# UNIVERSITY<sup>OF</sup> BIRMINGHAM University of Birmingham Research at Birmingham

# Human exposure to halogenated and organophosphate flame retardants through informal e-waste handling activities - A critical review

Ma, Yulong; Stubbings, William A.; Cline-Cole, Reginald; Harrad, Stuart

DOI: 10.1016/j.envpol.2020.115727

*License:* Creative Commons: Attribution-NonCommercial-NoDerivs (CC BY-NC-ND)

Document Version Peer reviewed version

Citation for published version (Harvard):

Ma, Y, Stubbings, WA, Cline-Cole, R & Harrad, S 2021, 'Human exposure to halogenated and organophosphate flame retardants through informal e-waste handling activities - A critical review', *Environmental Pollution*, vol. 268, no. Part A, 115727. https://doi.org/10.1016/j.envpol.2020.115727

Link to publication on Research at Birmingham portal

#### **General rights**

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

•Users may freely distribute the URL that is used to identify this publication.

•Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.

•User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?) •Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

#### Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

# Human exposure to halogenated and organophosphate flame retardants through informal e-waste handling activities - a critical review

Yulong Ma<sup>a</sup>, William A. Stubbings<sup>a</sup>, Reginald Cline-Cole<sup>b</sup>, Stuart Harrad<sup>a,\*</sup>

<sup>a</sup> School of Geography, Earth and Environmental Sciences, College of Life & Environmental Sciences, University

of Birmingham, Edgbaston, Birmingham, B15 2TT, UK

<sup>b</sup> Centre of West African Studies, Department of African Studies & Anthropology, School of History and Cultures,

University of Birmingham, Edgbaston, Birmingham, B15 2TT, UK

<sup>\*</sup> Corresponding author. E-mail address: S.J.Harrad@bham.ac.uk (S. Harrad)

## 1 Abstract

2 Informal electrical and electronic waste (e-waste) handling activities constitute a potentially important source of halogenated (HFRs) and organophosphate flame retardants (OPFRs) to the 3 environment and humans. In this review, two electronic databases (ScienceDirect and Web of 4 5 Science Core Collection) were searched for papers that addressed this topic. A total of 82 relevant 6 studies (including 72 studies selected from the two databases and 10 studies located from the 7 references of the first 72 selected studies) were identified that reported on human external and 8 internal exposure to HFRs and OPFRs arising as a result of informal e-waste handling activities. 9 Compared to the general population, higher levels of external exposure (i.e., inhalation, ingestion, 10 and dermal absorption) and internal exposure (i.e., blood serum, hair, breast milk, urine, and other 11 human matrices) to HFRs and OPFRs were identified for e-waste recyclers and residents inhabiting 12 e-waste dismantling and recycling zones, especially for younger adults and children. Food intake 13 and dust ingestion were the dominant exposure pathways for the majority of brominated flame 14 retardants (BFRs) and dechlorane plus (DP); while inhalation was identified as the most significant 15 pathway of human exposure to OPFRs in informal e-waste sites. The majority of research to date 16 has focused on China and thus future studies should be conducted in other regions such as Africa 17 and South Asia. Other suggested foci of future research are: examination of exposure via dermal 18 contact with e-waste, dietary exposure of local populations to OPFRs, confirmation of the existence 19 of and cause(s) of the higher body burdens of females compared with males amongst populations 20 impacted by informal e-waste handling, and characterisation of exposure of such populations to 21 chlorinated paraffins.

22

23 Keywords: WEEE; Brominated flame retardants; Organophosphate esters; Chlorinated paraffins

24

Main findings: Inhalation contributes most to human exposure to OPFRs, and dietary intake
 contributes most to human exposure to BFRs and DPs. Children and females are more exposed.

27

# 28 1. Introduction

Electrical and electronic waste (e-waste), also called waste electrical and electronic equipment
(WEEE), has become a global concern. In 2019, global generation of e-waste reached 53.6 million

tonnes (Mt), and this figure is estimated to reach 74 Mt by 2030 and 120 Mt by 2050, respectively 31 32 (Forti et al., 2020; World Economic Forum, 2019). Typically, household appliances like washing 33 machines, telecommunications, and IT equipment including computers, and consumer articles like TVs, comprise the majority of e-waste generated globally (Akram et al., 2019). Due to 34 35 environmental and economic considerations, much of the e-waste generated in high-income 36 countries has and continues to be exported for handling in low- and moderate-income countries 37 (Mihai et al., 2019; Pathak et al., 2017). Although measures outlined in the Basel Convention were 38 designed to prohibit the export of hazardous wastes to low- and moderate-income countries, these 39 are unlikely to be effective as some used electrical and electronic equipment is characterised as non-40 hazardous waste (Khan, 2016). Moreover, it was not until June 2019 when the EU clarified their 41 determinations on the regulation of persistent organic pollutants (POPs), that many non-hazardous 42 WEEE items would be reclassified as hazardous and become subject to the Hazardous Waste 43 Regulations 2005 (Official Journal of the European Union, 2019). Moreover, in low- and moderate-44 income countries, approximately 90 % of e-waste disposal and recycling activities are undertaken 45 by informal sector workers (Pathak et al., 2019). In India, for instance, 97 % of total e-waste 46 generated is handled by informal recycling yards in major cities like Delhi, Mumbai, Hyderabad, 47 and Bangalore (Rao et al., 2017).

48

49 This is concerning as various contaminants including halogenated flame retardants (HFRs) and 50 organophosphate flame retardants (OPFRs) can enter environmental media through different 51 activities reported to be undertaken at informal e-waste handling facilities. Informal e-waste 52 activities refer to unlicensed or unregulated e-waste dismantling and recycling activities that are 53 usually practiced by individuals and families using primitive techniques. These include: open 54 burning, acid leaching, and heating (Leung, 2019). The risks posed by such emissions are 55 compounded by the lack of effective personal protective equipment for use by those undertaking 56 such work, as well as by absence of other measures designed to protect the environment and human 57 health (Lundgren, 2012). HFRs and OPFRs have been widely used as additives at concentrations of 58 up to 10-15 % by weight in the plastic housing of various electric and electronic products to ensure 59 that these products meet fire safety regulations (Lassen and Havelund, 2006). However, from the 60 mid-2000s onwards, concerns about the adverse environmental and human health impacts of HFRs

61 like polybrominated diphenyl ethers (PBDEs) and hexabromocyclododecane (HBCDD) would lead 62 to restrictions and bans on their manufacture and use in new products in many jurisdictions, 63 including their listing under the Stockholm Convention on POPs of the United Nations Environment 64 Programme (UNEP) (Ge et al., 2020; Harrad, 2015; Huang et al., 2018; Ma et al., 2017b). As a 65 result, there has been increased demand for OPFRs and other HFRs such as tetrabromobisphenol-A (TBBP-A), novel brominated flame retardants (NBFRs), dechlorane plus (DP), and chlorinated 66 paraffins (CPs) (Chen et al., 2015; Ge et al., 2020; Gravel et al., 2020; He et al., 2017; Liu et al., 67 68 2016; Ma et al., 2017a; Zeng et al., 2018, 2020).

69

Previous studies have determined high concentrations of HFRs and OPFRs in environmental media 70 71 (Anh et al., 2017; Iqbal et al., 2017; Qin et al., 2019; Wang et al., 2018; Zheng et al., 2015b) and 72 foodstuffs (Anh et al., 2017; Huang et al., 2018; Tao et al., 2016; Zheng et al., 2015a) at various 73 informal e-waste dismantling and recycling sites such as Bui Dau (Vietnam), Guiyu and Qingyuan 74 (China), and Karachi (Pakistan). Such findings indicate high external exposure of local populations 75 (resident exposure) to HFRs and OPFRs through inhalation (Iqbal et al., 2017; Qin et al., 2019), 76 ingestion (Anh et al., 2017; Huang et al., 2018), and dermal absorption (Wu et al., 2016a). Elevated 77 internal exposure of local residents has also been demonstrated by reported concentrations of HFRs 78 or OPFRs in human matrices like serum (Guo et al., 2018, 2020; Lv et al., 2015), hair (Liang et al., 79 2016; Qiao et al., 2019), milk (Li et al., 2017), and urine (Lu et al., 2017; Shi et al., 2019) in informal 80 e-waste sites in South China (e.g., Qingyuan, Luqiao, and Wenling). Such evidence of elevated 81 human exposure has raised concerns about potential adverse health effects on populations impacted 82 by informal e-waste treatment (Akram et al., 2019; Asante et al., 2019; Awasthi et al., 2016, 2018; 83 Bakhiyi et al., 2018; Orisakwe et al., 2019; Shi et al., 2018). For instance, BFRs are endocrine 84 disruptors (Eguchi et al., 2015; Guo et al., 2018, 2019b; Zheng et al., 2017a, 2017b), and could exert 85 adverse effects on human semen quality (Yu et al., 2018); while OPFRs have been associated with 86 greater DNA damage (Lu et al., 2017). Moreover, health effects of HFR and OPFR exposure on 87 fetuses and infants are of particular concern due to their substantially weaker resistance and 88 immunity (Bai et al., 2019; Li et al., 2018; Xu et al., 2015; Zheng et al., 2017b).

89

90 The present review aims to: 1) summarise current research into human external exposure (i.e.,

91 inhalation, ingestion of food, dust, and soil, and dermal absorption) and internal exposure (i.e., blood
92 and serum, hair, breast milk, urine, and other human matrices) to HFRs and OPFRs through informal
93 e-waste dismantling and recycling activities for local residents; 2) identify potential health risks to
94 local residents in informal e-waste sites; and 3) highlight substantial research gaps that require
95 urgent investigation.

96

#### 97 2. Methods

98 Between 26/09/2019 and 07/09/2020, two electronic databases (ScienceDirect and Web of Science 99 Core Collection) were searched for research articles, reviews, book chapters, and other online resources. Terms searched were: "e-waste", "human exposure", and "flame retardants", with only 100 101 papers published between 2015 and 2020 selected. Using these search criteria, 1651 publications 102 were found on ScienceDirect, with a further 131 papers located on Web of Science Core Collection. 103 Further inspection by the authors comprising screening titles and abstracts identified relevant 104 publications (101 from ScienceDirect and 82 from Web of Science Core Collection). After removal 105 of duplicates (n=33), 150 publications remained for further screening.

106

107 These 150 publications were further rated for relevance by screening sampling methodology, 108 statistical data presented, and conclusions. As a result, 78 articles were excluded (including one 109 article not written in English), leaving 72 publications (consisting of 15 review articles, 4 book 110 chapters, and 53 research articles). In addition, references cited in these 72 selected publications 111 were also reviewed, adding a further 10 publications to the total included in this review. These 112 include: 1) four research papers published before 2015 but essential to this review in terms of data 113 interpretation (Ali et al., 2012; Labunska et al., 2014; Tue et al., 2013; Wang et al., 2010); 2) three 114 publications published between 2015 and 2020 but could not be located under current search 115 techniques due to their different foci (Abdallah and Harrad, 2018; Khan, 2016; Liu et al., 2016); and 116 3) three official reports (Lassen and Havelund, 2006; Lundgren, 2012; United States Environmental 117 Protection Agency, 2017).

118

# 3. Geographical distribution of studies into HFR and OPFR contamination from e-waste handling

It is notable that all the 53 selected research articles were conducted in lower- and upper-middle
income countries, according to the latest classification made by the World Bank (World Bank, 2020).
These include China (n=39), Vietnam (n=7), Thailand (n=2), Pakistan (n=1), Nigeria (n=1), South

- 124 Africa (n=1), Ghana (n=1), and Bangladesh (n=1) (Fig. 1).
- 125

## 126 4 Human external exposure to HFRs and OPFRs arising from informal e-waste activities

127 Previous studies have indicated 3 major pathways of human external exposure to HFRs and OPFRs 128 from informal e-waste activities, i.e., inhalation, ingestion of soil, dust, and food, and dermal 129 absorption from soil, dust, and e-waste articles (Anh et al., 2017; Huang et al., 2018; Iqbal et al., 130 2017; Qin et al., 2019; Wu et al., 2016a; Zheng et al., 2016). Once HFRs or OPFRs enter the 131 environment through informal e-waste activities, they can transfer between different environmental 132 media (Akram et al., 2019; Anh et al., 2017; Wang et al., 2018; Zheng et al., 2016). Particularly 133 important, HFRs and OPFRs can enter the food chain through bioconcentration and 134 biomagnification, thereby entering the human body through food ingestion resulting in exposure to 135 individuals beyond those directly undertaking e-waste handling, including young children (Anh et al., 2017; Sun et al., 2018; Wu et al., 2019; Zheng et al., 2016). Transfer of HFRs and OPFRs 136 137 between different environmental media and resultant human external exposure pathways are shown 138 in Fig.2, with recent studies (i.e. those published between 2015 and 2020) concerning human 139 external exposure to OPFRs and HFRs (including PBDEs, NBFRs, HBCDDs, TBBP-A, DPs, and 140 CPs) through informal e-waste handling activities summarised in Tables S1-S5 (Supplementary 141 Material).

142

#### 143 4.1 Human inhalation exposure to HFRs and OPFRs arising from informal e-waste activities 144 Inhalation of air has been highlighted as an important pathway of human exposure to HFRs and 145 OPFRs in informal e-waste dismantling and recycling sites, especially for OPFRs (Awasthi et al., 2016; Iqbal et al., 2017; Jiang et al., 2019; Luo et al., 2016). A study conducted in four e-waste 146 147 recycling sites (Jacob Lines, Surjani Town, Lyari, and Shershah) in Karachi City, Pakistan identified 148 human inhalation exposure to a range of HFRs and OPFRs. Specifically, these comprised: 8 PBDE congeners (BDE-28, -47, -99, -100, -153, -154, -183, and -209), 6 NBFRs (pentabromoethylbenzene 149 (PBEB), hexabromobenzene (HBB), 2-ethylhexyl-2,3,4,5-tetrabromobenzoate (TBB), 1,2-150

151 bis(2,4,6-tribromophenoxy) ethane (BTBPE), bis(2-ethylhexyl)-3,4,5,6-tetrabromophthalate (BEH-152 TBP), and decabromodiphenylethane (DBDPE)), 7 OPFRs (tris-(2-chloroethyl)-phosphate (TCEP), 153 tris-(2,3-dichloropropyl)-phosphate (TDCIPP), ethylhexyl diphenyl phosphate (EHDPP), tris(2-154 ethylhexyl)phosphate (TEHP), tri-n-butyl phosphate (TnBP), tris-(2-chloroisopropyl)-phosphate 155 (TCIPP), and tri-phenyl phosphate (TPHP)), as well as 2 isomers of DP (syn- and anti) (Iqbal et al., 2017). The authors reported human inhalation exposure to OPFRs to be the highest (2334 ng/kg 156 bw/day), with the average daily dose of TPHP, TEHP, TnBP, TCEP, EHDPP, TCIPP, and TDCIPP 157 158 being 1042 ng/kg bw/day, 308 ng/kg bw/day, 301 ng/kg bw/day, 230 ng/kg bw/day, 173 ng/kg 159 bw/day, 162 ng/kg bw/day, and 118 ng/kg bw/day, respectively (Iqbal et al., 2017). This was followed by PBDEs (19.1 ng/kg bw/day),  $\Sigma$ NBFRs (13.7 ng/kg bw/day), and DP (5.42 ng/kg bw/day) 160 161 (Iqbal et al., 2017). The calculated exposure doses were much lower than the reference doses suggested in previous literature for some of the target compounds (Table 1), indicating little health 162 163 risk caused by inhalation exposure. However, accurate assessment of risk is difficult since no 164 reference doses for mixtures of FRs (HFRs and OPFRs) are available, and the elevated exposures 165 ranging between 2876 and 12087 ng  $\Sigma$ FRs/kg bw/day are notable.

166

In comparison, human inhalation exposure to HFRs and OPFRs in Guiyu and Qingyuan, two major 167 168 e-waste recycling zones in China, was much lower (Luo et al., 2016; Wu et al., 2016a). In 2012, 169 samples of atmospheric particles were collected in Qingyuan and analysed for 12 OPFRs (2 isomers 170 of TnBP, TCEP, TCIPP, TDCIPP, TPHP, tris(2-butoxyethyl) phosphate (TBOEP), EHDPP, TEHP, 171 and 3 isomers of tris(2,4,6-trimethoxyphenyl)phosphine (TMPP)) (Luo et al., 2016). Of the 12 172 OPFRs, TEHP, and the 3 isomers of TMPP were not detected in any sample, and the average total 173 concentration of the remaining 8 OPFRs was 130±130 ng/m<sup>3</sup>. This indicates a mean daily inhalation 174 dose of  $12.1\pm4.1$  ng  $\Sigma$ OPFRs/kg bw/day for an average adult (TBOEP:  $6.8\pm4.1$  ng/kg bw/day, TnBP: 175 2.85±0.59 ng/kg bw/day, TIBP: 0.87±0.60 ng/kg bw/day, TCIPP: 0.64±0.28 ng/kg bw/day, TCEP: 176 0.33±0.20 ng/kg bw/day, TDCIPP: 0.25±0.12 ng/kg bw/day, TPHP: 0.18±0.06 ng/kg bw/day, 177 EHDPP: 0.172±0.049 ng/kg bw/day, respectively) (Luo et al., 2016). In Guiyu, daily inhalation 178 doses of BDE-47 and -99 were estimated to be 0.55 ng/kg bw/day and 0.33 ng/kg bw/day for adults, 179 and 1.77 ng/kg bw/day and 1.07 ng/kg bw/day for children, respectively (Wu et al., 2016a). Given 180 that this was much lower than the health based limit values (HBLVs) presented in previous studies

(Table 1), it is reasonable to conclude that human inhalation exposure to OPFRs and PBDEs inQingyuan and Guiyu presented a low health risk.

183

# 184 **4.2** Human ingestion exposure to HFRs and OPFRs arising from informal e-waste activities

## 185 *4.2.1 Dust ingestion*

186 Because of their relatively high octanol-air (K<sub>OA</sub>) and octanol-water partition coefficients (K<sub>OW</sub>), 187 many HFRs like high brominated PBDEs and NBFRs are likely to accumulate in atmospheric 188 particles and body lipids (Ji et al., 2017; Jiang et al., 2019; Luo et al., 2016; Ma et al., 2017a, 2017b; 189 Zheng et al., 2015a). An increasing number of studies have identified high concentrations of various 190 HFRs and OPFRs in indoor dust and foodstuffs, indicating that human exposure to HFRs and OPFRs 191 through indoor dust and food ingestion is non-negligible (Anh et al., 2017; Huang et al., 2018; Zeng 192 et al., 2016, 2018; Zheng et al., 2015b, 2016). For instance, Zheng et al. (2015b) determined 193 concentrations of 8 PBDE congeners (BDE-28, -47, -99, -100, -153, -154, -183, and -209), 4 NBFRs 194 (BEH-TBP, TBB, BTBPE, and DBDPE), 8 OPFRs (TEHP, TnBP, TCEP, TBOEP, TPHP, EHDPP, 195 TDCIPP, and TCIPP), and 2 isomers of DP (syn- and anti-) in indoor dust samples from some of the 196 largest e-waste dismantling and recycling sites in China (Longtang, Dali, and Guiyu), and calculated 197 human exposure to these contaminants through indoor dust ingestion. Assuming average dust 198 ingestion rates (20 mg/day for adults and 50 mg/day for children), and average FR concentrations 199 in dust; estimated daily intakes (EDIs) of PBDEs, NBFRs, OPFRs, and DPs for adults were: 1.11-200 24.1 ng/kg bw/day, 0.73-20.3 ng/kg bw/day, 1.36-23.5 ng/kg bw/day, and 0.08-1.73 ng/kg bw/day, 201 respectively, with the corresponding values for children 16-352 ng/kg bw/day, 11-296 ng/kg bw/day, 202 20-343 ng/kg bw/day, and 1.18-25.3 ng/kg bw/day, respectively (Zheng et al., 2015b). The highest 203 EDI values (assuming high ingestion of dust (50 mg/day for adults and 200 mg/day for children) contaminated at the 95<sup>th</sup> percentile concentration) for PBDEs, NBFRs, OPFRs, and DPs were: 168 204 ng/kg bw/day, 165 ng/kg bw/day, 226 ng/kg bw/day, and 12.8 ng/kg bw/day for adults, and 3915 205 206 ng/kg bw/day, 3844 ng/kg bw/day, 5282 ng/kg bw/day, and 298 ng/kg bw/day for children, 207 respectively (Zheng et al., 2015b). Although the calculated exposure doses were lower than the 208 reference doses suggested in previous studies (Table 1), the authors concluded that children had an 209 EDI of HFRs and OPFRs, that was 1 to 2 orders of magnitude higher than adults. Similar results 210 were reported in Vietnam (Anh et al., 2017) and Thailand (Muenhor et al., 2017, 2018).

211

212 Anh et al. (2017) measured concentrations of 8 PBDE congeners (BDE-28, -47, -99, -100, -153, -213 154, -183, and -209) in samples of home dust and fish from Bui Dau village, a major e-waste recycling site in Vietnam. Total PBDE concentrations in home dust were between 250-8650 ng/g, 214 215 with BDE-209 being the dominant congener (Anh et al., 2017). It is estimated that the EDI via dust 216 ingestion for adults and children was 0.71-2.47 ng/kg bw/day and 1.04-36.0 ng/kg bw/day under a 217 median dust ingestion scenario (20 mg/day for adults and 50 mg/day for children), and 0.18-6.19 218 ng/kg bw/day and 4.17-144 ng/kg bw/day under a high-end dust ingestion scenario (50 mg/day for 219 adults and 200 mg/day for children), respectively (applying an average body weight of 70 kg for 220 adults and 12 kg for children) (Anh et al., 2017). Specifically, it is notable that total PBDE 221 concentrations in home dust and the EDI reported by Anh et al. (2017) were similar to those reported 222 in a previous study conducted in the same area (Tue et al., 2013), indicating that PBDE 223 contamination in this area remained at the same level during the 7-year period (from 2008 to 2015). 224 Furthermore, the substantial increase in PBDE and NBFR concentrations in indoor dust in Qingyuan, 225 China from 2007 to 2013/2014 (Table 2) indicates that e-waste recycling workers and residents 226 became increasingly exposed to BFRs through dust ingestion over this period (He et al., 2017; Wang 227 et al., 2010; Zheng et al., 2015b). This is an interesting situation, seen from the perspective of other 228 areas where downward trends of PBDE contaminations have been reported (Harrad, 2015; Ma et 229 al., 2017b; Yu et al., 2016).

230

#### 231 4.2.2 Dietary intake

232 In addition to dust ingestion, food intake could also be an important exposure pathway of e-waste 233 recyclers and local residents to HFRs (Anh et al., 2017; Huang et al., 2018; Labunska et al., 2014, 234 2015; Tao et al., 2016; Wu et al., 2019; Zeng et al., 2016, 2018). In the same study conducted by 235 Anh et al. (2017), the EDI of 8 PBDE congeners through fish consumption (fish samples were 236 manually collected from ponds and canals located within the e-waste recycling area) for adults and 237 children in Bui Dau village, Vietnam was 0.72-46.4 ng/kg bw/day and 0.89-57.0 ng/kg bw/day, 238 respectively. In addition, the EDI of 5 NBFRs (pentabromobenzene (PBBz), HBB, BTBPE, BEH-239 TBP, and DBDPE), 2 DP isomers, and 3 HBCDD isomers through consumption of various foodstuffs was 36 ng/kg bw/day, 133 ng/kg bw/day, and 480 ng/kg bw/day for adults, and was 65 240

241 ng/kg bw/day, 350 ng/kg bw/day, and 1500 ng/kg bw/day for children, respectively, in Bui Dau village, Vietnam (Tao et al., 2016). It is notable that the EDI of  $\sum_{3}$ HBCDDs for adults and children 242 was 2.4 times and 7.5 times higher than the reference dose (200 ng/kg bw/day), respectively, 243 244 indicating potential health risk caused by dietary exposure to HBCDDs for local residents, 245 especially for children. In comparison, while human dietary exposure to NBFRs in informal e-waste dismantling and recycling sites in China (Huang et al., 2018; Labunska et al., 2014, 2015) was 246 247 comparable to that in Vietnam; much higher human exposure to PBDEs (1 to 2 orders of magnitude 248 higher) and much lower human exposure to HBCDDs (1 to 2 orders of magnitude lower) was 249 observed in China compared to Vietnam (Huang et al., 2018; Labunska et al., 2014, 2015; Wu et 250 al., 2019; Zeng et al., 2016). Furthermore, higher EDIs via food intake were identified for various 251 HFRs in informal e-waste dismantling and recycling sites in China between 2013 and 2016 (Tables 252 S1, S2, S3, and S5, Supplementary Material), especially for DPs, DBDPE, and CPs (Huang et al., 253 2018; Zeng et al., 2018), indicating potentially greater health concerns arising from exposure to 254 HFRs in these regions.

255

#### 256 4.3 Human dermal uptake of HFRs and OPFRs arising from informal e-waste activities

257 There is some debate about the contribution of dermal absorption to human external exposure to 258 HFRs and OPFRs. While some researchers state exposure risk through dermal contact is almost 259 negligible (Wu et al., 2016b), Wu et al. (2016a) suggest that the contribution of dermal absorption 260 to human external exposure to HFRs and OPFRs has been underestimated because previous 261 assessments of dermal exposure have addressed only inadvertent contact with contaminated dust or 262 soil and overlooked dermal absorption of both particulate and gaseous contaminants through air-to-263 skin transfer, as well as from direct skin contact with e-waste articles. Most studies to date have 264 focused on human exposure to HFRs and OPFRs through inhalation and ingestion (Anh et al., 2017; 265 Huang et al., 2018; Iqbal et al., 2017; Luo et al., 2016; Zeng et al., 2016; Zheng et al., 2015b), with 266 some consideration of exposure via dermal absorption limited. Abafe and Martincigh (2015) 267 estimated dermal absorption from dust of 8 PBDE congeners (BDE-28, -47, -99, -100, -153, -154, 268 -183, and -209) for e-waste dismantling and recycling workers from Durban, South Africa. Mean 269 and high-end exposure estimates were 0.87 ng/kg bw/day and 3.40 ng/kg bw/day, respectively, when 270 an equation adapted from the US Environmental Protection Agency was applied (Abafe and

271 Martincigh, 2015). The contribution of dermal absorption, defined as dermal absorption from dust 272 / (dust ingestion + dermal absorption from dust), was 40% under a mean dust ingestion scenario (20 273 mg/day for adults), and 20% under a high-end dust ingestion scenario (50 mg/day for adults). 274 However, this was likely underestimated since other dermal pathways such as air-to-skin transport, 275 especially those of lower brominated congeners considered more easily absorbed by human skin 276 due to their lower K<sub>OW</sub> (Abdallah et al., 2015; Wu et al., 2016a), was not considered. In Guiyu, for 277 example, the daily dermal intake of gaseous BDE-47 and BDE-99 through air-mediated transfer by 278 adults was estimated to be 0.65 ng/kg bw/day and 0.61 ng/kg bw/day, respectively, exceeding 279 exposure via inhalation of both gaseous and particle-bound BDE-47 (0.55 ng/kg bw/day) and BDE-280 99 (0.33 ng/kg bw/day) (Wu et al., 2016a). Similar results were also reported by Shen et al. (2019), 281 who identified higher EDIs of NBFRs, TBBP-A, and HBCDDs through dermal absorption rather 282 than dust ingestion for recyclers, local adults, and local children in Qingyuan, China (the same 283 equation adopted by Abafe and Martincigh (2015) was used in this study). Moreover, there appears 284 to date to have been no consideration of exposure via direct dermal contact with e-waste articles, 285 which may be a significant omission given recent data demonstrating the importance of dermal 286 exposure via direct contact with FR-treated fabrics (Abdallah and Harrad, 2018). Overall, dermal 287 absorption appears to be a potentially underestimated pathway of human external exposure to HFRs 288 (and perhaps, OPFRs) for e-waste dismantling and recycling workers or residents inhabiting e-waste 289 dismantling and recycling zones.

290

# 291 4.4 Variation in relative significance of different exposure pathways for HFRs and OPFRs

292 PBDEs are the most frequently reported HFRs globally in terms of human external exposure through 293 informal e-waste activities, especially through food and dust ingestion (Table S1, Supplementary 294 Material). Specifically, the EDI of PBDEs through food intake has been suggested to exceed other 295 exposure pathways such as inhalation (Wu et al., 2016a) and dust ingestion (Labunska et al., 2014). 296 In Wenling and Luqiao, China, for instance, the median EDI of PBDEs through food intake was 5 297 times and 2 times higher than that through dust ingestion for adults and children, respectively, with 298 high-end exposure via food intake approximately 10 and 30 times higher than that through dust 299 ingestion for adults and children, respectively (Labunska et al., 2014). Higher contribution to human 300 exposure to PBDEs via food consumption than dust ingestion was also identified for adults in two 11 / 32

301 Vietnamese informal e-waste sites when a medium dust ingestion rate (20 mg/day for adults and 50 302 mg/day for children) was applied (Anh et al., 2017). Moreover, for children, exposure through food 303 consumption was found roughly equivalent to that through dust ingestion (Anh et al., 2017). 304 However, the contribution of food intake to PBDE exposure was very likely underestimated by Anh 305 et al. (2017), since only fish consumption was included in that study. Furthermore, it is notable that 306 the relative contribution of different pathways varies for different PBDE congeners. In a study 307 conducted by Labunska et al. (2014), the relative contribution of dust ingestion to PBDE exposure 308 compared to that of food consumption (defined as EDI<sub>dust</sub> / (EDI<sub>food</sub> + EDI<sub>dust</sub>), using median exposure scenarios) was 1.0%, 2.1%, 0.2%, 0.2%, and 30.5% for BDE-47, -99, -153, -154, and -309 209, respectively, for adults, and 2.3%, 4.3%, 0.5%, 0.5%, and 49% for BDE-47, -99, -153, -154, 310 311 and -209, respectively, for children. Compared to lower brominated PBDE congeners, the 312 contribution of dust ingestion to overall exposure to BDE-209 is much higher.

313

314 Food intake has also been reported as the dominant pathway of human exposure to NBFRs (DBDPE, 315 BTBPE, etc), DPs, and HBCDD. In Bui Dau, Vietnam, the EDI for DBDPE, BTBPE, and 2 isomers of DP through food intake was 5 ng/kg bw/day, 31 ng/kg bw/day, and 133 ng/kg bw/day for adults, 316 317 and 2.7 ng/kg bw/day, 61 ng/kg bw/day, and 350 ng/kg bw/day for children, respectively (Tao et al., 318 2016). This figure was 2 to 4 orders of magnitude higher than exposure via dermal absorption of 319 soil and soil ingestion of DBDPE, BTBPE, and DPs in the same area (Someya et al., 2016). Similar 320 results were also reported in Qingyuan, China, where the estimated dietary intake of DBDPE, DPs, 321 as well as  $\alpha$ -,  $\beta$ -, and  $\gamma$ -HBCDD were about 1 to 2 orders of magnitude higher than dust ingestion 322 for both adults and children (He et al., 2017; Huang et al., 2018; Shen et al., 2019).

323

The main pathway of human exposure to OPFRs appears very different to that of BFRs and DPs, with inhalation identified as the most important exposure pathway in a recent study of an informal e-waste site in Pakistan. This study reported human exposure to OPFRs through inhalation to be 3 orders of magnitude higher than via soil ingestion (Iqbal et al., 2017). This result was in agreement with studies conducted in Qingyuan, China. Specifically, human exposure to OPFRs through inhalation was estimated to be 12.1±4.1 ng/kg bw/day (mean±SD) for adults in Qingyuan, China, which exceeded the EDI via dust ingestion reported by He et al. (2015) (median: 7.02 ng/kg bw/day), 331 Guo et al. (2019a) (mean: 5.85 ng/kg bw/day), and Zheng et al. (2015b) (mean: 1.36-2.11 ng/kg 332 bw/day) for adults. It is interesting that in both informal e-waste regions in Pakistan and China, nonchlorinated OPFRs comprised the majority of total OPFRs exposure via inhalation (Iqbal et al., 333 334 2017; Luo et al., 2016). This may be attributable to less extensive use of chlorinated OPFRs in 335 electrical and electronic products. Another interesting observation is that, despite the low 336 contribution of food intake to human exposure to OPFRs reported in informal e-waste sites to date, 337 chlorinated OPFRs were more frequently detected in chicken eggs sourced from an informal e-waste 338 site in China than non-chlorinated OPFRs (Zheng et al., 2016). This might be explained by the 339 relatively longer half-life of chlorinated OPFRs (Ma et al., 2017a).

340

341 Very limited data on human exposure to CPs through informal e-waste activities are available. To 342 the best of our knowledge, only 3 publications have reported human dietary exposure to CPs in 343 informal e-waste sites in China (Yuan et al., 2017; Zeng et al., 2016, 2018), with an increasing trend 344 of human dietary exposure identified in Longtang, China (Table S5, Supplementary Material). 345 Specifically, the EDI of short-chain chlorinated paraffins (SCCPs,  $C_{10}$ - $C_{13}$ ) through chicken egg 346 consumption increased by 4 times for adults and children between 2013 and 2016, and the EDI of median-chain chlorinated paraffins (MCCPs, C14-C17) also increased by nearly 30% (Zeng et al., 347 348 2018). No data were found about human exposure to CPs through inhalation, dermal contact, or dust 349 ingestion.

350

### **5** Human internal exposure to HFRs and OPFRs at informal e-waste handling sites

352 In addition to assessments of external exposure, internal human exposure to HFRs and OPFRs has 353 also been frequently examined in various informal e-waste dismantling and recycling areas, with 354 human blood and serum (Chen et al., 2015; Eguchi et al., 2015; Guo et al., 2020; Lv et al., 2015; 355 Schecter et al., 2018), human hair (Chen et al., 2015; Liang et al., 2016; Qiao et al., 2019), human 356 milk (Awasthi et al., 2016; Li et al., 2017; Shi et al., 2018), and human urine (Bai et al., 2019; Lu et 357 al., 2017; Shi et al., 2019; Yan et al., 2018) being the most commonly used biomarkers. An overview 358 of recent studies (i.e. those published 2015-2020) of internal human exposure to HFRs and OPFRs 359 through informal e-waste handling activities is provided as Tables S6 and S7 (Supplementary 360 Material).

361

# 362 5.1 Human blood and serum

363 Human blood and serum are frequently used indicators of human internal exposure to HFRs, since 364 HFRs are likely transported into multiple human organs and tissues through blood circulation (Chen et al., 2015; Eguchi et al., 2015; Kuo et al., 2019; Lv et al., 2015; Schecter et al., 2018). In a study 365 366 conducted in Baoai, an e-waste treatment site in northern Vietnam, serum samples from 40 female 367 e-waste recyclers were collected to assess their internal exposure to various contaminants including 368 7 PBDE congeners (BDE-47, -99, -100, -153, -154, -183, and -209) (Schecter et al., 2018). Among 369 the analyzed PBDE congeners, BDE-209 (median: 73.3 ng/g lipid; 95% confidence interval (CI): 32.4-138.2 ng/g lipid) was dominant, followed by BDE-153 (median: 13.0 ng/g lipid; 95% CI: 10.2-370 371 18.8 ng/g lipid) and BDE-183 (median: 7.3 ng/g lipid; 95% CI: 6.1-10.0 ng/g lipid) (Schecter et al., 372 2018). PBDE concentrations in serum from e-waste recyclers were 1 to 2 orders of magnitude higher 373 than those in non-recyclers whose PBDE concentrations in serum were frequently below limits of 374 detection (Schecter et al., 2018), indicating high occupational exposure of e-waste recyclers to 375 PBDEs. Similar conclusions have been reached in other studies (Eguchi et al., 2015; Guo et al., 376 2019b; Liang et al., 2016). For instance, in an e-waste recycling site in Wenling, China and an urban 377 area where no e-waste recycling activities were undertaken, 14 PBDE congeners (BDE-17, -28, -378 47, -66, -99, -100, -153, -154, -183, -203, -206, -207, -208, and -209) and DBDPE were determined 379 in serum samples taken from e-waste recyclers, non-occupationally-exposed residents of the e-waste 380 site, and urban residents (Liang et al., 2016). Mean concentrations of total PBDEs and DBDPE in 381 serum from e-waste recyclers were 656 ng/g lipid (range: 167-2530 ng/g lipid) and 125 ng/g lipid 382 (range: 26.7-439 ng/g lipid), respectively. These concentrations exceeded significantly those 383 detected in serum of non-occupationally-exposed residents (PBDEs: 123 ng/g lipid, range: 45.9-243 384 ng/g lipid; DBDPE: 56.1 ng/g lipid, range: 4.20-127 ng/g lipid), and urban residents (PBDEs: 24.6 385 ng/g lipid, range: 10.1-48.2 ng/g lipid; DBDPE: 13.8 ng/g lipid, range: nd-33.2 ng/g lipid) (Liang 386 et al., 2016). Moreover, Zheng et al. (2017b) identified higher PBDE concentrations in serum taken 387 from donors who lived in an e-waste site for over 20 years than those who lived there for less than 388 3 years (Table S6, Supplementary Material), indicating that PBDE concentrations were likely to 389 increase with increasing duration of residence in informal e-waste sites.

390

391 It is interesting to note that females have been reported to display greater HFR contamination in 392 serum than males in e-waste-impacted areas. Zheng et al. (2017a) reported that total PBDE 393 concentrations in female serum samples (mean: 2309 ng/g lipid; range: 206-35902 ng/g lipid) were 394 significantly higher (p < 0.05) than those in male serum samples (mean: 690 ng/g lipid; range: 105-1806 ng/g lipid) in Qingyuan, China. Specifically, females were found to have significantly higher 395 396 (p < 0.05) serum concentrations of congeners associated with the Deca-BDE commercial 397 formulation (sum of BDE-196, -197, -202, -203, -206, -207, -208, and -209; mean: 1896 ng/g lipid; 398 range: 100-34482 ng/g lipid) than did males (mean: 509 ng/g lipid; range: 64.5-1494 ng/g lipid). 399 Moreover, females displayed higher serum concentrations of congeners associated with the Penta-400 BDE formulation (sum of BDE-28, -47, -66, -85, -99, and -100; mean: 125 ng/g lipid; range: 44.6-401 853 ng/g lipid) and Octa-BDEs (sum of BDE-153, -154, and -183; mean: 287 ng/g lipid; range: 402 4.58-2667 ng/g lipid) than males (Penta-BDEs: mean: 104 ng/g lipid, range: 17.1-242 ng/g lipid; 403 Octa-BDEs: mean: 77.3 ng/g lipid, range: 8.02-283 ng/g lipid), but the difference was not significant 404 in this instance (p > 0.05) (Zheng et al., 2017a). Similar findings were also reported by (Chen et al., 405 2015), who found that concentrations of DPs in female serum (median: 230 ng/g lipid, range: 37-1400 ng/g lipid) were slightly higher than that in male serum (median: 180 ng/g lipid, range: 22-406 407 510 ng/g lipid). Unfortunately, no explanation for this difference was provided in either publication, 408 and the cause of this gender disparity is unclear.

409

#### 410 **5.2 Human hair**

411 As a non-invasive sampling matrix, human hair has been frequently used to measure human internal 412 exposure to HFRs and OPFRs (Chen et al., 2015; Liang et al., 2016; Qiao et al., 2019). A recent 413 study recruited 31 female e-waste dismantling workers from an e-waste recycling site in South 414 China, and measured concentrations of a wide variety of FRs (i.e., 8 PBDE congeners (BDE-28, -415 47, -99, -100, -153, -154, -183, and -209), 2 NBFRs (DBDPE and BTBPE), syn- and anti-DP, and 416 13 OPFRs (TnBP, TCEP, TDCIPP, TBOEP, TEHP, TPHP, EHDPP, triisopropyl phosphate (TiPrP), 417 tri-n-propyl phosphate (TPrP), 3 isomers of tricresyl phosphate (TCP), and TCIPP)) in hair samples (Qiao et al., 2019). The mean concentration of  $\Sigma$ OPFRs (the most abundant FRs) was 431 ng/g 418 419 (range: 189-1558 ng/g), with TBOEP, TCIPP, TEHP, and TPHP the dominant congeners, accounting 420 for 24.2%, 18.7%, 15.9%, and 11.1% of ΣOPFR concentrations, respectively (Qiao et al., 2019). 15 / 32

421 For  $\Sigma$ PBDEs, the mean concentration was 271 ng/g (range: 49.8-2104 ng/g), with BDE-209 422 dominant (accounting for 92.8% of \SigmaPBDEs) (Qiao et al., 2019). For ΣDPs and ΣNBFRs, mean 423 concentrations were 61.3 ng/g and 211 ng/g, with ranges of 1.64-360 ng/g and 16.4-991 ng/g, 424 respectively (Qiao et al., 2019). Notably, the study also identified an increasing trend in concentrations of PBDEs, DPs, and NBFRs from 2009 to 2015 (Qiao et al., 2019). Another study 425 426 conducted by Liang et al. (2016) also indicated high concentrations of PBDEs and DBDPE in human 427 hair, noting that exposure of e-waste recyclers (mean value: 292.9 ng/g for PBDEs and 82.5 ng/g 428 for DBDPE) was significantly higher than that of non-occupationally exposed residents (mean value: 429 55.8 ng/g for PBDEs and 29.4 ng/g for DBDPE) and urban residents (mean value: 12.9 ng/g for PBDEs and 10.9 ng/g for DBDPE). Significant correlations were reported between concentrations 430 431 of PBDEs and DBDPE in serum and hair, thereby indicating hair to be a useful matrix for 432 biomonitoring PBDEs and DBDPE exposure in humans (Liang et al., 2016). Similar conclusions 433 were also drawn regarding DPs in human hair, and as highlighted above for serum, it is notable that 434 female hair (median: 200 ng/g; range: 17-1100 ng/g) was more contaminated with DPs than male 435 hair (median: 19 ng/g; range: 6.3-150 ng/g). However, in this instance the gender difference was 436 attributed to the longer external exposure time of female hair (Chen et al., 2015).

437

#### 438 **5.3 Human milk**

Concentrations of HFRs in human milk can not only reflect internal exposure of female adults but 439 440 also dietary exposure of nursing infants (Shi et al., 2018; Tang and Zhai, 2017). Concentrations of 441 HFRs in human milk and the associated implications for human exposure, especially for infants, 442 have previously been reviewed for China and Africa (Shi et al., 2018; Orisakwe et al., 2019). 443 Specifically, a recent study conducted in an e-waste handling area in Wenling, China, collected 25 444 human milk samples from mothers who had lived there for over 20 years (defined as the  $R_{20}$  group) 445 and 21 human milk samples from mothers who had resided there for no more than 3 years (defined as the R<sub>3</sub> group), and determined concentrations of 8 PBDE congeners (BDE-28, -47, -99, -100, -446 447 153, -154, -183, and -209) in these samples (Li et al., 2017). It found that ΣPBDE concentrations in 448 the R<sub>20</sub> group (mean: 25.7±20.0 ng/g lipid; range: 7.89-90.6 ng/g lipid) exceeded significantly those in the R<sub>3</sub> group (mean: 6.68±5.61 ng/g lipid; range: 1.87-22.0 ng/g lipid) (Li et al., 2017). The two 449 450 groups had similar congener profiles of PBDEs, with BDE-209 and -153 being the most abundant 16 / 32

451 congeners in both groups (Li et al., 2017). Furthermore, the EDI of PBDEs for infants in the R<sub>20</sub>
452 group was in the range 15.8-243 ng/kg bw/day (median: 45.3 ng/kg bw/day), while in the R<sub>3</sub> group
453 the range was 5.43-43.0 ng/kg bw/day (median: 11.4 ng/kg bw/day) (Li et al., 2017). Although this
454 was much lower than the reference doses of the United States Environmental Protection Agency
455 (2017), the maximum EDI of BDE-47 (76.9 ng/kg bw/day) and BDE-153 (98.9 ng/kg bw/day) (Li
456 et al., 2017) for nursing infants in the R<sub>20</sub> group approached the corresponding reference doses
457 (BDE-47: 100 ng/kg bw/day; BDE-153: 200 ng/kg bw/day).

458

#### 459 **5.4 Human urine**

460 OPFR metabolites have been measured in human urine samples from e-waste recyclers, including 461 those of both chlorinated and non-chlorinated OPFRs (Bai et al., 2019; Lu et al., 2017; Shi et al., 462 2019; Yan et al., 2018). For instance, Lu et al. (2017) found urinary concentrations of chlorinated 463 and non-chlorinated OPFR metabolites in an e-waste impacted area in Qingyuan, China (mean: 4.0 464 ng/mL for chlorinated OPFR metabolites and 2.3 ng/mL for non-chlorinated OPFR metabolites) 465 exceeded significantly those in subjects from a rural area (mean: 2.1 ng/mL for chlorinated OPFR metabolites and 0.74 ng/mL for non-chlorinated OPFR metabolites), suggesting substantial human 466 exposure to OPFRs through e-waste handling activities. Negative correlations were determined 467 468 between age and urinary concentrations of each OPFR metabolite, and for bis(2-chloroethyl) phosphate (BCEP), bis(1-chloro-2-propyl) phosphate (BCIPP), bis(1,3-dichloro-2-propyl) 469 470 phosphate (BDCIPP), and diphenyl phosphate (DPHP), the negative correlations were significant 471 (Lu et al., 2017). These are similar to results reported by Yan et al. (2018) of significantly higher 472 concentrations of BCEP in urine samples from the 21-30 age group than the older age groups. This 473 might indicate higher exposure of younger people. However, a survey conducted by Shi et al. (2019) 474 generated different results, specifically that concentrations of OPFR metabolites in urine samples 475 from children were significantly lower than for adults who have been participating in e-waste 476 treatment for years. This discrepancy could be explained by the different sampling methodologies 477 adopted. In particular, only residents (and no e-waste workers) were sampled by Lu et al. (2017), 478 while only e-waste recyclers were sampled by Yan et al. (2018), which means the difference between occupational and non-occupational exposure was not considered. However, the results reported by 479 480 Shi et al. (2019) could be explained by high occupational exposure of adults since the children were 17 / 32

not involved in e-waste recycling activities. No data was found about concentrations of HFRs (or
metabolites) other than chlorinated OPFRs in urine samples from e-waste dismantling and recycling
areas.

484

#### 485 **5.5 Other human matrices**

Xu et al. (2015) compared 8 PBDE congeners (BDE-28, -47, -99, -100, -153, -154, -183, and -209) 486 487 in human placental tissue samples from an e-waste recycling site (Guiyu, China) and a reference 488 area (Haojiang, China). The study found that PBDE concentrations were much higher in samples 489 from the e-waste recycling site (mean: 61.39±85.42 ng/g/lipid; range: 0.89-516.97 ng/g lipid) than 490 those from the reference area (mean: 13.03±195.46 ng/g/lipid; range: 0.66-195.46 ng/g/lipid). This 491 could indicate higher exposure not only for mothers but also for fetuses in the e-waste recycling site 492 since partitioning of PBDEs from mothers to fetuses is considered as an important pathway of 493 exposure of fetuses to PBDEs (Xu et al., 2015; Zheng et al., 2017b). High concentrations of PBDEs 494 were also detected in human nails, abdominal subcutaneous adipose tissue, and umbilical cord tissue 495 from residents inhabiting informal e-waste sites in China (Li et al., 2018; Lv et al., 2015; Meng et 496 al., 2020). Apart from PBDEs, some OPFR metabolites (i.e., dibutyl phosphate (DBP) and DPHP) 497 were also frequently detected in amniotic fluid samples (Bai et al., 2019), indicating fetal exposure 498 to OPFRs.

499

#### 500 **5.6** Potential health risks originated from human internal exposure to HFRs and OPFRs

501 Health risk assessments of human exposure to HFRs and OPFRs have previously been reviewed in 502 Africa (Asante et al., 2019; Orisakwe et al., 2019), India (Awasthi et al., 2016), China (Awasthi et 503 al., 2018; Shi et al., 2018), and other regions on a global scale (Akram et al., 2019; Bakhiyi et al., 504 2018). Specifically, in informal e-waste sites, human exposure to TCIPP, TCEP, TNBP, and TPHP 505 were correlated with elevated DNA oxidative stress in e-waste sites, as the urinary concentrations 506 of BCIPP, BCEP, DBP, and DPHP were significantly increased as the concentration of 8-hydroxy-507 2'-deoxyguanosine (8-OHdG), a marker of DNA oxidative stress, increased (Lu et al., 2017). 508 Moreover, PBDEs and NBFRs are reported thyroid hormone (TH) disruptors (Eguchi et al., 2015; 509 Guo et al., 2019b; Zheng et al., 2017a, 2017b). They have strong binding affinity to thyroid-510 stimulating hormone (TSH), thyroglobulin, thyroxine-binding globulin (TBG), TH receptor  $\alpha$  (TR $\alpha$ ),

and iodothyronine deiodinase I (ID1), and therefore could disrupt TH-regulated proteins and gene expression (Guo et al., 2019b). It is also suggested that PBDEs, NBFRs (e.g., DBDPE, BTBPE, BEH-TBP, etc.), and DP could exert similar disrupting effects on female follicle-stimulating hormone (FSH) and male testosterone, with NBFRs showing stronger disrupting effects on human sex hormones than do PBDEs (Guo et al., 2018). Furthermore, PBDEs were found to have adverse effects on human semen quality measured by sperm concentration and count, sperm progressive motility, and sperm viability (Yu et al., 2018).

518

519 Specifically, health effects of HFR exposure on fetuses and infants are of particular concern due to 520 their substantially weaker resistance and immunity (Bai et al., 2019; Li et al., 2018; Xu et al., 2015; 521 Zheng et al., 2017b). Potential health risks for pregnant women and fetuses arising from OPFR 522 (especially TPHP and TnBP) exposure in an e-waste site (Qingyuan, China) were implied by Bai et 523 al. (2019), who determined high concentrations of OPFRs in paired amniotic fluid as well as in 524 maternal urine. Meanwhile, Xu et al. (2015), along with Li et al. (2018) reported that high prenatal 525 exposure to PBDEs in e-waste recycling areas may lead to adverse physiological development in 526 fetuses, in terms of reduced body-mass index, Apgar 1 score, and head circumference.

527

#### 528 6 Conclusions

529 The evidence reviewed in this study indicates 3 main pathways of human external exposure to HFRs 530 and OPFRs in informal e-waste handling sites, i.e., inhalation, ingestion of dust and food, and 531 dermal absorption. Current evidence suggests EDIs of OPFRs via inhalation exceed those via food 532 intake and dust ingestion, and thus inhalation could be the dominant pathway of human exposure to 533 OPFRs in informal e-waste sites; while for PBDEs, NBFRs, and DPs, food consumption and indoor 534 dust ingestion are likely more important contributors. An important factor emerging from our review 535 is that human exposure to HFRs and OPFRs through dermal absorption is insufficiently well-536 understood and may well be underestimated as pathways such as dermal absorption of both 537 particulate and gaseous contaminants through air-to-skin transfer, as well as from direct skin contact 538 with e-waste articles, have been largely overlooked. Children, infants, and fetuses, as well as e-539 waste recycling workers were found to experience higher HFR and OPFR exposure than did non-540 occupationally-exposed adults inhabiting e-waste handling zones, or those living in non-e-waste541 impacted locations. Gender differences in human internal exposure to HFRs in informal e-waste 542 locations were reported in some studies with higher HFR concentrations in serum and hair observed 543 in females compared to males. The cause of these higher HFR concentrations in females is unclear, 544 especially in serum.

545

Temporal changes in HFR concentrations in environmental media and humans have also been identified from previous studies. The evidence reviewed in this study shows increasing levels of HFRs in indoor dust, foodstuffs, and human hair with increasing duration of e-waste handling activity at a given site. Furthermore, PBDE concentrations in human serum and breast milk were likely to increase with increasing duration of residence in informal e-waste sites.

551

Most studies about human exposure to HFRs and OPFRs through informal e-waste handling activities were conducted in China, while studies in other countries or regions were limited. As the import of wastes from foreign countries was banned in China at the end of 2018, other low- and moderate-income countries are likely to receive more waste, and greater environmental and health effects could be caused by the improper treatment of this waste. More attention should therefore be paid to problems associated with informal e-waste treatment in low- and moderate-income countries in the future.

559

560 Based on the present review, we recommend that the following research gaps should be addressed 561 urgently. Firstly, very little is known about HFR and OPFR contamination of the environment and 562 humans arising from informal e-waste activities outside China, especially in Africa where such 563 activities appear to be growing substantially. Secondly, more detailed consideration of dermal 564 absorption as a pathway of human exposure to HFRs and OPFRs to those working in and inhabiting 565 e-waste handling areas is required. Thirdly, data on human dietary intake of OPFRs of residents of 566 informal e-waste sites is a priority for investigation. Moreover, the higher body burdens of HFRs in 567 females associated with informal e-waste recycling compared to males that have been reported 568 require further detailed study, both to verify such findings and to elucidate their cause(s). Finally, very little is understood about human exposure to CPs via all potential pathways as a result of 569 570 informal e-waste handling, and research to better understand the magnitude of such exposure and

571	the pathways	via	which	it	occurs	is	recommended.
-----	--------------	-----	-------	----	--------	----	--------------

# 573 **Declarations of interest**

- 574 The authors declare no conflict of interest.
- 575

# 576 Acknowledgments

- 577 The authors gratefully acknowledge provision of the Global Challenges PhD Scholarship to Yulong
- 578 Ma by the University of Birmingham.

579

582



583 Fig. 1: Global distribution of recent research studies reporting human exposure to HFRs and

584 **OPFRs through informal e-waste handling activities (2015-2020)** 



Fig. 2: Transfer of HFRs and OPFRs between various environmental media and associated human exposure pathways (Leung, 2019)

585

586

Compound	RfD	Compound	RfD
TnBP	24000	BDE-209	7000
TCEP	22000	Octa-BDE	3000
TCIPP	80000	Penta-BDE	2000
TBOEP	15000	BDE-47	100
TPHP	70000	BDE-99	100
TDCIPP	15000	BDE-153	200
ТСР	13000	DP	5000000(1)
BTBPE	243000	DP	2000000(2)
TBB	20000	DP	10000 <sup>(3)</sup>
BEH-TBP	20000	∑HBCDDs	200
DBDPE	333333	TBBP-A	600000

Table 1. Reference doses (RfD) values (ng/kg bw/day) for some HFRs and OPFRs

588 Notes: (1) chronic oral RfD; (2) dermal RfD; (3) inhalation RfDs

589 Source: Ali et al. (2012), except United States Environmental Protection Agency (2017) for PBDEs, Wang et al.

590 (2013) for DP, and Besis et al. (2017) for HBCDD and TBBP-A.

591

587

592

593

# Table 2. Concentrations of BDE-209, DBDPE, and BTBPE in indoor dust collected from

594							
sampling period	BDE-209		DBDPE		BTBPE		
	median	range	median	range	median	range	references
2007	988	105-140000	63.1	13.5-1144	20	n.d. <sup>(1)</sup> -998	(Wang et al., 2010)
2013	644-22500	146-195000	1160-26300	n.d181000	28-148	2.8-12700	(Zheng et al., 2015)
2013-2014	23800	8530-152000	2720	669-15000	n.a. <sup>(2)</sup>	n.a.	(He et al., 2017)
595		Notes: (1) n.d. = n	ot detected; (2) n.a	. = not available			

Notes: (1) n.d. = not detected; (2) n.a. = not available

## 596 References

- Abafe, O.A., Martincigh, B., 2015. An assessment of polybrominated diphenyl ethers and
  polychlorinated biphenyls in the indoor dust of e-waste recycling facilities in South Africa:
  implications for occupational exposure. Environmental Science and Pollution Research 22, 1407814086.
- Abdallah, M.A.E., Harrad, S., 2018. Dermal contact with furniture fabrics is a significant pathway of
   human exposure to brominated flame retardants. Environment International 118, 26-33.
- Abdallah, M.A., Pawar, G., Harrad, S., 2015. Effect of Bromine Substitution on Human Dermal
  Absorption of Polybrominated Diphenyl Ethers. Environmental Science & Technology 49, 1097610983.
- Akram, R., Fahad, S., Hashmi, M.Z., Wahid, A., Adnan, M., Mubeen, M., Khan, N., Rehmani, M.I.A.,
  Awais, M., Abbas, M., Shahzad, K., Ahmad, S., Hammad, H.M., Nasim, W., 2019. Trends of
  electronic waste pollution and its impact on the global environment and ecosystem. Environmental
  Science and Pollution Research 26, 16923-16938.
- Ali, N., Dirtu, A.C., Van den Eede, N., Goosey, E., Harrad, S., Neels, H., t Mannetje, A., Coakley, J.,
  Douwes, J., Covaci, A., 2012. Occurrence of alternative flame retardants in indoor dust from New
  Zealand: indoor sources and human exposure assessment. Chemosphere 88, 1276-1282.
- Anh, H.Q., Nam, V.D., Tri, T.M., Ha, N.M., Ngoc, N.T., Mai, P.T.N., Anh, D.H., Minh, N.H., Tuan, N.A.,
  Minh, T.B., 2017. Polybrominated diphenyl ethers in plastic products, indoor dust, sediment and
  fish from informal e-waste recycling sites in Vietnam: a comprehensive assessment of
  contamination, accumulation pattern, emissions, and human exposure. Environmental
  Geochemistry and Health 39, 935-954.
- Asante, K.A., Amoyaw-Osei, Y., Agusa, T., 2019. E-waste recycling in Africa: risks and opportunities.
   Current Opinion in Green and Sustainable Chemistry 18, 109-117.
- Awasthi, A.K., Wang, M., Awasthi, M.K., Wang, Z., Li, J., 2018. Environmental pollution and human
  body burden from improper recycling of e-waste in China: A short-review. Environmental Pollution
  243, 1310-1316.
- Awasthi, A.K., Zeng, X.L., Li, J.H., 2016. Relationship between e-waste recycling and human health risk
  in India: a critical review. Environmental Science and Pollution Research 23, 11509-11532.
- Bai, X.Y., Lu, S.Y., Xie, L., Zhang, B., Song, S.M., He, Y., Ouyang, J.P., Zhang, T., 2019. A pilot study
  of metabolites of organophosphorus flame retardants in paired maternal urine and amniotic fluid
  samples: potential exposure risks of tributyl phosphate to pregnant women. Environmental ScienceProcesses & Impacts 21, 124-132.
- Bakhiyi, B., Gravel, S., Ceballos, D., Flynn, M.A., Zayed, J., 2018. Has the question of e-waste opened
  a Pandora's box? An overview of unpredictable issues and challenges. Environment International
  110, 173-192.
- Besis, A., Christia, C., Poma, G., Covaci, A., Samara, C., 2017. Legacy and novel brominated flame
  retardants in interior car dust Implications for human exposure. Environmental Pollution 230, 871881.
- Chen, K., Zheng, J., Yan, X., Yu, L., Luo, X., Peng, X., Yu, Y., Yang, Z., Mai, B., 2015. Dechlorane Plus
  in paired hair and serum samples from e-waste workers: Correlation and differences. Chemosphere
  123, 43-47.
- Eguchi, A., Nomiyama, K., Minh Tue, N., Trang, P.T.K., Hung Viet, P., Takahashi, S., Tanabe, S., 2015.
  Residue profiles of organohalogen compounds in human serum from e-waste recycling sites in

- 640 North Vietnam: Association with thyroid hormone levels. Environmental Research 137, 440-449.
- Forti, V., Baldé, C.P., Kuehr, R., Bel, G., 2020. The global e-waste monitor 2020: quantities, flows, and
   the circular economy potential. <u>https://www.itu.int/en/ITU-</u>
   D/Environment/Documents/Toolbox/GEM 2020 def.pdf (Accessed 20 July 2020).
- Ge, X., Ma, S., Zhang, X., Yang, Y., Li, G., Yu, Y., 2020. Halogenated and organophosphorous flame
  retardants in surface soils from an e-waste dismantling park and its surrounding area: Distributions,
  sources, and human health risks. Environment International 139, 105741.
- Gravel, S., Lavoue, J., Bakhiyi, B., Lavoie, J., Roberge, B., Patry, L., Bouchard, M.F., Verner, M.A.,
  Zayed, J., Labreche, F., 2020. Multi-exposures to suspected endocrine disruptors in electronic waste
  recycling workers: Associations with thyroid and reproductive hormones. International Journal of
  Hygiene and Environmental Health 225, 113445.
- Guo, H., Zheng, X., Ru, S., Sun, R., Mai, B., 2019a. Size-dependent concentrations and bioaccessibility
  of organophosphate esters (OPEs) in indoor dust: A comparative study from a megacity and an ewaste recycling site. Science of the Total Environment 650, 1954-1960.
- Guo, L.C., Liu, T., Yang, Y., Yu,S., Gao, Y., Huang, W., Xiao, J., Ma, W., Rutherford, S., Zhang, Y., 2020.
  Changes in thyroid hormone related proteins and gene expression induced by polychlorinated
  biphenyls and halogen flame retardants exposure of children in a Chinese e-waste recycling area.
  Science of The Total Environment 742, 140597.
- Guo, L.C., Pan, S.X., Yu, S.B., Liu, T., Xiao, J.P., Zhu, B.H., Qu, Y.B., Huang, W.X., Li, M., Li, X., Zeng,
  W.L., Rutherford, S., Lin, L.F., Zhang, Y.H., Ma, W.J., 2018. Human Sex Hormone Disrupting
  Effects of New Flame Retardants and Their Interactions with Polychlorinated Biphenyls,
  Polybrominated Diphenyl Ethers, a Case Study in South China. Environmental Science &
  Technology 52, 13935-13941.
- Guo, L.C., Yu, S., Wu, D., Huang, J., Liu, T., Xiao, J., Huang, W., Gao, Y., Li, X., Zeng, W., Rutherford,
  S., Ma, W., Zhang, Y., Lin, L., 2019b. Disruption of thyroid hormone regulated proteins and gene
  expression by polychlorinated biphenyls, polybrominated diphenyl ethers and new flame retardants
  in residents of an e-waste region. Environmental Pollution 254, 112925.
- Harrad, S., 2015. A meta-analysis of recent data on UK environmental levels of POP-BFRs in an
  international context: Temporal trends and an environmental budget. Emerging Contaminants 1, 3953.
- He, C.T., Zheng, J., Qiao, L., Chen, S.J., Yang, J.Z., Yuan, J.G., Yang, Z.Y., Mai, B.X., 2015. Occurrence
  of organophosphorus flame retardants in indoor dust in multiple microenvironments of southern
  China and implications for human exposure. Chemosphere 133, 47-52.
- He, C.T., Zheng, X.B., Yan, X., Zheng, J., Wang, M.H., Tan, X., Qiao, L., Chen, S.J., Yang, Z.Y., Mai,
  B.X., 2017. Organic contaminants and heavy metals in indoor dust from e-waste recycling, rural,
  and urban areas in South China: Spatial characteristics and implications for human exposure.
  Ecotoxicology and Environmental Safety 140, 109-115.
- Huang, C.C., Zeng, Y.H., Luo, X.J., Tang, B., Liu, Y.E., Ren, Z.H., Mai, B.X., 2018. Level changes and
  human dietary exposure assessment of halogenated flame retardant levels in free-range chicken eggs:
  A case study of a former e-waste recycling site, South China. Science of the Total Environment 634,
  509-515.
- Iqbal, M., Syed, J.H., Breivik, K., Chaudhry, M.J.I., Li, J., Zhang, G., Malik, R.N., 2017. E-Waste Driven
  Pollution in Pakistan: The First Evidence of Environmental and Human Exposure to Flame
  Retardants (FRs) in Karachi City. Environmental Science & Technology 51, 13895-13905.

- Ji, X.W., Ding, J., Xie, X.C., Cheng, Y., Huang, Y., Qin, L., Han, C., 2017. Pollution Status and Human
   Exposure of Decabromodiphenyl Ether (BDE-209) in China. Acs Omega 2, 3333-3348.
- Jiang, Y., Yuan, L., Lin, Q., Ma, S., Yu, Y., 2019. Polybrominated diphenyl ethers in the environment and
  human external and internal exposure in China: A review. Science of the Total Environment 696,
  133902.
- Khan, S.A., 2016. E-products, E-waste and the Basel Convention: Regulatory Challenges and
  Impossibilities of International Environmental Law. Review of European, Comparative &
  International Environmental Law 25, 248-260.
- Kuo, L.J., Cade, S.E., Cullinan, V., Schultz, I.R., 2019. Polybrominated diphenyl ethers (PBDEs) in
  plasma from E-waste recyclers, outdoor and indoor workers in the Puget Sound, WA region.
  Chemosphere 219, 209-216.
- Labunska, I., Abdallah, M.A.E., Eulaers, I., Covaci, A., Tao, F., Wang, M., Santillo, D., Johnston, P.,
  Harrad, S., 2015. Human dietary intake of organohalogen contaminants at e-waste recycling sites in
  Eastern China. Environment International 74, 209-220.
- Labunska, I., Harrad, S., Wang, M., Santillo, D., Johnston, P., 2014. Human dietary exposure to PBDEs
  around E-waste recycling sites in Eastern China. Environmental Science & Technology 48, 55555564.
- Lassen, C., Havelund, S., 2006. Deca-BDE and Alternatives in Electrical and Electronic Equipment.
   Danish Environmental Protection Agency Report Environmental Project No. 1141.
   <u>https://www2.mst.dk/udgiv/publications/2007/978-87-7052-349-3/pdf/978-87-7052-350-9.pdf</u>
   (Accessed 05 Dec 2019).
- Leung, A.O.W., 2019. Environmental Contamination and Health Effects Due to E-waste Recycling, in:
  Prasad, M.N.V., Vithanage, M. (Eds.), Electronic Waste Management and Treatment Technology.
  Butterworth-Heinemann, pp. 335-362.
- Li, M., Huo, X., Pan, Y., Cai, H., Dai, Y., Xu, X., 2018. Proteomic evaluation of human umbilical cord
  tissue exposed to polybrominated diphenyl ethers in an e-waste recycling area. Environment
  International 111, 362-371.
- Li, X., Tian, Y., Zhang, Y., Ben, Y., Lv, Q., 2017. Accumulation of polybrominated diphenyl ethers in
  breast milk of women from an e-waste recycling center in China. Journal of Environmental Sciences
  52, 305-313.
- Liang, S., Xu, F., Tang, W.B., Zhang, Z., Zhang, W., Liu, L.L., Wang, J.X., Lin, K.F., 2016. Brominated
  flame retardants in the hair and serum samples from an e-waste recycling area in southeastern China:
  the possibility of using hair for biomonitoring. Environmental Science and Pollution Research 23,
  14889-14897.
- Liu, K., Li, J., Yan, S., Zhang, W., Li, Y., Han, D., 2016. A review of status of tetrabromobisphenol A
   (TBBPA) in China. Chemosphere 148, 8-20.
- Lu, S.Y., Li, Y.X., Zhang, T., Cai, D., Ruan, J.J., Huang, M.Z., Wang, L., Zhang, J.Q., Qiu, R.L., 2017.
   Effect of E-waste Recycling on Urinary Metabolites of Organophosphate Flame Retardants and
   Plasticizers and Their Association with Oxidative Stress. Environmental Science & Technology 51,
   2427-2437.
- Lundgren, K., 2012. The global impact of e-waste: Addressing the challenge.
   <u>https://www.ilo.org/wcmsp5/groups/public/---ed\_dialogue/---</u>
- 726 <u>sector/documents/publication/wcms\_196105.pdf</u> (Accessed 17 Feb 2020).
- 727 Luo, P., Bao, L.J., Guo, Y., Li, S.M., Zeng, E.Y., 2016. Size-dependent atmospheric deposition and

- inhalation exposure of particle-bound organophosphate flame retardants. Journal of HazardousMaterials 301, 504-511.
- Lv, Q.X., Wang, W., Li, X.H., Yu, L., Zhang, Y., Tian, Y., 2015. Polychlorinated biphenyls and
  polybrominated biphenyl ethers in adipose tissue and matched serum from an E-waste recycling
  area (Wenling, China). Environmental Pollution 199, 219-226.
- Ma, Y., Jin, J., Li, P., Xu, M., Sun, Y., Wang, Y., Yuan, H., 2017a. Organophosphate ester flame retardant
   concentrations and distributions in serum from inhabitants of Shandong, China, and changes
   between 2011 and 2015. Environmental Toxicology and Chemistry 36, 414-421.
- Ma, Y., Li, P., Jin, J., Wang, Y., Wang, Q., 2017b. Current halogenated flame retardant concentrations in
  serum from residents of Shandong Province, China, and temporal changes in the concentrations.
  Environmental Research 155, 116-122.
- Meng, H.J., Tang, B., Zheng, J., Ma, S.X., Cai, F.S., Zhuang, X., Wang, J.L., Yu, Y.J., 2020. Levels and
  sources of PBDEs and PCBs in human nails from e-waste, urban, and rural areas in South China.
  Environmental Science: Processes & Impacts 22, 1710-1717.
- Mihai, F.C., Gnoni, M.G., Meidiana, C., Ezeah, C., Elia, V., 2019. Waste Electrical and Electronic
  Equipment (WEEE): Flows, Quantities, and Management A Global Scenario, in: Prasad, M.N.V.,
  Vithanage, M. (Eds.), Electronic Waste Management and Treatment Technology. ButterworthHeinemann, pp. 1-34.
- Muenhor, D., Moon, H.B., Lee, S., Goosey, E., 2017. Polybrominated diphenyl ethers (PBDEs) in floor
  and road dust from a manual e-waste dismantling facility and adjacent communities in Thailand.
  Journal of Environmental Science and Health Part a-Toxic/Hazardous Substances & Environmental
  Engineering 52, 1284-1294.
- Muenhor, D., Moon, H.B., Lee, S., Goosey, E., 2018. Organophosphorus flame retardants (PFRs) and
  phthalates in floor and road dust from a manual e-waste dismantling facility and adjacent
  communities in Thailand. Journal of Environmental Science and Health Part a-Toxic/Hazardous
  Substances & Environmental Engineering 53, 79-90.
- Official Journal of the European Union, 2019. Regulation (EU) 2019/1021 of the European Parliament
  and of the Council of 20 June 2019 on persistent organic pollutants (Text with EEA relevance.).
  https://eur-lex.europa.eu/eli/reg/2019/1021/oj. (Accessed 29 May 2020).
- Orisakwe, O.E., Frazzoli, C., Ilo, C.E., Oritsemuelebi, B., 2019. Public Health Burden of E-waste in
  Africa. Journal of Health and Pollution 9, 1-12.
- Pathak, P., Srivastava, R.R., Ojasvi, 2017. Assessment of legislation and practices for the sustainable
   management of waste electrical and electronic equipment in India. Renewable and Sustainable
   Energy Reviews 78, 220-232.
- Pathak, P., Srivastava, R.R., Ojasvi, 2019. Environmental Management of E-waste, in: Prasad, M.N.V.,
  Vithanage, M. (Eds.), Electronic Waste Management and Treatment Technology. ButterworthHeinemann, pp. 103-132.
- Qiao, L., Zheng, X.B., Zheng, J., Chen, S.J., Zhong, C.Q., Chen, J.H., Yang, Z.Y., Mai, B.X., 2019.
  Legacy and Currently Used Organic Contaminants in Human Hair and Hand Wipes of Female EWaste Dismantling Workers and Workplace Dust in South China. Environmental Science &
  Technology 53, 2820-2829.
- Qin, Q.L., Xu, X.J., Dai, Q.Y., Ye, K., Wang, C.Y., Huo, X., 2019. Air pollution and body burden of
   persistent organic pollutants at an electronic waste recycling area of China. Environmental
   Geochemistry and Health 41, 93-123.

- Rao, M.N., Sultana, R., Kota, S.H., 2017. Electronic Waste, in: Rao, M.N., Sultana, R., Kota, S.H. (Eds.),
  Solid and Hazardous Waste Management. Butterworth-Heinemann, pp. 209-242.
- Schecter, A., Kincaid, J., Quynh, H.T., Lanceta, J., Tran, H.T.T., Crandall, R., Shropshire, W., Birnbaum,
  L.S., 2018. Biomonitoring of Metals, Polybrominated Diphenyl Ethers, Polychlorinated Biphenyls,
  and Persistent Pesticides in Vietnamese Female Electronic Waste Recyclers. Journal of
  Occupational and Environmental Medicine 60, 191-197.
- Shen, M.J., Ge, J.L., Lam, J.C.W., Zhu, M.S., Li, J., Zeng, L.X., 2019. Occurrence of two novel triazinebased flame retardants in an E-waste recycling area in South China: Implication for human exposure.
  Science of the Total Environment 683, 249-257.
- Shi, Y., Zheng, X., Yan, X., Wang, Y., Xu, R., Wang, M., Ren, M., Zheng, J., 2019. Short-term variability
   in levels of urinary phosphate flame retardant metabolites in adults and children from an e-waste
   recycling site. Chemosphere 234, 395-401.
- Shi, Z., Zhang, L., Li, J., Wu, Y., 2018. Legacy and emerging brominated flame retardants in China: A
  review on food and human milk contamination, human dietary exposure and risk assessment.
  Chemosphere 198, 522-536.
- Someya, M., Suzuki, G., Ionas, A.C., Tue, N.M., Xu, F., Matsukami, H., Covaci, A., Tuyen, L.H., Viet,
  P.H., Takahashi, S., Tanabe, S., Takigami, H., 2016. Occurrence of emerging flame retardants from
  e-waste recycling activities in the northern part of Vietnam. Emerging Contaminants 2, 58-65.
- Sun, R., Luo, X., Li, Q.X., Wang, T., Zheng, X., Peng, P., Mai, B., 2018. Legacy and emerging
  organohalogenated contaminants in wild edible aquatic organisms: Implications for
  bioaccumulation and human exposure. Science of the Total Environment 616-617, 38-45.
- Tang, J., Zhai, J.X., 2017. Distribution of polybrominated diphenyl ethers in breast milk, cord blood and
   placentas: a systematic review. Environmental Science and Pollution Research 24, 21548-21573.
- Tao, F., Matsukami, H., Suzuki, G., Tue, N.M., Viet, P.H., Takigami, H., Harrad, S., 2016. Emerging
  halogenated flame retardants and hexabromocyclododecanes in food samples from an e-waste
  processing area in Vietnam. Environmental Science-Processes & Impacts 18, 361-370.
- Tue, N.M., Takahashi, S., Suzuki, G., Isobe, T., Viet, P.H., Kobara, Y., Seike, N., Zhang, G., Sudaryanto,
  A., Tanabe, S., 2013. Contamination of indoor dust and air by polychlorinated biphenyls and
  brominated flame retardants and relevance of non-dietary exposure in Vietnamese informal e-waste
  recycling sites. Environment International 51, 160-167.
- United States Environmental Protection Agency, 2017. Technical Fact Sheet Polybrominated Diphenyl
  Ethers (PBDEs) (EPA 505-F-17-015). https://www.epa.gov/sites/production/files/201403/documents/ffrrofactsheet\_contaminant\_perchlorate\_january2014\_final\_0.pdf. (Accessed 31
  October 2019).
- Wang, D.G., Alaee, M., Byer, J.D., Brimble, S., Pacepavicius, G., 2013. Human health risk assessment
  of occupational and residential exposures to dechlorane plus in the manufacturing facility area in
  China and comparison with e-waste recycling site. Science of The Total Environment 445-446, 329336.
- Wang, J., Ma, Y.J., Chen, S.J., Tian, M., Luo, X.J., Mai, B.X., 2010. Brominated flame retardants in
  house dust from e-waste recycling and urban areas in South China: Implications on human exposure.
  Environment International 36, 535-541.
- Wang, T., Ding, N., Wang, T., Chen, S.J., Luo, X.J., Mai, B.X., 2018. Organophosphorus esters (OPEs)
  in PM2.5 in urban and e-waste recycling regions in southern China: concentrations, sources, and
  emissions. Environmental Research 167, 437-444.

- World Economic Forum, 2019. A New Circular Vision for Electronics: Time for a Global Reboot.
  www3.weforum.org/docs/WEF\_A\_New\_Circular\_Vision\_for\_Electronics.pdf. (Accessed 20 July
  2020).
- World Bank, 2020. World Bank Country and Lending Groups.
   <u>https://datahelpdesk.worldbank.org/knowledgebase/articles/906519-world-bank-country-and-</u>
   <u>lending-groups</u>. (Accessed 22 July 2020).
- Wu, C.C., Bao, L.J., Tao, S., Zeng, E.Y., 2016a. Dermal Uptake from Airborne Organics as an Important
   Route of Human Exposure to E-Waste Combustion Fumes. Environmental Science & Technology
   50, 6599-6605.
- Wu, Q., Leung, J.Y.S., Du, Y., Kong, D., Shi, Y., Wang, Y., Xiao, T., 2019. Trace metals in e-waste lead
  to serious health risk through consumption of rice growing near an abandoned e-waste recycling
  site: Comparisons with PBDEs and AHFRs. Environmental Pollution 247, 46-54.
- Wu, Y., Li, Y., Kang, D., Wang, J., Zhang, Y., Du, D., Pan, B., Lin, Z., Huang, C., Dong, Q., 2016b.
  Tetrabromobisphenol A and heavy metal exposure via dust ingestion in an e-waste recycling region
  in Southeast China. Science of the Total Environment 541, 356-364.
- Xu, L., Huo, X., Zhang, Y., Li, W., Zhang, J., Xu, X., 2015. Polybrominated diphenyl ethers in human
  placenta associated with neonatal physiological development at a typical e-waste recycling area in
  China. Environmental Pollution 196, 414-422.
- Yan, X., Zheng, X., Wang, M., Zheng, J., Xu, R., Zhuang, X., Lin, Y., Ren, M., 2018. Urinary metabolites
  of phosphate flame retardants in workers occupied with e-waste recycling and incineration.
  Chemosphere 200, 569-575.
- Yu, G., Bu, Q., Cao, Z., Du, X., Xia, J., Wu, M., Huang, J., 2016. Brominated flame retardants (BFRs):
  A review on environmental contamination in China. Chemosphere 150, 479-490.
- Yu, Y.J., Lin, B.G., Liang, W.B., Li, L.Z., Hong, Y.D., Chen, X.C., Xu, X.Y., Xiang, M.D., Huang, S.,
  2018. Associations between PBDEs exposure from house dust and human semen quality at an ewaste areas in South China-A pilot study. Chemosphere 198, 266-273.
- Yuan, B., Fu, J., Wang, Y., Jiang, G., 2017. Short-chain chlorinated paraffins in soil, paddy seeds (Oryza sativa) and snails (Ampullariidae) in an e-waste dismantling area in China: Homologue group
  pattern, spatial distribution and risk assessment. Environmental Pollution 220, 608-615.
- Zeng, Y., Ding, N., Wang, T., Tian, M., Fan, Y., Wang, T., Chen, S.J., Mai, B.X., 2020. Organophosphate
  esters (OPEs) in fine particulate matter (PM2.5) in urban, e-waste, and background regions of South
  China. Journal of hazardous materials 385, 121583.
- Zeng, Y., Huang, C., Luo, X., Liu, Y., Ren, Z., Mai, B., 2018. Polychlorinated biphenyls and chlorinated
  paraffins in home-produced eggs from an e-waste polluted area in South China: Occurrence and
  human dietary exposure. Environment International 116, 52-59.
- Zeng, Y.H., Luo, X.J., Tang, B., Mai, B.X., 2016. Habitat- and species-dependent accumulation of
  organohalogen pollutants in home-produced eggs from an electronic waste recycling site in South
  China: Levels, profiles, and human dietary exposure. Environmental Pollution 216, 64-70.
- Zheng, J., He, C.T., Chen, S.J., Yan, X., Guo, M.N., Wang, M.H., Yu, Y.J., Yang, Z.Y., Mai, B.X., 2017a.
  Disruption of thyroid hormone (TH) levels and TH-regulated gene expression by polybrominated
  diphenyl ethers (PBDEs), polychlorinated biphenyls (PCBs), and hydroxylated PCBs in e-waste
  recycling workers. Environment International 102, 138-144.
- Zheng, M.Y., Li, X.H., Zhang, Y., Yang, Y.L., Wang, W.Y., Tian, Y., 2017b. Partitioning of
   polybrominated biphenyl ethers from mother to fetus and potential health-related implications.

860 Chemosphere 170, 207-215.

- Zheng, X.B., Luo, X.J., Zheng, J., Zeng, Y.H., Mai, B.X., 2015a. Contaminant sources, gastrointestinal
  absorption, and tissue distribution of organohalogenated pollutants in chicken from an e-waste site.
  Science of the Total Environment 505, 1003-1010.
- Zheng, X., Xu, F., Chen, K., Zeng, Y., Luo, X., Chen, S., Mai, B., Covaci, A., 2015b. Flame retardants
  and organochlorines in indoor dust from several e-waste recycling sites in South China:
  Composition variations and implications for human exposure. Environment International 78, 1-7.
- Zheng, X., Xu, F., Luo, X., Mai, B., Covaci, A., 2016. Phosphate flame retardants and novel brominated
  flame retardants in home-produced eggs from an e-waste recycling region in China. Chemosphere
  150, 545-550.