

The bandwidth theorem for locally dense graphs

Staden, Katherine; Treglown, Andrew

DOI:

[10.1017/fms.2020.39](https://doi.org/10.1017/fms.2020.39)

License:

Creative Commons: Attribution (CC BY)

Document Version

Publisher's PDF, also known as Version of record

Citation for published version (Harvard):

Staden, K & Treglown, A 2020, 'The bandwidth theorem for locally dense graphs', *Forum of Mathematics, Sigma*, vol. 8, e42. <https://doi.org/10.1017/fms.2020.39>

[Link to publication on Research at Birmingham portal](#)

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.



RESEARCH ARTICLE

The bandwidth theorem for locally dense graphs

Katherine Staden¹ and Andrew Treglown²

¹Mathematical Institute, University of Oxford, Oxford, OX2 6GG United Kingdom; E-mail: staden@maths.ox.ac.uk.

²School of Mathematics, University of Birmingham, Edgbaston, Birmingham, B15 2TT United Kingdom;
E-mail: a.c.treglown@bham.ac.uk.

Received: 6 June 2019; **Revised:** 18 June 2020; **Accepted:** 16 June 2020

2020 Mathematics Subject Classification: Primary – 05C35

Keywords and phrases: bandwidth; embedding; regularity method

Abstract

The *bandwidth theorem* of Böttcher, Schacht, and Taraz [*Proof of the bandwidth conjecture of Bollobás and Komlós, Mathematische Annalen*, 2009] gives a condition on the minimum degree of an n -vertex graph G that ensures G contains every r -chromatic graph H on n vertices of bounded degree and of bandwidth $o(n)$, thereby proving a conjecture of Bollobás and Komlós [*The Blow-up Lemma, Combinatorics, Probability, and Computing*, 1999]. In this paper, we prove a version of the bandwidth theorem for *locally dense* graphs. Indeed, we prove that every locally dense n -vertex graph G with $\delta(G) > (1/2 + o(1))n$ contains as a subgraph any given (spanning) H with bounded maximum degree and sublinear bandwidth.

Contents

1	Introduction and results	2
2	Overview of the proof of Theorem 1.2	3
3	Preliminaries	5
3.1	Notation	5
3.1.1	Named graphs	6
3.2	Properties of locally dense graphs	7
4	The regularity and blow-up lemmas and associated tools	9
4.1	Regularity	9
4.2	Embedding lemmas	10
5	Finding the power of a Hamilton cycle	11
6	Lemmas for H	15
6.1	Partitioning a graph of low bandwidth: the basic lemma for H	15
6.2	Covering exceptional vertices: the second lemma for H	18
7	The lemma for G: adjusting cluster sizes	22
8	The proof of Theorem 1.2	28
9	Concluding remarks	34

1. Introduction and results

One of the fundamental topics in extremal graph theory is the study of minimum degree conditions that force a graph to contain a given spanning substructure. Perhaps the best known result in the area is Dirac's theorem [13], which states that any graph G on $n \geq 3$ vertices with minimum degree $\delta(G) \geq n/2$ contains a Hamilton cycle. The Pósa–Seymour conjecture (see [15] and [33]) states that any graph G on n vertices with $\delta(G) \geq rn/(r+1)$ contains the r th power of a Hamilton cycle. (The r th power of a Hamilton cycle C is obtained from C by adding an edge between every pair of vertices of distance at most r on C .) Komlós, Sárközy, and Szemerédi [28] proved this conjecture for sufficiently large graphs.

A decade ago, Böttcher, Schacht, and Taraz [9] proved a very general minimum degree result, the so-called *bandwidth theorem*. A graph H on n vertices is said to have *bandwidth at most b* if there exists a labelling of the vertices of H by the numbers $1, \dots, n$ such that for every edge $ij \in E(H)$, we have $|i-j| \leq b$. Clearly, every graph H has bandwidth at most $|H|-1$. Further, a Hamilton cycle has bandwidth 2, and in general the r th power of a Hamilton cycle has bandwidth at most $2r$. Böttcher, Preussmann, Taraz, and Würfl [7] proved that every planar graph H on n vertices with bounded maximum degree has bandwidth at most $O(n/\log n)$. The bandwidth theorem gives a condition on the minimum degree of a graph G on n vertices that ensures G contains every r -chromatic graph on n vertices of bounded degree and of bandwidth $o(n)$.

Theorem 1.1 (The bandwidth theorem, Böttcher, Schacht, and Taraz [9]). *Given any $r, \Delta \in \mathbb{N}$ and any $\gamma > 0$, there exist constants $\beta > 0$ and $n_0 \in \mathbb{N}$ such that the following holds. Suppose that H is an r -chromatic graph on $n \geq n_0$ vertices with $\Delta(H) \leq \Delta$ and bandwidth at most βn . If G is a graph on n vertices with*

$$\delta(G) \geq \left(\frac{r-1}{r} + \gamma \right) n,$$

then G contains a copy of H .

We remark that Theorem 1.1 had been conjectured by Bollobás and Komlós [26]. Since the bandwidth theorem was proven, a number of variants of the result have been obtained, including for arrangeable graphs [10] and degenerate graphs [30] and in the setting of random and pseudorandom graphs [1, 5, 23], as well as for robustly expanding graphs [24]. Very recently, a bandwidth theorem for approximate decompositions was proven by Condon, Kim, Kühn, and Osthus [12], whilst Glock and Joos [20] proved a μn -bounded edge colouring extension of Theorem 1.1. A general embedding result of Böttcher, Montgomery, Parczyk, and Person [6] also implies a bandwidth theorem in the setting of randomly perturbed graphs.

For many graphs H , the minimum degree condition in Theorem 1.1 is best-possible up to the term γn . For example, suppose that H is a K_r -factor (that is, we seek a collection of vertex-disjoint copies of K_r in G that together cover all the vertices in G). So $\chi(H) = r$, $\Delta(H) = r-1$, and H has bandwidth $r-1$. Suppose that G is obtained from two disjoint vertex classes A and B of sizes $n/r+1$ and $(r-1)n/r-1$, respectively, so that G contains all edges other than those with both endpoints in A . Then it is easy to see that G does not contain a K_r -factor; however, $\delta(G) = ((r-1)/r)n - 1$. In fact, note that the famous Hajnal–Szemerédi theorem [21] asserts that an n -vertex graph G contains a K_r -factor, provided $r|n$ and $\delta(G) \geq ((r-1)/r)n$. Thus, this extremal example is sharp. (Note, though, that for many r -partite graphs F , a significantly lower minimum degree condition than that in Theorem 1.1 ensures an F -factor; see [29].)

As for many other problems in the area, this extremal example has the characteristic that it contains a *large* independent set. There has thus been significant interest in seeking variants of classical results in extremal graph theory, where one now forbids the host graph from containing a large independent set. Indeed, nearly 50 years ago, Erdős and Sós [18] initiated the study of the Turán problem under the additional assumption of a small independence number. That is, they considered the number of edges in an n -vertex K_r -free graph with independence number $o(n)$. This topic is now known as *Ramsey–Turán theory* and has been extensively studied by numerous authors (see, for example, [2, 17, 31, 34]). More

recently, there has been interest in similar questions but where now one seeks a K_r -factor in an n -vertex graph with independence number $o(n)$ and large minimum degree (see [3, 4, 22]).

A stronger notion of a graph not containing a large independent set is that of being *locally dense*. More precisely, given $\rho, d > 0$, we say that an n -vertex graph G is (ρ, d) -dense if every $X \subseteq V(G)$ satisfies $e(G[X]) \geq d \binom{|X|}{2} - \rho n^2$. Note that the property of being locally dense is weaker than being dense and (pseudo)random and stronger than having a sublinear independence number. Locally dense graphs have been considered in a number of previous papers. For example, there have been several papers on a question of Erdős, Faudree, Rousseau, and Schelp [16]; there they considered a variant of the notion of (ρ, d) -dense and asked for the values of ρ and d guaranteeing that a (ρ, d) -dense graph contains a triangle. One can view the notion of *locally dense* as a parameter that ensures a graph is in some sense ‘random-like’. Therefore, there has been interest in determining the number of (homomorphic) copies of a fixed graph H in a (ρ, d) -dense graph G , and in particular whether this count is close to the value obtained if G were a random graph; the study of this topic (for graphs and hypergraphs) was initiated by Kohayakawa, Nagle, Rödl, and Schacht [25].

The aim of this paper is to prove the following locally dense version of the bandwidth theorem.

Theorem 1.2. *For all $\Delta \in \mathbb{N}$ and $d, \eta > 0$, there exist constants $\rho, \beta, n_0 > 0$ such that for every $n \geq n_0$, the following holds. Let H be an n -vertex graph with $\Delta(H) \leq \Delta$ and bandwidth at most βn . Then any (ρ, d) -dense graph G on n vertices with $\delta(G) \geq (1/2 + \eta)n$ contains a copy of H .*

In the case when H corresponds to a K_r -factor, Theorem 1.2 had been proven by Reiher and Schacht (see [4]). Note that in the case when H is connected, the minimum degree in Theorem 1.2 is best-possible up to the ηn term. Indeed, if G consists of two vertex-disjoint cliques, each of size $n/2$, then G trivially does not contain H , although G is locally dense and $\delta(G) = n/2 - 1$.

A striking feature of Theorem 1.2 is that, unlike Theorem 1.1, the minimum-degree condition does not depend on the chromatic number of H . In particular, when $\chi(H) = 2$, the minimum-degree condition in Theorem 1.2 is the same as that in Theorem 1.1. Thus, in the case of bipartite H , there is no benefit in adding the condition that G is locally dense. However, when $\chi(H) > 2$, the minimum degree in Theorem 1.2 is substantially reduced compared to the bandwidth theorem.

It would also be extremely interesting to prove a version of Theorem 1.2 for graphs of sublinear independence number. Note, though, that examples in [4] show that the statement of Theorem 1.2 is far from true if we require that G has a sublinear independence number instead of the locally dense condition. Indeed, the minimum degree necessary for the existence of a K_r -factor in such a graph is at least $(\frac{r-2}{r} + o(1))n$ for every $r \geq 4$. So, these two problems are genuinely different.

The proof of Theorem 1.2 draws on ideas from [8, 9], and our approach makes use of the regularity–blow-up method. We also employ several new ideas (particularly with regard to dealing with so-called *exceptional vertices*). In the next section, we give an overview of the proof of Theorem 1.2. In Section 3, we introduce some notation as well as several fundamental properties of locally dense graphs. The regularity and blow-up lemmas are presented in Section 4. A key step in the proof of Theorem 1.2 is to show that the hypothesis of this theorem ensures that G contains the r th power of a Hamilton cycle; we prove this in Section 5. The proof of Theorem 1.2 then breaks into two main parts: the proof of two so-called lemmas for H (presented in Section 6) and the lemma for G (presented in Section 7). In Section 8, we combine all these results to prove Theorem 1.2. We give some concluding remarks in Section 9.

Additional note. Since this paper was first submitted, Ebsen, Maesaka, Reiher, Schacht, and Schülke [14] have built on our work to generalise Theorem 1.2. Indeed, they replace the minimum degree condition on G with an inseparability condition.

2. Overview of the proof of Theorem 1.2

The overall strategy follows in the same spirit as the proof of the bandwidth theorem in [9], although the precise details of the proofs of the key steps in the argument turn out to be quite different. In particular, the setting of locally dense graphs both smooths over some aspects of the proof and introduces additional

difficulties. Often, in problems involving embedding a spanning structure, the most challenging aspect of the proof is dealing with so-called *exceptional vertices* (that is, trying to cover either the remaining last few vertices in the host graph or those few vertices that do not fit in some general structure in the host graph). In this paper, we take a novel approach to dealing with such vertices. Below we outline the key steps in our proof and highlight some of the main novelties in our approach.

Obtaining structure in G . Suppose that H and G are as in the statement of the theorem where $\chi(H) = r$. The first step in the proof is to apply the regularity lemma (Lemma 4.1) to G to obtain the reduced graph R of G . The reduced graph R is locally dense (with somewhat different parameters compared to G) and ‘inherits’ the minimum degree of G (that is, $\delta(R) > (1/2 + o(1))|R|$). These properties ensure that R contains an almost spanning subgraph Z_ℓ^{2r} that has the following properties:

- Z_ℓ^{2r} covers all but at most $2r$ of the vertices in R .
- Z_ℓ^{2r} consists of ℓ vertex-disjoint copies K^1, \dots, K^ℓ of K_{2r} (in particular, $|Z_\ell^{2r}| = 2r\ell$).
- For each $1 \leq i \leq \ell$, there are all possible edges between K^i and K^{i+1} except that we miss a perfect matching between the two. (Note here that $K^{\ell+1} := K^1$.)

The existence of Z_ℓ^{2r} in R can be guaranteed by finding a sufficiently large power of a Hamilton cycle in R . This is achieved in Section 5 (see Theorem 5.1). Using this, one can easily deduce that G contains an *almost* spanning structure \mathcal{C} that looks like the ‘blow-up’ of Z_ℓ^{2r} . More precisely, if $V(Z_\ell^{2r}) = \{1, \dots, 2r\ell\}$ and $V_1, \dots, V_{2r\ell}$ are the corresponding clusters in G ; then

- (i) $V(\mathcal{C}) = V_1 \cup \dots \cup V_{2r\ell}$.
- (ii) $\mathcal{C}[V_i, V_j]$ is ε -regular whenever $ij \in E(Z_\ell^{2r})$.
- (iii) If jk is an edge in one of the cliques K^i , then $\mathcal{C}[V_j, V_k]$ is superregular.

We refer to \mathcal{C} as a *cycle structure*.

Suppose that in fact \mathcal{C} is a spanning subgraph of G . In this case, ideally, one would now like to take the following approach. Let x_1, \dots, x_n denote the bandwidth ordering of H . Partition $V(H)$ into ℓ classes C_1, \dots, C_ℓ so that

- $c_i := |C_i| = |\cup_{j \in V(K^i)} V_j|$ for all $1 \leq i \leq \ell$.
- C_1 contains the vertices x_1, \dots, x_{c_1} , C_2 contains the vertices $x_{c_1+1}, \dots, x_{c_1+c_2}$, and so forth.

Then embed the vertices from C_1 into the clusters in G corresponding to the clique K^1 , embed the vertices from C_2 into the clusters in G corresponding to the clique K^2 , and so on.

At first sight, this seems like a plausible strategy: since the partition of $V(H)$ respected the bandwidth ordering of H (and as H has small bandwidth), most edges in H lie in the induced subgraphs $H[C_i]$; all remaining edges lie in the bipartite graphs $H[C_i, C_{i+1}]$. Suppose one could map each C_i onto the clusters corresponding to K^i , so that each such cluster V_j receives precisely $|V_j|$ vertices from C_i and, crucially, all edges xy in $H[C_i]$ are such that x and y are mapped to different clusters in K^i . That is, suppose we have a graph homomorphism ϕ_i between $H[C_i]$ and K^i that maps precisely $|V_j|$ vertices to each V_j . Further, suppose the ϕ_i together combine to give a graph homomorphism f from H to Z_ℓ^{2r} (so the edges in each $H[C_i, C_{i+1}]$ are mapped to edges in $R[V(K^i), V(K^{i+1})]$). Set $G_i := G[\cup_{j \in V(K^i)} V_j]$. Then (iii) above ensures that we could apply the blow-up lemma to each graph G_i so as to embed $H[C_i]$ into G_i . Further, (ii) ensures that we can achieve this embedding so all edges in the graphs $H[C_i, C_{i+1}]$ are also present. That is, we would obtain an embedding of H into G .

This naive approach fails, though, as there is no guarantee one can map each C_i onto the clusters corresponding to K^i so that each such cluster V_j receives precisely $|V_j|$ vertices from C_i . Furthermore, in the above approach, we assumed that \mathcal{C} is a spanning subgraph of G ; in reality, we have a small exceptional set V_0 of vertices in G uncovered by \mathcal{C} .

The basic lemma for H and the lemma for G . Instead of the above, we prove the so-called basic lemma for H (Lemma 6.1). Here we show that one can find a graph homomorphism f from H into Z_ℓ^{2r} so that for every cluster V_i of R , *approximately* $|V_i|$ vertices are mapped to it. This therefore ‘almost’

gives us the desired graph homomorphism f from H into Z_ℓ^{2r} . In the proof of Lemma 6.1, we rely on the fact that the K^i in Z_ℓ^{2r} are copies of K_{2r} ; note that in the analogous structure in the proof of the bandwidth theorem [9], the K^i are copies of K_r . To see why our condition is helpful for us, note that whilst in general, an r -partite graph H' does not have an ‘almost balanced’ graph homomorphism into K_r (since H' may have colour classes of wildly different sizes), for r -partite graphs H' of bounded degree and sublinear bandwidth, one can always find an almost balanced graph homomorphism from H' into K_{2r} .

Next, in the lemma for G (Lemma 7.1), we prove that if one does not have an exceptional set V_0 , then we can move vertices around the cycle structure \mathcal{C} in such a way as to ensure that now each cluster V_i in \mathcal{C} has size precisely corresponding to the number of vertices mapped to V_i by f . This is at the expense of weakening condition (iii): after applying Lemma 7.1, we only have that each clique K^i splits into two cliques K_1^i and K_2^i of size r such that if jk is an edge in one of the cliques K_1^i or K_2^i , then $\mathcal{C}[V_j, V_k]$ is superregular. However, this is still good enough to apply the blow-up lemma to find our desired embedding of H into G .

Special lemma for H . So far, we have been assuming that there is no exceptional set V_0 ; further, in the the proof of the bandwidth theorem [9], Böttcher, Schacht, and Taraz were able to utilise the large minimum degree to incorporate exceptional vertices into (their analogue of the cycle structure) \mathcal{C} . We have a significantly smaller minimum degree and so are unable to do this in our setting.

Instead, given the bandwidth ordering x_1, \dots, x_n of H , we reserve a short initial segment x_1, \dots, x_t ; and let H' denote the subgraph of H induced by x_1, \dots, x_t . Here, t will be significantly bigger than βn (recall that H has bandwidth at most βn), but H' will still only be a small fraction of H . Via the special lemma for H (Lemma 6.2), we will embed H' into G in such a way that, crucially, all of V_0 is covered by H' and, equally importantly, we do not cover more than a small proportion of each cluster V_i in \mathcal{C} .

In the proof of Lemma 6.2, since V_0 may contain only very few (or even no) edges, we must assign an independent set I in H' of size $|V_0|$ to be embedded onto V_0 . We then must connect up I through the rest of G to obtain a copy of H' . In particular, since H' is much smaller than H , the distance between two vertices $x, y \in I$ in H' may also be small. So it is crucial that G is ‘highly connected’. The connecting lemma (Lemma 3.3) ensures that this is the case. (Lemma 3.3 is also applied in the proof of Theorem 5.1.)

Care is also needed to ensure that Lemma 6.2 is compatible with the basic lemma for H (Lemma 6.1). That is, we use Lemma 6.2 to embed H' in G and Lemma 6.1 to embed $H \setminus H'$ in G . Thus, we need to ensure that the copies of H' and $H \setminus H'$ can be positioned in G in such a way that they ‘glue’ together to form a copy of H .

Note that the reader should view the above overview as an idealisation of the proof. Indeed, when we prove Theorem 1.2 in Section 8, some of the details will be a little different. For example, for technical reasons, it is important that we find a spanning copy of $Z_\ell^{r^*}$ in R for some $r^* \gg r$ rather than Z_ℓ^{2r} .

3. Preliminaries

3.1. Notation

Given a set X and $k \leq |X|$, write $\binom{X}{k}$ for the set of k -element subsets of X . Given $r \in \mathbb{N}$, write $[[2r]]^2 := [r]^2 \cup ([2r] \setminus [r])^2$. Given a function $f : X \rightarrow Y$ and $A \subseteq X$, we write $f|_A$ for the restriction of f to A and $f(A) := \{f(a) : a \in A\}$.

Given a graph G , we write $V(G)$ and $E(G)$ for the vertex and edge sets respectively, and $|G| := |V(G)|$ and $e(G) := |E(G)|$. The *degree* of a vertex $x \in V(G)$ is denoted by $d_G(x)$ and its neighbourhood by $N_G(x)$. The *degree* of a subset $X \subseteq V(G)$ is $d_G(X) := |\bigcap_{x \in X} N_G(x)|$. A subgraph $H \subseteq G$ is *s-extendable* if $d_G(V(H)) \geq s$. Given vertices x_1, \dots, x_k , we write $N_G(x_1, \dots, x_k) := \bigcap_{1 \leq i \leq k} N_G(x_i)$. If $A \subseteq V(G)$, we write $N_G(x, A) := N_G(x) \cap A$ and $d_G(x, A) := |N_G(x) \cap A|$. We say that A is *k-independent* if every vertex in A is at distance at least $k + 1$ in G ; that is, the shortest path in G between any pair of elements in A has length at least $k + 1$. Given $X, Y \subseteq V(G)$ (not necessarily disjoint), define $e_G(X, Y)$ to be the number of edges $xy \in E(G)$ with $x \in X$ and $y \in Y$. If X and Y are disjoint, then

$G[X, Y]$ is the bipartite graph with vertex classes X and Y whose edge set consists of all those edges in G with one endpoint in X and the other in Y .

Given two graphs G and H , we say that $f : V(H) \rightarrow V(G)$ is a *graph homomorphism* if $f(x)f(y) \in E(G)$ whenever $xy \in E(H)$. If f is additionally injective, we say that f is an *embedding (of H into G)*. Then $H \subseteq G$.

Throughout the paper, we will ignore floors and ceilings wherever they do not affect the argument. The constants in the hierarchies used to state our results are chosen from right to left. For example, if we claim that a result holds whenever $0 < 1/n \ll a \ll b \ll c \leq 1$ (where n is the order of the graph), then there are non-decreasing functions $f : (0, 1] \rightarrow (0, 1]$, $g : (0, 1] \rightarrow (0, 1]$, and $h : (0, 1] \rightarrow (0, 1]$ such that the result holds for all $0 < a, b, c \leq 1$ and all $n \in \mathbb{N}$ with $b \leq f(c)$, $a \leq g(b)$, and $1/n \leq h(a)$. Note that $a \ll b$ implies that we may assume in the proof that, for example, $a < b$ or $a < b^2$.

Given numbers a, b, c , we write $a = b \pm c$ to mean $a \in [b - c, b + c]$.

3.1.1. Named graphs

Given a graph H , the graph H^r , called the *r th power of H* , is obtained from H by adding an edge between every pair of vertices of distance at most r in H . In particular:

- $P_k^r = P = v_1 \dots v_k$ is an *r -path* if $V(P) = \{v_1, \dots, v_k\}$ and $E(P) = \bigcup_{j \in [r]} \{v_i v_{i+j} : 1 \leq i \leq k-j\}$.
- $C_k^r = C = w_1 \dots w_k$ is an *r -cycle* if $V(C) = \{w_1, \dots, w_k\}$ and $E(C) = \bigcup_{j \in [r]} \{w_i w_{i+j} : 1 \leq i \leq k\}$, where addition is modulo k .

Additionally,

- F is an *r -trail (of length s)* if there exists an ordered sequence of not necessarily distinct vertices v_1, \dots, v_s such that $V(F) = \{v_1, \dots, v_s\}$ and $E(F) = \bigcup_{j \in [r]} \{v_i v_{i+j} : 1 \leq i \leq s-j\}$. Observe that P_k^r is an r -trail, and $F \cong P_s^r$ if and only if $|F| = s$.
- A *K -tiling* is a collection of vertex disjoint copies of K . If it contains t copies, we denote it by $t \cdot K$. If $H \subseteq G$ is a K -tiling that is also spanning, we say that H is a *K -factor* of G .

Define

- Z_ℓ^r to be the graph with vertex set $[\ell] \times [r]$ in which $(i, j)(i', j')$ is an edge whenever (i) $|i - i'| \leq 1$ and $j \neq j'$ and when (ii) $i = \ell, i' = 1$, and $j \neq j'$.

Thus, Z_ℓ^r is obtained from a cycle on ℓ vertices by replacing each vertex with a clique on r vertices and replacing every edge with a complete bipartite graph minus a certain perfect matching. As indicated in Section 2, Z_ℓ^{2r} will be used in the proof of Theorem 1.2 as a framework for embedding (most of) H into G . Note that Böttcher, Schacht, and Taraz [9] used a very similar structure in their proof of the bandwidth theorem.

Observe that

$$2\ell \cdot K_r \subseteq C_{2r\ell}^{r-1} \subseteq Z_{2\ell}^r \subseteq C_{2r\ell}^{2r-1} \subseteq Z_\ell^{2r}, \quad (3.1)$$

and the lexicographic ordering of $V(Z_\ell^r)$ (that is, $(1, 1)(1, 2), \dots, (1, r), (2, 1), \dots, (\ell, r)$) is an $(r-1)$ -cycle ordering of $C_{r\ell}^{r-1}$.

Given $A, B \subseteq V(G)$ and $x_1, \dots, x_k \in V(G)$, when we say that, for example, $ABx_1 \dots x_k$ is an r -path (respectively, r -trail, r -cycle), we mean that any ordering $a_1, \dots, a_{|A|}$ of A and any ordering $b_1, \dots, b_{|B|}$ of B are such that $a_1 \dots a_{|A|} b_1 \dots b_{|B|} x_1 \dots x_k$ is an r -path (respectively, r -trail, r -cycle). An r -path (respectively, r -trail, r -cycle), $Ax_1 \dots x_k B$ or $x_1 \dots x_k AB$ is defined analogously.

Suppose X and Y are disjoint sets of vertices of size r . We say that an r -path P is *between X and Y* if $P = Xx_1 \dots x_k Y$ for some vertices x_1, \dots, x_k . Observe that $P[X], P[Y] \cong K_r$. Further, P *avoids* a set $W \subseteq V(G)$ if $V(P) \cap W = \emptyset$.

3.2. Properties of locally dense graphs

In this section we prove some simple properties of locally dense graphs G : that G induced on a large vertex subset is still locally dense; after removing a small set of vertices, G is still locally dense; and G contains many copies of cliques of a fixed size that additionally have a large common neighbourhood.

A fact that we shall use often throughout the paper is that if $0 < \rho < \rho'$ and $0 < d' < d$, then a (ρ, d) -dense graph is also (ρ', d') -dense.

Lemma 3.1. *Let $r, n \in \mathbb{N}$ and $0 < 1/n \ll \rho \ll d, 1/r$, and $0 < d, \alpha < 1$. Let G be a (ρ, d) -dense graph on n vertices, and let $U \subseteq V(G)$ where $|U| = \alpha n$. Then*

- (i) $G[U]$ is $(\rho/\alpha^2, d)$ -dense.
- (ii) $G \setminus U$ is $(\rho/(1-\alpha)^2, d)$ -dense.
- (iii) G contains at least $dn/2$ vertices of degree at least $dn/2$.
- (iv) G contains at least $(d/2)^{\binom{r+1}{2}} n^r / r!$ copies of K_r , each of which is $d^r n/2^r$ -extendable.

Proof. The proof of (i) is clear, and (ii) follows immediately from (i). For (iii), let $Y := \{v \in V(G) : d_G(v) \geq dn/2\}$. Then

$$2d \binom{n}{2} - 2\rho n^2 \leq 2e(G) \leq (n - |Y|) \frac{dn}{2} + |Y|n$$

and so

$$|Y| \geq \frac{dn - 2d - 4\rho n}{2 - d} \geq \frac{dn}{2},$$

proving (iii).

It remains to prove (iv). We claim that for each $i \leq r$, there is a set \mathcal{T}_i of (ordered) tuples $\mathbf{x} = (x_1, \dots, x_i)$ such that $G[\{x_1, \dots, x_i\}] \cong K_i$ and $d_G(\{x_1, \dots, x_i\}) \geq d^i n/2^i$ for all $\mathbf{x} \in \mathcal{T}_i$, and $|\mathcal{T}_i| \geq (d/2)^{\binom{i+1}{2}} n^i$. This will immediately imply (iv) as \mathcal{T}_r gives rise to at least $(d/2)^{\binom{r+1}{2}} n^r / r!$ (unlabelled) copies of K_r , each of which is $d^r n/2^r$ -extendable.

We will prove this by induction on i . Part (iii) implies that G contains a set \mathcal{T}_1 of $dn/2$ copies of K_1 that are all $dn/2$ -extendable. Suppose we have obtained \mathcal{T}_{i-1} with the required properties for some $2 \leq i \leq r$.

Fix $\mathbf{x} = (x_1, \dots, x_{i-1}) \in \mathcal{T}_{i-1}$. The graph $G_{\mathbf{x}} := G[N_G(x_1, \dots, x_{i-1})]$ induced by its neighbourhood contains at least $d^{i-1} n/2^{i-1}$ vertices, so (i) implies that it is $(2^{2i-2} \rho / d^{2i-2}, d)$ -dense and hence $(\sqrt{\rho}, d)$ -dense. Now, using the fact that $1/n \ll \sqrt{\rho} \ll d, 1/r$, (iii) implies that $G_{\mathbf{x}}$ contains at least $(d/2) \cdot d^{i-1} n/2^{i-1} = d^i n/2^i$ vertices, each of degree at least $d^i n/2^i$. Each such vertex y gives rise to an r -tuple $\mathbf{x}(y) := (x_1, \dots, x_{i-1}, y)$. Certainly $G[\{x_1, \dots, x_{i-1}, y\}] \cong K_i$; and further, $d_G(\{x_1, \dots, x_{i-1}, y\}) \geq d^i n/2^i$ since y has at least this many neighbours in the common neighbourhood of \mathbf{x} . Let \mathcal{T}_i be the collection of all these tuples $\mathbf{x}(y)$ formed from each $\mathbf{x} \in \mathcal{T}_{i-1}$. Observe that they have the required properties and are all distinct, so

$$|\mathcal{T}_i| \geq d^i n/2^i \cdot |\mathcal{T}_{i-1}| \geq (d/2)^{\binom{i}{2} + i} n^i = (d/2)^{\binom{i+1}{2}} n^i.$$

This completes the proof of the lemma. \square

We will need a *connecting lemma* to find a short r -path between two ‘extendable’ copies of K_r in a locally dense graph G with $\delta(G) > (1/2 + o(1))n$. The heart of the proof is the following lemma, which is the only part of the proof of Theorem 1.2 that requires $\delta(G) > (1/2 + o(1))n$ (elsewhere, linear minimum degree suffices). Somewhat similar lemmas have been used elsewhere in other settings: for example, [19, 35].

Lemma 3.2. *Let $0 < 1/n \ll \rho \ll d, \eta, 1/r < 1$, where $n, r \in \mathbb{N}$. Let G be an n -vertex graph, and let $U \subseteq V(G)$ be a subset of size $n' \geq \eta n/2$ such that $G[U]$ is (ρ, d) -dense and $d_G(x, U) \geq (1/2 + \eta)n'$*

for all $x \in V(G)$. Let X, Y, W be pairwise disjoint subsets of $V(G)$ such that $|X| = |Y| = \lceil 4r/\eta \rceil$ and $|W| \leq \eta n'/2$. Then there is $Z \subseteq U$ such that

- (i) $G[Z] \cong K_r$.
- (ii) $Z \cap (X \cup Y \cup W) = \emptyset$.
- (iii) There exist $X' \subseteq X$ and $Y' \subseteq Y$ with $|X'| = |Y'| = r$ such that $N_G(Z) \supseteq X' \cup Y'$.

Proof. Let $C := \lceil 4r/\eta \rceil$, and let $U' := U \setminus (X \cup Y \cup W)$. Then

$$e_G(U', X \cup Y) \geq (|X| + |Y|)((1/2 + \eta)n' - |X| - |Y| - |W|) \geq 2C(1/2 + \eta/4)n' \geq (1 + \eta/2)Cn'.$$

Let U'' be the collection of those vertices in U' that each have at least $C + r$ neighbours in $X \cup Y$. Then $(1 + \eta/2)Cn' \leq (C + r - 1)(n' - |U''|) + 2C|U''|$, and so

$$|U''| \geq \frac{(1 + \eta/2)Cn' - (C + r - 1)n'}{C - r + 1} \geq \frac{\eta n'}{4},$$

where the final inequality follows from the fact that $C \geq 4r/\eta$. There are not more than 2^{2C} ways a vertex can attach to $X \cup Y$, so there is $U^* \subseteq U''$ such that $N_G(v, X \cup Y)$ is identical for all $v \in U^*$ and $|U^*| \geq \eta n'/2^{2C+2}$. Note further that, since each such v has at least $C + r$ neighbours in $X \cup Y$, there are $X' \subseteq X$ and $Y' \subseteq Y$ such that $|X'| = |Y'| = r$ and $X' \cup Y' \subseteq N_G(U^*)$. Lemma 3.1(i) now implies that $G[U^*]$ is $(2^{4C+4}\rho/\eta^2, d)$ -dense and hence $(\sqrt{\rho}, d)$ -dense. But then, by Lemma 3.1(iv), there is $Z \subseteq U^*$ that spans a K_r . The desired properties (i)–(iii) are immediate. \square

As well as being applied in the proof of the connecting lemma below, Lemma 3.2 is also a key tool in the proof of Theorem 5.1 in Section 5, which in turn is a crucial tool for the proof of Theorem 1.2.

Lemma 3.3 (Connecting lemma). *Let $0 < 1/n \ll \rho \ll d, \eta \leq 1/r$, where $r \in \mathbb{N}$, and let G be a (ρ, d) -dense graph on n vertices with $\delta(G) \geq (1/2 + \eta)n$. Let W, X, Y be subsets of $V(G)$ such that $|W| \leq \eta n/4$ and X, Y induce r -cliques in G and each one either*

- *lies in a copy of $K_{\lceil 9r/\eta \rceil}$ that is disjoint from W or*
- *is ηn -extendable.*

Then G contains a copy of $P_{3r}^r = x_1 \dots x_{3r}$ avoiding W such that $Xx_1 \dots x_{3r}$ induces a copy of P_{4r}^r , and $x_1 \dots x_{3r}Y$ induces a copy of P_{4r}^r .

The connecting lemma will ensure that the reduced graph R of a graph G (as in Theorem 1.2) is ‘highly connected’. This property will be exploited when embedding a part of H into G so as to cover all of the exceptional set V_0 (specifically, we make use of Lemma 3.3 in Section 6.2).

Proof. Suppose that X is ηn -extendable. Let $C := \lceil 9r/\eta \rceil$ and $c := \lceil 4r/\eta \rceil$, and also let $G_X := G[N_G(X) \setminus W]$. Then Lemma 3.1(i) implies that G_X is a $(16\rho/(9\eta^2), d)$ -dense graph on at least $3\eta n/4$ vertices. But $4/(3\eta n) \ll 16\rho/(9\eta^2) \ll d, 1/C$, so Lemma 3.1(iv) implies that G_X contains a copy of K_C . Therefore, X lies in a copy of K_{C+r} that does not intersect W .

This implies that we may assume both X, Y lie in a copy of K_C that does not intersect W . Let X^* be the vertex set of the K_C containing X , and define Y^* analogously for Y . Choose $X' \subseteq X^*$ of size c that is disjoint from X . Since $|Y^*| - |Y| - |X| - |X'| = C - 2r - c \geq c$, we can choose $Y' \subseteq Y^*$ of size c that is disjoint from X, Y, X' . Apply Lemma 3.2 with $n, r, \eta, V(G), X', Y', X \cup Y \cup W$ playing the roles of n, r, η, U, X, Y, W to obtain $Z \subseteq V(G)$ that induces a copy of K_r and is disjoint from $X' \cup Y' \cup X \cup Y \cup W$; and there exist $X'' \subseteq X'$ and $Y'' \subseteq Y'$ such that $|X''| = |Y''| = r$ and $X'' \cup Y'' \subseteq N_G(Z)$. Notice that, by construction, each of $X \cup X'', X'' \cup Z, Z \cup Y'',$ and $Y'' \cup Y$ induce cliques, and the overlap of each consecutive pair induces a clique of size at least r . Further, none of these sets intersect with W . Thus $XX''ZY''Y$ induces an r -path. Thus there is an r -path with vertex set $X'' \cup Z \cup Y''$ (of length $3r$) that has the required property. \square

4. The regularity and blow-up lemmas and associated tools

4.1. Regularity

We will apply Szemerédi's regularity lemma in the proof of Theorem 1.2. For this, we need the following definitions. Given a bipartite graph G with vertex classes A and B and parameters $\varepsilon, \delta \in (0, 1)$,

- let $d_G(A, B) := \frac{e_G(A, B)}{|A||B|}$ be the *density* of G , and say that G is
- ε -*regular* if, for every $X \subseteq A$ and $Y \subseteq B$ with $|X| \geq \varepsilon|A|$ and $|Y| \geq \varepsilon|B|$, we have that $|d_G(A, B) - d_G(X, Y)| \leq \varepsilon$;
- (ε, δ) -*regular* if G is ε -regular and additionally $d_G(A, B) \geq \delta$;
- (ε, δ) -*superregular* if G is (ε, δ) -regular and additionally $d_G(a, B) \geq \delta|B|$ for every $a \in A$ and $d_G(b, A) \geq \delta|A|$ for every $b \in B$.

It will be convenient to use the degree form of the regularity lemma; this can be derived from the standard version [37].

Lemma 4.1 (Degree form of the regularity lemma). *For all $\varepsilon \in (0, 1)$ and $M' \in \mathbb{N}$, there exist $M, n_0 \in \mathbb{N}$ such that the following holds for all graphs G on $n \geq n_0$ vertices and $\delta \in (0, 1)$. There is a partition $V(G) = V_0 \cup V_1 \cup \dots \cup V_L$ and a spanning subgraph $G' \subseteq G$ such that*

- (i) $M' \leq L \leq M$.
- (ii) $|V_0| \leq \varepsilon n$.
- (iii) $|V_1| = \dots = |V_L| =: m$.
- (iv) $d_{G'}(x) \geq d_G(x) - (\delta + \varepsilon)n$ for all $x \in V(G)$.
- (v) For all $i \in [L]$, the graph $G'[V_i]$ is empty.
- (vi) For all $i \in [L]$, the graph $G'[V_i, V_j]$ is either empty or (ε, δ) -regular.

We call V_1, \dots, V_L the *clusters* of G and the vertices in V_0 the *exceptional vertices*. The graph G' is the *pure graph*. Note that the (ε, δ) -regular pairs may have very different densities. The *reduced graph* R of G with parameters ε, δ , and M' has vertex set $[L]$ and contains ij as an edge precisely when $G'[V_i, V_j]$ is (ε, δ) -regular.

The next lemma states that the reduced graph R of a locally dense graph G is still locally dense (with worse parameters) and, further, R inherits the minimum degree of G .

Lemma 4.2. *Let $0 < 1/n \ll 1/M' \ll \varepsilon \ll \delta \ll d \leq 1$; $1/M' \ll \rho \ll d$; $\delta \ll \eta$. Define $\rho^* := \max\{3\rho, 3\delta\}$. Let G be a (ρ, d) -dense graph of order n with $\delta(G) \geq (1/2 + \eta)n$. Apply Lemma 4.1 with parameters ε, δ , and M' to obtain a pure graph G' and a reduced graph R of G with $V(R) = [L]$. Then R is (ρ^*, d) -dense with $\delta(R) \geq (1/2 + \eta/2)L$.*

Proof. Here, (i)–(vi) will refer to the conclusions of Lemma 4.1. Parts (ii) and (iii) imply that

$$(1 - \varepsilon)n \leq mL \leq n. \quad (4.1)$$

Let $X \subseteq [L]$, and let $Y := \bigcup_{i \in X} V_i \subseteq V(G)$. So $|Y| = m|X|$. Then

$$d\left(\frac{|Y|}{2}\right) - \rho n^2 - |Y|(\delta + \varepsilon)n \leq e(G[Y]) - |Y|(\delta + \varepsilon)n \stackrel{(iv)}{\leq} e(G'[Y]) \stackrel{(v)}{\leq} e(R[X]) \cdot m^2$$

and so, dividing by m^2 ,

$$e(R[X]) \stackrel{(4.1)}{\geq} d \cdot \frac{|X|^2 - \frac{|X|}{m}}{2} - \rho \left(\frac{L}{1 - \varepsilon}\right)^2 - |X|(\delta + \varepsilon) \frac{L}{1 - \varepsilon} \geq d\left(\frac{|X|}{2}\right) - \rho^* L^2,$$

as required.

Let $i \in [L]$ and $x_i \in V_i$. Then $d_{G'}(x_i) \geq d_G(x_i) - (\delta + \varepsilon)n \geq (1/2 + \eta - \delta - \varepsilon)n$ by (iv). The number of clusters V_k of G containing some $y \in N_{G'}(x_i)$ is therefore at least

$$\frac{(1/2 + \eta - \delta - \varepsilon)n - |V_0|}{m} \geq \frac{(1/2 + \eta/2)n}{m} \geq (1/2 + \eta/2)L.$$

But then (vi) implies that i is adjacent to each of the vertices corresponding to these clusters in R . So $d_R(i) \geq (1/2 + \eta/2)L$, as required. \square

Note that in the case when $\rho \ll \delta$ in Lemma 4.2, R only inherits the property being locally dense with a significantly worse parameter playing the role of ρ . That is, now R is $(3\delta, d)$ -dense rather than (ρ, d) -dense.

The next well-known proposition states that (super)regular pairs are robust in the sense of adding or removing a small number of vertices. This version appears as Proposition 8 in [8].

Proposition 4.3. *Let G be a graph with $A, B \subseteq V(G)$ disjoint. Suppose that $G[A, B]$ is (ε, δ) -regular, and let $A', B' \subseteq V(G)$ be disjoint such that $|A \Delta A'| \leq \alpha|A'|$ and $|B \Delta B'| \leq \alpha|B'|$ for some $0 \leq \alpha < 1$. Then $G[A', B']$ is (ε', δ') -regular, with*

$$\varepsilon' := \varepsilon + 6\sqrt{\alpha} \quad \text{and} \quad \delta' := \delta - 4\alpha.$$

If, moreover, $G[A, B]$ is (ε, δ) -superregular and each vertex $x \in A'$ has at least $\delta'|B'|$ neighbours in B' and each vertex $x \in B'$ has at least $\delta'|A'|$ neighbours in A' , then $G[A', B']$ is (ε', δ') -superregular with ε' and δ' as above.

The following lemma is well known in several variations. The version here follows immediately from [36, Lemma 4.6].

Lemma 4.4. *Let $L \in \mathbb{N}$, and suppose that $0 < 1/m \ll \varepsilon \ll \delta, 1/\Delta, 1/L \leq 1$. Let R be a graph with $V(R) = [L]$ and $\Delta(R) \leq \Delta$. Let G be a graph with vertex partition V_1, \dots, V_L such that $|V_i| = m$ for all $1 \leq i \leq L$ and in which $G[V_i, V_j]$ is (ε, δ) -regular whenever $ij \in E(R)$. Then for each $i \in V(R)$, V_i contains a subset V'_i of size $(1 - \sqrt{\varepsilon})m$ such that for every edge ij of R , the graph $G[V'_i, V'_j]$ is $(4\sqrt{\varepsilon}, \delta/2)$ -superregular.*

4.2. Embedding lemmas

The next lemma is similar to a *partial embedding lemma* from [8, Lemma 10], which in turn is similar to an embedding lemma due to Chvátal, Rödl, Szemerédi, and Trotter [11]. Given a homomorphism from a graph H into the reduced graph R of G such that every pre-image is small, the lemma yields an embedding of some vertices of H into G while finding large candidate sets for the remaining vertices. Further (deviating from [8]), we would like to ensure that certain vertices of H are embedded into given target sets of large size.

Lemma 4.5 (Embedding lemma with target sets). *Let $0 < 1/n \ll 1/L \ll \varepsilon \ll c \ll \delta \ll 1/\Delta$, where $n, L \in \mathbb{N}$. Let G be an n -vertex graph, R an L -vertex graph, and H a graph on at most εn vertices such that*

- G has partition $\{V_a : a \in V(R)\}$, where $|V_a| \geq m \geq (1 - \varepsilon)n/L$ for all $a \in V(R)$ and $G[V_a, V_{a'}]$ is (ε, δ) -regular whenever $aa' \in E(R)$.
- $\Delta(H) \leq \Delta$, and there is a graph homomorphism $\phi : V(H) \rightarrow V(R)$ such that $|\phi^{-1}(a)| \leq 2\varepsilon m$ for all $a \in V(R)$.
- Let $X \cup Y$ be a partition of $V(H)$, and suppose that there is $W \subseteq X$ such that for each $w \in W$, there is a set $S_w \subseteq V_{\phi(w)}$ with $|S_w| \geq cm$.

Then there is an embedding f of $H[X]$ into G such that

- (i) $f(x) \in V_{\phi(x)}$ for all $x \in X$.
- (ii) $f(w) \in S_w$ for all $w \in W$.
- (iii) For all $y \in Y$, there exists $C_y \subseteq V_{\phi(y)} \setminus f(X)$ such that $C_y \subseteq N_G(f(x))$ for all $x \in N_H(y) \cap X$, and $|C_y| \geq cm$.

Since the proof of Lemma 4.5 is essentially identical to that of Lemma 10 from [8], we omit the proof.

We will also use the blow-up lemma of Komlós, Sárközy, and Szemerédi [27], which states that, for the purposes of embedding a spanning k -partite graph H of bounded degree, a graph G with a vertex partition into k classes, each pair of which is superregular, in fact behaves like a complete k -partite graph. Further, as in Lemma 4.5, one can ensure that a small fraction of the vertices of H are embedded into some given target sets.

Lemma 4.6 (Blow-up lemma [27]). *For every $d, \Delta, c > 0$ and $k \in \mathbb{N}$, there exist constants ε_0 and α such that the following holds. Let n_1, \dots, n_k be positive integers, $0 < \varepsilon < \varepsilon_0$, and G be a k -partite graph with vertex classes V_1, \dots, V_k where $|V_i| = n_i$ for $i \in [k]$. Let J be a graph on vertex set $[k]$ such that $G[V_i, V_j]$ is (ε, d) -superregular whenever $ij \in E(J)$. Suppose that H is a k -partite graph with vertex classes W_1, \dots, W_k of size at most n_1, \dots, n_k , respectively, with $\Delta(H) \leq \Delta$. Suppose further that there exists a graph homomorphism $\phi : V(H) \rightarrow V(J)$ such that $|\phi^{-1}(i)| \leq n_i$ for every $i \in [k]$. Moreover, suppose that in each class W_i , there is a set of at most αn_i special vertices y , each equipped with a set $S_y \subseteq V_i$ with $|S_y| \geq cn_i$. Then there is an embedding of H into G such that every special vertex y is mapped to a vertex in S_y .*

5. Finding the power of a Hamilton cycle

The next result states that for every $r \in \mathbb{N}$, every large locally dense n -vertex graph G with minimum degree at least $(1/2 + o(1))n$ contains the r th power of a Hamilton cycle. This is a very special case of our main result, Theorem 1.2.

Theorem 5.1. *For all $r, s \in \mathbb{N}$ and $d, \eta > 0$, there exist $\rho, n_0 > 0$ such that every (ρ, d) -dense graph G on $n \geq n_0$ vertices with $\delta(G) \geq (1/2 + \eta)n$ contains the r th power of a Hamilton cycle. In fact, for every $n' \in \mathbb{N}$ such that $n - s \leq n' \leq n$, G contains the r th power of a cycle covering precisely n' vertices.*

Note that Theorem 5.1 is an important tool in the proof of Theorem 1.2, in the same way that (an approximate version of) the result in [28] was used in the proof of Theorem 1.1. Indeed, Theorem 5.1 ensures that the reduced graph R of a graph G (as in Theorem 1.2) will contain a spanning $(4r - 1)$ -cycle. By (3.1), this implies R contains a spanning copy of Z_{ℓ}^{2r} . As outlined in Section 2, this copy of Z_{ℓ}^{2r} will be used as a ‘guide’ for embedding H into G .

We remark that one can give a significantly shorter proof of Theorem 5.1 if one only seeks the r th power of a cycle covering (say) at least $(1 - \eta)n$ vertices in G . However, for our application to Theorem 1.2, we (rather subtly) require that we have a $(4r - 1)$ -cycle in R covering all but a very small number of vertices (much fewer than $\rho|R|$ vertices in R can be left uncovered). So, such a weaker version of Theorem 5.1 is not sufficient.

The proof of Theorem 5.1 is an application of the *connecting–absorbing method*, a technique first developed by Rödl, Ruciński, and Szemerédi [32]. The first step in the proof is to find a short *absorbing* $2r$ -path P_{abs} in G that has the property that $V(P_{abs}) \cup Z$ spans an r -path in G (with the same start- and endpoints as P_{abs}) for any very small set of vertices Z . We then reserve a small pot of vertices V' (known as a *reservoir*), which will allow us to connect up pairs of paths into longer paths. Next we (via an application of the regularity lemma) find a collection \mathcal{P} of a constant number of vertex-disjoint $2C$ -paths that together cover almost all of the remaining vertices in G (here, C is chosen to be significantly bigger than r). Using vertices from the reservoir, we are then able to connect all the paths in \mathcal{P} together with P_{abs} to form a single r -cycle covering almost all the vertices in G . The remaining uncovered

Proof of Theorem 5.1. Note that if n is sufficiently large, then any n' -vertex induced subgraph G' of an n -vertex graph G as in the theorem must be $(2\rho, d)$ -dense with $\delta(G') \geq (1/2 + \eta/2)n'$. So as the r th power of a Hamilton cycle in G' corresponds to an r -cycle of length n' in G , it suffices to prove the first part of the statement of the theorem.

Further, it suffices to prove the theorem under the additional assumption that $d \ll \eta, 1/r$. Define constants $\rho, \varepsilon, \delta, d_1, \eta_0, \eta_1, \eta_2, \eta_3 > 0$, and $M' \in \mathbb{N}$, and apply the regularity lemma (Lemma 4.1) with inputs ε and M' to obtain some $M = M(\varepsilon, M')$ so that we have

$$0 < 1/M \leq 1/M' \ll \varepsilon \ll \delta \ll \rho \ll \eta_3 \ll \eta_2 \ll \eta_1 \ll \eta_0 \ll d_1 \ll d \ll \eta, 1/r. \quad (5.1)$$

Let n be sufficiently large, and consider any n -vertex graph G that is (ρ, d) -dense with $\delta(G) \geq (1/2 + \eta)n$.

Our initial aim is to construct a small absorbing $2r$ -path P_{abs} . The next claim provides the building blocks for this absorbing path.

Claim 5.2. *There exists a collection \mathcal{K} of at most $\eta_0 n/8r$ vertex-disjoint copies of K_{2r} in G such that:*

- (i) *Each $K \in \mathcal{K}$ is $d_1 n$ -extendable in G .*
- (ii) *Given any vertex $x \in V(G)$, there are at least $2\eta_2 n$ copies K of K_{2r} in \mathcal{K} so that $V(K) \subseteq N_G(x)$.*

Proof (of claim). Let \mathcal{C} denote the set of all copies of K_{2r} that are $d_1 n$ -extendable in G . So, certainly, $|\mathcal{C}| \leq n^{2r}$. Consider any $x \in V(G)$. Since $d_G(x) \geq n/2$, Lemma 3.1(i) implies that $G[N(x)]$ is $(4\rho, d)$ -dense. Thus, Lemma 3.1(iv) implies that there are at least $(d/2)^{\binom{2r+1}{2}} (n/2)^{2r} / (2r)!$ copies K of K_{2r} in \mathcal{C} so that $V(K) \subseteq N_G(x)$. (Here we use the property that $d^{2r}/2^{2r} \geq d_1$ by (5.1).) Let L_x denote the set of these copies of K_{2r} .

Let \mathcal{C}_p be obtained from \mathcal{C} by selecting each $K \in \mathcal{C}$ independently with probability

$$p := \frac{\eta_1}{n^{2r-1}}.$$

Hence,

$$\mathbb{E}(|\mathcal{C}_p|) \leq \eta_1 n \quad \text{and} \quad \mathbb{E}(|\mathcal{C}_p \cap L_x|) \geq (d/2)^{\binom{2r+1}{2}} \frac{(n/2)^{2r}}{(2r)!} \times \frac{\eta_1}{n^{2r-1}} \stackrel{(5.1)}{\geq} 2d_1 \eta_1 n$$

for each $x \in V(G)$. Thus, a Chernoff bound implies that, with high probability,

$$|\mathcal{C}_p| \leq 2\eta_1 n \quad \text{and} \quad |\mathcal{C}_p \cap L_x| \geq d_1 \eta_1 n \quad (5.2)$$

for all $x \in V(G)$. Let Y denote the number of pairs of copies of K_{2r} from \mathcal{C}_p that share at least one vertex. Then

$$\mathbb{E}(Y) \leq p^2 \binom{n}{2r} 2r \binom{n}{2r-1} \leq \eta_1^2 n.$$

By Markov's inequality, the probability that $|Y| \leq 2\eta_1^2 n$ is at least $1/2$. Therefore, there is a choice of \mathcal{C}_p such that this condition holds together with (5.2). Fix such a choice of \mathcal{C}_p ; then for each intersecting pair of cliques in \mathcal{C}_p , remove one of them to obtain a new collection \mathcal{K} . Note that the definition of \mathcal{C}_p and (5.2) implies that \mathcal{K} is a collection of at most $\eta_0 n/8r$ vertex-disjoint copies of K_{2r} in G . Further, since $d_1 \eta_1 n - 2\eta_1^2 n \geq d_1 \eta_1 n/2 \geq 2\eta_2 n$, we see that (ii) is satisfied, as desired. \square

With Claim 5.2 at hand, it is straightforward to obtain our desired absorbing $2r$ -path P_{abs} .

Claim 5.3. *G contains a $2r$ -path P_{abs} on at most $\eta_0 n$ vertices such that the following conditions hold.*

- (i) *Both the sets of the first and last $2r$ vertices on P_{abs} induce K_{2r} s in G that are $d_1 n$ -extendable. (Denote these sets by S and E , respectively.)*
- (ii) *Given any set $Z \subseteq V(G) \setminus V(P_{abs})$ of size at most $\eta_2 n$, there is an r -path P in G with vertex set $V(P_{abs}) \cup Z$ whose first $2r$ vertices are the elements of S (ordered as in P_{abs}) and last $2r$ vertices are the elements of E (ordered as in P_{abs}).*

Proof (of claim). Let \mathcal{K} be as in Claim 5.2, and enumerate its elements by K^1, \dots, K^t (so $t \leq \eta_0 n / 8r$). Apply Lemma 3.3 to G with $d_1, 2r, V(K^1), V(K^2), V(\mathcal{K})$ playing the roles of η, r, X, Y, W . (Note we can indeed apply this lemma by Claim 5.2(i) and as $|V(\mathcal{K})| \leq d_1 n / 4$.) We thus obtain a copy $P_1 = x_1^1 \dots x_{6r}^1$ of P_{6r}^{2r} in G , avoiding $V(\mathcal{K})$ such that $V(K^1)x_1^1 \dots x_{6r}^1$ and $x_1^1 \dots x_{6r}^1 V(K^2)$ both induce copies of P_{8r}^{2r} . Repeating this process iteratively, we obtain a collection P_1, \dots, P_{t-1} of vertex-disjoint copies of P_{6r}^{2r} in G so that $V(K^i)x_1^i \dots x_{6r}^i V(K^{i+1})$ induces a copy of P_{8r}^{2r} in G for each $1 \leq i \leq t-1$. (Here we have written $P_i = x_1^i \dots x_{6r}^i$.) Note that to ensure the P_i s are vertex-disjoint, at every step we update W ; so at step i , W contains $V(\mathcal{K})$ and the vertices from P_1, \dots, P_{i-1} (so $|W| \leq d_1 n / 4$).

Let P_{abs} denote the $2r$ -path obtained by the following concatenation:

$$P_{abs} := V(K^1)P_1V(K^2)P_2V(K^3) \dots V(K^{t-1})P_{t-1}V(K^t).$$

Notice that P_{abs} contains $(t-1)8r + 2r \leq 8rt \leq \eta_0 n$ vertices. Further, (i) follows since both K^1 and K^t are $d_1 n$ -extendable in G by definition of \mathcal{K} . Consider any set $Z = \{z_1, \dots, z_\ell\} \subseteq V(G) \setminus V(P_{abs})$ of size at most $\eta_2 n$. For each $1 \leq i \leq \ell$, by Claim 5.2(ii), there are at least $\eta_2 n$ choices for j_i such that:

- $2 \leq j_i \leq t-1$.
- $V(K^{j_i}) \subseteq N_G(z_i)$.

In particular, writing $V(K^{j_i}) = \{y_1, \dots, y_{2r}\}$, notice that

$$P_{j_i-1}y_1 \dots y_{2r}z_i y_{r+1} \dots y_{2r}P_{j_i} \quad (5.3)$$

is an r -path in G .

Since we have at least $\eta_2 n$ choices, we may define j_1, j_2, \dots, j_ℓ to be distinct. We can then insert each z_i into P_{abs} , as indicated by (5.3), to obtain the desired r -path P on $V(P_{abs}) \cup Z$. \square

Let S be as in Claim 5.3. Then $|N_G(S) \setminus V(P_{abs})| \geq d_1 n / 2$. Lemma 3.1(i) implies that $G_S := G[N_G(S) \setminus V(P_{abs})]$ is $(4\rho/d_1^2, d)$ -dense and therefore $(\rho^{1/2}, d)$ -dense. Set

$$C := \lceil 4r/\eta_3 \rceil. \quad (5.4)$$

Note that $\rho^{1/2} \ll d, 1/C$. Thus, Lemma 3.1(iv) implies that G_S contains a copy K_{2C+1}^S of K_{2C+1} . Similarly, we find a copy K_{2C+1}^E of K_{2C+1} in G that is disjoint from K_{2C+1}^S and P_{abs} so that $V(K_{2C+1}^E) \subseteq N_G(E)$. We will view both K_{2C+1}^S and K_{2C+1}^E as $2C$ -paths of length $2C+1$.

Set $G_0 := G \setminus (V(P_{abs}) \cup V(K_{2C+1}^S) \cup V(K_{2C+1}^E))$. Certainly, $|G_0| \geq (1 - 2\eta_0)n$ and

$$d_G(x, V(G_0)) \geq (1/2 + 3\eta/4)n \text{ for all } x \in V(G).$$

By selecting vertices randomly (and applying a Chernoff bound), one can obtain a set $V' \subseteq V(G_0)$ of $n' := \eta_3 n$ vertices such that

$$d_G(x, V') \geq (1/2 + \eta/2)n' \text{ for all } x \in V(G). \quad (5.5)$$

Set $G_1 := G[V']$ and $G_2 := G_0 \setminus V'$. Lemma 3.1(i) implies that G_1 is $(\rho/\eta_3^2, d)$ -dense and thus $(\rho^{1/2}, d)$ -dense. Similarly, G_2 is $(2\rho, d)$ -dense.

Apply Lemma 4.1 to G_2 with parameters ε, δ , and M' to obtain a partition V_0, V_1, \dots, V_ℓ of $V(G_2)$, pure graph G'_2 , and the reduced graph R of G_2 . Here, V_0 is the exceptional set on at most εn vertices, and $M' \leq \ell \leq M$. Set $m := |V_1| = \dots = |V_\ell|$. Then Lemma 4.2 implies that R is $(6\rho, d)$ -dense. In particular, Lemma 3.1(i) implies that R' is $(6\rho/\eta_3^2, d)$ -dense for any $R' \subseteq R$ on $\eta_3 \ell$ vertices.

Note that $1/\ell \ll 6\rho/\eta_3^2 \ll d, 1/C$. Thus, Lemma 3.1(iv) implies that every $R' \subseteq R$ on $\eta_3 \ell$ vertices contains a copy of K_{2C+1} . In particular, R contains a K_{2C+1} -tiling \mathcal{T} covering all but at most $\eta_3 \ell$ vertices.

Consider any copy K of K_{2C+1} in \mathcal{T} . The vertices of K correspond to clusters $V_{i_1}, \dots, V_{i_{2C+1}}$ in G_2 ; let G_K denote the subgraph of G'_2 induced by the vertices in these clusters combined. Every tuple (V_{i_j}, V_{i_k}) of such clusters forms an ε -regular pair of density at least δ in G_K . Moreover, Lemma 4.4 implies that for each such cluster V_{i_j} , there is a subset $V'_{i_j} \subseteq V_{i_j}$ of size $(1 - \varepsilon^{1/2})m$ so that (V'_{i_j}, V'_{i_k}) forms an $(4\varepsilon^{1/2}, \delta/2)$ -superregular pair in G_K (for each $1 \leq j \neq k \leq 2C+1$). The blow-up lemma (Lemma 4.6) now implies that G_K contains a $2C$ -path covering all but at most $(2C+1)\varepsilon^{1/2}m$ vertices in G_K .

Overall, this implies that G_2 contains a collection \mathcal{P} of at most $\ell/(2C+1) \leq M$ vertex-disjoint $2C$ -paths that together cover all but at most

$$\left((2C+1)\varepsilon^{1/2}m \times \frac{\ell}{2C+1} \right) + (\eta_3\ell \times m) + |V_0| \leq \varepsilon^{1/2}n + \eta_3n + \varepsilon n \stackrel{(5.1)}{\leq} 2\eta_3n \quad (5.6)$$

vertices in G_2 .

We will now use vertices in G_1 to connect all of the $2C$ -paths in $\mathcal{P} \cup \{K_{2C+1}^S, K_{2C+1}^E\}$ to obtain an r -path in G whose first $2C+1$ vertices are the vertices of K_{2C+1}^E and whose last $2C+1$ vertices are the vertices of K_{2C+1}^S . Note that we will have to reorder some of the vertices in the $2C$ -paths in \mathcal{P} , which is one reason we ‘drop’ from $2C$ -paths to an r -cycle. Label the $2C$ -paths in $\mathcal{P} \cup \{K_{2C+1}^S, K_{2C+1}^E\}$ by P_1, \dots, P_t , where $P_1 := K_{2C+1}^E$ and $P_t := K_{2C+1}^S$. In particular, note $M'/4C \leq t \leq M+2$.

For each P_i , let S_i denote the copy of K_C induced by the first C vertices on P_i ; let E_i denote the copy of K_C induced by the last C vertices on P_i ; and let P'_i denote the $2C$ -path obtained from P_i by deleting all vertices from S_i and E_i . (Note that P'_i is certainly non-empty.)

Claim 5.4. *Let $W \subseteq V(G_1)$ be arbitrary so that $|W| \leq \varepsilon n'$. Given any $1 \leq i \leq t-1$, there is an r -path P in G so that:*

- (i) $V(P) \cap V(G_2) = E_i \cup S_{i+1}$.
- (ii) $|V(P) \cap V(G_1)| = r$.
- (iii) *The first C vertices on P are precisely the vertices from E_i .*
- (iv) *The last C vertices on P are precisely the vertices from S_{i+1} .*
- (v) P is disjoint from W .

Proof (of claim). Apply Lemma 3.2 with $G, V', n', \eta_3, \sqrt{\rho}, d, E_i, S_{i+1}, W, r$ playing the roles of $G, U, n', \eta, \rho, d, X, Y, W, r$ to obtain a copy K of K_r in $G_1 = G[V']$ such that $V(K) \cap W = \emptyset$ (recall that $E_i \cup S_{i+1}$ is disjoint from V') and there exist $E'_i \subseteq E_i$ and $S'_{i+1} \subseteq S_{i+1}$ such that $|E'_i| = |S'_{i+1}| = r$ and $E'_i \cup S'_{i+1} \subseteq N_G(K)$.

Altogether, this implies that G_1 contains the desired r -path P . Indeed, we construct P so that the first $C-r$ vertices on P are those vertices in $E_i \setminus E'_i$ (in an arbitrary order); the next r vertices are the elements from E'_i ; after that, we take the vertices from K and then from S'_{i+1} ; the final $C-r$ vertices on P are from $S_{i+1} \setminus S'_{i+1}$. \square

With Claim 5.4 to hand, it is now easy to complete the proof of the theorem. Suppose that for some $j < t-1$, we have defined vertex-disjoint r -paths P_1^*, \dots, P_j^* such that, for each $i \leq j$, $P = P_i^*$ satisfies (i)–(iv) in Claim 5.4. Then define W to be all those vertices in an r -path P_1^*, \dots, P_j^* that lie in G_1 . So $|W| = jr \leq (M+2)r \leq \varepsilon n'$. Claim 5.4 then implies there is an r -path P_{j+1}^* in G that satisfies the conclusion of Claim 5.4 (where $j+1$ plays the role of i and P_{j+1}^* the role of P).

Thus, we obtain vertex-disjoint r -paths P_1^*, \dots, P_t^* such that, for each $i \leq t$, $P = P_i^*$ satisfies (i)–(iv) in Claim 5.4. Consider the concatenation

$$P^* := S_1 P'_1 P_1^* P'_2 P_2^* \dots P'_{t-1} P_{t-1}^* P'_t E_t.$$

This induces an r -path in G (with many additional edges). Further, note that by the definition of P_1 (and thus S_1), the first C vertices on P^* lie in K_{2C+1}^E and so are adjacent in G to every vertex in E .

Similarly, the last C vertices on P^* lie in K_{2C+1}^S and so are adjacent in G to every vertex in S . Thus, if we concatenate P^* together with P_{abs} , we obtain an r -cycle C^* in G (with many additional edges).

Note that, by (5.6), C^* covers every vertex in G except for at most $2\eta_3 n$ vertices in G_2 and at most $n' = \eta_3 n$ vertices in G_1 . Since $3\eta_3 n < \eta_2 n$, we may use the absorbing property (Claim 5.3(ii)) of P_{abs} to obtain the r th power of a Hamilton cycle in G , as required. \square

6. Lemmas for H

Our rough aim is to find ‘compatible’ partitions of the vertex sets of G and H that allow us to apply the embedding lemmas (Lemmas 4.5 and 4.6) to complete the embedding of H into G . In this section, we state and prove the so-called *lemmas for H* , whose input is some information about the structure of G and whose output is a suitable partition of H .

6.1. Partitioning a graph of low bandwidth: the basic lemma for H

At some stage of the proof, G will return some ‘ideal’ part sizes $\{m_{i,j} : (i,j) \in [\ell] \times [2r]\}$, where $\chi(H) \leq r$. We would then like to find a suitable partition of H , the parts of which are close to these ideal sizes (equivalently, a mapping f from $V(H)$ into $[\ell] \times [2r]$ whose pre-images have controlled size). This is the purpose of the next lemma. It guarantees that f is a graph homomorphism into Z_ℓ^{2r} and produces a small set B such that f restricted to $V(H) \setminus B$ is a graph homomorphism into a K_{2r} -factor (this is (B3)). Further, (B4) says that for the first few vertices of H (with respect to the bandwidth ordering of H), we have control of their images.

Before stating and proving Lemma 6.1, we would like to compare it to Lemma 8 in [9], the lemma for H in the bandwidth theorem. There, the assumptions on H are the same (in fact, slightly weaker), and the graph Z_ℓ^{2r} mentioned above is replaced by a given graph R of large minimum degree that contains a spanning subgraph S (very similar to Z_ℓ^r), which in turn contains a K_r -factor. Most edges are (and must be) mapped to the K_r -factor, which is much sparser than the K_{2r} -factor we have at our disposal. This means the proof of Lemma 8 in [9] is much harder to prove than our Lemma 6.1. Despite this, our lemma does not follow from the statement of Lemma 8 in [9], so we prove it here.

Lemma 6.1 (Basic lemma for H). *Let $n, r, \ell, \Delta \geq 1$ be integers, and let $\beta > 0$ be such that $0 < 1/n \ll 1/r, 1/\ell, 1/\Delta, \beta$. Let H be a graph on n vertices with $\Delta(H) \leq \Delta$, and assume that H has a labelling x_1, \dots, x_n of bandwidth at most βn and $\chi(H) \leq r$. Furthermore, suppose $\{m_{i,j} : (i,j) \in [\ell] \times [2r]\}$ is such that $\sum_{(i,j) \in [\ell] \times [2r]} m_{i,j} = n$; $m_{i,j} \geq 10\beta n$ for all $(i,j) \in [\ell] \times [2r]$; and $|m_{i,j} - m_{i,j'}| \leq 1$ whenever $i \in [\ell]$ and $j, j' \in [2r]$. Let $\chi : V(H) \rightarrow [r]$ be a proper colouring of H . Then there exist a mapping $f : V(H) \rightarrow [\ell] \times [2r]$ and a set of special vertices $B \subseteq V(H)$ with the following properties:*

- (B1) $B \cap \{x_1, \dots, x_{\beta n}\} = \emptyset$ and $|B| \leq 2\ell\beta n$.
- (B2) $||f^{-1}(i,j)| - m_{i,j}| \leq 10\beta n$ for every $(i,j) \in [\ell] \times [2r]$.
- (B3) For every edge $uv \in E(H)$, writing $f(u) = (i,j)$ and $f(v) = (i',j')$, we have $|i - i'| \leq 1$ and $j \neq j'$. If, additionally, $u, v \notin B$, then $i = i'$.
- (B4) For all $s \leq \beta n$, we have $f(x_s) = (1, \chi(x_s))$.

In particular, f yields a homomorphism from H to Z_ℓ^{2r} .

Note that the graph Z_ℓ^{2r} that appears in Lemma 6.1 will be found in the reduced graph R of G : since G is locally dense, R is also locally dense (see Lemma 4.2); and thus, by Theorem 5.1, we can find a spanning $(4r - 1)$ -cycle in R , which contains Z_ℓ^{2r} (see (3.1)).

Recall that each vertex in R corresponds to a unique cluster in G . In the proof of Theorem 1.2, the homomorphism f from H to $Z_\ell^{2r} \subseteq R$ will be a guide as to which cluster in G we should embed a vertex x into for most vertices $x \in V(H)$. That is, roughly speaking, if $f(x) = (i,j) \in V(R)$, we embed x into the cluster in G corresponding to (i,j) . Note, though, that f does not ‘guide’ us as to which vertices from

H we should embed into the exceptional set V_0 of G . So in the proof of Theorem 1.2, we in fact apply Lemma 6.1 to an almost spanning subgraph of H , rather than H itself; the remaining part of H is then embedded into G via an additional lemma for H (Lemma 6.2 in Section 6.2). In particular, Lemma 6.2 governs which vertices from H are embedded into V_0 . Property ($\mathcal{B}4$) of the homomorphism f is used to ensure that we can ‘fit’ the two Lemmas for H together to complete the embedding of H into G .

The idea of the proof of Lemma 6.1 is to first obtain a proper $2r$ -colouring χ' of H such that in any initial segment x_1, \dots, x_t of the bandwidth ordering of H , every colour is used roughly the same number of times in χ' . This then allows us to define f in a sequential way. That is, for some t_1 , we map each x_j in $\{x_1, \dots, x_{t_1}\}$ to $(1, \chi'(x_j))$; then, for some t_2 , we map each x_j in $\{x_{t_1+1}, \dots, x_{t_2}\}$ to $(2, \chi'(x_j))$, and so on.

Proof of Lemma 6.1. Let $N := \lceil 1/(2\beta) \rceil$, and partition the ordered vertices x_1, \dots, x_n into consecutive intervals A_1, A_2, \dots, A_{2N} , each of length βn (except possibly A_{2N} , which could be smaller). We view each interval as being ordered with the inherited bandwidth ordering.

We will first define a (proper) $2r$ -colouring $\chi' : V(H) \rightarrow [2r]$ by iteratively defining colourings χ'_i for $i \in [N]$ with the following properties:

$\mathcal{P}_1(i)$ $\chi'_i : \bigcup_{2 \leq t \leq 2i} A_t \rightarrow [2r]$ is a proper colouring of $H[\bigcup_{2 \leq t \leq 2i} A_t]$.

$\mathcal{P}_2(i)$ For all odd $2 \leq t \leq 2i$, we have $\chi'_i(A_t) \subseteq [r]$; and for all even $2 \leq t \leq 2i$, we have $\chi'_i(A_t) \subseteq [2r] \setminus [r]$.

$\mathcal{P}_3(i)$ Writing $b_i^j(s) := |\{x \in \bigcup_{2 \leq t \leq 2s} A_t : \chi'_i(x) = j\}|$ for all $j \in [2r]$ and $s \in [i]$, we have $|b_i^j(s) - b_i^{j'}(s)| \leq \beta n$ for all $(j, j') \in [[2r]]^2$ and $s \in [i]$.

For $\mathcal{P}_3(i)$, recall that $[[2r]]^2 := [r]^2 \cup ([2r] \setminus [r])^2$. Define $\chi'_1 : A_2 \rightarrow [2r]$ by setting $\chi'_1(x) = \chi(x) + r$. Clearly this satisfies $\mathcal{P}_1(1)$ – $\mathcal{P}_3(1)$, in particular as $|A_2| \leq \beta n$. Suppose we have defined χ'_i for some $i < N$ satisfying $\mathcal{P}_1(i)$ – $\mathcal{P}_3(i)$. By permuting the sets of colours $[r]$ and $[2r] \setminus [r]$, we can obtain a new proper $2r$ -colouring c_1 of $H[\bigcup_{2 \leq t \leq 2i} A_t]$ satisfying $\mathcal{P}_1(i)$ – $\mathcal{P}_3(i)$ and with the additional property that

$$|c_1^{-1}(1)| \geq \dots \geq |c_1^{-1}(r)| \quad \text{and} \quad |c_1^{-1}(r+1)| \geq \dots \geq |c_1^{-1}(2r)|. \quad (6.1)$$

Define $k : A_{2i+1} \cup A_{2i+2} \rightarrow [2r]$ by setting

$$k(x) = \begin{cases} \chi(x) & \text{if } x \in A_{2i+1} \\ \chi(x) + r & \text{if } x \in A_{2i+2}. \end{cases}$$

Clearly, k is a proper colouring of $H[A_{2i+1} \cup A_{2i+2}]$ since χ is. By permuting the sets of colours $[r]$ and $[2r] \setminus [r]$, we can obtain a new proper colouring c_2 of $H[A_{2i+1} \cup A_{2i+2}]$ from k such that

$$|c_2^{-1}(1)| \leq \dots \leq |c_2^{-1}(r)| \quad \text{and} \quad |c_2^{-1}(r+1)| \leq \dots \leq |c_2^{-1}(2r)| \quad (6.2)$$

(note that the ordering is reversed compared to (6.1)). Finally, define χ'_{i+1} by setting

$$\chi'_{i+1}(x) = \begin{cases} c_1(x) & \text{if } x \in \bigcup_{2 \leq t \leq 2i} A_t \\ c_2(x) & \text{if } x \in A_{2i+1} \cup A_{2i+2}. \end{cases} \quad (6.3)$$

The fact that $\mathcal{P}_2(i+1)$ holds is clear from $\mathcal{P}_2(i)$ and the definitions of c_1 , k , c_2 , and χ'_{i+1} .

To see that $\mathcal{P}_1(i+1)$ holds, let $x, y \in \bigcup_{2 \leq t \leq 2i+2} A_t$, where $xy \in E(H)$. We need to show that $\chi'_{i+1}(x) \neq \chi'_{i+1}(y)$. Let $2 \leq t, t' \leq 2i+2$ be such that $x \in A_t$ and $y \in A_{t'}$. Then $|t - t'| \leq 1$ since the intervals A_j respect the bandwidth ordering and each one (except perhaps A_{2N}) has size βn . If $|t - t'| = 1$, then $\mathcal{P}_2(i+1)$ implies that one of $\chi'_{i+1}(x), \chi'_{i+1}(y)$ lies in $[r]$ and the other in $[2r] \setminus [r]$, as required. So we may assume that $t = t'$. If $2 \leq t \leq 2i$, then $(\chi'_{i+1}(x), \chi'_{i+1}(y)) = (c_1(x), c_1(y))$. But c_1 is a proper colouring since it was obtained from χ'_i by permuting colours, and χ'_i is a proper colouring by $\mathcal{P}_1(i)$.

Suppose that $t \in \{2i+1, 2i+2\}$. Then, similarly, $(\chi'_{i+1}(x), \chi'_{i+1}(y)) = (c_2(x), c_2(y))$, and c_2 is a proper colouring since it was obtained from the proper colouring k by permuting colours. Thus $\mathcal{P}_1(i+1)$ holds.

For $\mathcal{P}_3(i+1)$, define for $j \in [2r]$ and $s \in [i+1]$

$$b_{i+1}^j(s) := \left| \left\{ x \in \bigcup_{2 \leq t \leq 2s} A_t : \chi'_{i+1}(x) = j \right\} \right|$$

and let $b_{i+1}^j := b_{i+1}^j(i+1) = |(\chi'_{i+1})^{-1}(j)|$. Then (6.3) implies that $b_{i+1}^j = |c_1^{-1}(j)| + |c_2^{-1}(j)|$ for all $j \in [2r]$. Now let $(j, j') \in [[2r]]^2$. Clearly, $|b_{i+1}^j(s) - b_{i+1}^{j'}(s)| \leq \beta n$ for all $s \in [i]$ since this is true for χ'_i and hence c_1 . So it remains to show that $|b_{i+1}^j - b_{i+1}^{j'}| \leq \beta n$. Equations (6.1) and (6.2) imply that the quantities $|c_1^{-1}(j)| - |c_1^{-1}(j')|$ and $|c_2^{-1}(j)| - |c_2^{-1}(j')|$ are never both positive and never both negative, since j and j' are in different orders. This implies that

$$\begin{aligned} |b_{i+1}^j - b_{i+1}^{j'}| &= ||c_1^{-1}(j)| - |c_1^{-1}(j')|| + ||c_2^{-1}(j)| - |c_2^{-1}(j')|| \\ &\leq \max \{ ||c_1^{-1}(j)| - |c_1^{-1}(j')||, ||c_2^{-1}(j)| - |c_2^{-1}(j')|| \}. \end{aligned}$$

Note that $||c_2^{-1}(j)| - |c_2^{-1}(j')|| \leq \beta n$. Further, c_1 was obtained from χ'_i by permuting colours in $[r]$ and in $[2r] \setminus [r]$, so there is some $(q, q') \in [[2r]]^2$ for which $c_1^{-1}(j) = (\chi'_i)^{-1}(q)$ and $c_1^{-1}(j') = (\chi'_i)^{-1}(q')$. Thus $||c_1^{-1}(j)| - |c_1^{-1}(j')|| = |b_i^q(i) - b_i^{q'}(i)|$, which is at most βn by $\mathcal{P}_3(i)$. Thus $\mathcal{P}_3(i+1)$ holds.

Therefore, we can obtain a colouring $\chi'_N : V(H) \setminus A_1 \rightarrow [2r]$ satisfying $\mathcal{P}_1(N) - \mathcal{P}_3(N)$. Finally, define $\chi' : V(H) \rightarrow [2r]$ by setting

$$\chi'(x) = \begin{cases} \chi'_N(x) & \text{if } x \in V(H) \setminus A_1 \\ \chi(x) & \text{if } x \in A_1. \end{cases} \quad (6.4)$$

The following properties hold:

- (i) $\chi' : V(H) \rightarrow [2r]$ is a proper colouring.
- (ii) For all odd $t \in [2N]$, we have $\chi'(A_t) \subseteq [r]$; and for all even $t \in [2N]$, we have $\chi'(A_t) \subseteq [2r] \setminus [r]$.
- (iii) Writing $d^j(s) := |\{x \in \bigcup_{t \in [s]} A_t : \chi'(x) = j\}|$ for all $j \in [2r]$ and $s \in [2N]$, we have $|d^j(s) - d^{j'}(s)| \leq 2\beta n$ for all $(j, j') \in [[2r]]^2$ and $s \in [2N]$.

Let $M_0 = n_0 := 0$. For all $i \in [\ell]$, let $M_i := \sum_{j \in [2r]} m_{i,j}$; and $n_i := \sum_{t \in [i]} M_t$. (Note that $n_\ell = n$.) Let $B_i := \{x_{n_{i-1}+1}, \dots, x_{n_i}\}$. So B_1, \dots, B_ℓ is a partition of $V(H)$ that respects the bandwidth ordering, and each interval inherits the bandwidth ordering. Let

$$B := \bigcup_{2 \leq i \leq \ell} \{x_{n_{i-1}+1}, \dots, x_{n_{i-1}+\beta n}\} \cup \bigcup_{1 \leq i \leq \ell-1} \{x_{n_i-\beta n+1}, \dots, x_{n_i}\},$$

and define $f : V(H) \rightarrow [\ell] \times [2r]$ by setting

$$f(x) := (i, \chi'(x)) \text{ if } x \in B_i. \quad (6.5)$$

We claim that f is the required mapping. Note $|B| = 2(\ell-1)\beta n$, and if $t \leq n_1 - \beta n$, then $x_t \notin B$. But $n_1 - \beta n \geq 9\beta n$, so certainly $\{x_1, \dots, x_{\beta n}\} \cap B = \emptyset$. Hence, $(\mathcal{B}1)$ holds. To show $(\mathcal{B}2)$, fix $i \in [\ell]$. Choose the smallest $p^- \in [2N]$ such that the first element of A_{p^-} lies in B_i and the largest $p^+ \in [2N]$ such that the last element of A_{p^+} lies in B_i . So B_i is the union of $\bigcup_{p^- \leq t \leq p^+} A_t$ together with a proper subset of $A_{p^- - 1}$ and a proper subset of $A_{p^+ + 1}$. Thus,

$$|f^{-1}(i, j)| = |\{x \in B_i : \chi'(x) = j\}| = d^j(p^+) - d^j(p^- - 1) \pm (|A_{p^- - 1}| + |A_{p^+ + 1}|) \quad (6.6)$$

for all $j \in [2r]$. Let $(j, j') \in [[2r]]^2$. Then the sizes of $f^{-1}(i, j)$ and $f^{-1}(i, j')$ do not differ much:

$$\begin{aligned} |f^{-1}(i, j) - f^{-1}(i, j')| &\stackrel{(6.6)}{\leq} \left| d^j(p^+) - d^{j'}(p^+) \right| + \left| d^j(p^- - 1) - d^{j'}(p^- - 1) \right| + 4\beta n \\ &\stackrel{(iii)}{\leq} 8\beta n. \end{aligned} \quad (6.7)$$

For any fixed $i \in [\ell]$,

$$\begin{aligned} S_1 &:= \sum_{j \in [r]} |f^{-1}(i, j)| \stackrel{(ii), (6.5)}{=} \sum_{t \text{ odd}} |B_i \cap A_t| \quad \text{and} \\ S_2 &:= \sum_{j \in [2r] \setminus [r]} |f^{-1}(i, j)| = \sum_{t \text{ even}} |B_i \cap A_t|. \end{aligned}$$

Therefore, $S_1 + S_2 = |B_i| = M_i$ and $|S_1 - S_2| \leq \beta n$. So $S_1, S_2 = M_i/2 \pm \beta n$. By definition of $m_{i,j}$ and M_i , we have that $|m_{i,j} - M_i/(2r)| \leq 1$ for all $j \in [2r]$. Now let $j \in [r]$. We have

$$\begin{aligned} ||f^{-1}(i, j)| - m_{i,j}| &\leq ||f^{-1}(i, j)| - M_i/(2r)| + 1 \leq \frac{1}{r} |r|f^{-1}(i, j)| - S_1| + 2\beta n \\ &\leq \frac{1}{r} \sum_{j' \in [r]} ||f^{-1}(i, j)| - |f^{-1}(i, j')|| + 2\beta n \stackrel{(6.7)}{\leq} 10\beta n, \end{aligned}$$

as required. The case when $j \in [2r] \setminus [r]$ is almost identical. Thus $(\mathcal{B}2)$ holds.

Now let $uv \in E(H)$, and write $f(u) = (i, j)$ and $f(v) = (i', j')$ for $i, i' \in [\ell]$ and $j, j' \in [2r]$. Since $|B_t| > \beta n$ for all $t \in [\ell]$ and $u \in B_i$ and $v \in B_{i'}$, we have that $|i - i'| \leq 1$ by consideration of the bandwidth ordering. We also have $j = \chi'(u)$ and $j' = \chi'(v)$, and χ' is a proper colouring of H , so $j \neq j'$. Suppose additionally that $u, v \notin B$. If $i \neq i'$, then u and v are separated by at least $2\beta n$ in the bandwidth ordering, so $uv \notin E(H)$, a contradiction.

Finally, if $s \leq \beta n$, then $x_s \in A_1 \cap B_1$. So $f(x_s) = (1, \chi'(x)) = (1, \chi(x))$ by (6.4). So $(\mathcal{B}4)$ holds. \square

6.2. Covering exceptional vertices: the second lemma for H

The second lemma for H will be used to find an embedding of a short initial segment of H (in bandwidth ordering) into G such that the exceptional set V_0 , obtained after applying the regularity lemma, lies in the image of this embedding. In fact the pre-image of V_0 will be a 2-independent set, which exists because H has small maximum degree and bandwidth. As well as embedding this initial segment, we would like to find target sets for its neighbours so that eventually we can extend this embedding to the whole of H .

Lemma 6.2 (Special lemma for H). *Let $n, r, L \geq 1$ be integers, and let $0 < 1/n \ll \beta \ll 1/L \ll \varepsilon \ll \rho \ll \eta \ll d, 1/r, 1/\Delta$. Let G be an n -vertex graph, R be an L -vertex graph, and $\{b_1, \dots, b_r\} \subseteq V(R)$ be such that*

- ($\mathcal{G}1$) G has vertex partition $\{V_0\} \cup \{V_a : a \in V(R)\}$ where $|V_0| \leq \varepsilon n$ and $|V_a| =: m$ for all $a \in V(R)$.
- ($\mathcal{G}2$) Each $v \in V_0$ is equipped with a subset $N_v \subseteq V(R)$ with $|N_v| \geq \eta L$.
- ($\mathcal{G}3$) R is (ρ, d) -dense, and $\delta(R) \geq (1/2 + \eta)L$.
- ($\mathcal{G}4$) $R[\{b_1, \dots, b_r\}] \cong K_r$, and $\{b_1, \dots, b_r\}$ lies in a copy of K_{18r/η^2} in R .

Then there exists an integer $s \leq \varepsilon^{1/4}n$ such that the following holds. Let H be a graph on $s + \beta n$ vertices with $\Delta(H) \leq \Delta$, and assume that H has a labelling $x_1, \dots, x_{s+\beta n}$ of bandwidth at most βn and $\chi(H) \leq r$. Let $X := \{x_1, \dots, x_s\}$ and $Y := \{x_{s+1}, \dots, x_{s+\beta n}\}$. Let $\chi : V(H) \rightarrow [r]$ be a proper colouring of H . Then there exists a mapping $f : V(H) \rightarrow V(R) \cup V_0$ with the following properties:

- ($\mathcal{D}1$) Setting $I := f^{-1}(V_0)$, we have that I is a subset of X that is 2-independent in H , and each vertex in V_0 is mapped onto from a unique vertex in H (so $|I| = |V_0|$).

- (D2) For all $v \in V_0$, setting $W_v := N_H(f^{-1}(v))$, we have $W_v \subseteq X$ and $f(W_v) \subseteq N_v$.
 (D3) $|f^{-1}(a)| \leq \varepsilon^{1/4}m$ for every $a \in V(R)$.
 (D4) For every edge $uv \in E(H)$ such that $f(u), f(v) \notin V_0$, we have $f(u)f(v) \in E(R)$.
 (D5) For all $y \in Y$, we have $f(y) = b_{\chi(y)}$.

To prove Lemma 6.2, we will need an auxiliary result, Lemma 6.3, which produces a ‘framework’ F in the reduced graph that we will later use to find f . This framework F is a $2r$ -trail such that for every $v \in V_0$, there is a copy T of K_{2r} in F such that $V(T) \subseteq N_v$.

Lemma 6.3. *Let $0 < 1/n \ll 1/L \ll \varepsilon \ll \rho \ll \eta \ll d, 1/r \leq 1$. Let G be an n -vertex graph, R be an L -vertex graph, and $\{b_1, \dots, b_r\} \subseteq V(R)$ be such that*

- (G1) G has vertex partition $\{V_0\} \cup \{V_a : a \in V(R)\}$ where $|V_0| \leq \varepsilon n$ and $|V_a| = m$ for all $a \in V(R)$.
 (G2) Each $v \in V_0$ is equipped with a subset $N_v \subseteq V(R)$ with $|N_v| \geq \eta L$.
 (G3) R is (ρ, d) -dense, and $\delta(R) \geq (1/2 + \eta)L$.
 (G4) $R[\{b_1, \dots, b_r\}] \cong K_r$, and $\{b_1, \dots, b_r\}$ lies in a copy of K_{18r/η^2} in R .

Then there exist an integer $K \leq L^{2r}$ and a subgraph $F \subseteq R$ such that

- (F1) F is a $2r$ -trail with ordering a_1, \dots, a_t where $t = (8K + 1)r$.
 (F2) There is a partition $V_0 = V_0^1 \cup \dots \cup V_0^K$ such that $N_v \supseteq \{a_{8(i-1)r+1}, \dots, a_{8(i-1)r+2r}\}$ for all $v \in V_0^i$ and $|V_0^i| \leq \sqrt{\varepsilon}m/L^{2r-1}$ for all $i \in [K]$.
 (F3) $(a_{t-r+1}, \dots, a_t) = (b_1, \dots, b_r)$.
 (F4) Every $a \in V(R)$ appears at most $L^{2r-1}/\varepsilon^{1/12}$ times in the sequence a_1, \dots, a_t .

Proof. Assume without loss of generality that $V(R) = [L]$. We first prove the following claim.

Claim 6.4. *There is a $K \leq L^{2r}$ and a set $\mathcal{T} = \{T_1, \dots, T_K\}$ of $((d/2)^{2r}\eta L)$ -extendable copies of K_{2r} in R such that there is a partition $V_0 = V_0^1 \cup \dots \cup V_0^K$ with the property that, for all $k \in [K]$, we have $|V_0^k| \leq \sqrt{\varepsilon}m/L^{2r-1}$ and $T_k \subseteq R[N_v]$ for all $v \in V_0^k$.*

Proof (of claim). By Lemma 3.1(i), we see that $R_v := R[N_v]$ is $(\rho L^2/|N_v|^2, d)$ -dense and hence $(\sqrt{\rho}, d)$ -dense, where we used (G2) and the fact that $\rho/\eta^2 < \sqrt{\rho}$. Lemma 3.1(iv) implies that R_v contains at least $(d/2)^{\binom{2r+1}{2}}\eta^{2r}L^{2r}/(2r)!$ copies of K_{2r} , each of which is $((d/2)^{2r}\eta L)$ -extendable in R_v (and thus R).

Let T_1, \dots, T_K be the set of $((d/2)^{2r}\eta L)$ -extendable copies of K_{2r} in R . So

$$K \leq \binom{L}{2r} \leq L^{2r}. \quad (6.8)$$

Then there is a partition $V_0^1 \cup \dots \cup V_0^K$ of V_0 into subsets (some of which may be empty) such that for all $k \in [K]$ and $v \in V_0^k$, we have that $R_v \supseteq T_k$ and

$$|V_0^k| \leq \frac{|V_0|}{(d/2)^{\binom{2r+1}{2}}\eta^{2r}L^{2r}/(2r)!} \stackrel{(G1)}{\leq} \frac{\varepsilon m}{(d/2)^{\binom{2r+1}{2}}\eta^{2r}(1-\varepsilon)L^{2r-1}/(2r)!} \leq \frac{\sqrt{\varepsilon}m}{L^{2r-1}},$$

as desired. \square

Let $\mathcal{T} := \{T_i : i \in [K]\}$ be obtained from the claim. To complete the proof, we will use the connecting lemma (Lemma 3.3) to join the K_{2r} s in \mathcal{T} into a $2r$ -trail. In so doing, we have to be careful not to visit any $a \in [L]$ too many times so as to ensure that (F4) holds.

Suppose, for some $0 \leq i < K - 1$ and all $j \in [i]$, we have obtained a copy $P_j = x_j^1 \dots x_j^{6r}$ of $P_{6r}^{2r} \subseteq R$ such that

$$\mathcal{P}_1(i) \quad V(T_j)x_j^1 \dots x_j^{6r} \text{ induces a copy } P'_j \text{ of } P_{8r}^{2r}.$$

$\mathcal{P}_2(i) \ x_j^1 \dots x_j^{6r} V(T_{j+1})$ induces a copy P_j'' of P_{8r}^{2r} .

$\mathcal{P}_3(i)$ Each $a \in [L]$ lies in at most $\varepsilon^{-1/12} L^{2r-1}/2$ of the $2r$ -paths P_1, \dots, P_i .

We would like to find P_{i+1} such that $\mathcal{P}_1(i+1) - \mathcal{P}_3(i+1)$ hold. We will say that $a \in [L]$ is *bad* if it appears in at least $\varepsilon^{-1/12} L^{2r-1}/3$ of P_1, \dots, P_i . Let D be the set of bad a . Since each P_j contains $6r$ vertices, we have

$$|D| \leq \frac{6ir}{\varepsilon^{-1/12} L^{2r-1}/3} \leq \frac{18\varepsilon^{1/12} Kr}{L^{2r-1}} \stackrel{(6.8)}{\leq} 18\varepsilon^{1/12} rL.$$

Recall from Claim 6.4 that T_{i+1} and T_{i+2} are both $((d/2)^{2r} \eta L)$ -extendable copies of K_{2r} in R . Since $\eta \ll d, 1/r$, they are $\eta^2 L$ -extendable copies. Apply Lemma 3.3 with $R, V(T_{i+1}), V(T_{i+2}), D, 2r, \eta^2$ playing the roles of G, X, Y, W, r, η to obtain a copy P_{i+1} of $P_{6r}^{2r} = x_{i+1}^1 \dots x_{i+1}^{6r}$ that avoids D and such that $V(T_{i+1})x_{i+1}^1 \dots x_{i+1}^{6r}$ induces a copy P'_{i+1} of P_{8r}^{2r} , and $x_{i+1}^1 \dots x_{i+1}^{6r} V(T_{i+2})$ induces a copy P''_{i+1} of P_{8r}^{2r} . So $\mathcal{P}_1(i+1)$ and $\mathcal{P}_2(i+1)$ hold. Now let $a \in [L]$. If $a \notin V(P_{i+1})$, then a lies in at most $\varepsilon^{-1/12} L^{2r-1}/2$ of P_1, \dots, P_{i+1} by $\mathcal{P}_3(i)$. Otherwise, since P_{i+1} avoids D , a lies in at most $\varepsilon^{-1/12} L^{2r-1}/3 + 1 < \varepsilon^{-1/12} L^{2r-1}/2$ of P_1, \dots, P_{i+1} . So $\mathcal{P}_3(i+1)$ holds. Therefore, we can find P_1, \dots, P_{K-1} satisfying $\mathcal{P}_1(K-1) - \mathcal{P}_3(K-1)$.

Next we want to find a $2r$ -path between T_K and $\{b_1, \dots, b_r\}$. Let $\{b'_1, \dots, b'_r\}$ be such that $\{b_1, \dots, b_r, b'_1, \dots, b'_r\}$ lies in a copy of K_{18r/η^2} in R (such vertices exist by (G4)). Apply Lemma 3.3 with $R, V(T_K), \{b_1, \dots, b_r, b'_1, \dots, b'_r\}, \emptyset, 2r, \eta^2$ playing the roles of G, X, Y, W, r, η to obtain a copy P_K of $P_{6r}^{2r} = x_K^1 \dots x_K^{6r}$ such that $V(T_K)x_K^1 \dots x_K^{6r}$ induces a copy P'_K of P_{8r}^{2r} , and furthermore $x_K^1 \dots x_K^{6r} b_1 \dots b_r b'_1 \dots b'_r$ induces a copy of P_{8r}^{2r} ; thus $x_K^1 \dots x_K^{6r} b_1 \dots b_r$ induces a copy P''_K of P_{8r}^{2r} . (Note that the vertices b'_1, \dots, b'_r were introduced only so that we could apply Lemma 3.3.) Clearly, $\mathcal{P}_3(K-1)$ implies that each $a \in [K]$ lies in at most $L^{2r-1} \varepsilon^{-1/12}/2 + 1$ of P_1, \dots, P_K .

Writing $V(T_i) = \{y_i^1, \dots, y_i^{2r}\}$ for all $i \in [K]$, R contains a $2r$ -trail $F' := \bigcup_{i \in [K]} (P'_i \cup P''_i)$ of length $(8K+1)r = t$, with ordering given by

$$(a_1, \dots, a_t) := (y_1^1, \dots, y_1^{2r}, x_1^1, \dots, x_1^{6r}, y_2^1, \dots, y_2^{2r}, \dots, y_K^1, \dots, y_K^{2r}, x_K^1, \dots, x_K^{6r}, b_1, \dots, b_r).$$

By construction, (F1) and (F3) hold.

We have that $V(T_i) = \{a_{8(i-1)r+1}, \dots, a_{8(i-1)r+2r}\}$ for all $i \in [K]$, which together with Claim 6.4 implies that (F2) holds. Now let $a \in [L]$. Then $\mathcal{P}_3(K-1)$ implies that a plays the role of some x_i^j with $(i, j) \in [K] \times [6r]$ at most $\varepsilon^{-1/12} L^{2r-1}/2 + 1$ times. Since each T_i with $i \in [K]$ is a distinct copy of K_{2r} in R , we see that a plays the role of some y_i^j with $(i, j) \in [K] \times [2r]$ at most $\binom{L-1}{2r-1} \leq L^{2r-1}$ times. Clearly, a plays the role of at most one of b_1, \dots, b_r . Thus the number of times a appears in the sequence a_1, \dots, a_t is at most $\varepsilon^{-1/12} L^{2r-1}/2 + L^{2r-1} + 2 \leq \varepsilon^{-1/12} L^{2r-1}$. So (F4) holds. \square

Armed with Lemma 6.3, we can now prove Lemma 6.2. The proof proceeds by splitting $V(H)$ into segments and assigning each one to a copy of K_r in R , according to the framework F . For example, the first segment of $V(H)$ will be assigned to $\{a_1, \dots, a_r\}$; and, more specifically, those vertices coloured i by χ will be mapped to a_i . In those special segments assigned to vertex sets of K_r s that lie in N_v for $v \in V_0^i$, we choose $|V_0^i|$ special vertices to be the pre-images of vertices in V_0^i . The property (F4) of F will ensure that not too many vertices are mapped to the same cluster of R .

Proof of Lemma 6.2. Let G and R be as in the statement of the lemma. Without loss of generality, we will assume that $V(R) = [L]$. Apply Lemma 6.3 to obtain $K \leq L^{2r}$ and $F \subseteq R$ such that

(F1) F is a $2r$ -trail with ordering a_1, \dots, a_t where $t = (8K+1)r$.

(F2) There is a partition $V_0 = V_0^1 \cup \dots \cup V_0^K$ such that $N_v \supseteq \{a_{8(i-1)r+1}, \dots, a_{8(i-1)r+2r}\}$ for all $v \in V_0^i$ and $|V_0^i| \leq \sqrt{em}/L^{2r-1}$ for all $i \in [K]$.

(F3) $(a_{t-r+1}, \dots, a_t) = (b_1, \dots, b_r)$.

($\mathcal{F}4$) Every $a \in [L]$ appears at most $L^{2r-1}/\varepsilon^{1/12}$ times in the sequence a_1, \dots, a_t .

Let

$$s := 8K\varepsilon^{1/3}m/L^{2r-1} \leq 8L\varepsilon^{1/3}m \stackrel{(\mathcal{F}1)}{\leq} 8\varepsilon^{1/3}n \leq \varepsilon^{1/4}n.$$

For all $i \in [K]$, let

$$u_i := |V_0^i| \stackrel{(\mathcal{F}2)}{\leq} \sqrt{\varepsilon}m/L^{2r-1} \quad \text{and} \quad b := \varepsilon^{1/3}m/L^{2r-1} > 100\beta mL \stackrel{(\mathcal{F}1)}{>} 99\beta n. \quad (6.9)$$

Let H, X, Y be as in the statement of the lemma. Define a partition of $X \cup Y = \{x_1, \dots, x_{s+\beta n}\}$ into $8K + 1$ intervals

$$B_1^1, B_1^2, \dots, B_1^8, B_2^1, B_2^2, \dots, B_2^8, B_3^1, \dots, B_K^1, B_K^2, \dots, B_K^8, B_{K+1}^1$$

where $|B_i^j| = b$ for all $(i, j) \in [K] \times [8]$; $|B_{K+1}^1| = \beta n$; and the first b vertices x_1, \dots, x_b in $X \cup Y$ form B_1^1 , the next b vertices in $X \cup Y$ form B_1^2 , and so on. In particular, $B_{K+1}^1 = Y$, and each interval comes equipped with the ordering inherited from the bandwidth ordering of H . The first claim identifies a set $I \subseteq X$ that will be the pre-image of V_0 in our desired mapping. Recall that given a graph J and $A \subseteq V(J)$, we say that A is *2-independent* if every pair of vertices in A is at distance at least 3 in J . In other words, A is an independent set and, additionally, the neighbourhoods of different vertices in A are disjoint.

Claim 6.5. *For each $i \in [K]$, there exists a 2-independent set $I_i \subseteq B_i^1$ (with respect to H) of size u_i such that $W(i) := \bigcup_{y \in I_i} N_H(y) \subseteq B_i^1$. Further, $I := \bigcup_{i \in [K]} I_i$ is a 2-independent set in H .*

Proof (of claim). Obtain A_i from B_i^1 by removing the first $2\beta n$ and last $2\beta n$ elements (which is possible by (6.9)). Suppose we have obtained a 2-independent set $I^j \subseteq A_i$ of size $0 \leq j < u_i$. Then for any $y \in A_i$, the set $I^j \cup \{y\}$ is a 2-independent set in H of size $j + 1$ if $y \notin I^j \cup N_H(y') \cup N_H(N_H(y'))$ for any $y' \in I^j$. The number of excluded y is at most

$$\begin{aligned} |I^j| + \sum_{x \in I^j} d_H(x) + \sum_{x \in I^j} \sum_{z \in N_H(x)} d_H(z) &\leq |I^j|(1 + \Delta + \Delta^2) \leq 2\Delta^2 u_i \\ &\stackrel{(6.9)}{\leq} 2\Delta^2 \sqrt{\varepsilon}m/L^{2r-1} \stackrel{(6.9)}{<} b - 4\beta n = |A_i|. \end{aligned}$$

Therefore, we can find a 2-independent set $I_i := I^{u_i}$ of size u_i in A_i . This, together with the bandwidth property and the definition of A_i , implies that $W(i) \subseteq \bigcup_{y \in I_i} (N_H(y) \cup N_H(N_H(y))) \subseteq B_i^1$. Thus there is no edge between $N_H(I_i)$ and $N_H(I_{i'})$ for $i \neq i'$. So $I = \bigcup_{i \in [K]} I_i$ is a 2-independent set in H , proving the claim. \square

Let $\chi : V(H) \rightarrow [r]$ be the given proper colouring of H . A second claim finds a suitable homomorphism $\phi : V(H) \rightarrow V(F)$ on which f will be based.

Claim 6.6. *For each $(i, j) \in [K] \times [8] \cup \{(K + 1, 1)\}$, let*

$$\phi(x) := a_{(8(i-1)+(j-1))r+\chi(x)} \quad \text{if } x \in B_i^j.$$

Then $\phi : V(H) \rightarrow V(F)$ is a graph homomorphism such that $|\phi^{-1}(a)| \leq \varepsilon^{1/4}m$ for all $a \in [L]$.

Proof (of claim). Note first that if a_k is in the image of ϕ for some $k \in \mathbb{N}$, then, recalling ($\mathcal{F}1$), we have that $k \in [t]$, so $V(F) \supseteq \phi(V(H))$. Let us check that ϕ is a homomorphism. Let $xy \in E(H)$. Let $(i, j), (i', j') \in [K] \times [8] \cup \{(K + 1, 1)\}$ be such that $x \in B_i^j$ and $y \in B_{i'}^{j'}$. Since H has bandwidth

at most βn and $|B_i^j|, |B_{i'}^{j'}| > \beta n$, we must have $(i', j') \in \{(i, j-1), (i, j), (i, j+1)\}$, where we let $(i, 9) := (i+1, 1)$ and $(i, 0) := (i-1, 8)$. So, writing

$$T_i := \{a_{(8(i-1)+(j-1))r+p} : p \in [r]\} \quad \text{and} \quad T_{i'} := \{a_{(8(i'-1)r+(j'-1))+p} : p \in [r]\}$$

we either have $T_i = T_{i'}$, or T_i and $T_{i'}$ are consecutive intervals in a_1, \dots, a_t , each of length r . In both cases, we have $\phi(x) \neq \phi(y)$ (in the first case, this follows from the fact that $\chi(x) \neq \chi(y)$). But $(\mathcal{F}1)$ now implies that $F[T_i \cup T_{i'}]$ is a clique, so since $\phi(x) \in T_i$ and $\phi(y) \in T_{i'}$ are distinct, $\phi(x)\phi(y) \in E(F)$, as required.

For the final assertion, each $a \in V(F)$ appears at most $L^{2r-1}/\varepsilon^{1/12}$ times in the sequence a_1, \dots, a_t by $(\mathcal{F}4)$. So, writing $\theta : V(H) \rightarrow [t]$ where $\phi(x) = a_{\theta(x)}$, we have

$$|\phi^{-1}(a)| \leq \frac{L^{2r-1}}{\varepsilon^{1/12}} \cdot \max_{k \in [t]} |\theta^{-1}(k)| \leq \frac{L^{2r-1}b}{\varepsilon^{1/12}} \stackrel{(6.9)}{=} \varepsilon^{1/4}m,$$

as desired. \square

Now let $H' := H \setminus I$, where $I := \bigcup_{k \in [K]} I_k$ and $W := \bigcup_{k \in [K]} W(k)$, where $W(k)$ is defined in Claim 6.5. Note also that $W \subseteq V(H')$ since, by Claim 6.5, I is an independent set.

Let $g : I \rightarrow V_0$ be a bijection such that $g(I_i) = V_0^i$ for all $i \in [K]$ (which is clearly possible by Claim 6.5 and (6.9)). Since I_i is a 2-independent set in H , the set of neighbourhoods $N_H(y)$ is pairwise disjoint over all $y \in I_i$. So for each $w \in W(i)$, there is a unique $y \in I_i$ for which $w \in N_H(y)$. Claim 6.6 implies that $|\phi^{-1}(a)| \leq \varepsilon^{1/4}m$ for all $a \in [L]$.

We claim that $f : V(H) \rightarrow [L] \cup V_0$ given by

$$f(x) = \begin{cases} \phi(x) & \text{if } x \in V(H) \setminus I \\ g(x) & \text{if } x \in I \end{cases} \quad (6.10)$$

is the required mapping. Note that $f(V(H) \setminus I) \subseteq [L]$ and $f(I) \subseteq V_0$. For $(\mathcal{D}1)$, note that, by Claim 6.5 and (6.10), $f(I) = g(I) = V_0$, $|I| = |V_0|$ and I is a 2-independent subset of X . For $(\mathcal{D}2)$, let $v \in V_0$ and $W_v := N_H(f^{-1}(v))$. Let $k \in [K]$ be such that $v \in V_0^k$. Then $f^{-1}(v) = g^{-1}(v) \in I_k$. So $W_v \subseteq W(i) \subseteq B_i^1 \subseteq X$. Let $x \in W_v \subseteq B_i^1$. By Claim 6.6 and $(\mathcal{F}2)$, we have that $f(x) = \phi(x) = a_{8(i-1)+\chi(x)} \in N_v$. This completes the proof of $(\mathcal{D}2)$.

For $(\mathcal{D}3)$, let $a \in V(R)$. Then $f^{-1}(a) \subseteq \phi^{-1}(a)$ has size at most $\varepsilon^{1/4}m$ by Claim 6.6. For $(\mathcal{D}4)$, let $uv \in E(H)$ be such that $f(u), f(v) \notin V_0$. So $u, v \in V(H) \setminus I$ and $f(u) = \phi(u)$ and $f(v) = \phi(v)$. By Claim 6.6, $\phi : V(H) \rightarrow V(F)$ is a homomorphism, so $f(u)f(v) \in E(F) \subseteq E(R)$. Finally, for $(\mathcal{D}5)$, we have that $Y \cap I = \emptyset$ by Claim 6.5, so for any $y \in Y$, we have $f(y) = \phi(y) = a_{t-r+\chi(y)} = b_{\chi(y)}$ by $(\mathcal{F}3)$. \square

7. The lemma for G : adjusting cluster sizes

Recall the definition of Z_ℓ^r from Section 3.1.1 and in particular that it contains a K_r -factor. Our goal in this section is to prove Lemma 7.1. Roughly speaking, it supposes that the reduced graph R of G contains a spanning copy of Z_ℓ^{2r} , its clusters V_1, \dots, V_L are equally sized, and pairs of clusters corresponding to the K_{2r} -factor $\ell \cdot K_{2r}$ in Z_ℓ^{2r} are superregular. Then we can adjust V_1, \dots, V_L slightly by reallocating a small number of vertices so that they have given sizes, at the expense of now having superregular pairs corresponding to a K_r -factor $2\ell \cdot K_r$.

To formalise the structural properties we need from G , we make the following definition (very similar to Definition 8.1 in [35]).

Definition 1 (r -Cycle structure). Given integers n, ℓ, r , a graph G on n vertices, and constants $\varepsilon, \delta > 0$, we say that G has an $(R, \ell, r, \mathcal{V}, \varepsilon, \delta)$ -cycle structure \mathcal{C} if the following hold:

- ($\mathcal{C}1$) $\mathcal{V} = \{V_0\} \cup \{V_{i,j} : (i, j) \in [\ell] \times [r]\}$ is a partition of $V(G)$, where $|V_0| \leq \varepsilon n$.
- ($\mathcal{C}2$) R has vertex set $[\ell] \times [r]$ and $R \supseteq Z_\ell^r$ and $G[V_{i,j}, V_{i',j'}]$ is (ε, δ) -regular whenever $(i, j)(i', j') \in E(R)$.
- ($\mathcal{C}3$) $G[V_{i,j}, V_{i,j'}]$ is (ε, δ) -superregular whenever $i \in [\ell]$ and $1 \leq j < j' \leq r$.

We say that \mathcal{V} induces \mathcal{C} . If $V_0 = \emptyset$, we say that \mathcal{C} is *spanning*.

The next definition concerns a convenient relabelling of the vertex set of a graph, which we will use for the reduced graph R .

Definition 2 (Bijection ϕ_ℓ^{2r}). Given integers r, ℓ , define $\phi_\ell^{2r} : [\ell] \times [2r] \rightarrow [2\ell] \times [r]$ by setting

$$\phi_\ell^{2r}(i, j) = \left((2i-1) + \left\lfloor \frac{j}{r} \right\rfloor, j - \left(\left\lfloor \frac{j}{r} \right\rfloor - 1 \right) r \right), \quad \text{for all } (i, j) \in [\ell] \times [2r]. \quad (7.1)$$

It is easy to check that ϕ_ℓ^{2r} is a bijection and

$$\begin{aligned} & \phi_\ell^{2r}(1, 1) \dots \phi_\ell^{2r}(1, r) \phi_\ell^{2r}(1, r+1) \dots \phi_\ell^{2r}(1, 2r) \dots \phi_\ell^{2r}(\ell, r+1) \dots \phi_\ell^{2r}(\ell, 2r) \\ &= (1, 1) \dots (1, r) (2, 1) \dots (2, r) \dots (2\ell, 1) \dots (2\ell, r). \end{aligned}$$

This implies that for all $a \in [2\ell]$ and distinct $b, b' \in [r]$, there are $i \in [\ell]$ and $(j, j') \in [[2r]]^2$ such that $(\phi_\ell^{2r}(i, j), \phi_\ell^{2r}(i, j')) = ((a, b), (a, b'))$.

Given a graph R and a bijection $\phi : V(R) \rightarrow V$ to some set V , we write $\phi(R)$ for the graph with vertex set $\{\phi(x) : x \in V(R)\}$ and edge set $\{\phi(x)\phi(y) : xy \in E(R)\}$. So $\phi(R) \cong R$.

In the language of Definition 1, the main result of this section states that, given a graph with a (spanning) $2r$ -cycle structure, we can obtain from it an r -cycle structure that is almost balanced, but the exact deviation from perfect balancedness can be controlled.

Lemma 7.1 (Lemma for G). Let $n, \ell, m, r \in \mathbb{N}$ and $0 < 1/n \ll \xi \ll 1/\ell \ll \varepsilon \ll \delta < 1/r$. Suppose that G is a graph on n vertices with a spanning $(R, \ell, 2r, \mathcal{V}, \varepsilon, \delta)$ -cycle structure, where $\mathcal{V} = \{V_{i,j} : (i, j) \in [\ell] \times [2r]\}$ and $|V_{i,j}| = m$ for all $(i, j) \in [\ell] \times [2r]$. Let $\{\tau_{a,b} \in \mathbb{Z} : (a, b) \in [2\ell] \times [r]\}$ be such that $0 \leq \tau_{a,b} \leq \varepsilon m$ for all $(a, b) \in [2\ell] \times [r]$. Then there exist positive integers $\{m_{a,b} : (a, b) \in [2\ell] \times [r]\}$ such that

- ($\mathcal{L}1$) $\sum_{(a,b) \in [2\ell] \times [r]} (m_{a,b} + \tau_{a,b}) = n$ and $m_{a,b} \geq (1 - \sqrt{\varepsilon})m$ and $|m_{a,b} - m_{a,b'}| \leq 1$ for all $a \in [2\ell]$ and $b, b' \in [r]$.
- ($\mathcal{L}2$) Given any $\{n_{a,b} \in \mathbb{N} : (a, b) \in [2\ell] \times [r]\}$ with $\sum_{(a,b) \in [2\ell] \times [r]} (n_{a,b} + \tau_{a,b}) = n$ and $|m_{a,b} - n_{a,b}| \leq \xi n$, there is a partition $\mathcal{X} = \{X_{a,b} : (a, b) \in [2\ell] \times [r]\}$ of $V(G)$ with $|X_{a,b}| = n_{a,b} + \tau_{a,b}$ and $|X_{a,b} \Delta V_{(\phi_\ell^{2r})^{-1}(a,b)}| \leq \sqrt{\varepsilon} m$ for all $(a, b) \in [2\ell] \times [r]$ such that G has a spanning $(\phi_\ell^{2r}(R), 2\ell, r, \mathcal{X}, \varepsilon^{1/3}, \delta/2)$ -cycle structure.

Proof. Note that

$$2r\ell m = n. \quad (7.2)$$

For each $(i, j) \in [\ell] \times [2r]$, choose $A_{i,j} \subseteq V_{i,j}$ satisfying

$$|A_{i,j}| = \tau_{\phi_\ell^{2r}(i,j)} \quad (7.3)$$

and let

$$Y_{i,j} := V_{i,j} \setminus A_{i,j}, \quad \text{so } (1 - \varepsilon)m \leq |Y_{i,j}| \leq m. \quad (7.4)$$

Let $\mathcal{Y} := \{Y_0\} \cup \{Y_{i,j} : (i,j) \in [\ell] \times [2r]\}$, where

$$Y_0 := V(G) \setminus \bigcup_{(i,j) \in [\ell] \times [2r]} Y_{i,j} = \bigcup_{(i,j) \in [\ell] \times [2r]} A_{i,j}.$$

Given a vertex $v \in V(G)$ and $(i,j) \in [\ell] \times [2r]$, we will say that $v \rightarrow Y_{i,j}$ is *valid* if

- $j \in [r]$ and $d_G(v, Y_{i,j'}) \geq (\delta - 2\varepsilon)m$ for all $j' \in [r] \setminus \{j\}$; or
- $j \in [2r] \setminus [r]$ and $d_G(v, Y_{i,j'}) \geq (\delta - 2\varepsilon)m$ for all $j' \in ([2r] \setminus [r]) \setminus \{j\}$.

The first claim furnishes us with many pairs $(v, Y_{i',j'})$ such that $v \in Y_{i,j}$ and $v \rightarrow Y_{i',j'}$ is valid.

Claim 7.2. *Let $i \in [\ell]$, and suppose that $1 \leq j \leq r < t \leq 2r$ or $1 \leq t \leq r < j \leq 2r$. Then every vertex $v \in Y_{i,j}$ is such that $v \rightarrow Y_{i,j}, Y_{i,t}$ is valid, and at least $(1 - \sqrt{\varepsilon})m$ are such that $v \rightarrow Y_{i+1,j}, Y_{i+1,t}$ are also valid. (Here, for example, $Y_{\ell+1,j} := Y_{1,j}$.)*

Proof (of claim). Let t, j be as in the statement. Since, by $(\mathcal{C}3)$, $G[V_{i,j}, V_{i,j'}]$ is (ε, δ) -superregular for all $j' \in [2r] \setminus \{j\}$, we have that every vertex $v \in V_{i,j}$ has at least $\delta|V_{i,j'}|$ neighbours in $V_{i,j'}$. Thus every vertex $v \in Y_{i,j} \subseteq V_{i,j}$ has at least $\delta m - \varepsilon m \geq (\delta - 2\varepsilon)m$ neighbours in $Y_{i,j'}$. In particular, $v \rightarrow V_{i,j}, V_{i,t}$ is valid.

From the definition of regularity, one can see the following. If $G[A, B]$ is an (ε, δ) -regular graph, then there are fewer than $\varepsilon|A|$ vertices with fewer than $(\delta - \varepsilon)|B|$ neighbours in B . Thus, if $S_{i,j}$ is a subset of $N_{i,j} := \{(i', j') \in V(R) : G[V_{i,j}, V_{i',j'}] \text{ is } (\varepsilon, \delta)\text{-regular}\}$, we see that there are at least $(1 - \varepsilon|S_{i,j}|)|V_{i,j}|$ vertices in $V_{i,j}$ with at least $(\delta - \varepsilon)m$ neighbours in $V_{i',j'}$ for all $(i', j') \in S_{i,j}$, and hence at least $(\delta - 2\varepsilon)m$ neighbours in $Y_{i',j'}$.

Recall that, since $Z_\ell^{2r} \subseteq R$ by $(\mathcal{C}2)$, we have that $N_{i,j} \supseteq \{(i, j'), (i+1, j') : j' \in [2r] \setminus \{j\}\}$. Thus the second assertion of the claim follows by taking $S_{i,j} := \{(i+1, j') : j' \in [2r] \setminus \{j, t\}\}$ and using the fact that $(1 - |S_{i,j}|\varepsilon)|V_{i,j}| - |A_{i,j}| \geq (1 - (2r-2)\varepsilon)m - \varepsilon m \geq (1 - \sqrt{\varepsilon})m$. \square

Next we prove the following claim, which will give us a ‘balanced’ partition.

Claim 7.3. *$V(G)$ has a partition $\{Y_0\} \cup \{U_{i,j} : (i,j) \in [\ell] \times [2r]\}$ such that the following hold for all $i \in [\ell]$:*

- (U1) $||U_{i,j}| - |U_{i,j'}|| \leq 1$ for all $(j, j') \in [[2r]]^2$.
- (U2) $|Y_{i,j} \triangle U_{i,j}| \leq r\varepsilon m$ for all $j \in [2r]$.
- (U3) If $j \in [r]$, then $U_{i,j} \setminus Y_{i,j} \subseteq \bigcup_{k \in [2r] \setminus [r]} Y_{i,k}$; and if $k \in [2r] \setminus [r]$, then $U_{i,k} \subseteq Y_{i,k}$.

Proof (of claim). Fix an $i \in [\ell]$, and, to simplify notation, let $A_j := Y_{i,j}$, $a_j := |A_j|$, $B_j := Y_{i,r+j}$, and $b_j := |B_j|$ for all $j \in [r]$. Suppose, without loss of generality, that $a_1 \geq \dots \geq a_r$ and $b_1 \geq \dots \geq b_r$. Let

$$S := \max \left\{ \sum_{j \in [r]} (a_1 - a_j), \sum_{j \in [r]} (b_j - b_r) \right\} \stackrel{(7.4)}{\leq} r\varepsilon m. \quad (7.5)$$

Now let $A_j(0) := A_j$ and $B_j(0) := B_j$, and $a_j(0) := |A_j(0)|$ and $b_j(0) := |B_j(0)|$ for all $j \in [r]$. Do the following for each $0 \leq s < S$. Fix $t^-, t^+ \in [r]$ such that $a_{t^-}(s) \leq a_j(s)$ and $b_{t^+}(s) \geq b_j(s)$ for all $j \in [r]$. Choose $x \in B_{t^+} \cap B_{t^+}(s)$, and let

$$A_j(s+1) := \begin{cases} A_j(s) \cup \{x\} & \text{if } j = t^- \\ A_j(s) & \text{if } j \in [r] \setminus \{t^-\}; \end{cases}$$

$$B_j(s+1) := \begin{cases} B_j(s) \setminus \{x\} & \text{if } j = t^+ \\ B_j(s) & \text{if } j \in [r] \setminus \{t^+\}; \end{cases}$$

Let $a_j(s+1) := |A_j(s+1)|$ and $b_j(s+1) := |B_j(s+1)|$ for all $j \in [r]$. The following properties are clear:

- (i) For all $0 \leq s < S$ and $j \in [r]$, we have $A_j(s) \supseteq A_j$ and $A_j(s) \setminus A_j \subseteq \bigcup_{k \in [r]} B_k$, and $B_j(s) \subseteq B_j$. Furthermore, for all $j \in [r]$, we have $\sum_{j \in [r]} |A_j(s) \setminus A_j| = \sum_{k \in [r]} |B_k \setminus B_k(s)| = s$.
- (ii) Letting $s_1 := \sum_{j \in [r]} (a_1 - a_j)$, we have that $a_1(s_1) = \dots = a_r(s_1) = a_1$; and for each $s > s_1$, we have $|a_j(s) - a_{j'}(s)| \leq 1$.
- (iii) Letting $s_2 := \sum_{j \in [r]} (b_j - b_r)$, we have that $b_1(s_2) = \dots = b_r(s_2) = b_r$; and for each $s > s_2$, we have $|b_j(s) - b_{j'}(s)| \leq 1$.

Now let $U_{i,j} := A_j(S)$ if $j \in [r]$ and $U_{i,j} := B_{j-r}(S)$ if $j \in [2r] \setminus [r]$. For $(\mathcal{U}1)$, the fact that $S = \max\{s_1, s_2\}$ together with (ii) and (iii) implies that $|a_j(S) - a_{j'}(S)| \leq 1$ and $|b_j(S) - b_{j'}(S)| \leq 1$ for all $j, j' \in [r]$. So $(\mathcal{U}1)$ holds. For $(\mathcal{U}2)$, we have by (i) that

$$\begin{aligned} |U_{i,j} \triangle Y_{i,j}| &= |U_{i,j} \setminus Y_{i,j}| = |A_j(S) \setminus A_j| \leq S \stackrel{(7.5)}{\leq} r\epsilon m \quad \text{if } j \in [r], \quad \text{and} \\ |U_{i,j} \triangle Y_{i,j}| &= |Y_{i,j} \setminus U_{i,j}| = |B_{j-r} \setminus B_{j-r}(S)| \leq S \stackrel{(7.5)}{\leq} r\epsilon m \quad \text{if } j \in [2r] \setminus [r]. \end{aligned}$$

Finally, $(\mathcal{U}3)$ follows immediately from (i). \square

The next claim shows that we can modify $\{U_{i,j}\}$ further to obtain a new partition with clusters of given sizes (each of which does not differ much from $|U_{i,j}|$).

Claim 7.4. *Let $\{Y_0\} \cup \{U_{i,j} : (i,j) \in [\ell] \times [2r]\}$ be any partition of $V(G)$ satisfying $(\mathcal{U}1)$ – $(\mathcal{U}3)$. Let $\{n'_{i,j} : (i,j) \in [\ell] \times [2r]\}$ be such that $\sum_{(i,j) \in [\ell] \times [2r]} n'_{i,j} = \sum_{(i,j) \in [\ell] \times [2r]} |U_{i,j}|$ and $||U_{i,j}| - n'_{i,j}| \leq \xi n$ for all $(i,j) \in [\ell] \times [2r]$. Then $V(G)$ has a partition $\{Y_0\} \cup \{W_{i,j} : (i,j) \in [\ell] \times [2r]\}$ such that the following hold for all $(i,j) \in [\ell] \times [2r]$:*

- $(\mathcal{W}1)$ $|W_{i,j}| = n'_{i,j}$.
- $(\mathcal{W}2)$ $|W_{i,j} \triangle U_{i,j}| \leq \epsilon m$.
- $(\mathcal{W}3)$ For every $v \in W_{i,j}$, we have that $v \rightarrow Y_{i,j}$ is valid.

Proof (of claim). Let

$$K := 2r\ell\xi n \stackrel{(7.2)}{=} 4r^2\ell^2\xi m \leq \frac{\epsilon m}{2}. \quad (7.6)$$

Suppose that, for some $0 \leq k < K/2$, we have found for each $(i,j) \in [\ell] \times [2r]$ subsets $U_{i,j}^k \subseteq V(G)$ such that the following hold:

- $\mathcal{A}_1(k)$ $\{Y_0\} \cup \{U_{i,j}^k : (i,j) \in [\ell] \times [2r]\}$ is a partition of $V(G)$.
- $\mathcal{A}_2(k)$ For all $v \in U_{i,j}^k$, we have that $v \rightarrow Y_{i,j}$ is valid.
- $\mathcal{A}_3(k)$ For all $(i,j) \in [\ell] \times [2r]$, we have $|U_{i,j}^k \triangle U_{i,j}| \leq 2k$.
- $\mathcal{A}_4(k)$ $\sum_{(i,j) \in [\ell] \times [2r]} ||U_{i,j}^k| - n'_{i,j}| \leq 2(r\ell\xi n - k)$.

We claim that we can set $U_{i,j}^0 := U_{i,j}$ for all $(i,j) \in [\ell] \times [2r]$. Indeed, $\mathcal{A}_1(0)$ holds by Claim 7.3. For $\mathcal{A}_2(0)$, let $(i,j) \in [\ell] \times [2r]$, and let $v \in U_{i,j}$. If $v \in Y_{i,j}$, then $v \rightarrow Y_{i,j}$ is valid by Claim 7.2. Otherwise, $v \in U_{i,j} \setminus Y_{i,j}$. Note that by $(\mathcal{U}3)$, this implies $j \in [r]$ and, further, $v \in \bigcup_{k \in [2r] \setminus [r]} Y_{i,k}$. So $v \rightarrow Y_{i,j}$ is valid by Claim 7.2. Property $\mathcal{A}_3(0)$ vacuously holds, and $\mathcal{A}_4(0)$ holds since $||U_{i,j}| - n'_{i,j}| \leq \xi n$ for all $(i,j) \in [\ell] \times [2r]$.

If $|U_{i,j}^k| = n'_{i,j}$ for all $(i,j) \in [\ell] \times [2r]$, then we stop. Otherwise, we will obtain sets $U_{i,j}^{k+1}$ from $U_{i,j}^k$. There must exist $(i^-, j^-), (i^+, j^+) \in [\ell] \times [2r]$ for which $|U_{i^-, j^-}^k| \leq n'_{i^-, j^-} - 1$ and $|U_{i^+, j^+}^k| \geq n'_{i^+, j^+} + 1$.

We will say that $(i_1, j_1) \rightarrow (i_2, j_2) \rightarrow \dots \rightarrow (i_s, j_s)$ is a *good chain (of length s)* if for all $p \in [s-1]$, there exist at least $(1 - \sqrt{\epsilon})m$ vertices $v \in Y_{i_p, j_p}$ such that $v \rightarrow Y_{i_{p+1}, j_{p+1}}$ is valid. Claim 7.2 implies that the following are good chains of length 3 (where here and for the remainder of the proof of Claim 7.4,

addition is modulo ℓ):

$$\begin{aligned}(i^+, j^+) &\rightarrow (i^+, j^- + r) \rightarrow (i^+ + 1, j^-) && \text{if } j^+, j^- \in [r] \\(i^+, j^+) &\rightarrow (i^+, j^- - r) \rightarrow (i^+ + 1, j^-) && \text{if } j^+, j^- \in [2r] \setminus [r] \\(i^+, j^+) &\rightarrow (i^+, j^-) \rightarrow (i^+ + 1, j^-) && \text{otherwise,}\end{aligned}$$

and further, in all cases and for all $t \geq 0$, the chain $(i^+ + t, j^-) \rightarrow (i^+ + t + 1, j^-)$ of length 2 is good. Together, this implies that in all cases, there is a good chain

$$(i^+, j^+) =: (i_1, j_1) \rightarrow \dots \rightarrow (i_S, j_S) := (i^-, j^-)$$

of some length S , where we choose the shortest such chain. As a crude estimate, we have, say, $S \leq 2\ell$, and $(i_s, j_s) \neq (i_{s'}, j_{s'})$ for any distinct $s, s' \in [S]$ (or we could find a shorter chain).

We will exchange vertices between successive clusters according to this chain. For each $s \in [S]$, there are by definition at least $(1 - \sqrt{\varepsilon})m$ vertices $v \in Y_{i_s, j_s}$ such that $v \rightarrow Y_{i_{s+1}, j_{s+1}}$ is valid. The number of these vertices that additionally lie in U_{i_s, j_s}^k is by $(\mathcal{U}2)$, $\mathcal{A}_3(0)$ and (7.6) at least $(1 - \sqrt{\varepsilon})m - 2k - r\varepsilon m > m/2$. So we can find $x_s \in U_{i_s, j_s}^k$ such that $x_s \rightarrow Y_{i_{s+1}, j_{s+1}}$ is valid. For each $(i, j) \in [\ell] \times [2r]$, set

$$U_{i,j}^{k+1} = \begin{cases} U_{i,j}^k \setminus \{x_1\} & \text{if } (i, j) = (i_1, j_1) \\ U_{i,j}^k \cup \{x_{s-1}\} \setminus \{x_s\} & \text{if } (i, j) = (i_s, j_s) \text{ for some } 2 \leq s < S \\ U_{i,j}^k \cup \{x_{S-1}\} & \text{if } (i, j) = (i_S, j_S) \\ U_{i,j}^k & \text{otherwise.} \end{cases}$$

Property $\mathcal{A}_1(k+1)$ holds by $\mathcal{A}_1(k)$, the definition of $U_{i,j}^{k+1}$, and the fact that each pair in the chain is distinct. Property $\mathcal{A}_2(k)$ and the choice of x_s imply that $\mathcal{A}_2(k+1)$ holds. We have

$$|U_{i,j}^{k+1} \triangle Y_{i,j}| \leq |U_{i,j}^{k+1} \triangle U_{i,j}^k| + |U_{i,j}^k \triangle Y_{i,j}| \stackrel{\mathcal{A}_3(k)}{\leq} 2(k+1),$$

proving $\mathcal{A}_3(k+1)$ (note here we are again using the fact that each pair in our chain is distinct). Finally, observe that $||U_{i^\pm, j^\pm}^{k+1}| - n'_{i^\pm, j^\pm}| = ||U_{i^\pm, j^\pm}^k| - n'_{i^\pm, j^\pm}| - 1$ and $|U_{i,j}^{k+1}| = |U_{i,j}^k|$ for all other pairs (i, j) . Therefore,

$$\sum_{(i,j) \in [\ell] \times [2r]} ||U_{i,j}^{k+1}| - n'_{i,j}| = \sum_{(i,j) \in [\ell] \times [2r]} ||U_{i,j}^k| - n'_{i,j}| - 2 \stackrel{\mathcal{A}_4(k)}{\leq} 2(r\ell\xi n - (k+1)),$$

proving $\mathcal{A}_4(k+1)$. So, for each $0 \leq k \leq K/2$, either the procedure has terminated or we are able to proceed to step $k+1$. Therefore, there is some $p \leq K/2$ such that $\sum_{(i,j) \in [\ell] \times [2r]} ||U_{i,j}^p| - n'_{i,j}| = 0$. Note that, by $\mathcal{A}_3(p)$, we have

$$|U_{i,j}^p \triangle U_{i,j}| \leq 2p \leq K \stackrel{(7.6)}{\leq} \varepsilon m$$

for all $(i, j) \in [\ell] \times [2r]$. Thus setting $W_{i,j} := U_{i,j}^p$ yields the required partition. \square

Apply Claim 7.3 to obtain $\{U_{i,j} : (i, j) \in [\ell] \times [2r]\}$ satisfying $(\mathcal{U}1)$ – $(\mathcal{U}3)$.

Let $\phi := \phi_\ell^{2r}$ as in Definition 2. Let

$$U'_{a,b} := U_{\phi^{-1}(a,b)} \quad \text{and} \quad m_{a,b} := |U'_{a,b}| \quad \text{for all } (a, b) \in [2\ell] \times [r].$$

We claim that $\{m_{a,b}\}$ satisfies $(\mathcal{L}1)$. Indeed, $(\mathcal{U}1)$ implies that $|m_{a,b} - m_{a,b'}| \leq 1$ for all $a \in [2\ell]$ and $b, b' \in [r]$, and further, writing $\phi^{-1}(a, b) = (i, j)$,

$$m_{a,b} = |U_{i,j}| \stackrel{(\mathcal{U}2)}{\geq} |Y_{i,j}| - r\epsilon m \stackrel{(7.3)}{\geq} |V_{i,j}| - (r+1)\epsilon m = (1 - (r+1)\epsilon)m \geq (1 - \sqrt{\epsilon})m.$$

Finally,

$$\sum_{(a,b) \in [2\ell] \times [r]} m_{a,b} = \sum_{(i,j) \in [\ell] \times [2r]} |U_{i,j}| = n - |Y_0| = n - \sum_{(a,b) \in [2\ell] \times [r]} \tau_{a,b},$$

so $(\mathcal{L}1)$ holds.

Now let $\{n_{a,b} \in \mathbb{N} : (a, b) \in [2\ell] \times [r]\}$ satisfy $\sum_{(a,b) \in [2\ell] \times [r]} (n_{a,b} + \tau_{a,b}) = n$ and $|m_{a,b} - n_{a,b}| \leq \xi n$. Let

$$n'_{i,j} := n_{\phi(i,j)} \quad \text{for all } (i, j) \in [\ell] \times [2r]. \quad (7.7)$$

Apply Claim 7.4 with input partition $\{Y_0\} \cup \{U_{i,j} : (i, j) \in [\ell] \times [2r]\}$ and input sizes $\{n'_{i,j} : (i, j) \in [\ell] \times [2r]\}$ to obtain a partition $\{Y_0\} \cup \{W_{i,j} : (i, j) \in [\ell] \times [2r]\}$ satisfying $(\mathcal{W}1)$ – $(\mathcal{W}3)$. Let

$$X_{a,b} := W_{\phi^{-1}(a,b)} \cup A_{\phi^{-1}(a,b)} \quad \text{for all } (a, b) \in [2\ell] \times [r]. \quad (7.8)$$

We claim that $\mathcal{X} := \{X_{a,b} : (a, b) \in [2\ell] \times [r]\}$ is the required partition for $(\mathcal{L}2)$. For all $(a, b) \in [2\ell] \times [r]$, we have

$$|X_{a,b}| \stackrel{(7.8)}{=} |W_{\phi^{-1}(a,b)}| + |A_{\phi^{-1}(a,b)}| \stackrel{(7.3), (\mathcal{W}1)}{=} n'_{\phi^{-1}(a,b)} + \tau_{a,b} \stackrel{(7.7)}{=} n_{a,b} + \tau_{a,b},$$

as required. Also, writing $(i, j) := \phi^{-1}(a, b) \in [\ell] \times [2r]$ and recalling that $A_{i,j} \subseteq V_{i,j}$, we have

$$\begin{aligned} |X_{a,b} \Delta V_{\phi^{-1}(a,b)}| &\stackrel{(7.8)}{=} |W_{i,j} \Delta V_{i,j}| \leq |W_{i,j} \Delta U_{i,j}| + |U_{i,j} \Delta V_{i,j}| \\ &\stackrel{(\mathcal{U}2), (\mathcal{W}2)}{\leq} 2r\epsilon m \leq 3r\epsilon |X_{a,b}| \leq \sqrt{\epsilon}m. \end{aligned} \quad (7.9)$$

Lastly, we need to check that \mathcal{X} induces a $(\phi(R), 2\ell, r, \mathcal{X}, \epsilon^{1/3}, \delta/2)$ -cycle structure. That is, we need to check that $(\mathcal{C}1)$ – $(\mathcal{C}3)$ hold. Property $(\mathcal{W}1)$ implies that $\mathcal{X} = \{X_{a,b} : (a, b) \in [2\ell] \times [r]\} = \{W_{i,j} \cup A_{i,j} : (i, j) \in [\ell] \times [2r]\}$ is a partition of $V(G)$. So $(\mathcal{C}1)$ holds. Now, by (3.1) and Definition 2, we see that $\phi(R)$ has vertex set $[2\ell] \times [r]$, and, since $Z_{\ell}^{2r} \subseteq R$, we have $Z_{2\ell}^r \subseteq \phi(R)$ (with the correct labelling). Let $(a, b), (a', b') \in E(\phi(R))$, and write $(i, j) := \phi^{-1}(a, b)$ and $(i', j') := \phi^{-1}(a', b')$. Then $(i, j)(i', j') \in E(R)$, so $G[V_{i,j}, V_{i',j'}]$ is (ϵ, δ) -regular by $(\mathcal{C}2)$ for \mathcal{V} . Then (7.9) implies that we can apply Proposition 4.3 with $\alpha := 3r\epsilon$ and $\epsilon' := \epsilon^{1/3} \geq \epsilon + 6\sqrt{\alpha}$ to see that $G[X_{a,b}, X_{a',b'}]$ is $(\epsilon^{1/3}, \delta/2)$ -regular. So $(\mathcal{C}2)$ holds.

For $(\mathcal{C}3)$, fix $a \in [2\ell]$, and let $b, b' \in [r]$ be distinct. Let $(i, j) := \phi^{-1}(a, b)$. Definition 2 implies that there exists j' such that $(j, j') \in [[2r]]^2$ and $\phi^{-1}(a, b') = (i, j')$. Let $x \in X_{a,b} \setminus V_{\phi^{-1}(a,b)} = W_{i,j} \setminus Y_{i,j}$. Then $(\mathcal{W}3)$ implies that $x \rightarrow Y_{i,j}$ is valid. Since $Y_{i,j} \subseteq V_{i,j}$, this means $d_G(x, V_{i,j^*}) \geq (\delta - 2\epsilon)m$ for all j^* such that $(j, j^*) \in [[2r]]^2$. So $d_G(x, V_{\phi^{-1}(a,b')}) \geq (\delta - 2\epsilon)m$, and hence (7.9) implies $d_G(x, X_{a,b'}) \geq (\delta - 2\epsilon)m - 2r\epsilon m \geq \delta|X_{a,b'}|/2$. Similarly, every $y \in X_{a,b'} \setminus V_{\phi^{-1}(a,b')}$ satisfies $d_G(y, X_{a,b}) \geq \delta|X_{a,b}|/2$. Moreover, $(\mathcal{C}3)$ for \mathcal{V} and (7.9) implies that $d_G(x, X_{a,b'}) \geq \delta|X_{a,b'}|/2$ for every $x \in V_{\phi^{-1}(a,b)}$ and $d_G(y, X_{a,b}) \geq \delta|X_{a,b}|/2$ for every $y \in V_{\phi^{-1}(a,b')}$. So Proposition 4.3 applied with $\alpha := 3r\epsilon$ and $\epsilon' := \epsilon^{1/3}$ implies that $G[X_{a,b}, X_{a,b'}]$ is $(\epsilon^{1/3}, \delta/2)$ -superregular. So $(\mathcal{C}3)$ holds. This completes the proof of $(\mathcal{L}2)$ and hence of the lemma. \square

8. The proof of Theorem 1.2

First, note that it suffices to prove the theorem under the additional assumption that $\eta \ll d, 1/\Delta$. Let $n_0, \beta, \rho, \varepsilon, c, \delta, \rho', L' > 0$ satisfy

$$0 < 1/n_0 \ll \beta \ll 1/L' \ll \rho \ll \varepsilon \ll c \ll \delta \ll \rho' \ll \eta \ll d, 1/\Delta. \quad (8.1)$$

Let G be a (ρ, d) -dense graph on $n \geq n_0$ vertices with $\delta(G) \geq (1/2 + \eta)n$. Let H be a graph on n vertices with $\Delta(H) \leq \Delta$ and bandwidth at most βn . Write $r := \chi(H)$; so as $\eta \ll 1/\Delta$, certainly $\eta \ll 1/r$.

Apply the regularity lemma (Lemma 4.1) with parameters $\varepsilon, (4r + 1)L'$ to obtain $L^* \in \mathbb{N}$. We may assume that $\beta \ll 1/L^*$.

Claim 8.1. *There exists $L' \leq \ell \leq L^*$, a partition $\mathcal{V} = \{V_0\} \cup \{V_{i,j} : (i, j) \in [\ell] \times [4r]\}$ of $V(G)$ with $|V_{i,j}| = m$ for all $(i, j) \in [\ell] \times [4r]$, a graph R on vertex set $[\ell] \times [4r]$ and a spanning subgraph G' of G , such that*

- (i) R is (ρ', d) -dense.
- (ii) $\delta(R) \geq (1/2 + \eta/3)|R|$.
- (iii) G' has an $(R, \ell, 4r, \mathcal{V}, 7\varepsilon^{1/4}, \delta/2)$ -cycle structure \mathcal{C} and $|V_0| \leq 2\varepsilon^{1/2}n$.
- (iv) $R[\{(1, 1), \dots, (1, 4r)\}] \cong K_{4r}$ and $\{(1, 1), \dots, (1, 4r)\}$ lies in a copy of K_{324r/η^2} in R .

Proof (of claim). Apply Lemma 4.1 to G with parameters $\varepsilon, \delta, (4r + 1)L'$ to obtain clusters V_1, \dots, V_L of size m' , an exceptional set V'_0 , a pure graph G' , and a reduced graph R' . So

$$Lm' \leq n \leq Lm' + \varepsilon n, \quad (8.2)$$

and $|R'| = L$, where

$$(4r + 1)L' \leq L \leq L^* \quad (8.3)$$

and $|V'_0| \leq \varepsilon n$,

$$\delta(G') \geq (1/2 + \eta)n - (\delta + \varepsilon)n \geq (1/2 + \eta/2)n \quad (8.4)$$

and $G'[V_i, V_j]$ is (ε, δ) -regular whenever $ij \in E(R')$. Lemma 4.2 implies that R' is $(3\delta, d)$ -dense and $\delta(R') \geq (1/2 + \eta/2)L$.

Let $r^* := 324r/\eta^2$. Apply Theorem 5.1 with $R', L, r^* - 1, 4r, 3\delta, d, \eta/2$ playing the roles of $G, n, r, s, \rho, d, \eta$ to obtain an $(r^* - 1)$ -cycle $C \cong C_{4r\ell}^{r^*-1} \subseteq R'$ of order $4r\ell$ where

$$(1 - \varepsilon)L \leq L - 4r \leq 4r\ell \leq L. \quad (8.5)$$

Relabel those clusters of R' corresponding to vertices of C so that they are now $\{V'_{i,j} : (i, j) \in [\ell] \times [4r]\}$, and

$$(1, 1)(1, 2) \dots (1, 4r)(2, 1) \dots (2, 4r) \dots (\ell, 1) \dots (\ell, 4r) = C_{4r\ell}^{r^*-1} \supseteq Z_\ell^{4r}. \quad (8.6)$$

Let $R := R'[V(C)]$. So $V(R) = [\ell] \times [4r]$. Observe that $\{(1, 1), \dots, (1, 4r)\}$ lies in a copy of K_{r^*} in R . For all $i \in [\ell]$, let

$$T(i) := R\left[\bigcup_{j \in [4r]} (i, j)\right] \stackrel{(8.6)}{\cong} K_{4r}.$$

Apply Lemma 4.4 with $G'[\bigcup_{j \in [4r]} V'_{i,j}], T(i), 4r - 1, 4r, V'_{i,1}, \dots, V'_{i,4r}, m', \varepsilon, \delta$ playing the roles of $G, R, \Delta, L, V_1, \dots, V_L, m, \varepsilon, d$ to obtain for each $j \in [4r]$ a subset $V_{i,j} \subseteq V'_{i,j}$ of size

$$|V_{i,j}| = m := (1 - \sqrt{\varepsilon})m' \quad (8.7)$$

such that for every distinct $j, j' \in [4r]$, the graph $G'[V_{i,j}, V_{i,j'}]$ is $(4\sqrt{\varepsilon}, \delta/2)$ -superregular. Let $V_0 := V(G) \setminus \bigcup_{(i,j) \in [\ell] \times [4r]} V_{i,j}$ and

$$\mathcal{V} := \{V_0\} \cup \{V_{i,j} : (i,j) \in [\ell] \times [4r]\}. \quad (8.8)$$

We have

$$\begin{aligned} n &\geq 4r\ell m \stackrel{(8.7)}{=} 4r\ell(1-\sqrt{\varepsilon})m' \stackrel{(8.5)}{\geq} (1-\sqrt{\varepsilon})(1-\varepsilon)Lm' \stackrel{(8.2)}{\geq} (1-\sqrt{\varepsilon})(1-\varepsilon)^2n \\ &\geq (1-2\varepsilon^{1/2})n. \end{aligned} \quad (8.9)$$

Since we will often compare m and βn in calculations, let us note here that

$$\beta n \stackrel{(8.2)}{\leq} \frac{\beta Lm'}{1-\varepsilon} \stackrel{(8.7)}{=} \frac{\beta Lm}{(1-\sqrt{\varepsilon})(1-\varepsilon)} \stackrel{(8.1),(8.3)}{\leq} 2\beta L^* \cdot m \leq \frac{\varepsilon^2 m}{L^*}. \quad (8.10)$$

We will now show that ℓ , R , and \mathcal{V} satisfy Claim 8.1(i)–(iv). We have that

$$4rL' \stackrel{(8.1)}{\leq} (1-\varepsilon)(4r+1)L' \stackrel{(8.3)}{\leq} (1-\varepsilon)L \stackrel{(8.5)}{\leq} 4r\ell \leq L \stackrel{(8.3)}{\leq} L^*.$$

So $L' \leq \ell \leq L^*$, as required. Note that (i) follows from Lemma 3.1(i) since $\rho' \gg \delta$. Further, $\delta(R) \geq \delta(R') - 4r \geq (1/2 + \eta/3)L$, so (ii) holds.

For (iii), we need to show that \mathcal{V} (see (8.8)) induces the required cycle structure \mathcal{C} . That is, we need to check that $(\mathcal{C}1)$ – $(\mathcal{C}3)$ hold with the desired parameters. The sets $V_{i,j}$ are pairwise-disjoint since the same is true for $V'_{i,j}$, so by the definition of V_0 , we have that \mathcal{V} is a partition of $V(G')$. Moreover,

$$|V_0| = n - 4r\ell m \stackrel{(8.9)}{\leq} 2\varepsilon^{1/2}n < 7\varepsilon^{1/4}n,$$

so $(\mathcal{C}1)$ holds. Certainly $V(R) = [\ell] \times [4r]$ and, by (8.6), $R \supseteq Z_\ell^{4r}$. Let $(i,j)(i',j') \in E(R)$. Then $(i,j)(i',j')$ has a corresponding edge in $R' \supseteq R$, so $G'[V'_{i,j}, V'_{i',j'}]$ is (ε, δ) -regular. Note that $\varepsilon + 6\sqrt{\varepsilon^{1/2}} \leq 7\varepsilon^{1/4}$ and $\delta - 4\varepsilon^{1/2} \geq \delta/2$. Thus Lemma 4.3 applied with $V'_{i,j}, V_{i,j}, V'_{i',j'}, V_{i',j'}, \varepsilon^{1/2}$ playing the roles of A, A', B, B', α implies that $G'[V_{i,j}, V_{i',j'}]$ is $(7\varepsilon^{1/4}, \delta/2)$ -regular, so $(\mathcal{C}2)$ holds. We have already seen, for every $i \in [\ell]$ and distinct $j, j' \in [4r]$, that $G'[V_{i,j}, V_{i,j'}]$ is $(4\sqrt{\varepsilon}, \delta/2)$ -superregular. Thus it is $(7\varepsilon^{1/4}, \delta/2)$ -superregular. So $(\mathcal{C}3)$ holds. Thus (iii) holds. We saw when we defined R that (iv) holds. This completes the proof of the claim. \square

Recall the definition of the bijection $\phi_\ell^{4r} : [\ell] \times [4r] \rightarrow [2\ell] \times [2r]$ given by

$$\phi_\ell^{4r}(i, j) = \left((2i-1) + \left\lfloor \frac{j}{2r} \right\rfloor, j - \left(\left\lfloor \frac{j}{2r} \right\rfloor - 1 \right) 2r \right), \quad \text{for all } (i, j) \in [\ell] \times [4r].$$

Recall also that $\phi_\ell^{4r}(R)$ is the graph with vertex set $\phi_\ell^{4r}(V(R)) = [2\ell] \times [2r]$ and edge set $E(\phi_\ell^{4r}(R)) = \{\phi_\ell^{4r}(x)\phi_\ell^{4r}(y) : xy \in E(R)\}$. For ease of notation, we will write

$$\begin{aligned} \phi &:= \phi_\ell^{4r}, \quad \text{so } \phi(1, b) = (1, b) \text{ for all } b \in [2r], \quad \text{and} \\ R^* &:= \phi(R), \quad \text{so } R^* \cong R, \quad V(R^*) = [2\ell] \times [2r], \end{aligned} \quad (8.11)$$

and $V(G')$ has partition $\mathcal{V} = \{V_0\} \cup \{V_{\phi^{-1}(a,b)} : (a,b) \in [2\ell] \times [2r]\}$.

Claim 8.2. *There exists a partition $\mathcal{X} = \{V_0\} \cup \{X_{a,b} : (a,b) \in [2\ell] \times [2r]\}$ of $V(G')$ and a surjective mapping $\psi : V(H) \rightarrow ([2\ell] \times [2r]) \cup V_0$ such that the following hold:*

- (i) $|\psi^{-1}(a, b)| = |X_{a,b}| \geq (1 - \varepsilon^{1/19})m$ for all $(a, b) \in [2\ell] \times [2r]$.
- (ii) G' has an $(R^*, 2\ell, 2r, \mathcal{X}, \varepsilon^{1/27}, \delta/4)$ -cycle structure \mathcal{C}' .
- (iii) $I := \psi^{-1}(V_0)$ is an independent set in H of size $|V_0|$; and for all $w \in W := \bigcup_{x \in I} N_H(x)$, there is a unique $u \in I$ such that $uw \in E(H)$ and $d_{G'}(\psi(u), X_{\psi(w)}) \geq cm/2$;
- (iv) $\psi|_{V(H) \setminus I} : V(H \setminus I) \rightarrow V(R^*)$ is a graph homomorphism.
- (v) There exists $X' \subseteq V(H) \setminus I$ with $W \subseteq X'$ and $|\psi^{-1}(a, b) \cap X'| \leq \varepsilon^{1/10}m$ for all $(a, b) \in [2\ell] \times [2r]$ such that, whenever $uv \in E(H)$ and $u, v \notin X' \cup I$, writing $\psi(u) =: (a, b)$ and $\psi(v) =: (a', b')$, we have $a = a'$ and $b \neq b'$. Moreover, writing

$$N := (\bigcup_{x \in X'} N_H(x)) \setminus (X' \cup I),$$

we have $|N| \leq \varepsilon m$.

Proof (of claim). For all $v \in V(G')$, write

$$N_{R^*}^c(v) := \{(a, b) \in [2\ell] \times [2r] : d_{G'}(v, V_{\phi^{-1}(a,b)}) \geq cm\} \quad (8.12)$$

and $d_{R^*}^c(v) := |N_{R^*}^c(v)|$. Then

$$(1/2 + \eta)n - (\delta + \varepsilon)n \stackrel{(8.4)}{\leq} d_{G'}(v) \leq d_{R^*}^c(v)m + (4r\ell - d_{R^*}^c(v))cm + |V_0|.$$

Claim 8.1(iii) implies that

$$4r\ell m \leq n \leq 4r\ell m + |V_0| \leq 4r\ell m + 2\varepsilon^{1/2}n.$$

Thus

$$d_{R^*}^c(v) \geq \frac{(1/2 + \eta - \delta - \varepsilon)n - 4r\ell cm - |V_0|}{(1 - c)m} \stackrel{(8.1)}{\geq} \frac{1/2 + \eta/2}{1 - c} \cdot 4r\ell \geq \frac{|R^*|}{2}. \quad (8.13)$$

We would like to apply Lemma 6.2 (special lemma for H) to obtain an integer s , with $G', R^*, 4r\ell, 2r, \eta/3, 2\varepsilon^{1/2}, \rho', d, n, m, N_{R^*}^c(v), (1, i)$ playing the roles of $G, R, L, r, \eta, \varepsilon, \rho, d, n, m, N_v, b_i$. For this, we need to check that $(\mathcal{G}1)$ – $(\mathcal{G}4)$ hold. For $(\mathcal{G}1)$, we know that G' has vertex partition $\{V_0\} \cup \{V_{i,j} : (i, j) \in [\ell] \times [4r]\} = \{V_0\} \cup \{V_{\phi^{-1}(a,b)} : (a, b) \in [2\ell] \times [2r]\}$, and $|V_0| \leq \varepsilon^{1/2}n$ and $|V_p| = m$ for all $p \in V(R^*)$. That $(\mathcal{G}2)$ holds follows from (8.13). Property $(\mathcal{G}3)$ follows from Claim 8.1(i) and (ii) and the fact that $R^* \cong R$. Finally, $(\mathcal{G}4)$ follows from (iv), noting that $324r/\eta^2 = 18 \cdot (2r) \cdot 1/(\eta/3)^2$, and the fact that $\phi(1, b) = (1, b)$ for all $b \in [2r]$ from (8.11). Therefore, we can apply Lemma 6.2 with the above parameters to obtain an integer

$$s \leq (2\varepsilon^{1/2})^{1/4}n \leq \varepsilon^{1/9}n. \quad (8.14)$$

Let $\chi : V(H) \rightarrow [r]$ be a proper colouring of H , let x_1, \dots, x_n be an ordering of $V(H)$ with bandwidth at most βn , and let

$$X := \{x_1, \dots, x_s\}, \quad Y := \{x_{s+1}, \dots, x_{s+\beta n}\} \subseteq Z := \{x_{s+1}, \dots, x_n\}, \quad (8.15)$$

$$H' := H[X \cup Y] \quad \text{and} \quad H'' := H[Z]. \quad (8.16)$$

Apply Lemma 6.2 (special lemma for H) with the above parameters and with $s, \beta, \Delta, H', X, Y, \chi$ playing the roles of $s, \beta, \Delta, H, X, Y, \chi$ to obtain a mapping

$$f : X \cup Y \rightarrow ([2\ell] \times [2r]) \cup V_0$$

with the following properties:

- ($\mathcal{D}1$) Setting $I := f^{-1}(V_0)$, we have that I is a subset of X that is 2-independent in H' , and each vertex in V_0 is mapped onto from a unique vertex in H (so $|I| = |V_0|$).
- ($\mathcal{D}2$) For all $v \in V_0$, setting $W_v := N_H(f^{-1}(v))$, we have $W_v \subseteq X$ and $f(W_v) \subseteq N_{R^*}^c(v)$.
- ($\mathcal{D}3$) $|f^{-1}(a, b)| \leq \varepsilon^{1/9}m$ for every $(a, b) \in [2\ell] \times [2r]$.
- ($\mathcal{D}4$) For every edge $uv \in E(H)$ such that $f(u), f(v) \notin V_0$, we have $f(u)f(v) \in E(R^*)$.
- ($\mathcal{D}5$) For all $y \in Y$, we have $f(y) = (1, \chi(y))$.

Let

$$\tau_{a,b} := |(f|_{X \setminus I})^{-1}(a, b)| = |(f|_X)^{-1}(a, b)| \quad \text{for all } (a, b) \in [2\ell] \times [2r]. \quad (8.17)$$

Then $0 \leq \tau_{a,b} \leq \varepsilon^{1/9}m$ for all $(a, b) \in [2\ell] \times [2r]$ by ($\mathcal{D}3$).

Apply Lemma 7.1 (the lemma for G) with $n - |V_0|, \ell, m, 2r, 11\beta, \varepsilon^{1/9}, \delta/2, G' \setminus V_0, R, \mathcal{V} \setminus \{V_0\}$ playing the roles of $n, \ell, m, r, \xi, \varepsilon, \delta, G, R, \mathcal{V}$ to obtain positive integers $\{m_{a,b} : (a, b) \in [2\ell] \times [2r]\}$ such that

- ($\mathcal{L}1$) $\sum_{(a,b) \in [2\ell] \times [2r]} (m_{a,b} + \tau_{a,b}) = n - |V_0|$ and $m_{a,b} \geq (1 - \varepsilon^{1/18})m$ and $|m_{a,b} - m_{a,b'}| \leq 1$ for all $a \in [2\ell]$ and $b, b' \in [2r]$.
- ($\mathcal{L}2$) Given any $\{n_{a,b} : (a, b) \in [2\ell] \times [2r]\}$ with $\sum_{(a,b) \in [2\ell] \times [2r]} (n_{a,b} + \tau_{a,b}) = n - |V_0|$ and $|m_{a,b} - n_{a,b}| \leq 11\beta(n - |V_0|)$, there is a partition $\mathcal{X} = \{V_0\} \cup \{X_{a,b} : (a, b) \in [2\ell] \times [2r]\}$ of $V(G')$ with $|X_{a,b}| = n_{a,b} + \tau_{a,b}$ and $|X_{a,b} \Delta V_{\phi^{-1}(a,b)}| \leq \varepsilon^{1/18}m$ for all $(a, b) \in [2\ell] \times [2r]$ such that G' has an $(R^*, 2\ell, 2r, \mathcal{X}, \varepsilon^{1/27}, \delta/4)$ -cycle structure.

Note that Lemma 7.1 yields a partition of $G' \setminus V_0$ into clusters, and the partition of $V(G')$ specified in ($\mathcal{L}2$) is simply this partition together with V_0 .

The next step is to apply Lemma 6.1 (basic lemma for H) to $H'' = H[Z]$ (which overlaps with H' in Y). Note that the number of vertices in H'' is $n - s \geq (1 - \varepsilon^{1/9})n$. Further,

$$\sum_{(a,b) \in [2\ell] \times [2r]} m_{a,b} \stackrel{(\mathcal{L}1)}{=} n - |V_0| - \sum_{(a,b) \in [2\ell] \times [2r]} \tau_{a,b} \stackrel{(8.17)}{=} n - |V_0| - |X \setminus I| \stackrel{(\mathcal{D}1)}{=} n - |X| \stackrel{(8.15)}{=} |Z| \quad (8.18)$$

and $m_{a,b} \geq (1 - \varepsilon^{1/18})m \geq 10\beta(n - s)$ by (8.1). Thus we can apply Lemma 6.1 with $n - s, r, 2\ell, \Delta, \beta, H'', (x_{s+1}, \dots, x_n), \chi, \{m_{a,b} : (a, b) \in [2\ell] \times [2r]\}$ playing the roles of $n, r, \ell, \Delta, \beta, H, (x_1, \dots, x_n), \chi, \{m_{a,b} : (a, b) \in [2\ell] \times [2r]\}$ to obtain a mapping

$$k : Z \rightarrow [2\ell] \times [2r] \quad (8.19)$$

and $B \subseteq Z$ with the following properties:

- ($\mathcal{B}1$) $B \subseteq Z \setminus Y$ and $|B| \leq 2\ell\beta n$.
- ($\mathcal{B}2$) $||k^{-1}(a, b)| - m_{a,b}| \leq 10\beta n$ for every $(a, b) \in [2\ell] \times [2r]$.
- ($\mathcal{B}3$) For every edge $uv \in E(H'')$, writing $k(u) =: (a, b)$ and $k(v) =: (a', b')$, we have $|a - a'| \leq 1$ and $b \neq b'$. If additionally $u, v \notin B$, then $a = a'$.
- ($\mathcal{B}4$) For all $y \in Y$, we have $k(y) = (1, \chi(y))$.

Let

$$n_{a,b} := |k^{-1}(a, b)| \quad \text{for all } (a, b) \in [2\ell] \times [2r]. \quad (8.20)$$

Then

$$\sum_{(a,b) \in [2\ell] \times [2r]} n_{a,b} \stackrel{(8.19), (8.20)}{=} |Z| \stackrel{(8.18)}{=} \sum_{(a,b) \in [2\ell] \times [2r]} m_{a,b} \stackrel{(\mathcal{L}2)}{=} n - |V_0| - \sum_{(a,b) \in [2\ell] \times [2r]} \tau_{a,b},$$

and for all $(a, b) \in [2\ell] \times [2r]$,

$$|n_{a,b} - m_{a,b}| \stackrel{(8.20)}{=} ||k^{-1}(a, b)| - m_{a,b}| \stackrel{(\mathcal{B}2)}{\leq} 10\beta n \leq 11\beta(n - |V_0|).$$

Thus $(\mathcal{L}2)$ implies that there is a partition $\mathcal{X} = \{V_0\} \cup \{X_{a,b} : (a, b) \in [2\ell] \times [2r]\}$ of $V(G')$ with, for all $(a, b) \in [2\ell] \times [2r]$,

$$|X_{a,b}| = |k^{-1}(a, b)| + |(f|_{X \setminus I})^{-1}(a, b)| \quad \text{and} \quad |X_{a,b} \triangle V_{\phi^{-1}(a,b)}| \leq \varepsilon^{1/18} m \quad (8.21)$$

such that G' has an $(R^*, 2\ell, 2r, \mathcal{X}, \varepsilon^{1/27}, \delta/4)$ -cycle structure.

Define a mapping $\psi : V(H) \rightarrow ([2\ell] \times [2r]) \cup V_0$ by setting

$$\psi(x) = \begin{cases} f(x) & \text{if } x \in X \\ k(x) & \text{if } x \in Z. \end{cases} \quad (8.22)$$

Finally, let $X' := (X \setminus I) \cup B$.

We need to check that \mathcal{X} , ψ , and X' satisfy Claim 8.2(i)–(v). For (i), we have

$$\begin{aligned} |\psi^{-1}(a, b)| &\stackrel{(8.22)}{=} |k^{-1}(a, b)| + |(f|_{X \setminus I})^{-1}(a, b)| \stackrel{(8.21)}{=} |X_{a,b}| \stackrel{(\mathcal{B}2)}{\geq} m_{a,b} - 10\beta n \\ &\stackrel{(\mathcal{L}1)}{\geq} (1 - \varepsilon^{1/18})m - 10\beta n \stackrel{(8.10)}{\geq} (1 - \varepsilon^{1/19})m. \end{aligned}$$

Further, we have already seen that (ii) holds.

Note that $I = f^{-1}(V_0) = \psi^{-1}(V_0)$ has size $|V_0|$ and is a 2-independent subset of X in H' by $(\mathcal{D}1)$. Let $w \in W := \bigcup_{x \in I} N_H(x)$. Since I is 2-independent, there is a unique $u \in I \subseteq X$ such that $uw \in E(H)$. So $w \in W_{f(u)} = W_{\psi(u)} \subseteq X$ in the notation of $(\mathcal{D}2)$. So $\psi(w) = f(w) \in N_{R^*}^c(f(u)) = N_{R^*}^c(\psi(u))$. Thus

$$d_{G'}(\psi(u), X_{\psi(w)}) \stackrel{(8.21)}{\geq} d_{G'}(\psi(u), V_{\phi^{-1}(\psi(w))}) - \varepsilon^{1/18} m \stackrel{(8.12)}{\geq} cm/2,$$

so (iii) holds.

For (iv), note that $k(y) = (1, \chi(y)) = f(y) = \psi(y)$ for all $y \in Y$ by $(\mathcal{D}5)$ and $(\mathcal{B}4)$. Observe that $\psi' := \psi|_{V(H) \setminus I}$ is a map into $V(R^*) = [2\ell] \times [2r]$. Let $xy \in E(H)$ where $x, y \notin I$. Suppose first that $x, y \in X \cup Y$. Then $\psi(x) = f(x)$ and $\psi(y) = f(y)$. Then $f(x), f(y) \notin V_0$, so $(\mathcal{D}4)$ implies that $f(x)f(y) \in E(R^*)$. Suppose now that $x, y \in Z$. Write $\psi(x) = k(x) = (a, b)$ and $\psi(y) = k(y) = (a', b')$, where $(a, b), (a', b') \in [2\ell] \times [2r]$. Then $(\mathcal{B}3)$ implies that $|a - a'| \leq 1$ and $b \neq b'$. Thus $\psi(x)\psi(y) \in E(Z_{2\ell}^{2r}) \subseteq E(R^*)$, as required. The only other possibility is that one of x, y is in X and the other is in $Z \setminus Y$. But then the distance between them in the bandwidth ordering of H is more than $|Y| = \beta n$, a contradiction to $xy \in E(H)$. Thus $\psi' : V(H \setminus I) \rightarrow V(R^*)$ is a graph homomorphism. So (iv) holds.

For (v), note that $B \subseteq Z$, so $X' \cap I = \emptyset$; and $W = \bigcup_{v \in V_0} W_v \subseteq X$; and $W \cap I = \emptyset$ since I is 2-independent in H' . So $W \subseteq X'$. Now let $(a, b) \in [2\ell] \times [2r]$. We have

$$\begin{aligned} |\psi^{-1}(a, b) \cap X'| &\leq |\psi^{-1}(a, b) \cap X| + |B| \leq |f^{-1}(a, b)| + |B| \\ &\stackrel{(\mathcal{D}3), (\mathcal{B}1)}{\leq} \varepsilon^{1/9} m + 2\ell\beta n \stackrel{(8.10)}{\leq} \varepsilon^{1/10} m. \end{aligned}$$

Now let $uv \in E(H)$, where $u, v \notin X' \cup I$. So $u, v \in Z \setminus B$. Write $\psi(u) = (a, b)$ and $\psi(v) = (a', b')$. Then $(a, b) = k(u)$ and $(a', b') = k(v)$, and $(\mathcal{B}3)$ implies that $a = a'$ and $b \neq b'$, as required.

Finally, define N as in (v). If $y \in N$, then either $y \in \bigcup_{x \in X} N_H(x) \setminus X \subseteq Y$ or $y \in \bigcup_{x \in B} N_H(x)$ (or both). So (8.15) and the fact that $\Delta(H) \leq \Delta$ imply that

$$|N| \leq |Y| + \Delta|B| \leq (2\Delta\ell + 1)\beta n \stackrel{(8.1), (8.10)}{\leq} \varepsilon m. \quad (8.23)$$

This completes the proof of (v) and hence of the claim. \square

In the final part of the proof, we will use the cycle structure \mathcal{C}' , mapping ψ and special set X' obtained in Claim 8.2 to find an embedding g of H into $G' \subseteq G$. We will do this in three stages: (1) define an embedding g_1 of I into V_0 , according to ψ ; (2) find an embedding g_2 of X using ψ as a framework,

such that there are large candidate sets for the neighbouring vertices N of X' ; (3) find an embedding g_3 of the remainder of H using the blow-up lemma, using the candidate sets obtained in (2) to ensure that g_2 is compatible with g_3 . Then set g to be the union of g_1, g_2, g_3 .

Stage (1) is easy; we simply define

$$g_1 : I \rightarrow V_0 \quad \text{where} \quad g_1(x) := \psi(x) \quad \text{for all } x \in I.$$

Since by Claim 8.2(iii), I is an independent set in H of size V_0 , we trivially have that g_1 is an embedding of $H[I]$ into $V(G')$.

For Stage (2), we will apply Lemma 4.5 (embedding lemma with target sets) to embed vertices in X' . Indeed, let $\psi^* := \psi|_{X' \cup N}$. Given $w \in W$, let u be the unique element of I such that $uw \in E(H)$, as guaranteed by Claim 8.2(iii). Let

$$S_w := N_{G'}(\psi(u), X_{\psi(w)}). \quad (8.24)$$

We will apply Lemma 4.5 with $G' \setminus V_0, R^*, H[X' \cup N], n - |V_0|, 4r\ell, \varepsilon^{1/27}, c/2, \delta/4, \Delta, \{X_{a,b}\}, (1 - \varepsilon^{19})m, \psi^*, X', N, W, S_w$ playing the roles of $G, R, H, n, L, \varepsilon, c, \delta, \Delta, \{V_a : a \in V(R)\}, m, \phi, X, Y, W, S_w$. To see why this is possible, note that, by Claim 8.1(iii), $G' \setminus V_0$ has $n - |V_0| \geq (1 - 2\varepsilon^{1/2})n$ vertices; and Claim 8.2(ii) (specifically ($\mathcal{C}2$)) implies that it has an $(\varepsilon^{1/27}, \delta/4)$ -regular partition $\{X_{a,b} : (a, b) \in V(R^*)\}$. Clearly, as a restriction of ψ , the function ψ^* is a suitable graph homomorphism, and by Claim 8.2(v) and (8.23), we have

$$|(\psi^*)^{-1}(a, b)| \leq |\psi^{-1}(a, b) \cap X'| + |N| \leq \varepsilon^{1/10}m + \varepsilon m \leq \varepsilon^{1/12}m. \quad (8.25)$$

Finally, $W \subseteq X'$ by Claim 8.2(v), and $|S_w| \geq cm/2$ by Claim 8.2(iii). So the above are suitable parameters for the application of Lemma 4.5.

Thus there is a mapping

$$g_2 : X' \rightarrow V(G') \setminus V_0$$

that is an embedding of $H[X']$ into G' such that

- ($\mathcal{T}1$) $g_2(x) \in X_{\psi^*(x)}$ for all $x \in X'$.
- ($\mathcal{T}2$) $g_2(w) \in S_w$ for all $w \in W$.
- ($\mathcal{T}3$) For all $y \in N$, there exists $C_y \subseteq X_{\psi^*(y)} \setminus g_2(X')$ such that $C_y \subseteq N_{G'}(g_2(x))$ for all $x \in N_H(y) \cap (X')$, and $|C_y| \geq cm/2$.

For Stage (3), we will do the following for each $a \in [2\ell]$. Let $U_{a,b} := X_{a,b} \setminus g_2(X')$ for all $b \in [2r]$. We want to show that $U_{a,b}$ has exactly the right size to embed the remaining vertices of H whose image under ψ is (a, b) . Indeed, let $\psi' := \psi|_{H \setminus (X' \cup I)}$. Then Claim 8.2(i) implies that

$$|U_{a,b}| = |X_{a,b}| - |g_2(X') \cap X_{a,b}| \stackrel{(\mathcal{T}1)}{=} |\psi^{-1}(a, b)| - |(\psi^*)^{-1}(a, b) \cap X'| = |(\psi')^{-1}(a, b)|$$

where we used the fact that $\psi^{-1}(a, b) \cap I = \emptyset$. This together with (8.25) implies that $|U_{a,b} \Delta X_{a,b}| = |(\psi^*)^{-1}(a, b) \cap X'| \leq \varepsilon^{1/10}m \leq 2\varepsilon^{1/10}|U_{a,b}|$. Let $b, b' \in [2r]$ be distinct. So $|U_{a,b}| \geq (1 - \varepsilon^{1/20})m$ by Claim 8.2(i). Recall from Claim 8.2(ii) (specifically ($\mathcal{C}3$)) that $G'[X_{a,b}, X_{a,b'}]$ is $(\varepsilon^{1/27}, \delta/4)$ -superregular. So given any $x \in U_{a,b}$, Claim 8.2(i) implies that

$$d_{G'}(x, U_{a,b'}) \geq \delta|X_{a,b}|/4 - \varepsilon^{1/10}m \geq (\delta/4 - \delta\varepsilon^{1/19} - \varepsilon^{1/10})m \geq \delta|U_{a,b}|/5.$$

Thus Proposition 4.3 with $G', X_{a,b}, U_{a,b}, X_{a,b'}, U_{a,b'}, \varepsilon^{1/27}, \delta/4, \varepsilon^{1/10}$ playing the roles of $G, A, A', B, B', \varepsilon, \delta, \alpha$ implies that $G'[U_{a,b}, U_{a,b'}]$ is $(2\varepsilon^{1/27}, \delta/5)$ -superregular for all distinct $b, b' \in [2r]$. The set N has size at most $\varepsilon m \leq 2\varepsilon|U_{a,b}|$ for any $b \in [2r]$; and for each $y \in N \cap (\psi')^{-1}(a, b)$, ($\mathcal{T}3$) guarantees a corresponding set $C_y \subseteq X_{\psi^*(y)} \setminus g_2(X') = U_{\psi'(y)} = U_{a,b}$ that has size at least $cm/2 \geq c|U_{a,b}|/3$. Let H_a denote the subgraph of H induced by the set of all $x \in V(H) \setminus (X' \cup I)$

such that $\psi'(x) = (a, b)$ for some $b \in [2r]$. Now apply, for each $a \in [2\ell]$, Lemma 4.6 (blow-up lemma) with $G'[\bigcup_{b \in [2r]} U_{a,b}]$ and H_a playing the roles of G and H and $2\varepsilon^{1/27}, 2\varepsilon, \delta/5, c/3, \Delta, 2r, \{U_{a,b} : b \in [2r]\}, \psi', C_y$ playing the roles of $\varepsilon, \alpha, \delta, c, \Delta, k, \{V_a : a \in [k]\}, \phi, S_y$. Altogether, this yields a mapping

$$g_3 : V(H) \setminus (X' \cup I) \rightarrow V(G') \setminus (V_0 \cup g_2(X'))$$

that is an embedding of $H \setminus (X' \cup I)$ into $V(G')$ such that every $y \in N$ is mapped to a vertex in C_y .

We claim that the mapping g given by

$$g(x) = \begin{cases} g_1(x) & \text{if } x \in I \\ g_2(x) & \text{if } x \in X' \\ g_3(x) & \text{otherwise} \end{cases} \quad (8.26)$$

is an embedding of H into G' (and hence into G).

Firstly, g is an injective map from $V(H)$ to $V(G')$ by the definitions of g_1, g_2, g_3 . So we just need to check that it is a graph homomorphism. Also, by their definitions, each of g_1, g_2, g_3 is an embedding of H induced on its respective domain into G' . So it suffices to check that whenever $xy \in E(H)$ and x, y are not both in I or in X' or in $V(H) \setminus (X' \cup I)$, $g(x)g(y) \in E(G')$.

Suppose first that $x \in I$ and $y \in V(H) \setminus I$. Then $g(x) = g_1(x) = \psi(x)$ and $y \in W \subseteq X'$ (here we used Claim 8.2(v)). So $g(y) = g_2(y)$. Claim 8.2(iii) implies that x is the only vertex in I that is a neighbour of y . Then

$$g(y) \stackrel{(8.26)}{=} g_2(y) \stackrel{(\mathcal{T}2)}{\in} S_y \stackrel{(8.24)}{=} N_{G'}(\psi(x), X_{\psi(y)}) = N_{G'}(g(x), X_{\psi(y)}).$$

So $g(x)g(y) \in E(G')$, as required.

Therefore, we may assume that $x \in X'$ and $y \in V(H) \setminus (X' \cup I)$. Then $g(x) = g_2(x)$, $y \in N$, and $g(y) = g_3(y) \in C_y$, where C_y was defined in $(\mathcal{T}3)$, which guarantees that $C_y \subseteq N_{G'}(g_2(x)) = N_{G'}(g(x))$. So $g(x)g(y) \in E(G')$, as required. This completes the proof of Theorem 1.2.

9. Concluding remarks

In this paper, we prove a version of the bandwidth theorem for locally dense graphs. As mentioned in the introduction, it is also of interest to seek minimum degree conditions that force a given spanning structure in a graph with sublinear independence number. In particular, it would be very interesting to obtain an analogue of the bandwidth theorem in this setting.

In a step in this direction, Balogh, Molla, and Sharifzadeh [4] proved the following result on triangle factors.

Theorem 9.1 (Balogh, Molla, and Sharifzadeh [4]). *For every $\varepsilon > 0$, there exist $\gamma > 0$ and $n_0 \in \mathbb{N}$ such that the following holds. For every n -vertex graph G with $n \geq n_0$ divisible by 3, if $\delta(G) \geq (1/2 + \varepsilon)n$ and G has independence number $\alpha(G) \leq \gamma n$, then G has a K_3 -factor.*

Perhaps the next natural step is to ascertain whether the conclusion of Theorem 9.1 can be strengthened to ensure the square of a Hamilton cycle.

Conjecture 9.2. *For every $\varepsilon > 0$, there exist $\gamma > 0$ and $n_0 \in \mathbb{N}$ such that the following holds. For every n -vertex graph G with $n \geq n_0$, if $\delta(G) \geq (1/2 + \varepsilon)n$ and $\alpha(G) \leq \gamma n$, then G contains the square of a Hamilton cycle.*

It is also natural to seek a version of Theorem 1.2 where now one replaces the condition of *locally dense* with a more restrictive *uniformly dense* condition: given $\rho, d > 0$, we say that an n -vertex graph G is (ρ, d) -uniformly dense if every $X, Y \subseteq V(G)$ satisfies $e_G(X, Y) \geq d|X||Y| - \rho n^2$. If one restricts to uniformly dense graphs, then one can substantially reduce the minimum degree condition in Theorem 1.2, as well as remove the bandwidth condition on H .

Theorem 9.3. For all $\Delta \in \mathbb{N}$ and $d, \eta > 0$, there exist constants $\rho, n_0 > 0$ such that for every $n \geq n_0$, the following holds. Let H be an n -vertex graph with $\Delta(H) \leq \Delta$. Then any (ρ, d) -uniformly dense graph G on n vertices with $\delta(G) \geq \eta n$ contains a copy of H .

Theorem 9.3 can be proven by a simple application of the blow-up lemma; a more general ‘rainbow’ version of Theorem 9.3 is given in [20, Corollary 1.3].

Acknowledgements. Katherine Staden was supported by ERC grant 306493, and Andrew Treglown was supported by EPSRC grant EP/M016641/1.

We would like to thank Maryam Sharifzadeh for many helpful conversations at the start of the project. Andrew Treglown is grateful to Stefan Glock and Felix Joos for a conversation on [20] that brought to light the version of the bandwidth theorem for uniformly dense graphs.

The authors are also grateful to the referees for their helpful and careful reviews.

Conflict of Interest: None.

References

- [1] P. Allen, J. Böttcher, J. Ehrenmüller, and A. Taraz, ‘The bandwidth theorem in sparse graphs’, *Advances Combin.* (2020):6, 60pp.
- [2] J. Balogh and J. Lenz, ‘Some exact Ramsey–Turán numbers’, *Bull. London Math. Soc.* **44** (2012), 1251–1258.
- [3] J. Balogh, A. McDowell, T. Molla, and R. Mycroft, ‘Triangle-tilings in graphs without large independent sets’, *Combin. Probab. Comput.* **27** (2018), 449–474.
- [4] J. Balogh, T. Molla, and M. Sharifzadeh, ‘Triangle factors of graphs without large independent sets and of weighted graphs’, *Random Structures Algorithms* **49** (2016), 669–693.
- [5] J. Böttcher, Y. Kohayakawa, and A. Taraz, ‘Almost spanning subgraphs of random graphs after adversarial edge removal’, *Combin. Probab. Comput.* **22** (2013), 639–683.
- [6] J. Böttcher, R. Montgomery, O. Parczyk, and Y. Person, ‘Embedding spanning bounded degree graphs in randomly perturbed graphs’, *Mathematika* **66** (2020), 422–447.
- [7] J. Böttcher, K. Preussmann, A. Taraz, and A. Würfl, ‘Bandwidth, expansion, treewidth, separators and universality for bounded-degree graphs’, *European J. Combin.* **31** (2010), 1217–1227.
- [8] J. Böttcher, M. Schacht, and A. Taraz, ‘Spanning 3-colourable subgraphs of small bandwidth in dense graphs’, *J. Combin. Theory B* **98** (2008), 752–777.
- [9] J. Böttcher, M. Schacht, and A. Taraz, ‘Proof of the bandwidth conjecture of Bollobás and Komlós’, *Math. Ann.* **343** (2009), 175–205.
- [10] J. Böttcher, A. Taraz, and A. Würfl, ‘Spanning embeddings of arrangeable graphs with sublinear bandwidth’, *Random Structures Algorithms*, **48** (2016), 270–289.
- [11] V. Chvátal, V. Rödl, E. Szemerédi, and W.T. Trotter Jr., ‘The Ramsey number of a graph with bounded maximum degree’, *J. Combin. Theory Ser. B* **34** (1983), 239–243.
- [12] P. Condon, J. Kim, D. Kühn, and D. Osthus, ‘A bandwidth theorem for approximate decompositions’, *Proc. London Math. Soc.* **118** (2019), 1393–1449.
- [13] G.A. Dirac, ‘Some theorems on abstract graphs’, *Proc. London Math. Soc.* **2** (1952), 69–81.
- [14] O. Ebsen, G.S. Maesaka, Chr. Reiher, M. Schacht, and B. Schülke, ‘Embedding spanning subgraphs in uniformly dense and inseparable graphs’, *Random Structures and Algorithms*, **57** (2020), 1077–1096.
- [15] P. Erdős, Problem 9, in: M. Fieldler (ed.), *Theory of Graphs and Its Applications* (Czech. Acad. Sci. Publ., Prague, 1964), 159.
- [16] P. Erdős, R.J. Faudree, C.C. Rousseau, and R.H. Schelp, ‘A local density condition for triangles’, *Discrete Math.* **127** (1994), 153–161.
- [17] P. Erdős, A. Hajnal, V.T. Sós, and E. Szemerédi, ‘More results on Ramsey–Turán type problems’, *Combinatorica* **3** (1983), 69–81.
- [18] P. Erdős and V.T. Sós, ‘Some remarks on Ramsey’s and Turán’s theorem’, in: *Combinatorial Theory and Its Applications* vol. **II** (North-Holland, Amsterdam, 1970), 395–404.
- [19] A. Ferber, R. Nenadov, A. Noever, U. Peter, and N. Skoric, ‘Robust hamiltonicity of random directed graphs’, *J. Combin. Theory Ser. B* **126** (2017), 1–23.
- [20] S. Glock and F. Joos, ‘A rainbow blow-up lemma’, *Random Structures Algorithms*, to appear.
- [21] A. Hajnal and E. Szemerédi, ‘Proof of a conjecture of Erdős’, in: *Combinatorial Theory and Its Applications* vol. **II** **4** (North-Holland, Amsterdam, 1970), 601–623.
- [22] J. Han, ‘On perfect matchings and tilings in uniform hypergraphs’, *SIAM J. Discrete Math.*, **32** (2018), 919–932.
- [23] H. Huang, C. Lee, and B. Sudakov, ‘Bandwidth theorem for random graphs’, *J. Combin. Theory B* **102** (2012), 14–37.

- [24] F. Knox and A. Treglown, 'Embedding spanning bipartite graphs of small bandwidth', *Combin. Probab. Comput.* **22** (2013), 71–96.
- [25] Y. Kohayakawa, B. Nagle, V. Rödl, and M. Schacht, 'Weak regularity and linear hypergraphs', *J. Combin. Theory B* **100** (2010), 151–160.
- [26] J. Komlós, 'The blow-up lemma', *Combin. Probab. Comput.* **8** (1999), 161–176.
- [27] J. Komlós, G.N. Sárközy, and E. Szemerédi, 'Blow-up lemma', *Combinatorica* **17** (1997), 109–123.
- [28] J. Komlós, G.N. Sárközy, and E. Szemerédi, 'Proof of the Seymour conjecture for large graphs', *Annals of Combinatorics* **2** (1998), 43–60.
- [29] D. Kühn and D. Osthus, 'The minimum degree threshold for perfect graph packings', *Combinatorica* **29** (2009), 65–107.
- [30] C. Lee, 'Embedding degenerate graphs of small bandwidth', submitted, arXiv:1501.05350.
- [31] D. Mubayi and V.T. Sós, 'Explicit constructions of triple systems for Ramsey–Turán problems', *J. Graph Theory* **52** (2006), 211–216.
- [32] V. Rödl, A. Ruciński, and E. Szemerédi, 'A Dirac-type theorem for 3-uniform hypergraphs', *Combin. Probab. Comput.* **15** (2006), 229–251.
- [33] P. Seymour, Problem section, in: T.P. McDonough and V.C. Mavron (eds.), *Combinatorics: Proceedings of the British Combinatorial Conference 1973* (Cambridge University Press, 1974), 201–202.
- [34] M. Simonovits and V. T. Sós, 'Ramsey–Turán theory', *Discrete Math.* **229** (2001), 293–340.
- [35] K. Staden and A. Treglown, 'On degree sequences forcing the square of a Hamilton cycle', *SIAM J. Disc. Math.* **31** (2017), 383–437.
- [36] K. Staden and A. Treglown, 'On degree sequences forcing the square of a Hamilton cycle', arXiv version, arXiv:1412.3498.
- [37] E. Szemerédi, 'Regular partitions of graphs', *Problèmes Combinatoires et Théorie des Graphes Colloques Internationaux CNRS* **260** (1978), 399–401.