

Comment on "A Modern Assessment of Cancer Risk in Adrenal Incidentalomas

Chortis, Vasileios; Sitch, Alice J; Bancos, Irina; Prete, Alessandro; Taylor, Angela E; Biehl, Michael; Deeks, Jonathan J; Arlt, Wiebke

DOI:

[10.1097/SLA.0000000000004804](https://doi.org/10.1097/SLA.0000000000004804)

License:

Creative Commons: Attribution (CC BY)

Document Version

Publisher's PDF, also known as Version of record

Citation for published version (Harvard):

Chortis, V, Sitch, AJ, Bancos, I, Prete, A, Taylor, AE, Biehl, M, Deeks, JJ & Arlt, W 2021, 'Comment on "A Modern Assessment of Cancer Risk in Adrenal Incidentalomas: Analysis of 2219 Patients" by Kahramangil B et al', *Annals of surgery*, vol. 274, no. 6, pp. e887-e888. <https://doi.org/10.1097/SLA.0000000000004804>

[Link to publication on Research at Birmingham portal](#)

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.



Annals of Surgery

DOI: 10.1097/SLA.0000000000004804

Letter to the Editor

Comment on “A Modern Assessment of Cancer Risk in Adrenal Incidentalomas: Analysis of 2219 Patients” by Kahramangil B et al.

Vasileios Chortis¹, Alice J. Sitch^{2,3}, Irina Bancos⁴, Alessandro Prete¹, Angela E. Taylor¹, Michael Biehl⁵, Jonathan J. Deeks^{2,3}, Wiebke Arlt^{1,3}

¹Institute of Metabolism and Systems Research and ²Institute of Applied Health Research, College of Medical and Dental Sciences, University of Birmingham, Birmingham, UK

³NIHR Birmingham Biomedical Research Centre, University Hospitals Birmingham NHS Foundation Trust and University of Birmingham, Birmingham, UK

⁴Division of Endocrinology, Diabetes, Metabolism and Nutrition³, Mayo Clinic, Rochester, MN, USA

⁵Bernoulli Institute for Mathematics, Computer Science and Artificial Intelligence, University of Groningen, Groningen, The Netherlands

Correspondence:

Professor Wiebke Arlt, Institute of Metabolism and Systems Research (IMSR), University of Birmingham, B15 2TT, Birmingham, UK

Email: w.arlt@bham.ac.uk

This is an open access article distributed under the Creative Commons Attribution License 4.0, (CCBY), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Dear Editor,

We have read with great interest the paper 'A Modern Assessment of Cancer Risk in Adrenal Incidentalomas – Analysis of 2219 Patients' by Kahramangil *et al*¹. The diagnostic work-up of incidentally discovered adrenal masses remains challenging, in particular with regard to the critically important differentiation of adrenocortical carcinoma (ACC) from benign adrenal lesions. This study on a large retrospective cohort of adrenal incidentalomas (AI) is a welcome contribution to the literature. We recently published a prospective, multi-center study validating a new biochemical diagnostic test for adrenal tumors, urine steroid metabolomics, in a similarly sized cohort of 2017 patients with newly discovered adrenal masses including 98 ACC (Evaluation of URINE steroid metabolomics in the differential diagnosis of AdrenoCortical Tumors, EURINE-ACT)². Notably, 84% of the tumors in our cohort were incidental findings (AI); patients with biochemical evidence of pheochromocytoma or recent extra-adrenal malignancy were excluded.

Our prospective study² and the retrospective study by Kahramangil *et al.*¹ concur on two important points of direct relevance to clinical practice. First, both studies report that ACCs are extremely rare in lesions measuring <4 cm (post-test probability of ACC 0.1% in Kahramangil *et al.*¹, 0.13% in the EURINE-ACT cohort). Second, both studies provide evidence that an unenhanced CT tumor attenuation value of >20 Hounsfield Units (HU) is a more specific diagnostic cut-off than the hitherto applied threshold of 10 HU. In the prospective EURINE-ACT study, we found a specificity of 64% (95%CI 61.4%-66.4%) for >10HU, which increased to 80% (95%CI 77.9%-82.0%) when applying >20HU, with similar sensitivities (100% and 99%, respectively). Kahramangil *et al.*¹ reported a similar specificity for >20HU (76.8%; sensitivity 94.1%, no 95% CIs reported).

However, we would like to voice important concerns about the implementation of the management algorithm proposed by Kahramangil *et al.*¹ into clinical practice. Their algorithm suggests that all adrenal masses with a size >4.6cm or with an attenuation >20HU on non-contrast CT should be surgically excised. Although these criteria indeed minimize the risk of missing an ACC, this comes at the very substantial cost of frequent unnecessary adrenalectomies in a much larger number of patients with benign adrenal tumors, with all attendant risks and costs^{3,4}. This is due to the poor positive predictive value (PPV) of tumor size and tumor attenuation, as a result of the low overall ACC prevalence in patients with AI^{1,2}. By our calculations, in the Kahramangil *et al.* cohort (1.7% overall rate of ACC), a tumor size >4.6 cm would correctly detect 35 out of 38 ACCs. However, this would also yield 231 false positive results, corresponding to patients who would end up undergoing adrenalectomy for non-ACC tumors. This is equivalent to 6.6 unnecessary adrenalectomies for each surgically removed ACC, excepting the rare cases of hormonally active adrenal tumors. Similarly, when applying a tumor attenuation of >20HU, only 6.3% of those lesions were actually confirmed as ACC by histopathology; the disproportionally high false-positive rate of 93.7% will result in 14.7 unnecessary adrenalectomies per excised ACC.

We also urge caution regarding their proposed algorithm from a methodological point of view. Kharangil *et al.* performed univariable logistic regression to identify risk factors, followed by multivariable logistic regression to present a prognostic model. However, the

proposed cut-points were derived from their retrospective dataset only, without any further internal or external validation. It is also problematic to use unadjusted models to select variables for adjusted models and the authors did not report confidence intervals in the main manuscript.

In our recently published prospective test validation study², we propose a management algorithm for patients with a newly diagnosed adrenal mass, based on a 'triple test', taking a stepwise approach, with sequential application of size criteria (diameter > 4 cm), imaging characteristics (non-contrast attenuation > 20 HU) and urine steroid metabolomics (USM)². USM is a new diagnostic biomarker test that combines mass spectrometry-based steroid metabolite profiling with steroid data analysis by a machine learning-based algorithm.^{2,5,6}. This strategy provided a positive predictive value of 76.4% and a negative predictive value of 99.7% for ACC diagnosis². This triple test approach has the great advantage of drastically curtailing the number of unnecessary adrenalectomies, while remaining highly sensitive to the detection of ACCs. In addition, it dramatically decreases the numbers of additional imaging procedures, which all have lower specificity than USM, as nicely demonstrated by Kahramangil et al.¹ for the widely used CT contrast wash-out studies, which they found to falsely predict malignancy in 91% of patients.

References

1. Kahramangil B, Kose E, Remer EM, et al. A Modern Assessment of Cancer Risk in Adrenal Incidentalomas: Analysis of 2219 Patients. *Ann Surg*. 2020.
2. Bancos I, Taylor AE, Chortis V, et al. Urine steroid metabolomics for the differential diagnosis of adrenal incidentalomas in the EURINE-ACT study: a prospective test validation study. *Lancet Diabetes Endocrinol*. 2020;8(9):773-781.
3. Palazzo F, Dickinson A, Phillips B, et al. Adrenal surgery in England: better outcomes in high-volume practices. *Clin Endocrinol (Oxf)*. 2016;85(1):17-20.
4. Park HS, Roman SA, Sosa JA. Outcomes from 3144 adrenalectomies in the United States: which matters more, surgeon volume or specialty? *Arch Surg*. 2009;144(11):1060-1067.
5. Arlt W, Biehl M, Taylor AE, et al. Urine steroid metabolomics as a biomarker tool for detecting malignancy in adrenal tumors. *J Clin Endocrinol Metab*. 2011;96(12):3775-3784.
6. Chortis V, Bancos I, Nijman T, et al. Urine Steroid Metabolomics as a Novel Tool for Detection of Recurrent Adrenocortical Carcinoma. *J Clin Endocrinol Metab*. 2020;105(3).