

## Age, BMI, and inflammation

Balter, Leonie JT; Raymond, Jane E.; Aldred, Sarah; Higgs, Suzanne; Bosch, Jos A.

DOI:

[10.1016/j.physbeh.2021.113324](https://doi.org/10.1016/j.physbeh.2021.113324)

License:

Creative Commons: Attribution-NonCommercial-NoDerivs (CC BY-NC-ND)

*Document Version*

Peer reviewed version

*Citation for published version (Harvard):*

Balter, LJT, Raymond, JE, Aldred, S, Higgs, S & Bosch, JA 2021, 'Age, BMI, and inflammation: associations with emotion recognition', *Physiology and Behavior*, vol. 232, 113324. <https://doi.org/10.1016/j.physbeh.2021.113324>

[Link to publication on Research at Birmingham portal](#)

### General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

### Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact [UBIRA@lists.bham.ac.uk](mailto:UBIRA@lists.bham.ac.uk) providing details and we will remove access to the work immediately and investigate.

1           **Age, BMI, and inflammation: associations with emotion recognition**

2           Leonie JT Balter<sup>\*A</sup> Jane E Raymond<sup>B</sup>, Sarah Aldred<sup>C</sup>, Suzanne Higgs<sup>B</sup> & Jos A Bosch<sup>D</sup>

4           <sup>A</sup> Stress Research Institute, Stockholm University, Stockholm, SE-10691

5           <sup>B</sup> School of Psychology, University of Birmingham, Birmingham, B15 2TT, UK

6           <sup>C</sup> School of Sport, Exercise, and Rehabilitation Sciences, University of Birmingham,  
7           Birmingham, B15 2TT, UK

8           <sup>D</sup> Psychology Department, Clinical Psychology, University of Amsterdam, Amsterdam, 1018  
9           WT, NL

11          \* Corresponding Author:

12          Stress Research Institute, Stockholm University

13          SE-10691 Stockholm, Sweden

14          Email: leonie.balter@su.se

17          Keywords: Inflammation; aging, body mass index; reading the mind in the eyes; emotion  
18          recognition; psychoneuroimmunology

## ABSTRACT

Experimental studies show that inflammation impairs the ability to interpret the mental state of another person, denoted theory of mind (ToM). The current study attempted a conceptual replication in states associated with elevated low-grade inflammation, i.e., high body weight and advanced age.

Ninety young ( $M = 26.3$  years,  $SD = 4.1$ ) or older ( $M = 70.7$  years,  $SD = 4.0$ ) participants with either a normal body mass index (BMI) ( $M = 22.4$ ,  $SD = 2.2$ ) or high BMI ( $M = 33.1$ ,  $SD = 3.8$ ) completed the Reading the Mind in the Eyes Test (RMET) to assess ToM. Plasma interleukin-6 (IL-6) level was measured to index low-grade inflammation.

As anticipated, elevated IL-6 levels were found with higher BMI, although not with increased age. IL-6 was associated with poorer task performance, independent of potential demographic and health confounders (e.g., sex, education, smoking status, alcohol intake, presence of illness symptoms, and medication intake). Analyses also revealed an interaction whereby young individuals with a high BMI showed worse RMET performance compared to their normal BMI counterparts, whereas the opposite pattern was found in older individuals.

The present observational study replicated experimental results showing that elevated low-grade inflammation is correlated with a lower ability to infer the mental states of others. These findings suggest that also naturalistic conditions of (protracted) low-grade inflammation may alter emotion recognition.

## 1. INTRODUCTION

Human and animal studies have identified acute inflammation as a powerful regulator of social behaviors [1]–[3]. Two recent studies showed that induced acute inflammation impaired the ability to recognize emotions expressed by others, a central component of social cognition denoted as Theory of Mind (ToM) [2], [3]. Impairment of emotion recognition is a transdiagnostic mechanism implicated in a number of mental health disorders, most notably depression [4], [5]. For example, impaired emotion recognition is thought to explain why depressed individuals tend to express more social difficulties and exhibit social withdrawal [4], [6]. Whether such inflammatory effects on emotion recognition are also observed in non-experimental, naturalistic conditions characterized by low-grade inflammation remains unanswered.

Protracted low-grade inflammation is seen among individuals with high BMI and in advanced age [7], [8]. In the former, inflammation is thought to emanate mainly from adipose cells and surrounding immune cells that produce copious amounts of inflammatory cytokines (Cancello & Clément, 2006). In aging, the sources of elevated inflammatory activity, denoted as “inflammaging”, are more diffuse and may involve factors such as oxidative stress, immunosenescence (i.e., age-related immune impairment), hormonal changes, and the gradual surge of inflammatory conditions such as atherosclerosis [10]–[12]. Age and BMI are therefore relatively independent determinants of low-grade inflammation. Considering that these states are rather stable (i.e., do not change from one week to the other), these factors may be utilized to study the relationship between emotion recognition and persistent low-grade inflammation.

High BMI and older age are associated with lower performance across multiple cognitive domains [13]–[18]. However, only a handful of studies have addressed the combined effects of BMI and age on cognitive functions, and none assessed emotion recognition. These isolated studies show both additive and synergistic effects, whereby obesity-related cognitive deficits are independent of age [19]–[22] or appear amplified with increased age [22]. Some reports also indicate a possible protective effect whereby the impairments associated with high BMI

disappear or reverse with increasing age; a phenomenon referred to as the “obesity paradox”. For example older adults with a relatively higher BMI exhibit less decline in visuospatial skills [23]–[26]. In a similar vein, poorer memory recall performance over 12 years was predicted by a decline in BMI during that period [27], and low BMI and weight loss in older age precede the onset of dementia [28]–[30]. Together these data suggest that in older age a lower BMI, rather than higher BMI, may be a risk factor for cognitive decline. However, age-dependent BMI effects on social cognition, and more specifically emotion recognition, have remained unexplored.

It is expected that both aging and obesity are linked to alterations in emotion recognition, although this may not apply to all emotions equally. E.g., recognition of disgust seems to be preserved with age whereas the emotions anger and sadness are particularly affected [31], [32]. A meta-analysis of emotion processing studies demonstrated that individuals with obesity have lower levels of emotional awareness and difficulty in using emotion regulation strategies [33]. The evidence for an impaired ability to recognize emotions appeared inconsistent [33]. However, this meta-analysis included only two studies that assessed emotion recognition. Of these two studies, one study reported no difference comparing women with obesity and normal-weight women [34], while the other study reported impairment among women with obesity [35]. Moreover, these inconclusive findings were based on studies with young to middle-aged individuals (between 28 and 49 years).

Therefore, the aim of the present cross-sectional study was twofold. First, to assess the association of BMI- and age-related low-grade inflammation (as measured by IL-6) with emotion recognition. Second, to determine if age and BMI show independent or interactive associations with emotion recognition.

## **2. METHOD**

### *2.1 Participants*

Ninety participants (60% female) were recruited through a database held by the University of Birmingham and via (online) advertisement. Inclusion criteria included an age

between 21 and 35 years ('young';  $M = 26.3$ ,  $SD = 4.1$ ) or between 63 and 80 years ('older';  $M = 70.7$ ,  $SD = 4.0$ ) and a BMI between 17 and 25 ('normal BMI';  $M = 22.4$ ,  $SD = 2.2$ ) or greater than 27 ('high BMI';  $M = 33.1$ ,  $SD = 3.8$ ). Height and body weight were confirmed in the laboratory. One participant had a BMI of 26.3 and thus failed to meet the BMI criterion for either the normal BMI or the high BMI group and was therefore removed from all analyses. Excluded were individuals who reported a history of gastric banding, eating disorders, neurological or inflammatory disorders (e.g., rheumatoid arthritis, inflammatory bowel disease, multiple sclerosis, periodontitis) or use of anti-depressant, anti-histamine, or anti-inflammatory (e.g., antibiotics) medication during the past 7 days were excluded. Participants reported normal or corrected-to-normal vision and stable body weight for at least six months (i.e., fluctuations  $< 7.5$  kg for individuals with high BMI,  $< 5$  kg for normal BMI individuals). The participants were paid a maximum of £25 to reimburse travel expenses. The study was conducted according to the guidelines laid down in the Declaration of Helsinki and all procedures were approved by the University of Birmingham Research Ethics Committee.

## 2.2 Procedures

Test sessions started throughout the day between 8:30 and 15:30 hours. Start times of test sessions were matched across the four groups to control for diurnal variations in IL-6 [36]. Written informed consent was obtained on arrival. Participants were instructed to have their breakfast/lunch as usual but avoid consumption of high-fat products (e.g., bacon, fries), because these foods may induce a short-lived inflammatory response [37], and to refrain from eating, drinking (except for water), and smoking for 1 hour before the start of the test session. Participants were also asked not to engage in strenuous physical exercise or consume alcohol within 12 hours before the test session, and reschedule their appointment if they had suspected infection symptoms on the day of testing. Self-reports were used to verify whether participants had complied with instructions. A blood sample was taken by venipuncture and questionnaires and cognitive tests were completed, including the reading the mind in the eyes test (see further below). Other tests included measures of attention and psychomotor speed

(published elsewhere [21]), memory, and reinforcement learning. The order in which the tasks were administered was fixed and the same across all the participants. Lastly, a measure of height and body composition was taken.

### **3. MATERIALS**

#### *3.1 Reading the Mind in the Eyes test*

The reading the mind in the eyes test (RMET) is considered an advanced test of theory of mind involving mental state attribution and complex emotion recognition from photographs of the eye region of the face [38], [39].

##### *3.1.1 Procedure*

The test display comprised a test eye image and four words placed in the center of the screen. The participant was instructed to select the word that best described what the person in the test image was thinking or feeling by pressing one of four computer keys (Q, P, A, L) that spatially corresponded to the position of each word. The correct (target) word had the same emotional valence as the accompanying three foil words. For example, the target word, 'panicked', was accompanied by 'arrogant', 'jealous' and 'hateful'. Target words were equally likely to appear in one of the four word locations on the screen. Each test display remained visible until a key response was made; the text test display was immediately presented thereafter. The test consisted of 36 different images, completed as one set. In line with previous studies, a glossary containing a definition of each word was available to the participant.

##### *3.1.2 Stimuli*

A grey-scale digital image (subtending 9 x° X 3.6 y° of visual angle) of the eye region of a face (including eyes and eyebrows) was presented in the middle of a grey field on a computer monitor. Four words describing mental states accompanied each test stimulus, presented in black Arial font (subtending 2.6 x° X 0.7 x° of visual angle).

### 3.2 Blood sampling

Blood (6 mL) was collected from an antecubital vein in the forearm into one vacutainer containing ethylenediaminetetraacetic acid (EDTA) as anticoagulant (Becton Dickinson Diagnostics, Oxford, United Kingdom). Samples were immediately centrifuged at 1500 x g for 10 min at 4 °C and plasma was aliquoted and stored at -80 °C for later assessment of interleukin-6 (IL-6), a marker of system low-grade inflammation. Plasma level of IL-6 was measured in duplicate using high-sensitivity enzyme-linked immunosorbent assay (ELISA) (Quantikine HS Human IL-6 ELISA, R&D Systems, UK) in accordance with the manufacturer's instructions. The limits of detection of this assay was 0.11 pg/mL, with intra- and inter-assay coefficients of variation (CVs) of 0.69-11.6%.

### 3.3 Anthropomorphic Measures

Participants were asked to remove footwear and coats and empty their pockets before a body composition measurement was taken using a TANITA BC-545N body composition analyzer (Tanita Europe, Amsterdam, The Netherlands). A stadiometer was used to measure height.

### 3.4 Questionnaires

To adjust for potential confounding factors, participants completed questionnaires about illness symptoms (modified version of self-administered comorbidity questionnaire (SCQ); Sangha, Stucki, Liang, Fossel, & Katz, 2003), sleep quality (Pittsburgh Sleep Quality Index (PSQI); Carpenter & Andrykowski, 1998), medication intake (number of medications), and demographic variables (i.e., age, sex, occupation status, education, smoking status (current smoker, ex-smoker, non-smoker), alcohol intake (units per week)).

## 4. STATISTICAL ANALYSES

Data from participants who disproportionately biased mean estimates were removed using Cook's Distance and data points that exceeded 3 SDs from means ( $N = 2$ , young high BMI individuals with an average accuracy < 30%).

For IL-6 analysis, log transformation was applied because of the skewed distribution of raw IL-6 values. There were six missing values (five young high BMI and one older high BMI participant). Outliers (values > 3 *SDs* from group means) were removed ( $n = 3$ , 2 = older normal BMI, 1 = young high BMI). The IL-6 results are also published elsewhere [21] with slight deviations due to missing data and test-specific exclusion criteria.

To assess a relationship between low-grade inflammation and RMET performance initial correlation analysis was performed. Multiple linear regression analysis was conducted to assess whether a significant correlation between low-grade inflammation and RMET performance was independent of age, BMI, time of day, demographic-, lifestyle- and health-factors, using: 1) an unadjusted model with IL-6, and 2) an adjusted model correcting for age, BMI, demographic-, lifestyle- and health-variables (i.e., illness symptoms, smoking, alcohol intake, sleep quality, medication intake, sex, education level) previously shown to be associated with inflammation and/or cognitive function. The results of the multiple regression models are presented as standardized coefficient estimates ( $\beta$ ),  $t$ -values, and 95% confidence intervals. In the event of a significant relationship between inflammation and overall RMET performance, mediation analysis was conducted. Variables were Z-transformed before linear regression and mediation analysis was conducting.

To assess possible additive and/or interactive effects of age and BMI on performance on the RMET, the percentage of total correct responses was calculated and Age group (young, older) and BMI group (normal BMI, high BMI) were entered as between-subject factors in an analysis of variance (ANOVA). To assess the effect of emotional valence, percentage correct was calculated for each emotional valence (positive, neutral, and negative expressions) [42], which was entered in a mixed model ANOVA. Previous studies reported sex differences in emotion recognition [43]–[46] but because the study was not designed to assess sex-dependent effects, sex was included as a covariate rather than a between-subjects factor (see Supplementary Materials 7.1 for exploratory analysis including sex as a between-subjects factor).

Alpha values were set at .05 throughout. For all analyses where appropriate, Levene's test of Equality of Variances and Mauchly's Test of Sphericity were used to test for assumption violations; adjustments were made as needed using the Greenhouse-Geisser correction. Bonferroni corrections were applied to post-hoc pairwise comparisons (two-tailed unless stated otherwise) to control for Type I error rate. The PROCESS macro (Hayes, 2013) was used to test possible mediation effects of IL-6 (Model 4 with 5000 bootstrap samples). In addition to traditional null hypothesis significance testing, Bayes factors were calculated using Bayesian ANOVAs, t-tests, and correlational analyses using default prior probabilities (see [48]) for guidelines on the interpretation of Bayes factor). All statistical analyses were conducted using SPSS v.24.0 (IBM-SPSS Inc., Chicago, IL, USA) and JASP version 0.13.

## 5. RESULTS

### *5.1 BMI, age, and inflammation*

Table 1 presents summary statistics of the included participants. The four groups did not differ in sex composition, age (i.e., between the two BMI groups), or BMI (i.e., between age groups). The IL-6 analysis and an extended table of descriptive statistics are also published elsewhere [21]. ANOVAs showed higher IL-6 levels in the high versus normal BMI group ( $F(1, 78) = 31.30, p < .001, \eta_p^2 = .29, BF_{10} = 39511.14$ ), and higher IL-6 levels in the older versus young group ( $F(1, 78) = 3.62, p = .061, \eta_p^2 = .04, BF_{10} = 1.10$ ) although the latter was non-significant. There was no evidence for an age x BMI group interaction ( $F(1, 76) = 2.38, p = .127, \eta_p^2 = .03; BF_{10} = 1.33$ ).

**Table 1.** Descriptive statistics of participant characteristics. Numbers in parenthesis indicate SD; ♦ indicates a significant main effect of age group, • indicates a significant main effect of BMI group.

	Young Normal BMI	Young High BMI	Older Normal BMI	Older High BMI
N	20	20	21	26
Age (years)				
Mean ♦	25	28	72	70
Range	21 – 32	21 – 35	66 – 79	63 – 76
Sex ( <i>n</i> Female)	11	14	13	15
IL-6 (pg/ml)•	1.04 ± 0.44	2.40 ± 1.24	1.67 ± 1.36	2.35 ± 1.03
Range	0.34 – 2.10	1.12 – 5.61	0.43 – 6.54	1.09 – 5.99
Weight Status				
BMI (kg/m <sup>2</sup> )•	21.7 ± 2.5	33.1 ± 3.4	23.0 ± 1.7	32.5 ± 3.8
Body fat %				
Females•	27.9 ± 3.9	45.9 ± 4.7	33.3 ± 5.6	44.9 ± 4.8
Males•	15.7 ± 4.4	27.6 ± 3.8	22.4 ± 5.0	31.8 ± 4.9
Number of illness symptoms (SCQ)♦	0.95 ± 1.3	1.3 ± 1.9	3.0 ± 2.3	5.1 ± 3.2

## 5.2 Inflammation and emotion recognition

As can be seen in Figure 1, mean performance on the RMET (averaged across valence) was significantly negatively correlated with IL-6 ( $r(80) = -.279, p = .012, BF_{10} = 3.06$ ). Multiple regression analysis confirmed that IL-6 remained a significant predictor of mean RMET performance when adjusting for age and BMI group, time of day, and demographic-, lifestyle- and health-variables (adjusted model) (see Table 2).

Repeating the regression analysis with age and BMI as a continuous factor produced similar results (see Supplementary Materials Table S1). Inflammation had no mediating role in the relationship between inflammation and emotion recognition (results reported in Supplementary Materials 7.2).

**Table 2.** Multiple regression analysis ( $N = 87$ ) of the relationship between inflammation (IL-6) and overall emotion recognition (model 1) adjusted for influences of age, BMI, time of day, and health- and demographic variables (adjusted model). Age group: 1 = young, 2 = older; BMI group: 1 = low BMI, 2 = high BMI; Sex: 1 = Female; 2 = Male; Smoke: 0 = never, 1 = ex-smoker, 2 = smoker; Alcohol intake in units; Illness symptoms = number of symptoms; Sleep quality: higher score is indicative of lower quality of sleep; Medication intake = number of medications; \*\*\*  $p < .001$ , \*\*  $p < .01$ , \*  $p < .05$ .


Overall emotion recognition					
Model $R^2 = .08^*$					
Adjusted model $R^2 = .25^*$					
	$\beta$	$t$	95% CI		
			Lower	Upper	
Model 1					
IL-6	-.280*	-2.558	-.484	-.060	
Adjusted model					
IL-6	-.375**	-2.902	-.616	-.114	
Age group	.007	0.049	-.258	.271	
BMI group	.070	0.530	-.185	.319	
Sex	-.132	-1.089	-.353	.104	
Education	.087	0.715	-.159	.337	
Smoking	.142	1.164	-.095	.361	
Alcohol	-.102	-0.787	-.334	.145	
Illness symptoms	.435*	2.304	.060	.834	
Sleep quality	.045	0.383	-.189	.279	
Medication intake	-.201	-1.144	-.549	.149	
Time of day	-.034	-0.308	-.239	.175	

### 5.3 Age-dependent BMI effects on emotion recognition


As can be seen in Figure 2a, an age x BMI group interaction ( $F(1, 82) = 12.01$ ,  $p < .001$ ,  $\eta_p^2 = .13$ ,  $BF_{10} = 45.61$ ) showed that young individuals with a high BMI ( $M = 62.7\%$ ,  $SE = 2.6\%$ ) performed worse than their normal BMI counterparts ( $M = 73.3\%$ ,  $SE = 2.6\%$ ) (controlling for sex) ( $t(38) = 2.74$ ,  $p = .010$ ,  $d = 0.85$ ,  $BF_{10} = 5.27$ ). In contrast, in the older group, a main effect of BMI group indicated that individuals with a high BMI ( $M = 73.1\%$ ,  $SE = 2.3\%$ ) outperformed those with a normal BMI ( $M = 66.1\%$ ,  $SE = 2.6\%$ ) (controlling for sex)

( $t(45) = -2.07$ ,  $p = .044$ ,  $d = -0.60$ ,  $BF_{10} = 1.65$ ). Moreover, additionally adjusting for health symptoms and time of day (age x BMI interaction  $F(1, 80) = 9.26$ ,  $p = .003$ ,  $\eta_p^2 = .10$ ) produced similar results. Using age- and sex-adjusted body fat percentages (see [49]) instead of BMI to define weight-groups did not yield a different pattern of results. RMET accuracy of young individuals with a body fat percentage indicative of overweight groups performed 11.1% worse ( $SE = 3.8\%$ ) as compared to their leaner counterparts. In older individuals, those with overweight body fat percentages performed 2.5% better ( $SE = 3.8\%$ ) as compared to their leaner counterparts.

Stimuli selected on valence (negative, neutral, positive) showed that performance was best on positive expressions ( $M = 78.2\%$ ,  $SE = 1.9\%$ ), followed by neutral ( $M = 69.4\%$ ,  $SE = 1.6\%$ ) and negative expressions ( $M = 63.5\%$ ,  $SE = 1.9\%$ ), with significant differences between all three valence types ( $F(2, 166) = 24.92$ ,  $p < .001$ ,  $\eta_p^2 = .23$ ,  $BF_{10} = 3.735e+7$ ). As shown in Figure 2b, the data provided no evidence for an age or BMI group x valence interaction ( $F$ 's  $< 1$ ,  $BF_{10} = 0.11$  and  $BF_{10} = 0.14$ , respectively).



**Figure 1.** Scatterplot depicting individual accuracy scores and IL-6 levels. For each subgroup linear regression lines are fitted. Please note that the y-axis starts at chance level (i.e., 25% accuracy).



**Figure 2.** Individual (dots) and group mean (horizontal bars) accuracy scores for RMET total score (A), and group means for each emotional valence for each BMI and Age group (B); Errors bars indicate standard error of the mean. Please note that the y-axes start at 25% accuracy (i.e., chance level).

## 6. DISCUSSION

The present study examined the relationship between inflammation and emotion recognition. As expected, BMI was linked to elevated inflammatory activity, assessed as plasma IL-6. The expected positive association with age was non-significant. In line with earlier experimental research [2], [3], low-grade inflammation correlated with impaired performance on the RMET. This association withstood full adjustment for demographic-, lifestyle- and health-variables. The direct effects of BMI appeared non-linear however, whereby young individuals with a high BMI performed worse on the RMET than their normal BMI counterparts, whereas the reverse was observed for older adults. Analyses could not confirm a mediating role of IL-6 in these group differences, although this may reflect low statistical power.

The current results are consistent with the hypothesis that low-grade inflammation in a non-experimental setting may be a biobehavioral pathway linked to impaired emotion recognition, and further may be taken to suggest that ‘normal’ variations in inflammatory activity within relatively healthy populations may be associated with impaired emotion recognition. The fact that the prevalence of overweight and obesity has reached epidemic proportions worldwide [50], and the majority of these individuals likely have elevated levels of inflammatory activity, adds relevance to this notion. Being less sensitive to such social cues may have direct consequences for the dynamics of social interactions. The present findings are also consistent with preliminary evidence that high BMI exerts negative effects on emotion recognition, which some studies have already established in children and adolescents [51]–[53] but also [54].

Evidence of an ‘obesity paradox’ is mostly limited to studies using global measures of cognition, such as the Mini Mental State Examination Test, and focused on so-called ‘cold’ cognitions (e.g., memory, attention) [24]–[26], [55]. The current study is the first to extend these observations to cognitive processes that have social-affective components (i.e., so-called ‘hot’ cognitions).

At present, it remains unclear how or under what conditions older individuals may be spared for BMI-related cognitive deficits [56]. For that reason, some have attributed this

“obesity paradox” to methodological issues. Several methodological limitations may indeed apply to the current analyses also. It has been argued that BMI may not represent a reliable index of adiposity in older individuals, because in aging adipose tissue increase is often without weight gain due to a parallel decrease in lean body mass (e.g., muscle mass) [57]. However, when the current analyses replaced BMI by using age- and sex-adjusted body fat percentages to define weight-groups (see [49]), analyses did not yield a different pattern of results (see Results sections). Related to this are findings that patients with cardiovascular disease who have abdominal obesity in combination with a low BMI, are at higher risk of mortality, suggesting that the fat distribution plays an important role in BMI-health relationships [58], [59]. A second caveat is potential of selection bias; i.e., it cannot be excluded that the older high BMI group recruited in the present study may represent a healthier subsample of high BMI adults than in the general population, because heavier individuals may have experienced more overweight-related diseases that prevented them from taking part in research (i.e., those available for research are a relatively healthy subsample) [60]. However, no stringent health-related exclusion criteria were applied in the current study, and accordingly the older high BMI group indeed reported significantly higher levels of illness symptoms (see Table 1). Moreover, adjusting for illness symptoms did not alter the pattern of results. Thirdly, while there are links between obesity and depression in middle-aged and older adults [61], [62], there are also reports suggesting that depression in older age is associated with weight loss rather than with obesity [63], [64]. These data suggest that a lower BMI in older age may be a risk factor for disturbances in emotional processes. To reduce potential influences of age-related weight loss, stable body weight for at least six months prior to study enrollment was an inclusion criterion in the current study. However, whether weight loss may have occurred before this six month period cannot be ruled out. A further limitation is the moderate sample size and null-findings, including the lack of mediation, should thus be interpreted with some caution. Notwithstanding, to address these potential issues, future research should strive for larger and representative samples. Lastly, the assessment of IL-6 primarily acted as an inflammation check, i.e., a correlate/marker of inflammation, and no causal assumptions about the role of

383 IL-6 in the observed effects can be made. Another inflammatory factor or process  
384 mechanistically linked to inflammation may be the causal factor.

385 In summary, the present observational study aimed to replicate experimental human  
386 results showing that elevated low-grade inflammation is negatively related to emotion  
387 recognition. In young participants, higher BMI was associated with poorer emotion recognition  
388 whereas the opposite was observed for older participants. However, these relationships did  
389 not appear to be mediated by IL-6. Protracted low-grade inflammation, in otherwise  
390 predominantly healthy individuals, may thus present a biobehavioral pathway influencing  
391 emotion recognition across age and body weight categories. A possible protective effect of  
392 high-body weight in older individuals warrants further scrutiny.

393

## 7. SUPPLEMENTARY MATERIALS

**Table S1.** Multiple regression analysis for the relationship between IL-6 and RMET accuracy with age and BMI as continuous predictors instead of groups (see Table 2 for the same regression analysis with age and BMI groups). Statistical significance is indicated as follows; \*  $p < .05$ , \*\*  $p < .01$ , \*\*\*  $p < .001$ .

Model $R^2 = .05^*$					
Adjusted model $R^2 = .27^*$					
	$\beta$	t	95% CI		
			Lower	Upper	
Model 1					
IL-6	-.280*	-2.558	-.484	-.060	
Adjusted model					
IL-6	-.304*	-2.191	-.565	-.026	
Age	-.074	-0.531	-.334	.194	
BMI	-.053	-0.383	-.322	.218	
Sex	-.141	-1.153	-.364	.097	
Education	.068	0.566	-.176	.315	
Smoking	.142	1.142	-.099	.364	
Alcohol	-.084	-0.659	-.314	.158	
Illness symptoms	.492*	2.624	.121	.890	
Sleep quality	.043	-0.367	-.191	.278	
Medication intake	-.200	-1.141	-.549	.150	
Time of day	-.044	-0.407	-.248	.164	

### 7.1 Sex-dependent effects BMI on emotion recognition

In light of prior research showing sex effects on emotion recognition, exploratory analyses were performed, confirming a significant effect of sex (males:  $M = 64.8\%$ ,  $SE = 2.0\%$ ; females:  $M = 71.2\%$ ,  $SE = 1.6\%$ ) ( $F(1, 79) = 6.27$ ,  $p = .014$ ,  $\eta_p^2 = .06$ ;  $BF_{10} = 1.28$ ), and therefore analyses included sex as a covariate. A BMI group x Age group x sex interaction ( $F(1, 79) = 4.02$ ,  $p = .048$ ,  $\eta_p^2 = .04$ ) indicated that the reduced emotion recognition in young

individuals with a high BMI was driven by lower performance of males ( $M = 53.2\%$ ,  $SE = 5.6\%$ ) (Females:  $M = 68.4\%$ ,  $SE = 4.6\%$ ) ( $F(1, 36) = 5.14$ ,  $p = .029$ ,  $\eta_p^2 = .13$ ;  $BF_{10} = 3.08$ ). Sex had no significant effect in the older group ( $F(1, 43) = 3.12$ ,  $p = .085$ ,  $\eta_p^2 = .07$ ;  $BF_{10} = 0.81$ ) and also no interaction of sex x BMI group was evident in the older group ( $F < 1$ ;  $BF_{10} = 0.41$ ).

## 7.2 Mediation analysis results

A mediating variable is a variable that is part of the pathway by which an independent variable affects a dependent variable. The main requirement for mediation is that the *indirect effect* of the independent variable (e.g., BMI) through the mediator (e.g., IL-6) on the dependent variable (e.g., accuracy) is significant. Mediation indirect effects can be interpreted as the strength of the relationship between the independent variable (BMI) and dependent variable (accuracy) when accounting for the mediating pathway (IL-6) [65]. Variables were Z-transformed before analysis yielding standardized regression coefficients.

### 7.2.1 Reading the Mind in the Eyes Test (overall accuracy)

*Young age groups.* BMI in the young group (controlling for sex) was associated with higher IL-6 ( $\beta = .713$ , 95% CI = .480 .947,  $p < .001$ ) and lower RMET accuracy ( $\beta = -.416$ , 95% CI = -.723 -.109,  $p = .009$ ). IL-6 was not significantly associated with RMET accuracy when adjusting for BMI and sex ( $\beta = -.202$ , 95% CI = -.671, .266,  $p = .386$ ) and BMI was not a significant predictor of accuracy when adjusting for IL-6 and sex ( $\beta = -.272$ , 95% CI = -.726, .183,  $p = .232$ ). The indirect effect of IL-6 on the association between BMI and performance was not significant ( $\beta = -.144$ , 95% CI = -.454, .215), suggesting that IL-6 did not have a mediating role.

*Older age groups.* BMI in the older group (controlling for sex) was associated with higher IL-6 ( $\beta = .475$ , 95% CI = .190 .760,  $p = .002$ ). RMET accuracy was not significantly associated with BMI ( $\beta = -.136$ , 95% CI = -.169 -.442,  $p = .373$ ). IL-6 was not significantly associated with RMET accuracy when adjusting for BMI and sex ( $\beta = -.304$ , 95% CI = -.632, .025,  $p = .069$ ) and BMI was not a significant predictor of accuracy when adjusting for IL-6 and sex ( $\beta = .281$ , 95% CI = -.055, .616,  $p = .099$ ). The indirect effect of IL-6 on the association

441 between BMI and performance was not significant ( $\beta = -.144$ , 95% CI =  $-.345, .074$ ), suggesting  
442 that IL-6 did not have a mediating role.

443

## 8. ACKNOWLEDGEMENTS

The authors would like to thank Anne Clemens, Greta Ontrup, and Marina Wissink for their valuable contribution in recruiting and testing participants.

## 9. FUNDING AND DISCLOSURE

This research was funded by internal funds of the University of Amsterdam and University of Birmingham. The authors report no conflicts of interest, financial or otherwise.

## REFERENCES

- [1] N. I. Eisenberger, M. Moieni, T. K. Inagaki, K. A. Muscatell, and M. R. Irwin, "In Sickness and in Health: The Co-Regulation of Inflammation and Social Behavior," *Neuropsychopharmacology*, vol. 42, no. 1. pp. 242–253, 2017.
- [2] L. J. T. Balter *et al.*, "Low-grade inflammation decreases emotion recognition – Evidence from the vaccination model of inflammation," *Brain, Behavior, and Immunity*, 2018.
- [3] M. Moieni, M. R. Irwin, I. Jevtic, E. C. Breen, and N. I. Eisenberger, "Inflammation impairs social cognitive processing: A randomized controlled trial of endotoxin.," *Brain. Behav. Immun.*, vol. 48, pp. 132–138, 2015.
- [4] E. Bora and M. Berk, "Theory of mind in major depressive disorder: A meta-analysis," *Journal of Affective Disorders*, vol. 191. pp. 49–55, 2016.
- [5] M. N. Dalili, I. S. Penton-Voak, C. J. Harmer, and M. R. Munafò, "Meta-analysis of emotion recognition deficits in major depressive disorder," *Psychological Medicine*. 2015.
- [6] R. M. Hirschfeld *et al.*, "Social functioning in depression: a review.," *J. Clin. Psychiatry*, vol. 61, no. 4, pp. 268–75, 2000.
- [7] L. Capuron, S. Geisler, K. Kurz, F. Leblhuber, B. Sperner-Unterweger, and D. Fuchs, "Activated Immune System and Inflammation in Healthy Ageing: Relevance for Tryptophan and Neopterin Metabolism.," *Curr. Pharm. Des.*, p. 24641220, 2014.
- [8] R. C. Shelton and A. H. Miller, "Eating ourselves to death (and despair): The contribution of adiposity and inflammation to depression," *Progress in Neurobiology*. 2010.
- [9] R. Cancellato and K. Clément, "Is obesity an inflammatory illness? Role of low-grade inflammation and macrophage infiltration in human white adipose tissue," *BJOG: An International Journal of Obstetrics and Gynaecology*, vol. 113, no. 10. pp. 1141–1147, 2006.
- [10] H. Y. Chung *et al.*, "Molecular inflammation: Underpinnings of aging and age-related diseases," *Ageing Research Reviews*. 2009.
- [11] C. Franceschi *et al.*, "Inflammaging and anti-inflammaging: A systemic perspective on aging and longevity emerged from studies in humans," *Mech. Ageing Dev.*, 2007.
- [12] D. Baylis, D. B. Bartlett, H. P. Patel, and H. C. Roberts, "Understanding how we age: insights into inflammaging," *Longev. Heal.*, 2013.
- [13] G. Coppin, S. Nolan-Poupart, M. Jones-Gotman, and D. M. Small, "Working memory and reward association learning impairments in obesity.," *Neuropsychologia*, vol. 65, pp. 146–55, 2014.
- [14] A. B. Fagundo *et al.*, "Executive functions profile in extreme eating/weight conditions: From Anorexia Nervosa to Obesity," *PLoS One*, 2012.
- [15] E. Glisky, "Changes in Cognitive Function in Human Aging," in *Brain aging: Models, methods, and mechanisms*, 2007.
- [16] J. Gunstad, R. H. Paul, R. A. Cohen, D. F. Tate, and E. Gordon, "Obesity is associated with memory deficits in young and middle-aged adults," *Eat. Weight Disord.*, 2006.
- [17] S. Higgs and M. S. Spetter, "Cognitive Control of Eating: the Role of Memory in Appetite and Weight Gain," *Current obesity reports*. 2018.
- [18] C. Prickett, L. Brennan, and R. Stolwyk, "Examining the relationship between obesity and cognitive function: a systematic literature review.," *Obes. Res. Clin. Pract.*, vol. 9, no. 2, pp. 93–113, 2015.
- [19] J. Gunstad *et al.*, "C-reactive protein, but not homocysteine, is related to cognitive dysfunction

- in older adults with cardiovascular disease.," *J. Clin. Neurosci.*, vol. 13, no. 5, pp. 540–6, 2006.
- [20] J. Gunstad, R. H. Paul, R. A. Cohen, D. F. Tate, M. B. Spitznagel, and E. Gordon, "Elevated body mass index is associated with executive dysfunction in otherwise healthy adults," *Compr. Psychiatry*, vol. 48, no. 1, pp. 57–61, 2007.
- [21] L. J. T. Balter, S. Higgs, S. Aldred, J. A. Bosch, and J. E. Raymond, "Inflammation Mediates Body Weight and Ageing Effects on Psychomotor Slowing," *Sci. Rep.*, 2019.
- [22] K. M. Stanek *et al.*, "Body mass index and neurocognitive functioning across the adult lifespan," *Neuropsychology*, 2013.
- [23] J. Gunstad, A. Lhotsky, C. R. Wendell, L. Ferrucci, and A. B. Zonderman, "Longitudinal examination of obesity and cognitive function: Results from the baltimore longitudinal study of aging," *Neuroepidemiology*, vol. 34, no. 4, pp. 222–229, 2010.
- [24] H. K. Kuo *et al.*, "Cognitive function in normal-weight, overweight, and obese older adults: An analysis of the advanced cognitive training for independent and vital elderly cohort," *J. Am. Geriatr. Soc.*, vol. 54, no. 1, pp. 97–103, 2006.
- [25] H. M. Noh *et al.*, "Relationships between cognitive function and body composition among community-dwelling older adults: A cross-sectional study," *BMC Geriatr.*, 2017.
- [26] D. H. Yoon, S. H. Choi, J. H. Yu, J. H. Ha, S. H. Ryu, and D. H. Park, "The relationship between visceral adiposity and cognitive performance in older adults," *Age Ageing*, 2012.
- [27] C. K. Suemoto, P. Gilsanz, E. R. Mayeda, and M. M. Glymour, "Body mass index and cognitive function: The potential for reverse causation," *Int. J. Obes.*, 2015.
- [28] D. S. Knopman, S. D. Edland, R. H. Cha, R. C. Petersen, and W. A. Rocca, "Incident dementia in women is preceded by weight loss by at least a decade," *Neurology*, 2007.
- [29] T. Sobów, W. Fendler, and R. Magierski, "Body mass index and mild cognitive impairment-to-dementia progression in 24 months: A prospective study," *European Journal of Clinical Nutrition*. 2014.
- [30] E. Pedditizi, R. Peters, and N. Beckett, "The risk of overweight/obesity in mid-life and late life for the development of dementia: A systematic review and meta-analysis of longitudinal studies," *Age Ageing*, 2016.
- [31] T. Ruffman, J. D. Henry, V. Livingstone, and L. H. Phillips, "A meta-analytic review of emotion recognition and aging: Implications for neuropsychological models of aging," *Neuroscience and Biobehavioral Reviews*. 2008.
- [32] S. Sullivan and T. Ruffman, "Emotion recognition deficits in the elderly," *Int. J. Neurosci.*, 2004.
- [33] J. Fernandes, F. Ferreira-Santos, K. Miller, and S. Torres, "Emotional processing in obesity: a systematic review and exploratory meta-analysis," *Obes. Rev.*, vol. 19, no. 1, pp. 111–120, 2018.
- [34] S. Bergmann *et al.*, "Emotional availability, understanding emotions, and recognition of facial emotions in obese mothers with young children," *J. Psychosom. Res.*, 2016.
- [35] D. Rommel, J. L. Nandrino, C. Ducro, S. Andrieux, F. Delecourt, and P. Antoine, "Impact of emotional awareness and parental bonding on emotional eating in obese women," *Appetite*, 2012.
- [36] G. Nilsson, M. Lekander, T. Åkerstedt, J. Axelsson, and M. Ingre, "Diurnal variation of circulating interleukin-6 in humans: A meta-analysis," *PLoS One*, vol. 11, no. 11, 2016.
- [37] M. Herieka and C. Erridge, "High-fat meal induced postprandial inflammation," *Molecular Nutrition and Food Research*, vol. 58, no. 1, pp. 136–146, 2014.
- [38] S. Baron-Cohen, S. Wheelwright, J. Hill, Y. Raste, and I. Plumb, "The 'Reading the Mind in the Eyes' Test revised version: a study with normal adults, and adults with Asperger syndrome or high-functioning autism," *J. Child Psychol. Psychiatry*, vol. 42, no. 2, pp. 241–251, 2001.
- [39] B. F. M. Oakley, R. Brewer, G. Bird, and C. Catmur, "Theory of mind is not theory of emotion: A cautionary note on the reading the mind in the eyes test," *J. Abnorm. Psychol.*, 2016.
- [40] O. Sangha, G. Stucki, M. H. Liang, A. H. Fossel, and J. N. Katz, "The self-administered comorbidity questionnaire: A new method to assess comorbidity for clinical and health services research," *Arthritis Rheum.*, vol. 49, no. 2, pp. 156–163, 2003.
- [41] J. S. Carpenter and M. A. Andrykowski, "Psychometric evaluation of the Pittsburgh Sleep Quality Index," *J. Psychosom. Res.*, vol. 45, no. 1, pp. 5–13, 1998.
- [42] P. Maurage *et al.*, "The 'Reading the Mind in the Eyes' test as a new way to explore complex emotions decoding in alcohol dependence," *Psychiatry Res.*, vol. 190, no. 2–3, pp. 375–378, 2011.
- [43] R. Saylik, E. Raman, and A. J. Szameitat, "Sex differences in emotion recognition and working memory tasks," *Front. Psychol.*, 2018.
- [44] T. S. H. Wingenbach, C. Ashwin, and M. Brosnan, "Sex differences in facial emotion

- recognition across varying expression intensity levels from videos," *PLoS One*, 2018.
- [45] K. L. Evans and E. Hampson, "Sex-dependent effects on tasks assessing reinforcement learning and interference inhibition," *Front. Psychol.*, 2015.
- [46] Z. Zhang, K. F. Manson, D. Schiller, and I. Levy, "Impaired associative learning with food rewards in obese women," *Curr. Biol.*, vol. 24, no. 15, pp. 1731–1736, 2014.
- [47] A. Hayes, "Introduction to mediation, moderation, and conditional process analysis," *New York, NY Guilford*, pp. 3–4, 2013.
- [48] E.-J. Wagenmakers *et al.*, "Bayesian inference for psychology. Part II: Example applications with JASP," *Psychon. Bull. Rev.*, 2017.
- [49] J. A. Pasco, K. L. Holloway, A. G. Dobbins, M. A. Kotowicz, L. J. Williams, and S. L. Brennan, "Body mass index and measures of body fat for defining obesity and underweight: A cross-sectional, population-based study," *BMC Obes.*, 2014.
- [50] WHO, "Obesity and overweight Key facts," *Fact sheet No. 311*, 2015. .
- [51] X. Caldú *et al.*, "Effect of the catechol-O-methyltransferase Val 158Met polymorphism on theory of mind in obesity," *Eur. Eat. Disord. Rev.*, 2019.
- [52] A. Koch and O. Pollatos, "Reduced facial emotion recognition in overweight and obese children," *J. Psychosom. Res.*, 2015.
- [53] I. Percinell, B. Ozbaran, S. Kose, D. G. Simsek, and S. Darcan, "Increased deficits in emotion recognition and regulation in children and adolescents with exogenous obesity," *World J. Biol. Psychiatry*, 2018.
- [54] P. Surcinelli, B. Baldaro, A. Balsamo, R. Bolzani, M. Gennari, and N. C. F. Rossi, "Emotion recognition and expression in young obese participants: preliminary study," *Percept. Mot. Skills*, 2007.
- [55] E. Smith, P. Hay, L. Campbell, and J. N. Trollor, "A review of the association between obesity and cognitive function across the lifespan: Implications for novel approaches to prevention and treatment," *Obes. Rev.*, vol. 12, no. 9, pp. 740–755, 2011.
- [56] J. C. D. Nguyen, A. S. Killcross, and T. A. Jenkins, "Obesity and cognitive decline: Role of inflammation and vascular changes," *Front. Neurosci.*, 2014.
- [57] A. S. Jackson, I. Janssen, X. Sui, T. S. Church, and S. N. Blair, "Longitudinal changes in body composition associated with healthy ageing: men, aged 20-96 years.," *Br. J. Nutr.*, 2012.
- [58] M. Zeller *et al.*, "Relation between body mass index, waist circumference, and death after acute myocardial infarction," *Circulation*, 2008.
- [59] M. B. Kadakia, C. S. Fox, B. M. Scirica, S. A. Murphy, M. P. Bonaca, and D. A. Morrow, "Central obesity and cardiovascular outcomes in patients with acute coronary syndrome: Observations from the MERLIN-TIMI 36 trial," *Heart*, 2011.
- [60] H. R. Banack and J. S. Kaufman, "The 'Obesity Paradox' Explained," *Epidemiology*, 2013.
- [61] L. de Wit, F. Luppino, A. van Straten, B. Penninx, F. Zitman, and P. Cuijpers, "Depression and obesity: A meta-analysis of community-based studies," *Psychiatry Res.*, 2010.
- [62] F. S. Luppino *et al.*, "Overweight, obesity, and depression: A systematic review and meta-analysis of longitudinal studies," *Archives of General Psychiatry*, vol. 67, no. 3. pp. 220–229, 2010.
- [63] V. L. Forman-Hoffman, J. W. Yankey, S. L. Hillis, R. B. Wallace, and F. D. Wolinsky, "Weight and depressive symptoms in older adults: Direction of influence?," *Journals Gerontol. - Ser. B Psychol. Sci. Soc. Sci.*, 2007.
- [64] J. Kim, J. W. Noh, J. Park, and Y. D. Kwon, "Body mass index and depressive symptoms in older adults: A cross-lagged panel analysis," *PLoS One*, 2014.
- [65] A. F. Hayes, "Beyond Baron and Kenny: Statistical mediation analysis in the new millennium," *Commun. Monogr.*, 2009.