

New results on quasi-subfield polynomials

Euler, Marie; Petit, Christophe

DOI:

[10.1016/j.ffa.2021.101881](https://doi.org/10.1016/j.ffa.2021.101881)

License:

Creative Commons: Attribution-NonCommercial-NoDerivs (CC BY-NC-ND)

Document Version

Peer reviewed version

Citation for published version (Harvard):

Euler, M & Petit, C 2021, 'New results on quasi-subfield polynomials', *Finite Fields and Their Applications*, vol. 75, 101881. <https://doi.org/10.1016/j.ffa.2021.101881>

[Link to publication on Research at Birmingham portal](#)

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

New Results on Quasi-Subfield Polynomials

Marie Euler^{a,1,*}, Christophe Petit^{b,c,1}

^a*DGA MI, France*

^b*Université libre de Bruxelles, Département d'informatique, Belgium*

^c*University of Birmingham, School of Computer Science, United Kingdom*

Abstract

¹ Quasi-subfield polynomials were introduced by Huang *et al.* together with a new algorithm to solve the Elliptic Curve Discrete Logarithm Problem (ECDLP) over finite fields of small characteristic. In this paper we provide both new quasi-subfield polynomial families and a new theorem limiting their existence. Our results do not allow to derive any speedup for the new ECDLP algorithm compared to previous approaches.

Keywords: linearized polynomials, cryptography, elliptic curve discrete logarithm problem,

2000 MSC: 11T06,11T71,94A60

1. Introduction

Let p be a prime and let n, n' be two positive integers. For any prime power q , let \mathbb{F}_q be the finite field with q elements. When n' divides n , the finite field $\mathbb{F}_{p^{n'}}$ is a subfield of \mathbb{F}_{p^n} . The polynomial

$$X^{p^{n'}} - X$$

splits over \mathbb{F}_{p^n} and its roots are exactly all the elements of $\mathbb{F}_{p^{n'}}$. Quasi-subfield polynomials, introduced by Huang *et al.* [1], naturally generalize this polynomial.

Definition 1 (informal). *A quasi-subfield polynomial (QSP) is a polynomial of the form*

$$L(X) := X^{p^{n'}} - \lambda(X)$$

*Corresponding author

Email addresses: marie.euler@hotmail.com (Marie Euler),
christophe.f.petit@gmail.com (Christophe Petit)

¹Part of this work was done in fulfillment of a master thesis requirement of the first author at the University of Oxford

²Supported in part by an EPSRC grant EP/S01361X/1.

¹Declarations of interest: none

such that “most” of its roots are distinct and defined over \mathbb{F}_{p^n} , and moreover $d := \deg \lambda$ is “small”.

When n' does not divide n , the degree of λ cannot be too small, as shown in the following lemma.

Lemma 1. [1, Lemma 4.1] *Let $L(X) = X^{p^{n'}} - \lambda(X) \in \mathbb{F}_{p^n}[X]$ completely splitting over \mathbb{F}_{p^n} , such that $\ell := \log_p \deg \lambda > 0$. Then we have*

$$\left\lfloor \frac{n}{n'} \right\rfloor \ell + (n \bmod n') \geq n'.$$

A similar result in the non split case is also provided in [1, Lemma C.2].

In light of this lemma, it is useful to associate to any QSP a “quality” parameter $\beta := \ell n/n'^2$, where $\ell := \log_p \deg \lambda$ as above. This leads to a more formal definition of QSP :

Definition 2. *A quasi-subfield polynomial (QSP) is a polynomial of the form*

$$L(X) := X^{p^{n'}} - \lambda(X) \in \mathbb{F}_{p^n}[X]$$

which splits completely (or at least has approximately $p^{n'}$ roots), for which moreover $\beta(L) := \frac{n\ell}{n'^2} \leq 1$ with $\ell := \log_p \deg \lambda$.

In their paper, Huang *et al.* provide a QSP family over $\mathbb{F}_{p^n}[x]$ with $n = p_{a+1}$ and $\beta = 1 - \frac{1}{p_a}(1 - 1/p^r + 1/(p^r p_a))$ with $p_a = 1 + p^r + p^{2r} + \dots + p^{ar}$. Most importantly, they show how quasi-subfield polynomials can be used to solve the Elliptic Curve Discrete Logarithm Problem (ECDLP), a problem of major importance for cryptography.

Our results. We expand the study of quasi-subfield polynomials initiated in [1], focusing on polynomials whose roots form a subgroup of either the additive or the multiplicative group of finite fields. In the additive case this amounts to searching for a *linearized polynomial*

$$L(X) = X^{p^{n'}} - (a_\ell X^{p^\ell} + a_{\ell-1} X^{p^{\ell-1}} + \dots + a_0 X)$$

splitting over \mathbb{F}_{p^n} , with “small” ℓ . In the multiplicative case this amounts to searching for a polynomial of the form

$$L(X) = X^{p^{n'}} - X^d$$

where $p^{n'} - d$ divides $p^n - 1$, and d is small.

Our main result is the following theorem on *linearized* QSPs.

Theorem 1. *Let $L(X) := X^{p^{n'}} - (a_\ell X^{p^\ell} + a_{\ell-1} X^{p^{\ell-1}} + \dots + a_0 X)$ be a linearized polynomial over \mathbb{F}_{p^n} , with $\ell \geq 1$. If L splits completely over \mathbb{F}_{p^n} then $\beta := \ell n/n'^2 \geq 3/4$.*

The equality is obtained for example with $X^{p^2} + X^p + X$ in $\mathbb{F}_{p^3}[X]$. The theorem improves on Lemma 1 for linearized polynomials with parameters n, n' satisfying $(n \bmod n') \geq n'/4$. In the special case of trinomials, a similar result was independently obtained with similar techniques by McGuire and Mueller [2].

We also introduce methods to generate new families of linearized QSP from known ones, and we exhibit new additive and multiplicative QSP families.

Finally, we apply our results to Huang *et al.*'s ECDLP algorithm.

Impact on ECDLP security. The complexity of Huang *et al.*'s ECDLP algorithm crucially relies on the quality parameter β of the quasi-subfield polynomial used. Based on their complexity estimations, a value of β at most 0.1 would be needed to obtain a complexity improvement over generic, state-of-the-art ECDLP algorithms. In contrast, Theorem 1 rules out any β smaller than $3/4$ in the case of linearized polynomials, and the best quasi-subfield polynomial we found so far has $\beta \simeq 0.7$. Further work will be needed to improve Huang *et al.*'s approach with new ideas and better QSP families, or to provide a definite proof that it will not improve on generic algorithms.

Outline. The remaining of this paper is organized as follows. Section 2 recalls our new lower bound for the β value of linearized QSPs and it provides its proof. Section 3 includes our new QSP families. Section 4 discusses the impact of our results on ECDLP and Section 5 concludes the paper.

2. A new lower bound on β for linearized quasi-subfield polynomials

In this section we first recall known properties of linearized polynomials, including a characterization of linearized polynomials that split completely over their field of definition. We then proceed to prove Theorem 1, and we compare the bound it provides with the bounds given in [1] and [2].

2.1. Linearized polynomials

Let p be a prime and n be positive integer. Let \mathbb{F}_{p^n} be the finite field with p^n elements. We write $\text{Gal}(\mathbb{F}_{p^n}/\mathbb{F}_p)$ for the Galois group of \mathbb{F}_{p^n} with respect to \mathbb{F}_p . For any automorphism $\sigma \in \text{Gal}(\mathbb{F}_{p^n}/\mathbb{F}_p)$, there exists $s \in \mathbb{Z}$ with $\gcd(s, n) = 1$ such that $\sigma(X) = X^{p^s}$. In the following we write X^σ for $\sigma(X)$.

Definition 3 (Linearized polynomials). *Let $\sigma \in \text{Gal}(\mathbb{F}_{p^n}/\mathbb{F}_p)$ and let $f = X^d + a_{d-1}X^{d-1} + \dots + a_1X + a_0 \in \mathbb{F}_{p^n}[X]$. The linearized polynomial related to f and σ is the polynomial*

$$L_{f,\sigma} = X^{\sigma^d} + a_{d-1}x^{\sigma^{d-1}} + \dots + a_1X^\sigma + a_0X \in \mathbb{F}_{p^n}[X].$$

Moreover d is called the σ -degree of $L_{f,\sigma}$.

Let $L := L_{f,\sigma} \in \mathbb{F}_{p^n}[X]$ be a linearized polynomial with coefficients in \mathbb{F}_{p^n} as above. Let C_L be the companion matrix of f , namely

$$C_L := \begin{bmatrix} 0 & 0 & \cdots & 0 & -a_0 \\ 1 & 0 & \cdots & 0 & -a_1 \\ 0 & 1 & \cdots & 0 & -a_2 \\ \vdots & \vdots & \ddots & \vdots & \\ 0 & 0 & \cdots & 1 & -a_{d-1} \end{bmatrix}.$$

We also define the matrix

$$A_L := C_L \cdot C_L^\sigma \cdot C_L^{\sigma^2} \cdots C_L^{\sigma^{n-1}},$$

where C_L^σ is the matrix obtained by applying σ coefficient-wise on C_L . Note that A_L and C_L are square matrices of dimension d .

The following result (independently due to McGuire and Sheekey [3] and Csajbók *et al.* [4]) characterizes linearized polynomials that split completely.

Proposition 1. *Let $L = X^{\sigma^d} + a_{d-1}x^{\sigma^{d-1}} + \dots + a_1X^\sigma + a_0X$ be a linearized polynomial with coefficients in \mathbb{F}_{p^n} . Then L has p^{n_1} roots defined over \mathbb{F}_{p^n} , where n_1 is the dimension of the eigenspace of A_L with eigenvalue 1. In particular, L splits completely over $\mathbb{F}_{p^n}[X]$ if and only if A_L is the identity matrix.*

A corollary of this proposition is that the maximum number of roots of L is p^d , and $L_{f,\sigma}$ splits completely in \mathbb{F}_{p^n} only if $\sigma(X) = X^p$. From now on, we will therefore only consider the case $\sigma(X) = X^p$ and thus write L_f instead of $L_{f,\sigma}$.

Another important property of completely splitting linearized polynomials is the following one.

Proposition 2. *Let $f = a_0 + a_1X + \dots + X^d \in \mathbb{F}_p[X]$. Then the following properties are equivalent.*

1. $L_f(X)$ splits completely over $\mathbb{F}_{p^n}[X]$
2. $L_f(X)$ divides $X^{p^n} - X$
3. f divides $X^n - 1$

This proposition directly follows from the fact that L_f divides L_g for the composition of polynomials if and only if $f|g$ for the multiplication of polynomials [5, Chapter 11].

2.2. Proof of Theorem 1

We will now prove Theorem 1, namely that for any linearized polynomial $L = X^{p^{\ell}} - (a_\ell X^{p^\ell} + a_{\ell-1}X^{p^{\ell-1}} + \dots + a_0X)$ with $\ell \geq 1$, $a_\ell \neq 0$ and $\forall i, a_i \in \mathbb{F}_{p^n}$, if L splits completely over \mathbb{F}_{p^n} then $\beta := \ell n / n^2 \geq 3/4$.

This result is in fact a consequence of the following lemma which highlights that the field has to be big enough to have completely splitting sparse linearized polynomials in it.

Lemma 2 (Lower bound on n). *Let $L_f = X^{p^{n'}} - (a_\ell X^{p^\ell} + a_{\ell-1} X^{p^{\ell-1}} + \dots + a_0 X)$ with $\ell \geq 1$, $a_\ell \neq 0$ and $\forall i, a_i \in \mathbb{F}_{p^n}$. If L_f splits completely over \mathbb{F}_{p^n} then*

$$n \geq n' + (n' - \ell) \left\lfloor \frac{n' - 1}{\ell} \right\rfloor$$

Let us first observe that this result is indeed enough to prove the theorem. Indeed one can notice that since n' and ℓ are integers, we have $\left\lfloor \frac{n' - 1}{\ell} \right\rfloor \geq \frac{n'}{\ell} - 1$. Therefore, by Lemma 2 we have $n \geq n' + (n' - \ell) \frac{n'}{\ell} - (n' - \ell) \geq \frac{n'^2}{\ell} - n' + \ell$. Thus, $\beta = \frac{\ell n}{n'^2} \geq 1 - \frac{\ell}{n'} + \frac{\ell^2}{n'^2} = 1 - \frac{\ell}{n'} \left(1 - \frac{\ell}{n'}\right) \geq 1 - 1/4 = 3/4$.

We will now show that Lemma 2 boils down to the proof of a result about the power of a matrix. By Proposition 1, $L := L_f$ splits completely over \mathbb{F}_{p^n} if and only if $A_L := C_L \cdot C_L^\sigma \cdots C_L^{\sigma^{n-1}} = I$ where C_L is the companion matrix of f .

Therefore we have to prove that

$$\text{If } n < n' + (n' - \ell) \left\lfloor \frac{n' - 1}{\ell} \right\rfloor, \text{ then } A_L \neq I$$

The remainder of the proof will consider matrices defined over the polynomial ring $\mathbb{F}_{p^n}[a_0, \dots, a_\ell]$, where a_i are the coefficients of L as above. However as our result only depends on the value of n' and ℓ but not on the specific coefficients a_i , we represent any element $x \in \mathbb{F}_{p^n}[a_i]$ by a symbol $\tilde{x} \in \{0, 1, a^\bullet, \otimes\}$, noting it $x \rightsquigarrow \tilde{x}$, according to the following rules :

- if $x \rightsquigarrow 0$ then $x = 0$;
- if $x \rightsquigarrow 1$ then $x = 1$;
- if $x \rightsquigarrow a^\bullet$ then x is a power of $(-a_\ell)$ and thus $x \neq 0$;
- ($x \rightsquigarrow \otimes$ implies no condition on x).

For instance $-a_0 \rightsquigarrow \otimes$, $-a_\ell \rightsquigarrow a^\bullet$, $a_0 + a_\ell \rightsquigarrow \otimes$, and also $-a_\ell \rightsquigarrow \otimes$. One can notice that if $x_1 \rightsquigarrow \tilde{x}_1$ and $x_2 \rightsquigarrow \tilde{x}_2$, then $x_1 + x_2 \rightsquigarrow z_1$ and $x_1 \cdot x_2 \rightsquigarrow z_2$ with z_1 and z_2 given by the following tables.

+	0	1	a^\bullet	\otimes
0	0	1	a^\bullet	\otimes
1	1	\otimes	\otimes	\otimes
a^\bullet	a^\bullet	\otimes	\otimes	\otimes
\otimes	\otimes	\otimes	\otimes	\otimes

\cdot	0	1	a^\bullet	\otimes
0	0	0	0	0
1	0	1	a^\bullet	\otimes
a^\bullet	0	a^\bullet	a^\bullet	\otimes
\otimes	0	\otimes	\otimes	\otimes

For instance, if $x_1 \rightsquigarrow a^\bullet$, $x_2 \rightsquigarrow a^\bullet$, $x_3 \rightsquigarrow \otimes$, then $x_1 + x_2 \rightsquigarrow \otimes$, $x_1 \cdot x_2 \rightsquigarrow a^\bullet$ and $x_1 \cdot x_3 \rightsquigarrow \otimes$.

We extend this notation to matrices over $\mathbb{F}_{p^n}[a_i]$: Let $M = (m_{i,j})$ be a square matrix of dimension n' with coefficients in \mathbb{F}_{p^n} . If for all i and j , $m_{i,j} \rightsquigarrow \tilde{m}_{i,j}$,

we denote $M \rightsquigarrow \tilde{M} := (\tilde{m}_{i,j})$. We observe that if $M \rightsquigarrow \tilde{M}$ and $N \rightsquigarrow \tilde{N}$ then $M + N \rightsquigarrow \tilde{M} + \tilde{N}$ and $MN \rightsquigarrow \tilde{M}\tilde{N}$ where the operations are done according to the above tables.

Let M and \tilde{M} such that $M \rightsquigarrow \tilde{M}$. If there is a zero on the main diagonal of \tilde{M} or there is a non zero coefficient (1 or a^\bullet) outside of this diagonal, then we see that \tilde{M} cannot represent the identity matrix (denoted $I \not\rightsquigarrow \tilde{M}$) and then $M \neq I$.

Therefore we will give \tilde{A}_L such that $A_L \rightsquigarrow \tilde{A}_L$ and prove that if $n < n' + (n' - \ell) \lfloor \frac{n' - 1}{\ell} \rfloor$ then $I \not\rightsquigarrow \tilde{A}_L$.

First of all, we can recall that $A_L := C_L \cdot C_L^\sigma \cdot \dots \cdot C_L^{\sigma^{n-1}}$ with

$$C_L = \begin{bmatrix} 0 & 0 & \dots & 0 & \dots & \dots & -a_0 \\ 1 & 0 & \dots & 0 & \dots & \dots & -a_1 \\ 0 & 1 & \dots & 0 & \dots & \dots & -a_2 \\ \vdots & \vdots & \ddots & & \ddots & & \vdots \\ 0 & 0 & \dots & 1 & \dots & \dots & -a_\ell \\ \vdots & \vdots & & \vdots & \ddots & & 0 \\ \vdots & \vdots & & \vdots & & \ddots & \vdots \\ 0 & 0 & \dots & & \dots & 1 & 0 \end{bmatrix}$$

Moreover, since σ acts on matrices coefficient-wise, we can observe that for all $k \geq 0$, $C_L^{\sigma^k} \rightsquigarrow M$ with

$$M := \begin{bmatrix} 0 & 0 & \dots & 0 & \dots & \dots & \circledast \\ 1 & 0 & \dots & 0 & \dots & \dots & \circledast \\ 0 & 1 & \dots & 0 & \dots & \dots & \circledast \\ \vdots & \vdots & \ddots & & \ddots & & \vdots \\ 0 & 0 & \dots & 1 & \dots & \dots & a^\bullet \\ \vdots & \vdots & & \vdots & \ddots & & 0 \\ \vdots & \vdots & & \vdots & & \ddots & \vdots \\ 0 & 0 & \dots & & \dots & 1 & 0 \end{bmatrix}$$

Therefore $A_L \rightsquigarrow M^n$. Our goal is then to study the powers of M , which is a companion matrix defined on $\{0, 1, a^\bullet, \circledast\}$ and to prove the following result.

Lemma 3 (Small powers are not the identity).

$$\text{If } n < n' + (n' - \ell) \lfloor \frac{n' - 1}{\ell} \rfloor, \text{ then } I \not\rightsquigarrow M^n.$$

More precisely, we will prove the following lemma which exhibits a non zero coefficients outside of the main diagonal. We will note $M_{i,j}^n$ the coefficient in the i -th row, j -th column of the matrix M^n .

Lemma 4 (A non zero coefficient). *The following two claims are true:*

- If $n < n'$, then $M_{n+1,1}^n = 1$.
- If $n' \leq n < n' + (n' - \ell) \left\lfloor \frac{n' - 1}{\ell} \right\rfloor$, then we have $M_{i_n,1}^n = a^\bullet \neq 0$
with $i_n = n - (n' - \ell) \left\lfloor \frac{n - \ell}{n' - \ell} \right\rfloor + 1 \in [2, n']$

Proof. Adapting a result of Chen and Louck [6] (see Appendix A for details) on powers of companion matrices we have $M_{i,j}^n = 1$ if $n = i - j$ and otherwise

$$M_{i,j}^n = \sum_{\substack{\mathbf{k}=(k_1,\dots,k_{n'}) \\ \sum_{i=1}^{n'} ik_i=n-i+j}} w_{\mathbf{k}} \cdot 0^{k_1+\dots+k_{n'-\ell-1}} \cdot (a^\bullet)^{k_{n'-\ell}} \cdot \otimes^{k_{n'-\ell+1}+\dots+k_{n'}}$$

where $\mathbf{k} = (k_i)_{1 \leq i \leq n'}$ are non-negative integers and

$$w_{\mathbf{k}} = \frac{k_{n'-i+1} + \dots + k_{n'}}{k_1 + \dots + k_{n'}} \left(k_1 + \dots + k_{n'} \right)_{k_1, \dots, k_{n'}}.$$

The first part of Lemma 4 follows from the case $n = i - j$.

When $n \neq i - j$, the contribution of any \mathbf{k} with $k_1 + \dots + k_{n'-\ell-1} > 0$ to the sum is null. Therefore we get

$$M_{i,j}^n = \sum_{\substack{\mathbf{k}=(k_{n'-\ell},\dots,k_{n'}) \\ \sum_{i=n'-\ell}^{n'} ik_i=n-i+j}} w_{\mathbf{k}} \cdot a^\bullet \cdot \otimes^{k_{n'-\ell+1}+\dots+k_{n'}}, \quad (1)$$

where $w_{\mathbf{k}}$ is now

$$w_{\mathbf{k}} = \frac{k_{\max(n'-i+1, n'-\ell)} + \dots + k_{n'}}{k_{n'-\ell} + \dots + k_{n'}} \left(k_{n'-\ell} + \dots + k_{n'} \right)_{k_{n'-\ell}, \dots, k_{n'}}.$$

Note that here we do not want to remove the exponents over the unknown symbol. Indeed when $k_{n'-\ell+1} + \dots + k_{n'} = 0$, we have $\otimes^{k_{n'-\ell+1}+\dots+k_{n'}} = 1$; so we get a term $w_{\mathbf{k}} \cdot a^\bullet$ which in the case $w_{\mathbf{k}} = 1$ is exactly what we need to prove that this coefficient is a^\bullet .

Let $n' \leq n \leq n' + (n' - \ell) \left\lfloor \frac{n' - 1}{\ell} \right\rfloor - 1$, and let $i_n := n - (n' - \ell) \left\lfloor \frac{n - \ell}{n' - \ell} \right\rfloor + 1$. We have

$$\begin{aligned} n - (n' - \ell) \frac{n - \ell}{n' - \ell} + 1 &\leq i_n \leq n - (n' - \ell) \left(\frac{n - \ell}{n' - \ell} - \frac{n' - \ell - 1}{n' - \ell} \right) + 1 \\ n - (n - \ell) + 1 &\leq i_n \leq n - (n - n' + 1) + 1 \\ 2 &\leq \ell + 1 \leq i_n \leq n' \end{aligned}$$

In particular, $M_{i_n,1}^n$ is well-defined and it is not in the top left corner.

Moreover, $n' - i_n + 1 \leq n' - (\ell + 1) + 1 = n' - \ell$ so $\max(n' - i_n + 1, n' - \ell) = n' - \ell$. Also, $n - i_n + 1 = (n' - \ell) \lfloor \frac{n - \ell}{n' - \ell} \rfloor$. We then have :

$$\begin{aligned} w_{\mathbf{k}} &= \frac{k_{n'-\ell} + \cdots + k_{n'}}{k_{n'-\ell} + \cdots + k_{n'}} \binom{k_{n'-\ell} + \cdots + k_{n'}}{k_{n'-\ell}, \dots, k_{n'}} = \binom{k_{n'-\ell} + \cdots + k_{n'}}{k_{n'-\ell}, \dots, k_{n'}} \\ &= \frac{(k_{n'-\ell} + \cdots + k_{n'})!}{k_{n'-\ell}! \cdots k_{n'}!} \end{aligned}$$

and

$$M_{i_n, 1}^n = \sum_{\substack{k_{n'-\ell}, \dots, k_{n'} \\ \sum_{i=n'-\ell}^{n'} i k_i = (n' - \ell) \lfloor \frac{n - \ell}{n' - \ell} \rfloor}} w_{\mathbf{k}} \cdot a^\bullet \cdot \otimes^{k_{n'-\ell+1} + \cdots + k_{n'}}$$

In order to show that $M_{i_n, 1}^n = a^\bullet$, we now show that this sum in fact only involves one term, and that this term is exactly a^\bullet .

Clearly $\mathbf{k} = (k_{n'-\ell}, \dots, k_{n'}) = \left(\lfloor \frac{n - \ell}{n' - \ell} \rfloor, 0, \dots, 0 \right)$ is a solution to

$$\sum_{i=n'-\ell}^{n'} i k_i = (n' - \ell) \lfloor \frac{n - \ell}{n' - \ell} \rfloor \quad (2)$$

and for this \mathbf{k} we have $w_{\mathbf{k}} \cdot a^\bullet \cdot \otimes^{k_{n'-\ell+1} + \cdots + k_{n'}} = 1 \cdot a^\bullet \cdot 1 = a^\bullet$. It remains to show that it is the only valid \mathbf{k} to prove that $M_{i_n, 1}^n = a^\bullet$.

By contradiction, let us assume that there exists a solution $(k_{n'-\ell}, \dots, k_{n'})$ to Equation (2) such that

$$\sum_{i=n'-\ell+1}^{n'} k_i = \sum_{i=0}^{\ell-1} k_{n'-i} > 0 \quad (3)$$

We recall that by definition, all the k_i are non-negative. From Equation (2), we have :

$$\begin{aligned} (n' - \ell) \lfloor \frac{n - \ell}{n' - \ell} \rfloor &= \sum_{i=n'-\ell}^{n'} i k_i = \sum_{i=0}^{\ell} (n' - i) k_{n'-i} \\ &= (n' - \ell) \sum_{i=0}^{\ell} k_{n'-i} + \sum_{i=0}^{\ell-1} (\ell - i) k_{n'-i} \quad (4) \end{aligned}$$

$$\begin{aligned} &\geq (n' - \ell) \frac{\sum_{i=0}^{\ell} (\ell - i) k_{n'-i}}{\ell} + \sum_{i=0}^{\ell-1} (\ell - i) k_{n'-i} \\ &\geq \left(\frac{n' - \ell}{\ell} + 1 \right) \sum_{i=0}^{\ell-1} (\ell - i) k_{n'-i} = \frac{n'}{\ell} \sum_{i=0}^{\ell-1} (\ell - i) k_{n'-i} \quad (5) \end{aligned}$$

From Equation (4), we have $(n' - \ell) \mid \sum_{\iota=0}^{\ell-1} (\ell - \iota) k_{n'-\iota}$. We also have, by hypothesis (3), that $\sum_{\iota=0}^{\ell-1} (\ell - \iota) k_{n'-\iota} \geq \sum_{\iota=0}^{\ell-1} k_{n'-\iota} > 0$ hence $\sum_{\iota=0}^{\ell-1} (\ell - \iota) k_{n'-\iota} \geq (n' - \ell)$.

Together with Equation (5) this implies $(n' - \ell) \lfloor \frac{n - \ell}{n' - \ell} \rfloor \geq \frac{n'}{\ell} (n' - \ell)$ so

$$\left\lfloor \frac{n - \ell}{n' - \ell} \right\rfloor \geq \frac{n'}{\ell}. \quad (6)$$

On the other hand, thanks to the condition on n being not too big, we have

$$\frac{n - \ell}{n' - \ell} \leq \frac{n' + (n' - \ell) \left\lfloor \frac{n' - 1}{\ell} \right\rfloor - 1 - \ell}{n' - \ell} \leq \left\lfloor \frac{n' - 1}{\ell} \right\rfloor + \left(1 - \frac{1}{n' - \ell}\right).$$

This implies $\left\lfloor \frac{n - \ell}{n' - \ell} \right\rfloor \leq \left\lfloor \frac{n' - 1}{\ell} \right\rfloor < \frac{n'}{\ell}$, contradicting Equation (6). We deduce that $\mathbf{k} = \left(\left\lfloor \frac{n - \ell}{n' - \ell} \right\rfloor, 0, \dots, 0\right)$ is the only solution to Equation (2) and thus

$$M_{i_n, 1}^n = a^\bullet \neq 0$$

□

2.3. Comparison with other lower bounds

Theorem 1 improves on the bound given in Lemma 4.1 of [1] whenever the QSP is linearized and $(n \bmod n') \geq n'/4$.

Our bound is similar to the one given by Daniela Mueller and Gary McGuire in [2], which was established during the completion of this article. Indeed, Theorem 1.1 in [2] shows (using also [3] and [4]) that a linearized trinomial $L = X^{q^d} - bX^q - aX \in \mathbb{F}_{p^n}$, with $b \neq 0$ and $q = p^k$ a power of p such that $n = k\tilde{n}$, splits completely only if

$$\tilde{n} \geq (d - 1)d + 1 = d^2 - d + 1.$$

Let us compare it with the bound given by Lemma 2. We write $n = k\tilde{n}$ so that $q = p^k$. Thus $L = X^{p^{k^d}} - bX^{p^k} - aX$. Lemma 2 gives:

$$k\tilde{n} = n \geq kd + (kd - k) \left\lfloor \frac{kd - 1}{k} \right\rfloor \geq k(d + (d - 1)(d - 1))$$

Thus, as Mueller and McGuire, we get $\tilde{n} \geq d + (d - 1)^2 = d^2 - d + 1$.

While their bound is less general than ours as it only applies to trinomials, they gave a more complete description of completely splitting linearized trinomials. Indeed, they also take into account the case $\ell = 0$ which we did not consider in this article, and they exhaustively describe all possible such polynomials when $\tilde{n} \leq (d - 1)d + 1$:

- either $\tilde{n} = id$ with $i \leq d - 1$, $b = 0$ and $a^{1+q^d+\dots+q^{(i-1)d}} = 1$,

Quasi-subfield polynomial	p	n	β
L_{f_a} with $f_0 = 1 + X^{q^{-1}} + \dots + X^{q^{d-1}}$, $f_a = a + X + X^q + \dots + X^{q^d}$ for any $a \neq 0 \in \mathbb{F}_q$ and q a power of p	any	$q^{d+1} - 1$	$1 - \frac{q^{d-1}}{(1+q+\dots+q^{d-1})^2}$
$L_{(X^{n-1})/f}$ with L_f a completely splitting QSP of degree n'	any	any integer	$1 - (\frac{n'}{n-n'})^2(1 - \beta(L_f))$
$X^{p^{n'}} - X^a$ with $n' = n - i$ and $a = p^{n'} \pmod{\frac{p^n-1}{p^{2i}-1}}$	any	$2ik,$ $k \geq 2$	$\leq 1 - \frac{1}{(2k-1)^2}$
$X^p - X^a$ with $k \geq 2$ and $a = p \pmod{\frac{p-k}{k-1}}$	$k^n + k - 1$	any integer	≤ 1
$X^{p^{n-1}} - X^a$ with $k \geq 2$ and $r = \frac{(p^n-1)(k-(-1)^n)}{(k^n-k)(k^n-(-1)^n)}$ $a = p^{n-1} \pmod r$	$k^n - k - (-1)^n$	$n > 2$ $k^n \gg 1$	≤ 1

Table 1: New families of quasi-subfield polynomials

- either $\tilde{n} = (d-1)d + 1$, $a^{1+q+\dots+q^{(d-1)d}} = (-1)^{d-1}$, $b = -a^{qe_1}$ where $e_1 = \sum_{i=0}^{d-1} q^{id}$ and $d-1$ is a power of p

See also [7] for the complete description of completely splitting linearized trinomials when $\tilde{n} \leq (d-1)d + d - 1$.

3. New families of quasi-subfield polynomials

In this section, we will prove that the additive polynomials of Proposition 5 and the multiplicative polynomials of Proposition 6 are quasi-subfield polynomials. We first provide general tools to deduce new linearized QSPs from known ones and we define equivalence classes among linearized quasi-subfield polynomials. We then successively focus on additive and multiplicative families, and we finish with the case where the extension degree is a Mersenne prime.

Since the case $\log_p \deg \lambda = 0$ corresponds to subfield polynomials, which are well-known, we will only consider the case $\log_p \deg \lambda > 0$.

3.1. Equivalent classes of linearized quasi-subfield polynomials

In order to simplify our search of quasi-subfield polynomials, we will first discuss transformations to deduce new linearized quasi-subfield polynomials from

known ones. We will introduce two types of transformations. The first transformation will change the value of β and thus potentially improve it. The second one will keep the same β , thus it will not produce more interesting linearized quasi-subfield polynomials, but it will allow us to group them by equivalence classes.

Both transformations will only concern completely splitting QSPs with coefficients in \mathbb{F}_p . We will write $\mathcal{Q}_{p,n}^L$ for the set of completely splitting linearized QSPs in $\mathbb{F}_{p^n}[X]$.

The first way to obtain a linearized quasi-subfield polynomial from another one is what we call the inversion process.

Proposition 3 (Inversion). *Let $f = X^{n'} + a_\ell X^\ell + \dots + a_0 \in \mathbb{F}_p[X]$ such that $L_f \in \mathcal{Q}_{p,n}^L$ and $n' < n$. Let $g = (X^n - 1)/f$. Then $L_g \in \mathcal{Q}_{p,n}^L$. We say that L_g is the inverse of L_f .*

Proof. By Proposition 2, we know that $f|X^n - 1$ so g is well-defined.

Let us write g as $X^{n-n'} + b_r X^r + \dots + b_0$. We will prove that $r+n' = n-n'+\ell$. Indeed, $X^n - 1 = f \cdot g = (X^{n'} + a_\ell X^\ell + \dots + a_0)(X^{n-n'} + b_r X^r + \dots + b_0) = X^n + b_r X^{r+n'} + a_\ell X^{\ell+n-n'} + \dots + a_0 b_0$. So, if $r+n' > \ell+n-n'$ then the coefficient of $X^{r+n'}$ in $f \cdot g$ comes exclusively from $(b_r X^r) \cdot X^{n'}$ and is therefore non zero. Conversely, if $r+n' < \ell+n-n'$ then the coefficient of $X^{\ell+n-n'}$ in $f \cdot g$ comes exclusively from $(a_\ell X^\ell) \cdot X^{n-n'}$ and thus is not zero. We deduce

$$\begin{aligned} \beta(L_g) &= \frac{n \cdot r}{(n-n')^2} = \frac{n \cdot (n-2n'+\ell)}{(n-n')^2} = 1 - \frac{n'^2 - \ell \cdot n}{(n-n')^2} \\ &= 1 - \left(\frac{n'}{n-n'} \right)^2 (1 - \beta(L_f)) \leq 1. \end{aligned}$$

□

Our second family of transformations keep the value of β unchanged.

Proposition 4 (Transformations preserving β). *Let $k \geq 1$ and $\gamma \in \mathbb{F}_{p^n}^*$. Let $f = X^{n'} + a_\ell X^\ell + \dots + a_0 \in \mathbb{F}_p[X]$. Then the following properties are equivalent:*

- (a) $L_f \in \mathcal{Q}_{p,n}^L$,
- (b) $L_{f(X^k)} = X^{p^{k \cdot n'}} + a_\ell X^{p^{k \cdot \ell}} + \dots + a_1 X^{p^k} + a_0 X \in \mathcal{Q}_{p,kn}^L$,
- (c) (when $n|p-1$) for any $\alpha \in \mathbb{F}_p$ with $\alpha^n = 1$, $\alpha^{-n'} L_{f(\alpha \cdot X)} = \alpha^{-n'} (\alpha^{n'} X^{p^{n'}} + \alpha^\ell a_\ell X^{p^\ell} + \dots + \alpha a_1 X^p + a_0 X) \in \mathcal{Q}_{p,n}^L$,
- (d) $\gamma^{-p^{n'}} L_f(\gamma \cdot X) = \gamma^{-p^{n'}} ((\gamma \cdot X)^{p^{n'}} + a_\ell (\gamma \cdot X)^{p^\ell} + \dots + a_1 (\gamma \cdot X)^p + a_0 \gamma X) \in \mathcal{Q}_{p,n}^L$.

Proof. One can observe that these four quasi-subfield polynomials have the same β . Therefore, we only have to show that their splitting conditions are equivalent. The equivalence between (a) and (b) directly comes from Proposition 2. Indeed,

$$\begin{aligned} L_f \in \mathcal{Q}_{p,n}^L &\Leftrightarrow f|X^n - 1 \Leftrightarrow f(X^k)|X^{kn} - 1 \text{ in } \mathbb{F}_p[X] \\ &\Leftrightarrow L_{f(X^k)} \in \mathcal{Q}_{p,kn}^L. \end{aligned}$$

Properties (a) and (c) are also equivalent as

$$\begin{aligned} L_f \in \mathcal{Q}_{p,n}^L &\Leftrightarrow f|X^n - 1 \Leftrightarrow f(\alpha \cdot X)|(\alpha \cdot X)^n - 1 \\ &\Leftrightarrow f(\alpha \cdot X)|X^n - 1 \text{ since } \alpha^n = 1 \\ &\Leftrightarrow \alpha^{-n'} f(\alpha \cdot X)|X^n - 1 \\ &\Leftrightarrow \alpha^{-n'} L_{f(\alpha \cdot X)} \in \mathcal{Q}_{p,n}^L. \end{aligned}$$

Finally, replacing X by $\gamma \cdot X$ clearly does not change the fact that the polynomial is split, therefore (a) \Leftrightarrow (d) is trivial. \square

Let us now reconsider the transformations (a) \Leftrightarrow (b) and (a) \Leftrightarrow (c). As these transformations do not change the value of β and they send completely splitting linearized quasi-subfield polynomials with coefficients in \mathbb{F}_p onto other ones, we can define equivalence classes by saying that two completely splitting linearized quasi-subfield polynomials with coefficients in \mathbb{F}_p are equivalent to each other if one can be obtained from the other with one through one of the previous transformations. Obviously, since the transformations leave the value of β - which determines the efficiency of the ECDLP algorithm - unchanged, we are only interested in finding one representative of each class.

3.2. Examples of completely splitting linearized QSPs

In order to find examples of linearized QSPs, we performed a systematic search of representatives of classes of equivalence of completely splitting QSPs. We will now explain how we did this search, and present our results.

From now on, we will only consider polynomials with coefficients in the base field \mathbb{F}_p . Then any L of the shape $X^{p^{kn'}} + a_\ell X^{p^{k\ell}} + \dots + a_0 \in \mathcal{Q}_{p,kn}^L$ is equivalent (in the above sense) to $X^{p^{n'}} + a_\ell X^{p^\ell} + \dots + a_0 \in \mathcal{Q}_{p,n}^L$. Whenever we have a non trivial factor d of all the element of the set $\{i \geq 1, a_i \neq 0\} \cup \{n\}$, we may use it to reduce the degree of the polynomial by a factor d . Therefore we may reduce the search of representatives of each class to polynomials of the shape $X^{p^{n'}} + a_\ell X^{p^\ell} + \dots + a_0 \in \mathbb{F}_p[X]$ with $\{i \geq 1, a_i \neq 0\} \cup \{n\}$ setwise coprime.

One can also notice that transformation (a) \Leftrightarrow (c) cannot often be used. Indeed if n is prime then $\alpha^n = 1$ implies $n|p-1$.

Since we restrict the search to polynomials in $\mathbb{F}_p[X]$, all the coefficients of C_L are in \mathbb{F}_p and thus A_L is merely C_L^n . Hence, Proposition 1 says that L splits completely over \mathbb{F}_{p^n} if and only if $C_L^n = I$.

Let $L = X^{p^{n'}} + a_\ell X^{p^\ell} + \dots + a_0$ be a fixed linearized polynomial in $\mathbb{F}_p[X]$. We can assume $a_0 \neq 0$ since if $a_0 = 0$ then 0 is a root of L with multiplicity

at least p so L does not split completely. We can search for the smallest n such that L splits completely over \mathbb{F}_{p^n} . This amounts to searching for n such that $C_L^n = I$, in other words finding the order of C_L . Note that C_L is in $\text{GL}_{n'}(\mathbb{F}_p)$ since $\det C_L = (-1)^{n'} a_0 \neq 0$, hence n exists. Moreover as we also want $\beta(L) = n \cdot \ell / (n')^2 \leq 1$, we only have to compute C_L^k with $k < n'^2 / \ell$. If we find such a k with $C_L^k = I$ then $L_f \in \mathcal{Q}_{p,k}^L$.

This naturally leads to an algorithm to produce a set of representatives of the previously defined equivalence classes. The results output by our algorithm when asking for representatives of the equivalence classes for $p \in \{2, 3, 5, 7\}$, $n' \leq 16$ and coefficients values restricted to $\{0, 1, -1\}$, are presented in appendix (Table B.2). Observing patterns in them allowed us to conjecture new types of quasi-subfield polynomials. We present one representative per equivalence class, as other quasi-subfield polynomials can be obtained by using the rules listed in Proposition 4.

Proposition 5 (Families of linearized QSPs). *The following types of linearized polynomials are quasi-subfield polynomials:*

Type 1 L_h with $h = X^{p^a} + \dots + X^{p^0} + 1$, where $q = p^r$, $r \geq 0$, $n = p_{a+1}$, $p_i = 1 + q + \dots + q^i$ and $a \geq 2$, $\beta = 1 - \frac{1}{p_a} (1 - \frac{p_a - 1}{q \cdot p_a})$. It is the family introduced in [1].

Type 1bis $X^{p^{n-1}} + \dots + X^{p^2} + X^p + X$, $n' = n - 1$, $\beta = 1 - \frac{1}{(n-1)^2}$.

Type 2 L_{f_a} with $f_a = \begin{cases} X^{q^d-1} + \dots + X^{q-1} + 1 & \text{if } a = 0 \\ X^{q^d} + \dots + X^q + X + a & \text{otherwise} \end{cases}$, $n = q^{d+1} - 1$,
 $q = p^r$, $r \geq 1$, $a \in \mathbb{F}_q$, $\beta = 1 - \frac{q^{d-1}}{(1+q+\dots+q^{d-1})^2}$

Type 3 *Inverses of Type 1 and inverses of Type 2.*

Proof. Type 1 is proven in [1, Lemma 4.3]. Moreover it is obvious that L_{X-1} is a quasi-subfield polynomial over \mathbb{F}_{p^n} for any p prime and n (tolerating here $\ell = 0$). By Proposition 3, its inverse $L_{(X^n-1)/(X-1)} = L_{X^{n-1} + \dots + X^2 + X + 1}$ is a quasi-subfield polynomial in \mathbb{F}_{p^n} . This proves the Type 1bis. One can notice that it is in fact a particular case of Type 1 (when $r = 0$).

For Type 2 polynomials, we need to show that f_a divides $X^n - 1$ and thus we look at the factorisation of $X^{q^{d+1}-1} - 1$. It appears easier to compute this by looking at $X(X^{q^{d+1}-1} - 1) = X^{q(d+1)} - X$ since the Frobenius is easy to compute in \mathbb{F}_{p^n} . One can observe with $g = X^{q^d} + \dots + X^q + X$, we have that $X^{q^{d+1}} - X = g^q - g$. Thus, as in Berlekamp's polynomial trace factorization algorithm, we get that $X^{q^{d+1}} - X = \prod_{a \in \mathbb{F}_q} \text{gcd}(X^{q^{d+1}} - X, g + a)$. But we also have $X^{q^{d+1}} - X = (g + a)^q - (g + a)$ so $g_a | X^{q^{d+1}} - X$ and $X^{q^{d+1}} - X = \prod_{a \in \mathbb{F}_q} (g + a) = X f_0 \prod_{a \in \mathbb{F}_q} f_a$

That is why, $\prod_{a \in \mathbb{F}_q} f_a = X^n - 1$. This gives that Type 2 polynomials split completely over \mathbb{F}_{p^n} . Thus it only remains to verify that $\beta \leq 1$.

If $a = 0$, then $n' = q^d - 1$ and $\ell = q^{d-1} - 1$ hence

$$\begin{aligned}\beta &= \frac{(q^{d-1} - 1)(q^{d+1} - 1)}{(q^d - 1)^2} = 1 - \frac{q^{d+1} + q^{d-1} - 2q^d}{(q^d - 1)^2} = 1 - \frac{q^{d-1}(q-1)^2}{(q^d - 1)^2} \\ &= 1 - \frac{q^{d-1}}{(1 + q + \dots + q^{d-1})^2} < 1,\end{aligned}$$

while if $a \neq 0$, then $n = q^{d+1} - 1$, $n' = q^d$, $\ell = q^{d-1}$ hence

$$\beta = \frac{q^{d-1}(q^{d+1} - 1)}{q^{2d}} = 1 - \frac{1}{q^{d+1}} < 1.$$

Proposition 3 addresses Type 3 polynomials. We know that for Type 1 we have $Xh^q = h + X^n - 1$ thus $X^n - 1 = h(Xh^{q-1} - 1)$ and the inverse of L_h is $L_{Xh^{q-1}-1}$. For Type 2, we have $\prod_{a \in \mathbb{F}_p} f_a = X^n - 1$, thus the inverse of L_{f_a} is $L_{\prod_{b \neq a} f_b}$. \square

Recall that this list does not cover all the equivalence classes. It was only conjectured from what was found with small n and small p and coefficients values in $\{0, 1, -1\}$. For example, when we launch the algorithm for very small n with coefficients allowed to be anything in \mathbb{F}_p , we get for $p = 5$ and $n = 4$, that $L_{(X^2+X+3)}$ is a linearized QSP. Indeed $(X^2 + X + 3)(X^2 - X + 3) = (X^2 + 3)^2 - X^2 = X^4 + X^2 - 1 - X^2 = X^4 - 1$, so $X^2 + X + 3$ splits in \mathbb{F}_{5^4} and $\beta = 4 \times 1/4 = 1$. Moreover, computing its equivalence class using Proposition 4, we observe that no element of its equivalence class has all its coefficients in $\{0, 1, -1\}$.

3.3. Examples of multiplicative quasi-subfield polynomials

We now study another family of quasi-subfield polynomials considered in [1], namely polynomials whose roots form a multiplicative group of \mathbb{F}_{p^n} .

More precisely, we consider quasi-subfield polynomials of the type

$$L = X^{p^{n'}} - X^a$$

together with an integer r such that $a = p^{n'} \pmod r$, $r | p^n - 1$ and $n' > \log_p(r)$. Indeed, L factors as $X^a(X^{p^{n'}-a} - 1)$, so the number of roots of L in \mathbb{F}_{p^n} is at most $1 + p^{n'} - a$. Moreover, there are $\gcd(k, p^n - 1)$ roots of $X^k - 1$ in \mathbb{F}_{p^n} . In order to have the maximal number of distinct roots, we must choose tuples (p, n, n', r) such that for $a = p^{n'} \pmod r$, and $\gcd(p^{n'} - a, p^n - 1) = p^{n'} - a$, i.e. $p^{n'} - a | p^n - 1$.

Proposition 6 (Multiplicative quasi-subfield polynomials). *Let p, n, n', r be defined in any of the following three ways:*

1. Let p prime and $k \geq 2$ and $i \geq 1$ integers. Let $n = 2ik$, $n' = i(2k - 1) = n - i$ and $r = \frac{p^n - 1}{p^{2i} - 1}$;

2. Let $p = k^n + k - 1$ prime and $k \geq 2$ an integer. Let $n' = 1$ and $r = (p - k)/(k - 1)$;
3. Let $p = k^n - k - (-1)^n$ be prime, $n > 2$ and $k > 1$ integers such that $k^n \gg 1$. Let $n' = n - 1$ and let $r = \frac{(p^n - 1)(k - (-1)^n)}{(k^n - k)(k^n - (-1)^n)}$.

Let $a = p^{n'} \bmod r$ and let $L = X^{p^{n'}} - X^a \in \mathbb{F}_{p^n}[X]$. Then L is a quasi-subfield polynomial.

We make a few observations before proving this proposition. For the first family when $p = 2$, $i = 1$ and $k = 2$, we get $r = (2^4 - 1)/(2^2 - 1) = 5$ and $a = 3$, and thus $L = X^8 - X^3$ is a quasi-subfield polynomial. On the other hand, $\beta(L) = \log_2(3) \cdot 4/3^2 \simeq 0.70 < 0.75$. This shows that Theorem 1 is not valid for multiplicative quasi-subfield polynomials.

In the second and last families, we can choose n prime as is the case for the subfield curves recommended by NIST. However, there are values of n which may not lead to any suitable parameter set for the second and third types. For example, with $n = 5$ and $k > 1$, the integer $k^5 + k - 1 = (k^3 + k^2 - 1)(k^2 - k + 1)$ so $k^5 + k - 1$ is not prime. More generally, for all $n \equiv 5 \pmod{6}$, $(k^2 - k + 1)|(k^n + k - 1)$ and thus any integer of the shape $k^{5+6i} + k - 1$ with $k > 1$ is composite. Similarly, for all $n \equiv 2 \pmod{6}$, $(k^2 - k + 1)|(k^n - k + 1)$ and thus there is no prime of the shape $k^{2+6i} - k + 1$ with $k > 1$ and $i > 0$.

The last two families overlap when $n = 2$ as $(k - 1)^2 + (k - 1) - 1 = k^2 - k - 1$. We excluded the case $n = 2$ in the last family, because such a choice of n' and r would lead to $\beta = 0$, which is not allowed in our definition of quasi-subfield polynomials. Yet, thanks to the last two families, we have a multiplicative quasi-subfield polynomial for any n and $p = k^n - k - (-1)^n$ prime.

Finally, it is worth noticing that the case $p = k^n - k - (-1)^n$ is the most promising one among the families introduced. Indeed, primes of the form $f(2^m)$, where $f(x)$ is a low-degree polynomial with small integer coefficients, are often used in cryptography since they were introduced in [8]. Indeed as well as for Mersenne primes, they allow fast modular reduction. They are called Solinas primes, or generalized Mersenne primes. Coming back to our example, $f(x) = x^n - x - (-1)^n$ verifies the constraint required about the weights of the coefficients, so the last family when applied with k a power of 2 corresponds to Solinas primes. It is then important to notice that Curve448, which is part of the approved elliptic curves for use by the US Federal Government, uses a prime exactly of this shape: $p = 2^{448} - 2^{224} - 1$. [9][10]. Moreover, four others curves that were recommended by NIST in 1999 [11] also uses Solinas primes: p-192 ($p = 2^{192} - 2^{64} - 1$), p-224 ($p = 2^{224} - 2^{96} + 1$) and p-256 ($p = 2^{256} - p^{224} + 2^{192} + 2^{96} - 1$) and p-384 ($p = 2^{384} - 2^{128} - 2^{96} + 2^{32} - 1$). Therefore, it may seem interesting to study more deeply multiplicative quasi-subfield polynomials when p is a Solinas prime. For a list of Solinas primes of the shape $2^n - 2^m \pm 1$, one can consult [8]. Of course, this approach is still far from threatening the security of these curves: they are defined on a prime field \mathbb{F}_p while we are considering an extension field \mathbb{F}_{p^n} with $n \geq 2$ and have $\beta \simeq 1$ so we obtain a complexity of $O(p^{0.95n})$. (See Remark 1 for the detail)

Proof. In each case, we show that for $a := p^{n'} \bmod r$, we have $p^{n'} - a | p^n - 1$ and $\beta := \frac{n \log_p a}{n'^2} \leq 1$.

1. We first note that $r = \frac{p^n - 1}{p^{2i} - 1}$ is an integer since $2i | n$.

We now show that $a = p^{n'} \bmod r = \frac{p^{i(2k-1)+1}}{p^i+1}$. Indeed, we have

$$p^{n'} - \frac{p^{i(2k-1)+1}}{p^i+1} = \frac{p^n - 1}{p^{2i} - 1} (p^i - 1) = r(p^i - 1)$$

and

$$\begin{aligned} \frac{p^{i(2k-1)+1}}{p^i+1} \frac{1}{r} &= \frac{(p^{i(2k-1)+1})}{(p^i+1)} \frac{(p^{2i}-1)}{(p^{2ik}-1)} = (p^{i(2k-1)+1}) \frac{(p^i-1)}{(p^{2ik}-1)} \\ &= \frac{p^{2ik} - p^{i(2k-1)} + p^i - 1}{(p^{2ik}-1)} \\ &= 1 - \frac{p^{i(2k-1)} - p^i}{p^{2ik}-1} < 1 \end{aligned}$$

since $2k - 1 > 1$.

Therefore, we have $p^{n'} - a = r(p^i - 1) = \frac{p^{2ik}-1}{p^i+1}$ and thus $p^{n'} - a | p^n - 1$. This implies that L splits completely over \mathbb{F}_{p^n} and it has $p^{n'} - a + 1 = \frac{p^{2ik}-1}{p^i+1} + 1 \approx p^{i(2k-1)} = p^{n'}$ roots.

Moreover, using $a = \frac{p^{i(2k-1)+1}}{p^i+1} = \sum_{j=0}^{2k-2} (-p^i)^j \leq p^{i(2k-2)}$, we get

$$\beta = \frac{\log_p(a) \cdot n}{n'^2} \leq \frac{i(2k-2) \cdot 2ik}{(i(2k-1))^2} = 1 - \frac{1}{(2k-1)^2} \leq 1.$$

2. We first note that $r = \frac{p-k}{k-1} = \frac{k^n-1}{k-1}$ is integer since $k-1 | k^n - 1$. Moreover, $p = \frac{p-k}{k-1}(k-1) + k = r(k-1) + k$ with $k < 1 + k + \dots + k^{n-1} = r$ so $a = (p \bmod r) = k$ and $p - a = k^n - 1$. Therefore,

$$\begin{aligned} p^n - 1 &= (k^n + k - 1)^n - 1 = \sum_{i=1}^n \binom{n}{i} (k^n - 1)^i k^{n-i} \\ &= (k^n - 1) \sum_{i=1}^n \binom{n}{i} (k^n - 1)^{i-1} k^{n-i} \\ &= (p - a) \sum_{i=1}^n \binom{n}{i} (k^n - 1)^{i-1} k^{n-i}. \end{aligned}$$

Consequently, we have $p - a | p^n - 1$, so $L = X^p - X^a$ splits in \mathbb{F}_{p^n} and it has $p^{n'} - a + 1 = p - a + 1 = (k^n + k - 1) - k + 1 = k^n$ roots. This is very

close to $p = p^{n'}$ if $k^n \gg 1$. Furthermore, since $k^n \leq k^n + k - 1 = p$, we have

$$\beta = \log_p(a) \cdot n/1 = \log_p(a^n) = \log_p(k^n) \leq 1.$$

3. The third proof is very similar to the first two proofs, and presented in Appendix C.

□

3.4. Quasi-subfield polynomials with n Mersenne

When $n = 2^k - 1$ is a Mersenne prime, $(X^n - 1)/(X - 1)$ has $(n - 1)/k$ irreducible factors of degree k over \mathbb{F}_2 , which gives a large number of potential candidates for linearized quasi-subfield polynomials in \mathbb{F}_{2^n} . We note that for $p = 2$, $p^{d+1} - 1$ is a Mersenne number, hence Type 2 of Proposition 5 gives examples of such polynomials.

The case of linearized quasi-subfield polynomials with n a Mersenne prime number is also treated in the appendix of [1]. Interestingly, [1] argued that such parameters were unlikely to exist. We now recall (and slightly extend) their heuristic argument, and we show that Type 2 polynomials from Proposition 5 give a counter-example to it.

Reasoning from [1]. Let us consider k such that $n = 2^k - 1$ is prime, and denote by $N(k, n')$ the number of distinct polynomials of degree n' that divide $X^n - 1$. Then [1] gives the following lemma:

Lemma 5. *We have $N(k, n') = \binom{\lfloor n/k \rfloor}{\lfloor n'/k \rfloor}$ if $n' \bmod k \in \{0, 1\}$, and $N(k, n') = 0$ otherwise. Moreover, $\log \left(\binom{\lfloor n/k \rfloor}{\lfloor n'/k \rfloor} \right) \simeq (n'/k) \log(n/n')$ when $n/k \gg 1$ and $n'/k \gg 1$.*

The argument in [1] relies on the following heuristic approximation: for n a Mersenne prime, we may assume that the density of “sparse enough” polynomials (*i.e.* polynomials of the shape $X^{n'} - \lambda(X)$ with $\deg(\lambda)$ small) is identical for factors of $X^n - 1$ as for random polynomials of the same degree.

Since in $\mathbb{F}_2[X]$, there are $2^{n'}$ monic polynomials of degree n' and 2^ℓ monic polynomials of degree at most ℓ , this assumption allows us to approximate the number of polynomials of degree n' that divide $X^n - 1$ and are sparse enough by $N(k, n')2^{\ell - n'}$. Accordingly, such polynomials a priori exist if and only if

$$\ell > n' - (n'/k) \log(n/n').$$

The case considered in the appendix of the article is when the quasi-subfield polynomial approach beats generic algorithms on ECDLP, which as we will prove in Lemma 8 requires $\alpha_\beta = \frac{1}{2\kappa\beta} \geq 1$ for some algorithmic constant κ .

We recall their argument in this case first, even if Type 2 does not fall in this category since its $\alpha_\beta \simeq \frac{1}{2\kappa}$ is not bigger than 1. To improve on generic algorithms, we want $\alpha_\beta = \frac{1}{2\kappa\ell n/n'^2} \geq 1$ hence $\ell \leq \frac{n'^2}{2\kappa n}$. With the previous

constraint on ℓ , we obtain $\frac{n'^2}{2\kappa n} > n' - (n'/k) \log(n/n')$. Thus, since $k \simeq \log(n)$, we get $\frac{n'}{2\kappa n} > 1 - \log(n/n')/\log(n) = \log(n')/\log(n)$. Therefore, $\frac{\log(n)}{2\kappa n} > \frac{\log(n')}{n'}$. Since $2\kappa \simeq 10$, and $n' < n$, this inequality can never be satisfied (except if $n' = 1$) so according to the above heuristic approximation, there should not be any linearized quasi-subfield polynomials with n Mersenne and big n and n' beating the generic algorithms.

The case of Type 2 polynomials. The same reasoning can be extended to quasi-subfield polynomials that do not verify $\alpha_\beta \geq 1$.

In this case, we only require $\beta = \ell \cdot n/n'^2 \leq 1$. Therefore, we have $\ell \leq n'^2/n$ (instead of $\ell \leq \frac{n'^2}{2\kappa n}$ before). This constraint added to the same heuristic as before gives: $\frac{n'^2}{n} > n' - (n'/k) \log(n/n')$ which similarly as in the previous paragraph gives $\frac{\log(n)}{n} > \frac{\log(n')}{n'}$. Since the function $\log(x)/x$ is decreasing for $x > e$, we deduce (following the same heuristic reasoning) that quasi-subfield polynomials are unlikely to exist for $n > n' \geq 3$.

This conclusion, however, is contradicted by the existence of Type 2 polynomials from Proposition 5.

On the heuristic approximations used in [1]. The above contradiction shows that the heuristic approximation used in [1] does not hold in general: when n is a Mersenne prime, there exists an ℓ such that the density of “sparse enough” polynomials (*i.e.* polynomials of the shape $X^{p^{n'}} - \lambda(X)$ with $\deg(\lambda) \leq \ell$) is bigger for factors of $X^n - 1$ than for random polynomials.

A similar heuristic in [1] says that there are only rare parameters for which we can have a quasi-subfield multiplicative polynomials. It uses really similar arguments to the ones introduced before for the case where n is a Mersenne prime. Property 6 shows that this heuristic about the distribution of completely splitting polynomials also fails.

4. Application to Cryptography

While quasi-subfield polynomials are mathematical objects of independent interest, the main motivation for their introduction in [1] is a cryptographic application. In this section we first recall the Elliptic Curve Discrete Logarithm Problem (ECDLP) and standard approaches to solve it. We then describe Huang *et al.*'s algorithm [1] using quasi-subfield polynomials and we explain how its complexity crucially depends on the parameter β of the polynomial. Next, we apply our results to this ECDLP algorithm, and discuss the resulting complexity. Finally, we introduce some aspects of coding theory where our results on linearized polynomials could be useful.

4.1. ECDLP and previous ECDLP algorithms

Let us consider an ECDLP instance: Let \mathcal{E} be an elliptic curve on $K = \mathbb{F}_{p^n}$, P a point on the curve \mathcal{E} and Q a point in $\langle P \rangle$, the group generated by P . We are looking for k such that $Q = kP$.

Before considering the algorithm using QSPs [1], we recall two algorithms for solving the ECDLP and their complexity.

- Exhaustive search (or brute-force algorithms) : it corresponds to the computation all the elements of $\langle P \rangle$ until finding Q . The cost is $O(\langle P \rangle) = O(p^n)$ for typical parameters.
- Generic algorithms such as Baby-Step-Giant-Step or Pollard-Rho [12]: The complexity is $O(\sqrt{|\langle P \rangle|}) \approx O(p^{n/2})$.

These will be used as benchmarks to assess the performance of our algorithm. For more information about these algorithms and other approaches to solve the ECDLP, the reader can consult *Recent progress on the elliptic curve discrete logarithm problem* [13] by Galbraith and Gaudry. It is also worth noticing that these two algorithms can solve the discrete logarithm problem in any group and we can hope that the new algorithm, which uses the structure of the group, has a better complexity.

When n is composite, we can write $n = \tilde{n}k$ and consider $q = p^k$ so that $\mathbb{F}_{p^n} = \mathbb{F}_{q^{\tilde{n}}}$. Better algorithms exist in this situation: Gaudry [14] succeeded in 2009 to find an algorithm solving the elliptic curve discrete logarithm problem on $\mathbb{F}_{q^{\tilde{n}}}$ in $O(q^{2-2/\tilde{n}})$. For $\tilde{n} = 2$, it leads to an algorithm with cost $O(p^{(n/2)(2-1)}) = O(p^{n/2})$ comparable with generic algorithms. For $\tilde{n} = 3$, it leads to an algorithm with cost $O(p^{(n/3)(2-2/3)}) = O(p^{4/9n})$, slightly better than generic algorithms. Diem also proved that there exists a sequence of prime powers $Q_i = q_i^{n_i}$ with $n_i \simeq \sqrt{\log(q_i)}$ such that the ECDLP in $\mathcal{E}(F_{Q_i})$ can be solved in subexponential time [15]. It works with any elliptic curve over \mathbb{F}_{Q_i} and uses an approach similar to the one introduced below but with subfield polynomials instead of quasi-subfield polynomials. This was one of the motivation of this new approach.

4.2. The quasi-subfield approach

We will now introduce the algorithm of [1], which uses quasi-subfield polynomials to solve elliptic curve discrete logarithm problems.

Let \mathcal{E} be an elliptic curve on $K = \mathbb{F}_{p^n}$, $P \in \mathcal{E}$ and $Q \in \langle P \rangle$. The elliptic curve discrete logarithm problem asks for computing k such that $Q = kP$. For simplicity and concreteness, we assume the curve is given in reduced Weierstrass coordinates.

Let $R \in \mathbb{F}_{p^n}[X]$ be a quasi-subfield polynomial. We define V as the set of the roots of R and $\mathcal{F} := \{(x, y) \in \mathcal{E} | x \in V\}$. The algorithm first computes more than $|V|$ relations of the shape:

$$a_j P + b_j Q = P_1 + \dots + P_m$$

with a_j, b_j random and $P_i \in \mathcal{F}$. Linear algebra on the relations then gives the value of k such that $Q = kP$.

In order to compute these relations, the algorithm uses Semaev's summation polynomials [16]: for an elliptic curve \mathcal{E} defined over a field K , the r th summation polynomial $S_r \in K[X]$ is such that

$$S_r(x_1, \dots, x_r) = 0 \Leftrightarrow \exists (x_1, y_1), \dots, (x_r, y_r) \in \mathcal{E}, (x_1, y_1) + \dots + (x_r, y_r) = 0$$

For given a_j, b_j , we compute $a_j P + b_j Q = (X_j, Y_j)$. Then, computing P_1, \dots, P_m such that

$$a_j P + b_j Q = P_1 + \dots + P_m = (x_1, y_1) + \dots + (x_m, y_m)$$

with $x_i \in V$ amounts to finding $x_1, \dots, x_m \in V$ such that $S_{m+1}(X_i, x_1, \dots, x_m) = 0$ and then finding the associated y_i .

The polynomial equation $S_{m+1}(X_i, x_1, \dots, x_m) = 0$ is solved as follows. Let \mathcal{M} be the set of monomials in $K[x_1, \dots, x_m]$, and let i be a positive integer. For $f = \sum_{M \in \mathcal{M}} a_M M \in K[x_1, \dots, x_m]$, let $F^i(f) = \sum_{M \in \mathcal{M}} a_M^i M$. Let also $\phi : K[x_1, \dots, x_m] \rightarrow K[x_1, \dots, x_m]$ defined by

$$f(x_1, \dots, x_m) \mapsto F^{n'}(f)(\lambda(x_1), \dots, \lambda(x_m)).$$

Note that $f^{p^{n'}} \equiv \phi(f) \pmod{(x_1^{p^{n'}} - \lambda(x_1), \dots, x_m^{p^{n'}} - \lambda(x_m))}$.

Finally, let $S^{(0)}(x_1, \dots, x_m) = S_{m+1}(X_i, x_1, \dots, x_m)$ and for $k \in \{1, \dots, m-1\}$, let $S^{(k)}(x_1, \dots, x_m) = \phi(S^{(k-1)}(x_1, \dots, x_m))$. One can then solve the polynomial equation $S_{m+1}(X_i, x_1, \dots, x_m) = 0$ by solving the system $\mathcal{S} = \{S^{(k)} = 0\}_{k=1}^{m-1}$. This is a sparse polynomial system with m equations and m variables.

4.3. Complexity of the quasi-subfield approach

We now recall the complexity estimations of this algorithm as given in [1].

We know that $\mathcal{S} = \{S^{(k)}\}_{k=1}^{m-1}$ is a sparse polynomial system of m equations and m variables. Therefore, it can be solved efficiently using Rojas' sparse resultant algorithm [17] and a univariate polynomial root finding algorithm such as BTA [5]. According to [1] (Lemma 3.1) the cost of this step is $\tilde{O}(m^{5.188}(3p^\ell)^{\kappa m^2})$. Here we introduce the notation κ as the numerical value 4.876 used in [1] may be suboptimal. Moreover, the system has solutions only with probability $\frac{|\mathcal{F}|^m/m!}{p^n}$ since (X_i, Y_i) is a random point on \mathcal{E} with $|\mathcal{E}| \simeq p^n$ and the number of sums of m points in \mathcal{F} is approximately $|\mathcal{F}|^m/m!$. Also, heuristically, half of the values in V are the x-coordinates of exactly two points on the curve so $|\mathcal{F}| \simeq |V| \simeq p^{n'}$. As we need $p^{n'}$ relations of this type, the cost of the relation search phase is $p^{n'} \frac{m! p^n}{p^{n'm}} \tilde{O}(m^{5.188}(3p^\ell)^{\kappa m^2})$.

Once all the $p^{n'}$ relations are gathered, each of them involves m points. Therefore, the system built from these relations is sparse. Thus, a sparse linear algebra algorithm can be used to finish the computation [18], at a cost approximately $mp^{2n'}$. This gives the complete cost of the algorithm:

$$m! p^{n-n'm+n'} \tilde{O}(m^{5.188}(3p^\ell)^{\kappa m^2}) + mp^{2n'}$$

Rewriting this expression to make β appear, we get the following estimation of the complexity:

Proposition 7 (Complexity of Huang *et al.*'s algorithm). *Let $P = X^{p^{n'}} - \lambda(X)$ be a β -quasi-subfield polynomial and let $\ell = \log_p(\deg \lambda)$. If $|\mathcal{F}| \simeq |\mathcal{V}| \simeq p^{n'}$, the*

complexity of Huang et al.'s algorithm is

$$\tilde{O}\left(m!p^{n\left(1+\kappa\beta\left(\frac{n'm}{n}\right)^2-\frac{n'm}{n}\right)+n'}m^{5.188}3^{\kappa m^2}\right)+mp^{2n'}$$

where κ is a constant involved in the cost of the resolution of the system \mathcal{S} currently majored by 4.876.

In the following we define $\alpha = n'm/n > 0$, and we try to find α which minimises the complexity. We will assume that m is fixed.

Proposition 8 (Best choice of parameters). *We can observe the following results in order to optimize the complexity:*

- The minimal complexity is obtained for $\alpha = \alpha_\beta$ with $\alpha_\beta := \frac{1}{2\kappa\beta}$. Then, the complexity becomes $\tilde{O}\left(p^{\max(2\alpha_\beta/m, 1-\alpha_\beta(1/2-1/m))n}\right)$.
- In order to beat brute force algorithms, $m > \max(2\alpha_\beta, 2)$ is required. So we have interest not to choose a very small integer for m .
- If $\alpha_\beta < 2$ and $m \gg 1$, then the complexity becomes $\tilde{O}\left(p^{(1-\alpha_\beta/2)n}\right)$
- Therefore, to beat generic algorithms, we need $\alpha_\beta > 1$

We remark that the condition $\alpha_\beta < 2$ is not really restrictive. Indeed for all the quasi-subfield polynomials exhibited in this paper, we have $\alpha_\beta < \alpha_{0.5} < 1$.

Proof. Let us now prove these four results. The complexity of the algorithm is bounded by $\tilde{O}\left(m!p^{n\left(1+\kappa\beta\left(\frac{n'm}{n}\right)^2-\frac{n'm}{n}\right)+n'}m^{5.188}3^{\kappa m^2}\right)+mp^{2n'}$ which with the α -notation and the fact that m is considered as a fixed integer, can be rewritten as $\tilde{O}\left(p^{n(\kappa\beta\alpha^2-\alpha+1)+\alpha n/m}+p^{2\alpha n/m}\right)$.

Since $\kappa\beta\alpha^2 - \alpha + 1$ is minimum for $\alpha = \frac{1}{2\kappa\beta} = \alpha_\beta$ (we recall that we only consider $\beta > 0$) and then has minimal value $\kappa\beta\frac{1}{(\kappa\beta)^2} - \frac{1}{2\kappa\beta} + 1 = 1 - \frac{1}{4\kappa\beta} = 1 - \frac{\alpha_\beta}{2}$, we get that the complexity can be rewritten as

$$\tilde{O}\left(p^{\max(2\alpha_\beta/m, 1-\alpha_\beta(1/2-1/m))n}\right).$$

In order to beat the brute force algorithms (which corresponds to a complexity of $O(p^n)$), what we need is to have on one side $2\alpha_\beta/m < 1$ which is true as soon as $m > 2\alpha_\beta$, and on the other side, $1 - \alpha_\beta(1/2 - 1/m) < 1$, namely $m > 2$. Hence $m > \max(2\alpha_\beta, 2)$.

Moreover, one can notice that $2\alpha_\beta/m \leq 1 - \alpha_\beta(1/2 - 1/m)$ if only if $\alpha_\beta(1/m + 1/2) \leq 1$. Therefore if $m \gg 1$ then $\alpha_\beta \leq 2$ implies $2\alpha_\beta/m \leq 1 - \alpha_\beta(1/2 - 1/m)$, so we can rewrite the complexity as $\tilde{O}\left(p^{(1-\alpha_\beta/2)n}\right)$.

Generic algorithms have a complexity of $O(p^{n/2})$, therefore we need $\alpha_\beta > 1$ to run faster than them. \square

The following table gives concrete complexity estimates for various values of β , assuming $\kappa = 4.876$.

β	1.0	0.8	0.6	0.4	0.2	0.15	0.1
$1 - \alpha_\beta/2$	0.949	0.936	0.915	0.872	0.744	0.658	0.487

Complexity estimates of Huang *et al.*'s algorithm for various values of β . By Proposition 8 the complexity of Huang *et al.*'s algorithm is $\tilde{O}\left(p^{(1-\alpha_\beta/2)n}\right)$.

Remark 1. We observe that for $\beta = 1$, we have $1 - \alpha_\beta/2 \approx 0.95$ so we get a complexity slightly better than the one of brute force algorithms. We can beat generic algorithms for $\alpha_\beta > 1$, which for this specific value of κ implies $\beta < 0.103$.

4.4. Impact of our results on ECDLP

We will now study the consequences of Theorem 1. Let L be a linearized quasi-subfield polynomial. Then by Theorem 1 we have $\beta(L) \geq 3/4$ and with $\kappa \simeq 4.876$ we get $\alpha_\beta = \frac{1}{2 \cdot \kappa \beta(L)} \leq \frac{2}{3 \cdot \kappa} < 1/7$. This shows that $a_\beta < 1$ and it is not possible to beat generic algorithms with L . The best complexity we can hope is indeed $\tilde{O}(p^{(1-\alpha_\beta(1/2-1/m))n})$, which is bigger than $\tilde{O}(p^{(1-1/14)n})$.

The previous estimation uses the approximation $\kappa \simeq 4.876$. If we succeeded to have $\kappa < 1.5$ then, when $\beta(L) = 3/4$, we would have $\alpha_\beta > 1$, so such a polynomial L could allow us to have an algorithm running faster than generic algorithms.

All the quasi-subfield polynomials exhibited in this article have $\beta > 0.7$ and thus $\alpha_\beta < 1$. In particular, none of them currently leads to an algorithm running faster than generic algorithms.

4.5. Links with coding theory

Linearized polynomials have attracted considerable interest, and our results can therefore be used in other contexts as well.

For example, linearized polynomials occur in rank-metric codes. The characterisation of completely splitting linearized trinomials given in [7] is used in the same article to study codes of the shape $C_{3,n,\sigma} = \langle x, x^\sigma, x^{\sigma^3} \rangle_{\mathbb{F}_{q^n}}$, with both their result on existence and non existence of such trinomials being used. In [19], codes of the shape

$$\mathcal{C}_T := \left\{ a_0 X^{q^{t_0}} + a_1 X^{q^{t_1}} + \dots + a_{k-1} X^{q^{t_{k-1}}}, a_0, a_1, \dots, a_{k-1} \in \mathbb{F}_{q^n} \right\}$$

for sets $T = \{t_0 < t_1 < \dots < t_{k-1}\} \subset \{0, \dots, n-1\}$ are studied. Maybe, Theorem 1, which gives wider results than [7] on completely splitting linearized polynomials could help to study such codes.

This notion also appears in cyclic subspace codes. For instance in [7], families of cyclic subspace codes are exhibited via linearized polynomials. Interestingly, the paper uses a parameter called *gap* which characterizes the sparsity of a polynomial and implies bounds on the minimal distance of an associated code. The gap is defined as $n' - \ell$ for a linearized polynomial $P = X^{p^{n'}} - \lambda(X) \in \mathbb{F}_{p^n}[X]$ with λ of degree p^ℓ . It is therefore similar to our parameter $\beta := \frac{\ell \cdot n}{n'^2}$.

5. Conclusion

We studied the existence of quasi-subfield polynomials (QSP) introduced by Huang *et al.* [1]. We proved a new lower bound on the β parameter of *linearized* QSP, and we introduced several new QSP families. We leave as an open problem the classification of all the QSP.

The main motivation underlying [1] is a new algorithm to solve the elliptic curve discrete logarithm problem, with a complexity depending on the β parameter of the QSP used. We showed that this algorithm is currently outperformed by other algorithms even with our new QSP families. Moreover, our new bound suggests that Huang *et al.*'s algorithm will remain worse if only linearized QSPs are used.

Acknowledgements

We would like to thank the reviewers for their helpful comments, especially for pointing out the link between this work and coding theory.

References

- [1] M.-D. Huang, M. Kusters, C. Petit, S. L. Yeo, Y. Yun, Quasi-subfield polynomials and the elliptic curve discrete logarithm problem, *Journal of Mathematical Cryptology* 14 (1) (2020) 25–38.
- [2] G. McGuire, D. Mueller, Some results on linearized trinomials that split completely, *Finite Fields and their Applications* (2020) 149.
- [3] G. McGuire, J. Sheekey, A characterization of the number of roots of linearized and projective polynomials in the field of coefficients, *Finite Fields and Their Applications* 57 (2019) 68–91.
- [4] B. Csajbók, G. Marino, O. Polverino, F. Zullo, A characterization of linearized polynomials with maximum kernel, *Finite Fields and Their Applications* 56 (2019) 109–130.
- [5] E. Berlekamp, *Algebraic coding theory*, World Scientific, 1968.

- [6] W. Y. C. Chen, J. D. Louck, The combinatorial power of the companion matrix, *Linear Algebra and its Applications* 232 (1996) 261–278. doi:10.1016/0024-3795(95)90163-9.
URL <http://www.sciencedirect.com/science/article/pii/S0024379595901639>
- [7] P. Santonastaso, F. Zullo, Linearized trinomials with maximum kernel, arXiv preprint arXiv:2012.14861.
- [8] J. A. Solinas, et al., Generalized mersenne numbers, Citeseer, 1999.
- [9] M. Hamburg, Ed448-Goldilocks, a new elliptic curve, Tech. Rep. 625 (2015). URL <https://eprint.iacr.org/2015/625>
- [10] I. T. L. Computer Security Division, Transition Plans for Key Establishment Schemes | CSRC (Oct. 2017).
URL <https://csrc.nist.gov/News/2017/Transition-Plans-for-Key-Establishment-Schemes>
- [11] NIST, Recommended elliptic curves for federal government use (1999).
- [12] P. C. van Oorschot, M. J. Wiener, Parallel collision search with cryptanalytic applications, *J. Cryptology* 12 (1999) 1–28. doi:10.1007/PL00003816.
- [13] S. D. Galbraith, P. Gaudry, Recent progress on the elliptic curve discrete logarithm problem, *Designs, Codes and Cryptography* 78 (1) (2016) 51–72. doi:10.1007/s10623-015-0146-7.
URL <http://link.springer.com/10.1007/s10623-015-0146-7>
- [14] P. Gaudry, Index calculus for abelian varieties of small dimension and the elliptic curve discrete logarithm problem, *Journal of Symbolic Computation* 44 (12) (2009) 1690–1702. doi:10.1016/j.jsc.2008.08.005.
URL <https://linkinghub.elsevier.com/retrieve/pii/S074771710800182X>
- [15] C. Diem, On the discrete logarithm problem in elliptic curves, *Compositio Mathematica* 147 (1) (2011) 75–104. doi:10.1112/S0010437X10005075.
URL https://www.cambridge.org/core/product/identifier/S0010437X10005075/type/journal_article
- [16] I. Semaev, Summation polynomials and the discrete logarithm problem on elliptic curves, Tech. Rep. 031 (2004).
URL <http://eprint.iacr.org/2004/031>
- [17] J. M. Rojas, Solving Degenerate Sparse Polynomial Systems Faster, *Journal of Symbolic Computation* 28 (1) (1999) 155–186. doi:10.1006/jsc.1998.0271.
URL <http://www.sciencedirect.com/science/article/pii/S0747717198902711>

- [18] D. Wiedemann, Solving sparse linear equations over finite fields, IEEE Transactions on Information Theory 32 (1) (1986) 54–62. doi:10.1109/TIT.1986.1057137.
- [19] B. Csajbok, G. Marino, O. Polverino, Y. Zhou, Mrd codes with maximum idealizers, Discrete Mathematics 343 (9) (2020) 111985.

Appendix A. Adaptation of the results from [6]

In [6], Chen and Louck give a formula for computing the powers of the following companion matrices :

$$C(u_1, \dots, u_m) := \begin{bmatrix} u_1 & u_2 & \cdots & \cdots & u_m \\ 1 & 0 & \cdots & \cdots & 0 \\ 0 & 1 & \cdots & \cdots & 0 \\ \vdots & \vdots & \ddots & & \vdots \\ 0 & 0 & \cdots & 1 & 0 \end{bmatrix}$$

Comparing it with our definition of companion matrices,

$$D(a_1, a_2, \dots, a_m) = \begin{bmatrix} 0 & 0 & \cdots & \cdots & a_1 \\ 1 & 0 & \cdots & \cdots & a_2 \\ 0 & 1 & \cdots & \cdots & \vdots \\ \vdots & \vdots & \ddots & & \vdots \\ 0 & 0 & \cdots & 1 & a_m \end{bmatrix},$$

we notice that $D(a_1, a_2, \dots, a_m)$ is the antitranspose of $C(a_m, \dots, a_2, a_1)$. Formally, we have :

$$D(a_1, a_2, \dots, a_m) = P \cdot C(a_m, \dots, a_2, a_1)^T \cdot P \text{ with } P = \begin{bmatrix} 0 & \cdots & 0 & 1 \\ \vdots & \cdots & 1 & 0 \\ \vdots & & & \vdots \\ 1 & \cdots & & 0 \end{bmatrix}.$$

Since $P^2 = Id$, we get that for all $n \geq 0$, $D(a_1, a_2, \dots, a_m)^n$ is the antitranspose of $C(a_m, \dots, a_2, a_1)^n$. Chen and Louck give the following formula for the coefficient (i, j) of $C(u_1, \dots, u_m)^n$:

$$c_{i,j}^{(n)} = \sum_{k_1, \dots, k_m} \frac{k_j + k_{j+1} + \cdots + k_m}{k_1 + \cdots + k_m} \binom{k_1, k_2, \dots, k_m}{k_1 + \cdots + k_m} u_1^{k_1} \cdots u_m^{k_m}$$

where the summation is over non-negative integers satisfying $\sum k_l = n - i + j$. Moreover, when the previous sum is not defined, *i.e.* when $n = i - j$, $c_{i,j}^{(n)} = 1$.

Since applying the antitranspose boils down to swapping the coefficients (i, j) and $(m+1-j, m+1-i)$ ³, we get the expression of the coefficient (i, j) of $D(a_1, a_2, \dots, a_m)^n$:

$$\begin{aligned} d_{i,j}^{(n)} &= c_{m+1-j, m+1-i}^{(n)} \\ &= \sum_{k_1, \dots, k_m} \frac{k_{m-i+1} + k_{m-i+2} + \dots + k_m}{k_1 + \dots + k_m} \binom{k_1, k_2, \dots, k_m}{k_1 + \dots + k_m} a_m^{k_1} \dots a_1^{k_m} \end{aligned}$$

where the summation is over non-negative integers satisfying

$$\sum \iota k_\iota = n - (m - j + 1) + m - i + 1 = n - i + j$$

Moreover, when the previous sum is not defined (*i.e.* when $n = j - i$) then $d_{i,j}^{(n)} = 1$.

In the proof of Lemma 4, we consider $M := D(\otimes, \dots, \otimes, a^\bullet, 0, \dots, 0)$ a matrix of dimension n' , which leads to $M_{i,j}^n = 1$ if $n = i - j$, and

$$M_{i,j}^n = \sum_{\substack{\mathbf{k}=(k_1, \dots, k_{n'}) \\ \sum_{\iota=1}^{n'} \iota k_\iota = n - i + j}} w_{\mathbf{k}} \cdot 0^{k_1 + \dots + k_{n'-\ell-1}} \cdot (a^\bullet)^{k_{n'-\ell}} \cdot \otimes^{k_{n'-\ell+1} + \dots + k_{n'}}$$

otherwise, where $\mathbf{k} = (k_\iota)_{1 \leq \iota \leq n'}$ are non-negative integers and

$$w_{\mathbf{k}} = \frac{k_{n'-i+1} + \dots + k_{n'}}{k_1 + \dots + k_{n'}} \binom{k_1 + \dots + k_{n'}}{k_1, \dots, k_{n'}}.$$

Appendix B. Some linearized QSP

In this section we provide the list of linearized QSP found through the systematic search described in Section 3.2. Recall that this search only covers representatives of equivalence classes for $p \in \{2, 3, 5, 7\}$, $n' \leq 16$, and coefficients values restricted to $\{0, 1, -1\}$.

For the sake of readability, we list polynomials f instead of their corresponding quasi-subfield polynomials L_f . We provide the values of n and p such that L_f is in $\mathcal{Q}_{p,n}^L$ and indicate the associated value β . We mark by a checkmark in the table when the linearized polynomial belongs to the category (as defined in Proposition 5), except for the last category where we give the value of the inverse.

³We number the rows, as well as the columns from 1 to m .

f	n	β	p	T1	T2	T3
$X^2 + X + 1$	3	0.75	2	✓	✓	
$X^2 + X + 1$	3	0.75	3,5,7	✓	✓	
$X^3 + X + 1$	7	0.78	2	✓	✓	
$X^3 + X + 1$	8	0.8	3	✓	✓	
$X^3 + X^2 + X + 1$	4	0.8	2,3,5,7	✓	✓	
$X^4 + X + 1$	15	0.9	2	✓	✓	
$X^4 + X + 1$	13	0.8	3	✓	✓	
$X^4 + X^2 + X + 1$	7	0.8	2	✓	✓	$X^3 + X + 1$
$X^4 + X^3 + X^2 + X + 1$	5	0.9	2,3,5,7	✓	✓	
$X^5 + X + 1$	21	0.8	2	✓	✓	
$X^5 + X + 1$	24	0.9	5	✓	✓	
$X^5 + X^4 + X^3 + X^2 + X + 1$	6	0.9	2,3,5,7	✓	✓	
$X^5 - X^3 - X^2 + X - 1$	8	0.9	3	✓	✓	$X^3 + X + 1$
$X^6 + X + 1$	31	0.8	5	✓	✓	
$X^6 + X^5 + \dots + X^2 + X + 1$	7	0.9	2,3,5,7	✓	✓	
$X^7 + X + 1$	48	0.9	7	✓	✓	
$X^7 + X^3 + X + 1$	15	0.9	2	✓	✓	
$X^7 + X^6 + \dots + X^2 + X + 1$	8	0.9	2,3,5,7	✓	✓	
$X^8 + X + 1$	63	0.9	2	✓	✓	
$X^8 + X + 1$	57	0.8	7	✓	✓	$X^7 + X^3 + X + 1$
$X^8 + X^4 + X^2 + X + 1$	15	0.9	2	✓	✓	
$X^8 + \dots + X^2 + X + 1$	9	0.9	2,3,5,7	✓	✓	
$X^9 + X + 1$	73	0.9	2	✓	✓	
$X^9 + X + 1$	80	0.9	3	✓	✓	
$X^9 + X^3 + X + 1$	26	0.9	3	✓	✓	
$X^9 - X^6 - X^5 + X^3 - X^2 + X - 1$	13	0.9	3	✓	✓	$X^4 + X + 1$
$X^9 + \dots + X^2 + X + 1$	10	0.9	2,3,5,7	✓	✓	

f	n	β	p	T1	T2	T3
$X^{10} + X + 1$	91	0.9	3	✓		
$X^{10} + \dots + X^2 + X + 1$	11	0.9	2,3,5,7	✓		
$X^{11} + X^8 + X^7 + X^5 + X^3 + X^2 + X + 1$	15	0.9	2	✓		$X^4 + X + 1$
$X^{11} + \dots + X^2 + X + 1$	12	0.9	2,3,5,7	✓		
$X^{12} + \dots + X^2 + X + 1$	13	0.9	2,3,5,7	✓		
$X^{13} + X^4 + X + 1$	40	0.9	3	✓		
$X^{13} + \dots + X^2 + X + 1$	14	0.9	2,3,5,7	✓		
$X^{14} + \dots + X^2 + X + 1$	15	0.9	2,3,5,7	✓		
$X^{15} + X^7 + X^3 + X + 1$	31	0.9	2	✓	✓	
$X^{15} + X^{14} + \dots + X^2 + X + 1$	16	0.9	2,3,5,7	✓		
$X^{16} + X + 1$	255	0.9	2		✓	
$X^{16} + X^4 + X + 1$	63	0.9	2		✓	
$X^{16} + X^8 + X^4 + X^2 + X + 1$	31	0.9	2	✓		$X^{15} + X^7 + X^3 + X + 1$
$X^{16} + X^{12} + X^{11} + X^8 + X^6 + X^4 + X^3 + X^2 + X + 1$	21	0.9	2	✓		$X^5 + X + 1$
$X^{16} + \dots + X^2 + X + 1$	17	0.9	2,3,5,7	✓		
\vdots						

Table B.2: Classification of the outputs of the algorithm

Appendix C. Proof of the third family of multiplicative QSPs

We will now demonstrate that the third family of Proposition 6 is a family of multiplicative QSPs. We recall that it is defined as $X^{p^{n'}} - X^a \in \mathbb{F}_{p^n}$ with

- $p = k^n - k - (-1)^n$ prime, $n > 2$ and $k > 1$ integers such that $k^n \gg 1$.
- $n' = n - 1$
- $r = \frac{(p^n - 1)(k - (-1)^n)}{(k^n - k)(k^n - (-1)^n)}$
- $a := p^{n'} \pmod r$

We will prove that $r|p^n - 1$, provide an explicit formula for a and show that $\beta := n \log_p a / (n')^2 \leq 1$.

We will first show that $r = \frac{(p^n - 1)(k - (-1)^n)}{(k^n - k)(k^n - (-1)^n)}$ is an integer dividing $p^n - 1$. We can notice that $(k^n - k)|p^n - 1$. Indeed we have :

$$\begin{aligned} (p^n - 1) &= ((k^n - k) + (-1)^{n+1})^n - 1 \\ &= \sum_{i=1}^n \binom{n}{i} (k^n - k)^i (-1)^{(n+1)(n-i)} + (-1)^{n(n+1)} - 1 \\ &= (k^n - k) \sum_{i=0}^{n-1} \binom{n}{i} (k^n - k)^i (-1)^{(n+1)(n-i)} \end{aligned}$$

Similarly, $(k^n - (-1)^n)|p^n - 1$ since

$$\begin{aligned} (p^n - 1) &= ((k^n - (-1)^n) - k)^n - 1 \\ &= \sum_{i=1}^n \binom{n}{i} (k^n - (-1)^n)^i (-k)^{n-i} + (-k)^n - 1 \\ &= (k^n - (-1)^n) \left(\sum_{i=0}^{n-1} \binom{n}{i} (k^n - (-1)^n)^i (-k)^{n-i} + (-1)^n \right) \end{aligned}$$

Therefore $(k^n - k)(k^n - (-1)^n) / \gcd(k^n - k, k^n - (-1)^n)$ is an integer dividing $p^n - 1$. Moreover,

$$\begin{aligned} \gcd(k^n - k, k^n - (-1)^n) &= \gcd((k^n - (-1)^n) - (k^n - k), k^n - (-1)^n) \\ &= \gcd(k - (-1)^n, k^n - (-1)^n) \\ &= k - (-1)^n \text{ since } k - (-1)^n | k^n - (-1)^n. \end{aligned}$$

Hence $p^n - 1 = r \cdot \frac{(k^n - k)(k^n - (-1)^n)}{\gcd(k^n - k, k^n - (-1)^n)}$ and thus $r|p^n - 1$

Since the value of r depends on the parity of n , we distinguish two cases for the remaining of the proof.

If n is even. We now show that $a := (p^{n'} \bmod r) = \frac{p^{n-1}+1}{k^n-k}$. Indeed, $\frac{p^{n-1}+1}{k^n-k} = p^{n-1} - \frac{p^n-1}{k^n-k} = p^{n'} - r \frac{k^n-(-1)^n}{k-(-1)^n}$. Moreover, $\frac{p^{n-1}+1}{k^n-k} \simeq k^{n(n-1)-n} = k^{n^2-2n}$ while $r \approx k^{n^2+1-2n}$ for $k^n \gg 1$, so $\frac{p^{n-1}+1}{k^n-k} \simeq r/k$ and thus for k big enough, we have $\frac{p^{n-1}+1}{k^n-k} < r$.

Furthermore, $p^{n'} - a = \frac{p^n-1}{k^n-k} | p^n - 1$ so L splits over \mathbb{F}_{p^n} and it has $p^{n'} - a + 1 = \frac{p^n-1}{k^n-k} + 1 \simeq k^{n(n-1)}$ roots. This is close to $p^{n'}$ roots if $k^n \gg 1$.

Finally, $\frac{a^n}{p^{(n-1)^2}} \simeq \frac{(k^{n^2-2n})^n}{k^{n(n-1)^2}} = (k^n)^{(n^2-2n-n^2+2n+1)} = k^{-n} < 1$ when $k^n \gg 1$ so $a^n < p^{(n-1)^2}$, thus

$$\beta = n \log_p a / (n-1)^2 = (\log_p a^n) / (\log_p p^{(n-1)^2}) \leq 1.$$

If n is odd. We now show that $a = \frac{p^{n-1}k+1}{k^n+1}$.

Indeed $\frac{p^{n-1}k+1}{k^n+1} = p^{n-1} - \frac{p^n-1}{k^n+1} = p^{n'} - r \frac{k^n-k}{k-(-1)^n}$ and we have

$$\begin{aligned} r &= \frac{(p^n-1)(k+1)}{(k^n-k)(k^n+1)} \\ &= \frac{p^{n-1}}{k^n+1} (p(k+1)) \frac{1}{k^n(1-k^{1-n})} - \frac{1}{(k^n-k)(k^n+1)} \\ &= \frac{p^{n-1}}{k^n+1} (k^{n+1} + k^n + o(k^3)) k^{-n} (1 + k^{1-n} + o(k^{1-n})) + o(1) \text{ for } k^n \gg 1 \\ &= \frac{p^{n-1}}{k^n+1} (k+1 + o(k^{3-n})) (1 + k^{1-n} + o(k^{1-n})) + o(1) \\ &= \frac{p^{n-1}}{k^n+1} (k+1 + o(k^{3-n})) + o(1) \\ &= \frac{p^{n-1}k+1}{k^n+1} - \frac{1}{k^n+1} + \frac{p^{n-1}}{k^n+1} (1 + o(k^{3-n})) + o(1) \\ &= \frac{p^{n-1}k+1}{k^n+1} + \frac{p^{n-1}}{k^n+1} (1 + o(k^{3-n})) + o(1) \\ &= \frac{p^{n-1}k+1}{k^n+1} + \frac{p^{n-1}}{k^n+1} (1 + o(1)) \text{ since } n \geq 3, \\ &> \frac{p^{n-1}k+1}{k^n+1} \end{aligned}$$

Moreover, $p^{n'} - a = \frac{p^n-1}{k^n+1} | p^n - 1$ so L splits over \mathbb{F}_{p^n} and it has $p^{n'} - a + 1 = \frac{p^n-1}{k^n+1} + 1 \approx k^{n(n-1)} \approx p^{n'}$ roots if $k^n \gg 1$.

Furthermore,

$$\begin{aligned}
a^n &= \left(\frac{p^{n-1}k + 1}{k^n + 1} \right)^n = \left(\frac{p^{n-1}k}{k^n + 1} + o(1) \right)^n \\
&= \frac{p^{(n-1)^2 + n-1} k^n}{(k^n + 1)^n} + o\left(\frac{p^{n(n-1)} k^n}{k^{n^2}} \right) \\
&= p^{(n-1)^2} \frac{p^{n-1} k^n}{(k^n + 1)^n} + o(k^{n(n-1)^2}) \\
&= p^{(n-1)^2} \frac{(k^{n(n-1)} - (n-1)k^{1+n(n-2)} + o(nk^{1+n(n-2)}))k^n}{(1 + 1/k^n)^n} k^{-n^2} \\
&\quad + o(k^{n(n-1)^2}) \\
&= p^{(n-1)^2} (1 - (n-1)k^{1-n} + o(nk^{1-n}))(1 - k^{-n} + o(k^{-n})) + o(k^{n(n-1)^2}) \\
&= p^{(n-1)^2} (1 - (n-1)k^{1-n} + o(nk^{1-n})) + o(k^{n(n-1)^2}) \\
&< p^{(n-1)^2}
\end{aligned}$$

Thus $\beta := n \log_p a / (n-1)^2 = (\log_p a^n) / (\log_p p^{(n-1)^2}) \leq 1$.