# UNIVERSITY<sup>OF</sup> BIRMINGHAM University of Birmingham Research at Birmingham

# Sedentary behaviour and bone health in older adults

McMichan, L.; Dick, M.; Skelton, D. A.; Chastin, S. F.M.; Owen, N.; Dunstan, D. W.; Fraser, W. D.; Tang, J. C.Y.; Greig, C. A.; Agyapong-Badu, S.; Mavroeidi, A.

DOI: 10.1007/s00198-021-05918-2

*License:* None: All rights reserved

Document Version Peer reviewed version

#### Citation for published version (Harvard):

McMichan, L, Dick, M, Skelton, DA, Chastin, SFM, Owen, N, Dunstan, DW, Fraser, WD, Tang, JCY, Greig, CA, Agyapong-Badu, S & Mavroeidi, A 2021, 'Sedentary behaviour and bone health in older adults: a systematic review', *Osteoporosis International*, vol. 32, no. 8, pp. 1487-1497. https://doi.org/10.1007/s00198-021-05918-2

Link to publication on Research at Birmingham portal

#### Publisher Rights Statement:

McMichan, L., Dick, M., Skelton, D. et al. Sedentary behaviour and bone health in older adults: a systematic review. Osteoporos Int 32, 1487–1497 (2021). https://doi.org/10.1007/s00198-021-05918-2

#### **General rights**

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

•Users may freely distribute the URL that is used to identify this publication.

•Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.

•User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?) •Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

#### Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

1	Sedentary behaviour and bone health in older adults: a systematic review
2	Lauren McMichan <sup>1</sup> , Michael Dick <sup>1</sup> , Dawn A. Skelton <sup>2</sup> , Sebastien F.M. Chastin <sup>2,9</sup> , Neville
3	Owen <sup>3,4</sup> , David W. Dunstan <sup>3,5</sup> , William D. Fraser <sup>6</sup> , Jonathan C.Y. Tang <sup>6</sup> , Carolyn A. Greig <sup>7,8</sup> ,
4	Sandra Agyapong-Badu <sup>7</sup> , Alexandra Mavroeidi <sup>1</sup>
5	
6	<sup>1</sup> Department of Physical Activity for Health, School of Psychological Sciences and Health,
7	University of Strathclyde, Glasgow, UK
8	<sup>2</sup> Centre for Living, Department of Physiotherapy and Paramedicine, School of Health and
9	Life Sciences, Glasgow Caledonian University, Glasgow, UK
10	<sup>3</sup> Baker Heart and Diabetes Institute, Melbourne, Australia
11	<sup>4</sup> Centre for Urban Transitions, Swinburne University, Melbourne Australia
12	<sup>5</sup> Mary MacKillop Institute for Health Research, Australian Catholic University, Melbourne,
13	Australia
14	<sup>6</sup> Norwich Medical School, University of East Anglia, Norwich, UK
15	<sup>7</sup> School of Sport, Exercise and Rehabilitation Sciences, University of Birmingham,
16	Birmingham, UK
17	<sup>8</sup> MRC-Versus Arthritis Centre for Musculoskeletal Ageing Research, University of
18	Birmingham, UK
19	<sup>9</sup> Department of Movement and Sports Science, Ghent University, Ghent, Belgium
20	Correspondence: <u>lauren.mcmichan@strath.ac.uk</u> / <u>alexandra.mavroeidi@strath.ac.uk</u>
21	

22	Decl	larati	ons

#### 23 Conflict of Interest

- 24 Dawn A. Skelton is a Director of a not for profit training company, Later Life Training,
- 25 which provides training to health and leisure professionals to deliver strength and balance

26 training.

- 27 Lauren McMichan, Michael Dick, Sebastien F.M. Chastin, Neville Owen, David W. Dunstan,
- 28 William D. Fraser, Jonathan C.Y. Tang, Carolyn A. Greig, Sandra Agyapong-Badu,
- 29 Alexandra Mavroeidi declare no conflict of interests.

#### 30 **Disclosure**

- 31 Neville Owen is supported by a National Health and Medical Research Council of Australia
- 32 (NHMRC) Centre of Research Excellence Grant (#1057608), NHMRC Senior Principal
- 33 Research Fellowship (#1118225), and the Victorian Government's Operational Infrastructure
- 34 Support Program. David W. Dunstan is supported by a NHMRC Senior Research Fellowship
- 35 #1078360 and the Victorian Government's Operational Infrastructure Support Program.

36 Funding

- 37 This study was supported by a grant from the Royal Osteoporosis Society.
- 38
- 39
- 40
- 41
- 42

# 44 Abstract:

# 45 <u>Purpose:</u>

- 46 Older adults spend more than 8 hours/day in sedentary behaviours. Detrimental effects of
- 47 sedentary behaviour (SB) on health are established, yet little is known about SB and bone
- 48 health (bone mineral density; BMD) in older adults. The purpose of this review is to examine
- 49 associations of SB with BMD in older adults.
- 50 <u>Methods:</u>
- 51 Five electronic databases were searched: Web of Science (Core Collection); PubMed;
- 52 EMBASE; Sports Medicine and Education; and PsycInfo. Inclusion criteria were: healthy
- older adults mean age  $\geq 65$  years; measured SB; measured BMD using dual-energy X-ray
- 54 absorptiometry. Quality was assessed using National Institute of Health Quality Assessment
- 55 Tool for Observational Cohort and Cross-Sectional Studies.
- 56 <u>Results:</u>
- 57 After excluding duplicates 17,813 papers were assessed; 17,757 were excluded on
- title/abstract, 49 at full text, resulting in two prospective and five cross-sectional
- 59 observational studies reviewed. Four were rated 'good' and three were rated 'fair' using the
- 60 quality assessment criteria. Findings varied across the studies and differed by gender. In
- 61 women, four studies reported significant positive associations of SB with BMD at different
- 62 sites, and two found significant negative associations. Five studies which examined both men
- and women, men reported negative or no associations of SB with femoral neck, pelvic, whole
- 64 body, spine or leg BMD.

# 65 <u>Conclusion:</u>

- 66 While these findings suggest differences between men and women in the associations of SB
- 67 with BMD, they may be due to the varying anatomical sections examined for BMD, the
- 68 different methods used to measure SB, the varied quality of the studies included and the
- 69 limited number of published findings.
- 70

# 71 Keywords:

72 Sedentary behaviour, older adults, bone health, bone mineral density

- 74
- 75

# 76 Abbreviations:

- 77 BMD = Bone Mineral Density
- 78 DXA = Dual-X-Ray Absorptiometry
- 79 FN = Femoral Neck
- 80 LPA = Light Physical Activity
- 81 LS = Lumbar Spine
- 82 MVPA = Moderate-to-Vigorous Physical Activity
- 83 PA = Physical Activity
- 84 SB = Sedentary Behaviour
- ST = Sedentary Time
- TB = Total Body
- TF = Total Femur
- TH = Total Hip
- TS = Total Spine
- 90
- 91
- 92
- 93
- 94
- 95
- 96
- 97
- \_ \_
- 98
- 99

#### Introduction

Sedentary behaviour (SB) can be defined as "any waking activity characterized by an 101 102 energy expenditure  $\leq 1.5$  metabolic equivalents (METs), while in a sitting, reclining or lying posture" [1]. High volumes of SB can be detrimental to health, particularly in people who do 103 not undertake sufficient amounts of moderate-vigorous physical activity (MVPA) [2, 3]. 104 105 Adverse health consequences include higher risk of cardiovascular disease [4], diabetes mellitus [5], and reduced cognitive function [6]. SB encompasses many behaviours 106 performed routinely throughout the day, for example, sitting at a chair/sofa, driving to and 107 from places, and watching television [7]. Older adults (aged 65+ years) can accumulate > 8 108 hours of time spent in SB daily [8, 9], with an average being 9.4 hours/day [10]. 109 110 The beneficial effects of daily weight bearing physical activities on bone health are 111 well established [11]. Aging is a natural process, within which bone mass deterioration occurs, including changes to the structure and composition [12] of bone tissue. Although 112 some bone loss is typical of the aging process, osteoporosis is not an inevitable disease of the 113 old, with many risk factors for osteoporosis and osteoporotic fracture being modifiable [13]; 114 low levels of PA have long been recognised as such a risk factor. A published consensus 115 statement on Exercise and Osteoporosis recommends meeting the PA guidelines 116 (accumulation of 150 mins/week of moderate PA) for health and reducing prolonged SB 117 118 alongside more specific recommendations for exercise (resistance training and impact) [14]. There is a high prevalence of fractures in those over the age of 50 years, with one in two 119 women and one in five men fracturing a bone [15]. It is estimated that 500,000 fragility 120 fractures occur in the United Kingdom every year [16], with hospital costs of hip fractures 121 alone estimated at £1.1 billion [17]. 122

Mechanical forces (through gravitation or muscular loading) are essential for the 123 maintenance of bone health, therefore reducing these forces can have a detrimental effect 124 [18]. Space flight and bed rest studies have shown that reducing mechanical forces leads to 125 substantial reductions in bone strength [19]. Although the space-flight and bed-rest evidence 126 is from extreme and unusual circumstances, SB also involves the reduction of mechanical 127 forces, and could have a detrimental effect of bone health. A recent systematic review 128 129 explored the effects of SB on bone health in children, adolescents and young adults [20]. The review yielded 17 studies. It was reported that there was a moderately negative association 130 131 between SB and bone health in the lower extremities. It was also reported that one less hour of sedentary time mimics the positive effect of 18 minutes of MVPA in femoral neck bone 132 mineral density (BMD); however, this finding was weighted heavily on one strong 133 longitudinal study in boys [21]. 134 Gender also appears to play an important role on bone quality, with men exhibiting up 135 136 to 20% higher BMD compared to women [22]. This was evident in a study which analysed

data from the National Health and Nutrition Examination Study where negative associations
between SB and hip BMD in adult women, but not men, were identified [23].

Despite the emerging evidence on the potential detrimental influence that SB may
have on skeletal health in younger populations, little is known about the associations of SB
and bone health (specifically, BMD) in older adults. Therefore, the purpose of this study was
to systematically review the evidence on associations of SB with BMD (total and sitespecific) in older adults.

144

145 Methods

# 146 <u>Protocol and Registration</u>

147	The protocol for this systematic review was registered on Prospero
148	[CRD42019138999] in June 2019. The review was modelled using the PRISMA guidelines
149	[24, 25].
150	
151	Eligibility Criteria
152	Studies which explored the associations of SB on BMD in healthy older adults (mean
153	age $\geq$ 65 years old) were included in the review. Other inclusion criteria were studies which
154	measured SB, and measured BMD using Dual-Energy X-ray Absorptiometry (DXA). Studies
155	were peer reviewed, and in the English language.
156	
157	Information Sources
158	Five electronic databases were searched: Web of Science (Core Collection), PubMed,
159	EMBASE, Sports Medicine and Education, and PsycInfo. The search strategy was originally
160	conducted in March 2019. The search strategy was then repeated using additional search
161	terms to broaden the search results. This was conducted in June 2019.
162	
163	Search
164	The search strategy used for the databases is shown in Table 1. Note adaptations to
165	truncations and limiting factors were made based on the individual databases.
166	

Population	(Adult* OR "Older Adult*" OR Elderly OR Geriatric OR Ageing OR					
	Aged)					
Search Operator	AND					
Outcome	(Bone OR "Bone Health" OR "Bone Mineral Density" OR "Bone					
	Mineral" OR "Bone Mass" OR "Bone Fracture*" OR "Bone Strength"					
	OR Osteoporosis OR "Bone Mineral Content")					
Search Operator	AND					
Exposure	(Sedentar* OR "sedentary behavior" OR "Sedentary Behaviour" OR					
	"Sedentary Time" OR "Sitting Time" OR Sitting OR "screen time" OR					
	"television viewing" OR inactiv* OR "activity restriction" OR					
	"Computer use" OR "stationary behaviour" OR "stationary behavior"					
	OR lying OR reclining OR "non-screen based behaviour" OR "non-					
	screen based behavior")					
Limits	English only. Humans only.					
Study Selev	Study Selection					
<u>Study Selec</u>						
Articles ret	rieved from the search strategy were imported into EndNote Reference					
Manager, version 2	X8.2 (Thomson Reuters, Philadelphia, PA) and duplicates were removed.					

\_

Articles were then exported to a Microsoft Excel, version 2016 (Microsoft Corp, Redmond, 172 WA) spreadsheet where titles were screened. Articles included based on title screening were 173 reviewed at abstract level, and then reviewed as full text. We also reviewed the bibliography 174 of full text papers to identify any additional related papers. All articles were reviewed by the 175 first author (LM) and a sample (10%) was double checked by another reviewer (AM) as per 176 PROPSERO protocol. Any articles where there was uncertainty at abstract and full text level 177 178 were also checked by the senior author (AM). If there were any discrepancies, a discussion between the two authors was conducted until an agreement was reached. Exclusion of 179 180 articles were based on criteria and were excluded if they did not assess SB and BMD in older adults (mean age  $\geq 65$  years). Studies which included multiple age ranges but performed a 181 sub-analysis on older adults were included in the review. 182

183

184

#### Data Collection Process and Data Items

Data were extracted and imported into a standardised Microsoft Word, version 2016 (Microsoft Corp, Redmond, WA) table. Data extracted were: author(s)/year of publication; study design; sample size; gender; age range; SB measurement method and outcomes; BMD measurement methods and outcomes; overall results. Were pivotal data was missing we aimed to contact the authors and request such data.

190

# 191 <u>Quality Assessment</u>

192 Quality of studies included in the review was assessed using the 'Quality Assessment
193 Tool for Observational Cohort and Cross-Sectional Studies [26]. The tool consists of a 14194 item checklist: clearly stated research question; specific study population; rate of
195 participation of eligible persons; subject selection process; justification of sample size;

exposure measured prior to outcome(s); suitable timeframe between exposure and outcome;
levels of exposure; exposure measures clearly defined; exposure(s) assessed more than once
over time; outcome measures defined valid, reliable and consistent; blinding of outcome
assessors; loss to follow-up; and, adjustment for key confounders. Studies were then awarded
a rating of good, fair or poor. Quality was assessed by the first and last authors (LM/AM) to
ensure agreement. It should be noted that one of the co-authors (SFMC) was the co-author of
3 of the included studies.

203

#### 204 Summary of Measures

Primary exposure measures were SB (self-reported by questionnaire or objectively measured) and BMD measures were the outcomes. Studies included objective and subjective methods of assessing SB. The anatomical sites that were evaluated were grouped into three separate categories; lower extremities (including the femoral neck (FN), total femur (TF), hip, legs), trunk (including ribs, lumbar spine (LS)) and total body (TB). Markers of bone health were measured using DXA in all included studies since comparison of bone health is not possible if different assessments methods are used.

212

213 **Results** 

#### 214 <u>Study Selection</u>

The initial search strategy yielded 19,194 potentially relevant studies (Figure 1.). Following deduplication, this number was reduced to 17,813. Seventeen thousand seven hundred fifty seven articles were excluded based on title and abstract. Forty-nine articles were excluded at full text (see supplementary material for complete list of excluded studies),

219	leaving seven included for review [23, 27-32]. One of those studies (Chastin et al [23]) did
220	not present data for the over 65s separately in their published manuscript. However, as SFMC
221	is a co-author in the current review, he was able to repeat the main paper analysis from the
222	NHANES database refining it to those aged over 65 years; this sub analysis was included in
223	the current review.

232

### 225 <u>Study Characteristics</u>

Five studies were cross-sectional [23, 27, 29-32] and two were longitudinal [28, 29]. Sample size ranged from 112 [30] to 1134 [29] participants. Two studies included women only [27, 29]. Mean age of participants ranged from  $64.5 \pm 7.2$  years [28] to  $76.9 \pm 5.3$  years (men) and  $76.7 \pm 4.7$  years (women) [31]. A full summary of study characteristics and results are shown in Table 2. Four studies were rated good [23, 28-30] for quality, whilst three were rated fair [27, 31, 32].



Fig 1 PRISMA diagram [25] of the screening process

#### 234

# Measurement of BMD and SB

235 All studies measured BMD using DXA. Six studies used Hologic models [23, 27, 28, 30-32], whilst one used a Lunar model [29]. Measures of SB differed between studies. Two studies 236 used questionnaires to assess SB [27, 29]. The questionnaires asked how many hours per day 237 participants were sedentary. Five studies assessed SB using objective measures 238 (accelerometry) [23, 28, 30-32]. Two studies [31, 32] used ActiTrainer/ActiGraph wGT3X-239 240 BT, two used used ActiGraph GT1M [23, 28], and one used GENEActiv Action [30]. Accelerometer placement was different between studies, with the majority of the studies 241 [23, 31, 32] using hip mounded accelerometers and one study [30] using a leg mounded 242 243 accelerometer. All studies asked participants to wear their accelerometers for 6-7 consecutive 244 days but some participants had to remove their accelerometer during waking hours when engaging in water based activities [23, 31, 32]. Every study had its own unique wear-time 245 protocol. For example, data were excluded from the analysis if participants did not wear 246 device for at least 5 days and wore the device for less than 10 hours per day [23, 28] but these 247 criteria were different for the studies by Rodrigues- Gomez et al who included only results 248 with at least 4 valid days that included at least 8h/day of wear time [31, 32]. 249

250

#### 251 <u>Total Body BMD</u>

Four studies measured total BMD [28, 30-32]. McMillan et al [28] found no significant
associations over time between SB and total BMD in either men or women, using prospective
linear regression analyses.

Rodriguez-Gomez et al [31] reported a borderline significant positive association
between SB and total body BMD when both genders were analysed together. There were

gender differences when analyses were separated. There were no significant associations 257 between SB and total body BMD for men. For women, significant positive associations were 258 found between SB and total body BMD ( $\gamma = 0.022$ ; p = 0.00). Gender differences were also 259 reported in Rodriguez-Gomez et al [32] who found a significantly negative association 260 between SB and total body BMD ( $\gamma = -0.015$ ; p = 0.041) in robust healthy men, but reported 261 significant positive associations between SB and total body BMD ( $\gamma = 0.020$ ; p = 0.003) in 262 263 robust healthy women. It is important to note that the two Rodriguez-Gomez [31, 32] analyses are based on the same cohort and models from both studies were adjusted for age, 264 265 gender, BMI, fat and lean mass, alcohol, smoking, nutrition, calcium, education, level of income, marital status, frailty, arthritis, and thyroid disease. 266

Onambele-Pearson et al [30] also found differences between men and women. For 267 daily SB, there were no significant associations with total BMD for men, whilst there was a 268 significant positive association (r = 0.317, p < 0.01) for women. However, when analyses 269 270 were adjusted (age, total fat mass, general anthropometry) this was non-significant. Onambele-Pearson et al [30] also explored the association between SB bouts and BMD. 271 Whilst breaks in SB did not have a significant impact on total BMD in women, there was a 272 positive association in men (r = 0.330, p < 0.01). There was also positive association in men 273 when SB bouts were  $\geq 5 \text{ mins}$  (r = 0.373, p < 0.01). 274

275

# 276 <u>Lower Extremities</u>

In a prospective study, Nguyen et al [29] reported that, over time, sedentary lifestyle significantly reduced BMD in the femoral neck (FN) (-1.5  $\pm$  0.2%, p < 0.001) in women. Adjusted analyses (accounting for age, PA, baseline weight, weight change over time and baseline BMD) also suggested a significant reduction in femoral neck BMD (-1.35  $\pm$  0.8%, p 281 < 0.05). The other longitudinal study included in the review (McMillan et al [28]) also</p>
282 conducted analyses separately by gender. Their adjusted (model 2) prospective multivariate
283 linear regression analyses, showed a significantly positive association between SB and total
284 hip BMD ( $\beta = 0.199$ , p = 0.046) in women. For adjusted (model 2) prospective multivariate
285 linear regression analyses in men, there were significantly negative associations between SB
286 and femoral neck BMD ( $\beta = -0.232$ , p = 0.047). Model 2 analyses were adjusted for age,
287 height, lean mass and smoking.

Onambele-Pearson et al [30] reported no significant associations for daily SB for men and BMD ; in women, there were significant positive associations between daily SB and lower limb BMD (r = 0.272,  $r_{adj}^n = 0.260$ , p < 0.05). Similar findings were reported for breaks in SB (r = 0.299,  $r_{adj}^n =$  non-significant, p < 0.05), and for bouts of SB < 5 mins (r = 0.334,  $r_{adj}^n =$  non-significant, p < 0.01). There was also a positive association between breaks in SB and pelvic BMD in women only (r = 0.232,  $r_{adj}^n =$  non-significant, p < 0.05).

Braun et al [27] reported a negative association in older women between ST and femoral BMD (b (SE) = -0.0028 (0.0001);p = 0.027). These analyses were adjusted for race/ethnicity, milk consumption or supplement use, BMI, smoking, osteoporosis history, prednisone or cortisone use, and menopausal status.

Rodriguez-Gomez et al [31] reported significant positive associations between SB and leg leg BMD ( $\gamma = 0.028$ , p = 0.00). There were no significant associations reported for any femoral region assessed. When men and women were examined separately, there were no associations between SB and leg/femoral region BMD in men. In women, there were significantly positive associations for leg BMD ( $\gamma = 0.063$ , p = 0.00), but no associations for femoral regions. In the pelvic region, a significant negative association was reported between SB and BMD ( $\gamma = -0.027$ , p = 0.05) in men only.

Rodriguez-Gomez et al [32] reported significant positive associations between SB and leg BMD ( $\gamma = 0.035$ , p = 0.000) in robust (those who do not exhibit any of the frailty criteria set out by Fried et al [33]) older adults. No associations were reported for femoral neck. When analyses were conducted separately for men and women, men were found to have negative associations for leg BMD ( $\gamma = -0.018$ , p = 0.036). However, in women there were positive associations between SB and leg BMD ( $\gamma = 0.066$ , p = 0.000). No associations were reported for femoral neck in either gender.

312 <u>Trunk</u>

Only two studies reported significant associations between SB and areas of the trunk 313 [23, 30]. In women, Onambele-Pearson et al [30] reported significant positive association 314 315 between daily SB and spine BMD (r = 0.233, p < 0.05), although this was non-significant 316 when adjusted for confounders. There were significant positive associations between SB breaks and SB bouts of < 5 minutes, and BMD of the ribs (r = 0.266, p < 0.05; r = 0.328, p <317 318 0.01, respectively). The association between breaks in SB and BMD were non-significant when adjusted for confounders. There was a negative association between W50% min 319 (defined as "the bout duration below which half of all sedentary time is accrued" [30]) and rib 320 BMD (r = -0.224, p < 0.05; non-significant after adjusting). For men, breaks in SB and SB 321 bouts  $\geq$  5 minutes were positively associated with rib BMD (r = 0.282, p < 0.05; r = 0.349, p 322 323 < 0.01, respectively).

A significant negative association with sedentary time (ST – total duration of daily SB bouts) and spine BMD with (p = 0.05) and a significant positive association between fragmented ST and spine BMD (p = 0.05) were reported for the unpublished sub-analysis of the NHANES data by Chastin et al [23], yet there were non-significant differences between

328	women and men. Analyses were adjusted for age, BMI, ethnicity, parathyroid hormones,
329	smoking, alcohol (men only) and prednisone use (women only). (Table 2).
330	
331	Quality Assessment
332	Of the seven studies included in the review, four were rated good [23, 28-30] and
333	three were rated fair [27, 31, 32]. Table 3 provides a full summary of each study and the
334	criteria which the assessment was based on.

Author	Sample size, gender (♂/♀), age (years)	SB outcomes measured	SB assessment method	Bone assessment method	Sites of anatomical assessment	Conclusions/results			
	CROSS-SECTIONAL STUDIES								
Chastin et al (sub analysis of ref [23]	591 (n =259 ♀) Age = 75.2 ± 6.7	Total ST (Daily) Bouts of SB	Accelerometry (ActiGraph GT1M)	DXA	TF, FN, Trochanter Ward triangle, Intertrochanter, Spine	No significant association for men or women between ST or bouts of SB with TF, FN, Trochanter Ward triangle, Intertrochanter BMD. Significant negative association with spine BMD for ST (p =0.05), sub analysis per gender leads to non-significant results. Significant positive association between more fragmented ST			
						(shorter bout duration) and spine BMD (p=0.05), sub analysis per gender leads to non-significant results.			
Onambele- Pearson et al [30]	112 (n = 61 $\bigcirc$ ) Age = 72.5 ± 6.4	ST (hours/day), Bouts of SB	Accelerometry (GENEActiv Action)	DXA	Ribs, Spine, Pelvis, Upper Limbs, Lower Limbs, Total Body	$\delta$ = Significant positive association between breaks in SB and BMD for ribs (p < 0.05) and total BMD (p < 0.01). SB bouts > 5 minutes were positively associated with lower limbs (p < 0.05), ribs and total BMD (p < 0.01). No significant associations reported for total ST.			
						Q = Significantly positive association between total ST and spine, lower limb (p < 0.05) and total (p < 0.01) BMD. Significant positive correlation between breaks in SB and ribs, pelvis and lower limbs (p < 0.05). For bouts < 5 minutes, there were positive associations for ribs and lower limbs (p < 0.01). There was a negative association between W50% and rib BMD (p < 0.05).			
Rodriguez-Gomez et al [31]	871 (n =476 ♀) ♂ age = 76.9 ± 5.3 ♀ age = 76.7 ± 4.7	Total ST, separately for $\vec{c}$ and $\hat{c}$	Accelerometry (ActiTrainer & Actigraph wGT3X-BT)	DXA	TB, LS (L1-L4), FN, TH (greater trochanter, inter trochanter, Ward's	$^{\diamond}$ ♀: SB positively associated with leg /BMD (p = 0.00) and whole body BMD (p = 0.05). $^{\diamond}$ : SB negatively associated with pelvic BMD (p = 0.05)			
					triangle) and FN), BMD &	$\mathbb{Q}$ : SB was positively associated with whole body /BMD (;p = 0.00) and leg /BMD (p = 0.00).			
						$\bigcirc$ : Reduce time spent in SB to reduce fracture risk.			

# **Table 2** Overview of study attributes and findings on sedentary behaviour with bone outcomes

Rodriguez-Gomez et al [32]	540 (n = 289 ♀), ♂&♀ Age = 76.0 ± 4.4 (robust individuals only)	Total ST	Accelerometry (ActiTrainer & Actigraph wGT3X-BT)	DXA	TB, LS (L1-L4), FN, TH, Leg BMD &	SB significantly positively associated with leg BMD/ (p = 0.000) for whole sample. $\Im$ SB significantly negatively associated with TB BMD (p = 0.041) and leg BMD/ (p = 0.036) in robust men. $\Im$ SB significantly positively associated with TB BMD/ (p = 0.000) and leg BMD/ (p = 0.000) in robust women.
Braun et al [27]	327 ♀ only Age ≥ 65	SB (mins/day)	Questionnaire	DXA	FN, LS (L1-L4) (trochanter, inter trochanter, Ward's triangle, TF, TS)	SB significant predictor of decreased Femoral BMD ( $p = 0.027$ ), yet not a significant association in spinal BMD ( $p > 0.05$ ).
			LONGITUDINA	L PERSPECTI	VE STUDIES	
McMillan et al [28]	209, (n =111 ♀), Age = 64.5 ± 7.2	SB (mins/day), separately for $and$	Accelerometry (ActiGraph GT1M)	DXA	TH, LS, FN, Pelvis, Legs & TB	$\Diamond$ : Negative association between SB and FN /BMD ( p = 0.047) over 2.2 years. $\Diamond$ : SB was positively associated with TH /BMD (; p = 0.046) over 2.2 years.
Nguyen et al [29]	1134 $\bigcirc$ only n = 366 sedentary Age = 69.9 $\pm$ 7.4 <sup>†</sup> (N=1134; N = 827 at follow up)	ST (hours/day)	Questionnaire	DXA	FN	ST was significantly associated with reduced BMD $(-1.5 \pm 0.2\%, p < 0.001)$ . 47% of individual's with significant bone loss (n = 163) were categorised within the sedentary group.

336 Abbreviations: BMD bone mineral density, DXA dual-X-ray-absorptiometry, FN femoral neck, LPA light physical activity, LS lumbar spine, MVPA moderate-to-vigorous physical activity, PA

337 338 physical activity, SB sedentary behaviour, ST sedentary time, TB total body, TF total femur, TH total hip, TS total spine. † Mean age was calculated by as follows ((age of sedentary group x n

sedentary group) + (age of moderately active group x n moderately active group) + (age active group x n active group )/(total N of participants)).

Table 3 Summary of the quality assessment for each study using the National Institute of Health Quality Assessment Tool for Observational Cohort and Cross-Sectional
 Studies\*.

	Chastin et al [23]	Braun et al [27]	McMillan et al [28]	Nguyen et al [29]	Onambele- Pearson et al [30]	Rodriguez- Gomez et al [31]	Rodriguez- Gomez et al [32]
1) Was the research question or objective in this paper clearly stated?	Y	Y	Y	Y	Y	Y	Y
2) Was the study population clearly specified and defined?	Y	Y	Y	Y	Y	Y	Y
3) Was the participation rate of eligible persons at least 50%?	Y	Y	Y	Y	CD	CD	CD
4) Were all the subjects selected or recruited from the same or similar populations (including the same time period)? Were inclusion and exclusion criteria for being in the study prespecified and applied uniformly to all participants?	Y	Y	Y	Y	Y	Y	Y
5) Was a sample size justification, power description, or variances and effect estimates provided?	Ν	Ν	Ν	Ν	Y	Ν	Ν
6) For the analyses in the paper, were the exposure(s) of interest measured prior to the outcome(s) being measured?	CD	CD	CD	CD	NA	NA	NA
7) Was the timeframe sufficient so that one could reasonably expect to see an association between exposure and outcome if it existed?	NA	NA	Y	Y	NA	NA	NA
8) For exposures that can vary in amount or level, did the study examine different levels of exposure as related to the outcome (e.g. categories of exposure, or exposure measured as continuous variable)?	Y	N	NA	NA	Y	Y	Y
9) Were the exposure measures (independent variables) clearly defined, valid, reliable, and implemented consistently across all study participants?	Y	Y	Y	Y	Y	Y	Y
10) Was the exposure(s) assessed more than once over time?	NA	Ν	Y	Y	NA	NA	NA
11) Were the outcome measures (dependent variables) clearly defined, valid, reliable, and implemented consistently across all study participants?	Y	Y	Y	Y	Y	Y	Y
12) Were the outcome assessors blinded to the exposure status of participants?	NA	NA	NA	NA	NA	NA	NA
13) Was loss to follow up after baseline 20% or less?	NA	NA	Y	Ν	NA	NA	NA
14) Were key potential confounding variables measured and adjusted statistically for their impact on the relationship between exposure(s) and outcome(s)?	Y	Y	Y	Y	Y	Y	Y
Rating	Good	Fair	Good	Good	Good	Fair	Fair

342 \* Y = Yes, N = No, CD = Cannot Determine, NA = Not Applicable

#### Discussion

This review examined the associations between sedentary behaviour (SB) and BMD
in older adults (mean age ≥ 65 years). Following the screening of potentially relevant articles,
seven were included for review. Studies varied with respect to: measurement of SB
(subjective vs objective measures); anatomical measures of BMD; and whether they were
cross-sectional or longitudinal in nature. All studies used DXA as a method to assess BMD.

#### 349 <u>Summary of Evidence</u>

Longitudinal studies [28, 29] that explored the associations of SB on BMD over time, reported negative associations between higher levels of SB with BMD for both sexes and at all sites measured (except for total hip BMD in the female subgroup of the McMillan study [28]). Disparities may have arisen due to the larger sample size of women in Nguyen's [29] study in comparison to McMillan et al [28] and both studies measured SB differently (subjectively vs objectively, respectively).

The results from the cross-sectional studies however revealed a different pattern 356 between men and women. Some verified the deleterious associations of higher levels of SB 357 on BMD [27] in women, while others failed to identify a significant association when sub 358 analysis by gender took place [23] (possibly due to the smaller numbers of participants in 359 each subgroup). However; the majority of the studies [30-32] found a disparity in results 360 between men and women. In women, positive associations were observed between SB and 361 BMD (at different measuring sites) while the opposite was true for their male counterparts. It 362 appears that, as discussed by Rodriguez-Gomez et al [31], other movement behaviours could 363 364 have an impact on the SB and BMD association. It is common to find concomitant higher SB and higher MVPA levels, thus the MVPA could have a positive association on BMD in the 365 presence of high levels of SB [31]. In addition, it was also highlighted that greater sedentary 366

time could result in more frequent breaks in sitting time. More frequent postural changes
could result in higher volumes of mechanical load bearing, thus resulting in a positive
association on BMD [31].

Likewise, Onambele-Pearson et al [30] reported higher BMD in women who reported 370 more frequent sedentary bouts and identified that this frequency of interruption to SB could 371 372 contribute the higher BMD. This is consistent with other research in post-menopausal women, whereby a greater number of breaks in SB resulted in a 10% reduction in the odds of 373 being diagnosed with osteoporosis/osteopenia [35]. A small study in frailer older adults found 374 that breaking SB on a roughly hourly basis throughout the day improved physical function 375 (timed up and go and 30 second chair stand) over a 10-week period, with no significant 376 change in total ST or PA [36]. Similarly, Aunger et al [37] reported clinically significant 377 improvement in physical function with non-significant increases in daily steps and time spent 378 379 upright, despite non-significant decreases in SB. This modest body of emerging evidence 380 suggests that regular interruptions to ST may be beneficial to bone health in older adults. Further investigation is warranted, with a particular emphasis being on the wider application 381 of thigh-worn accelerometers which have been shown to have higher accuracy for detecting 382 postural changes than wrist and waist-worn accelerometers. 383

384 Measurement of SB also varied between the studies included in this review. The two 385 studies that measured SB subjectively using questionnaires, reported negative associations between femoral BMD and SB [27, 29] in women, which appears to contradict the findings 386 of objectively measured studies. This could be attributed to the underestimation of SB and 387 388 overestimation of PA; a bias which is commonly acknowledged when subjectively measuring PA [39]. It is reported that a random error of 2.5 hours per day is observed in subjective 389 390 assessment of total SB and that subjective measures are not valid in assessing SB bouts [40]. Studies that used objective measures of SB reported more positive associations between SB 391

392	and BMD in women [28, 30-32]. It has been recently suggested, that moving to a single SB
393	question assessing the whole day (via means of a visual analogue scale) might be worth
394	considering for future studies, in cases where the use of a device-based measurement of PA is
395	not practically possible [41].

#### 397 <u>Strengths and Limitations</u>

This is the first review to explore the associations between SB and BMD in healthy 398 399 older adults. However, there are a number of limitations influencing interpretation of the 400 study findings. We appreciate that in order to reduce the risk of bias ideally two independent reviewers should have carried out all the steps of study selection and data extraction. In this 401 402 study and due to time and resource limits the primary author (LM) screened the titles and abstracts, and excluded any irrelevant articles and only a sample (10%) of the studies were 403 checked by one other reviewer (AM). As per PROSPERO protocol, when there were 404 discrepancies in the inclusion or exclusion criteria discussion took place until an agreement 405 406 was made.

407 Of the studies reviewed, two were prospective, and five were cross-sectional in design, as such any associations found here are not of a causal inference, and the possibility 408 of bi-directional associations in the cross sectional studies cannot be ruled out. It should also 409 be noted that the reported significant associations between SB and BMD, do not necessarily 410 translate to clinically important associations and thus caution should be applied when 411 interpreting these for such use. The generalisability of the results from this study should also 412 413 be considered in light to the moderate quality of the studies included in the review, and the low numbers of participants in the subgroup (gender specific) analysis. In addition only 414 healthy adults were assessed (clinical population were excluded from this review), therefore 415

there was no analysis on different populations and different health status in older adults. Thisincluded omitting the analysis of those who were deemed as 'frail' [32].

Likewise, the role of body weight as a confounder was not analysed in detail as part 418 of this review. There is extensive body of findings suggesting that a higher body weight or 419 body mass index can be associated with higher BMD [42, 43, 44] and reduced fracture risk 420 421 [45]. These associations are probably attributable to the accentuated mechanical loading on the skeleton due to the increase in body mass although the exact mechanism is still not fully 422 understood [46]. Indeed future studies should interpret data in the context of a number of 423 confounders (including body weight and/or BMI) but also comorbidities, which are common 424 in this older population and can induce sedentary behaviour. 425

426 There were various anatomical sites assessed using DXA and various methods of monitoring 427 objectively and subjectively SB (different accelerometer types and questionnaires); in the absence of standardised assessments what may be concluded from the findings of the studies 428 429 is limited. Although BMD measurement remains the most useful diagnostic tool for identifying patients with osteoporosis other technologies (e.g. ultra-high-resolution peripheral 430 QCT, and 3D magnetic resonance imaging [MRI]) could noninvasively assess bone cross-431 sectional geometry and trabecular architecture. The combination of these, as well the 432 433 assessment of number of fractures, may provide a more comprehensive picture of bone 434 strength/health, compared with 2-dimensional BMD measurements in future studies. In addition this review included studies that used different densitometers to assess BMD, which 435 is inevitably a limitation due to the well-established inherent measurement differences between 436 scanners. In order to make progress in this field, we need well-designed longitudinal studies 437 in this age group, with objective measures of SB and PA, and assessment of bone outcomes 438 beyond just DXA. 439

#### Conclusion

This systematic review aimed to determine the associations between SB and BMD in 441 healthy older adults (mean age  $\geq 65$  years). In conclusion, the research suggests there are 442 gender difference in the associations of SB with BMD, with SB seemingly positive 443 association on BMD in older women, but having a negative or no association in older men. 444 445 However, there were only seven studies included in the review, with men being assessed in five of those studies, thus limiting the conclusions that can be drawn and thus these gender 446 specific results should be treated with caution though due to the inherited issues with the 447 relative small numbers of participants in subgroup analyses In order to better understand the 448 associations of SB on BMD in older adults, there is a particular need to examine variations in 449 patterns of sedentary time, using objective measures, including sit-stand transitions and how 450 451 these might vary between men and women.

452

#### 453 **References**

Tremblay MS et al (2017) Sedentary Behavior Research Network (SBRN) –
 Terminology Consensus Project process and outcome. Int J Behav Nutr Phys Act
 14:75-91.

457 2. Tremblay MS, Colley RC, Saunders TJ, Healy GN, Owen N (2010) Physiological and
458 health implications of a sedentary lifestyle. Appl Physiol Nutr Metab 35:725-740.

3. Biswas A, Oh PI, Faulkner GE et al (2015) Sedentary time and its association with
risk for disease incidence, mortality, and hospitalization in adults: a systematic review
and meta-analysis. Ann Intern Med 162:123-132.

# 462 4. Chrysant SG, Chrysant GS (2015) The cardiovascular consequences of excess sitting 463 time. J Clin Hypertens 17:528-531.

464	5.	Hu FB, Li TY, Colditz GA, Willett WC, Manson JE (2003) Television watching and
465		other sedentary behaviours in relation to risk of obesity and type 2 diabetes mellitus in
466		women. J Am Med Assoc 289:1785-1791.
467	6.	Falck RS, Davis JC, Liu-Ambrose T (2017) What is the association between
468		sedentary behaviour and cognitive function? A systematic review. Br J Sports Med
469		51:800-811.
470	7.	Chastin SFM, Schwarz U, Skelton DA (2013) Development of a consensus taxonomy
471		of sedentary behaviors (SIT): Report of Delphi Round 1. Plos One 8:e82313.
472	8.	Copeland J, et al (2017) Sedentary time in older adults: a critical review of
473		measurement, associations with health, and interventions. Br J Sports Med 51:1539-
474		1539.
475	9.	Harvey JA, Chastin SFM, Skelton DA (2013) Prevalence of sedentary behavior in
476		older adults: A systematic review. Int J Environ Res Public Health 10:6645-6661.
477	10.	Harvey JA, Chastin SF, Skelton DA (2015) How sedentary are older people? A
478		systematic review of the amount of sedentary behavior. J Aging Phys Act 23:471-87.
479	11.	Bonjour JP, Chevalley T, Ferrari S, Rizzoli R (2009) The importance and relevance of
480		peak bone mass in the prevalence of osteoporosis. Salud Publica Mex 51:S5-S17.
481	12.	Demontiero O, Vidal C, Duque G (2012) Aging and bone loss: new insights for the
482		clinician. Ther Adv Musculoskel 4:61-76.
483	13.	Abrahamsen B, Brask-Lindemann D, Rubin KH, Schwarz P (2014) A review of
484		lifestyle, smoking and other modifiable risk factors for osteoporotic fractures. Bone
485		Key Rep 3:574-580.
486	14.	National Osteoporosis Society (2018) Strong, Steady and Straight: An expert
487		consensus statement on physical activity and exercise for Osteoporosis.
488		https://www.bgs.org.uk/sites/default/files/content/attachment/2019-02-

489	20/FINAL%20Consensus%20Statement_St	rong%20Steady%20and%20Straight_DE
490	C18.pdf Accessed 21 May 2020.	
491	15. van Staa TP, Dennison EM, Leufkens HGM	I, Cooper C (2001) Epidemiology of
492	fractures in England and Wales. Bone 29:51	7-522.
493	16. Svedbom A, Hernlund E, Ivergard M, et al	(2013) Osteoporosis in the European
494	Union: a compendium of country-specific re	eports. Arch Osteoporos 8:137-355.
495	17. Leal J, Gray AM, Prieto-Alhambra D, et al	(2016) Impact of hip fracture on hospital
496	care costs: a population-based study. Osteop	poros Int 27:549-558.
497	18. Kohrt WM, Barry DW, Schwartz RS (2009)	) Muscle forces or gravity: What
498	predominates mechanical loading on bone?	Med Sci Sports Exerc 41: 2050–2055.
499	19. LeBlanc AD, Spector ER, Evans HJ, Sibong	ga JD (2007) Skeletal responses to space
500	flight and bed rest: A review. J Musculoske	let Neuronal Interact 7:33-47.
501	20. Keodijk JB, van Rijswijk J, Oranje, WA, et	al (2017) Sedentary behaviour and bone
502	health in children, adolescents and young ad	lults: a systematic review. Osteoporos Int
503	28:2507–2519.	
504	21. Vaitkeviciute D, Latt E, Maestu J, Jurimae	Г, Saar M, Purge P, Maasalu K, Jurimae J
505	(2014) Physical activity and bone mineral a	ccrual in boys with different body mass
506	parameters during puberty: a longitudinal st	udy. PLoS One 9:e107759
507	22. Naganathan V, Sambrook P (2003) Gender	differences in volumetric bone density: a
508	study of opposite-sex twins. Osteoporos Int	14:564–569.
509	23. Chastin S, Mandrichenko O, Helbostadt, J,	Skelton DA (2014) Associations between
510	objectively-measured sedentary behaviour a	and physical activity with bone mineral
511	density in adults and older adults, the NHA	NES study. Bone, 64:254-262.

512	24. Liberati A, Altman DG, Tetzlaff J, et al (2009) The PRISMA statement for reporting
513	systematic reviews and meta-analyses of studies that evaluate health care
514	interventions: Explanation and elaboration. BMJ 339:b2700.
515	25. Moher D, Liberati A, Tetzlaff, J Altman, DG, Grp P (2009) Preferred Reporting Items
516	for Systematic Reviews and Meta-Analyses: The PRISMA Statement. BMJ
517	339:b2535.
518	26. National Institutes of Health National Heart, Lung and Blood Institute (n.d.) Study
519	quality assessment tools. <u>https://www.nhlbi.nih.gov/health-topics/study-quality-</u>
520	assessment-tools. Accessed 30 September 2019.
521	27. Braun SI, Kim Y, Jetton AE, Kang M, Morgan DW (2015) Prediction of bone mineral
522	density and content from measures of physical activity and sedentary behaviour in
523	younger and older females. Prev Med Rep 2:300-305.
524	28. McMillan LB, Aitken D, Ebeling P, Jones G, Scott D (2018) The relationship
525	between objectively assessed physical activity and bone health in older adults differs
526	by sex and mediated by lean mass. Osteoporosis Int 29:1379-1388.
527	29. Nguyen TV, Sambrook PN, Eisman JA (1998) Bone loss, physical activity, and
528	weight change in elderly women: The Dubbo osteoporosis epidemiology study. J
529	Bone Miner Res 13:1458-1467.
530	30. Onambele-Pearson G, Wullems J, Doody C, Ryan D, Morse C, Degens H (2019)
531	Influence of habitual physical behavior – sleeping, sedentarism, physical activity – on
532	bone health in community-dwelling older people. Front Physiol 10:1-15.
533	31. Rodriguez-Gomez I, Manas A, Losa-Reyna J et al (2018) Associations between
534	sedentary time, physical activity and bone health among older people using
535	compositional data analysis. Plos One 13:1-17.

536	32. Rodriguez-Gomez I, Manas A, Losa-Reyna J et al (2019) The impact of movement
537	behaviours on bone health in elderly with adequate nutritional status: Compositional
538	data analysis depending on frailty status. Nutr 11:582-594.
539	33. Fried LP, Tangen CM, Walston J et al (2001) Frailty in older adults: Evidence for a
540	phenotype. J Gerontol Ser A Med Sci 56: M146-M156.
541	34. Braun SI, Kim Y, Jetton AE, Kang M, Morgan DW (2017) Sedentary behaviour,
542	physical activity, and bone health in postmenopausal women. J Aging Phys Activ
543	25:173-181.
544	35. Harvey JA, Chastin SFM, Skelton DA (2018) Breaking sedentary behaviour has the
545	potential to increase / maintain function in frail older adults. J Frailty Sarcop Falls
546	4:26-34.
547	36. Aunger, JA, Greaves, CJ, Davis, ET, Asamane, EA, Whittaker, AC, Greig, CA (2020)
548	A novel behavioural INTErvention to REduce Sitting Time in older adults undergoing
549	orthopaedic surgery (INTEREST): results of a randomised-controlled feasibility
550	study. Aging Clin Exp Res. <u>https://doi.org/10.1007/s40520-020-01475-6</u> .
551	37. Shephard RJ (2003) Limits to the measurement of habitual physical activity by
552	questionnaires. Br J Sport Med 37:197-206.
553	38. Chastin SFM, Dontje ML, Skelton DA, Čukić I, Shaw RJ, Gill JMR, Greig CA, Gale
554	CR, Deary IJ, Der G, Dall PM (2018) Systematic comparative validation of self-
555	report measures of sedentary time against an objective measure of postural sitting
556	(activPAL). Int J Behav Nutr Phys Act 15:21-32.
557	39. Strain T, Milton K, Dall P, Standage M, Mutrie N (2019) How are we measuring
558	physical activity and sedentary behaviour in the four home nations of the UK? A
559	narrative review of current surveillance measures and future directions. Br J Sports
560	Med 0:1-9.

561	40. Kim SJ, Yang WG, Cho E, Park EC (2012) Relationship between weight, body mass
562	index and bone mineral density of lumbar spine in women. J Bone Metab 19:95-102.
563	41. Felson DT, Zhang Y, Hannan MT, Anderson JJ (1993) Effects of weight and body
564	mass index on bone mineral density in men and women: the Framingham study. J
565	Bone Miner Res 8:567-573.
566	42. Reid IR (2002) Relationships among body mass, its components, and bone. Bone
567	31:547-555.
568	43. Nielson CM, Srikanth P, Orwoll ES (2012) Obesity and fracture in men and women:
569	an epidemiologic perspective. J Bone Miner Res 27:1-10.
570	44. Iwaniec UT, Turner RT (2016) Influence of body weight on bone mass, architecture,

and turnover. J Endocrinol 230:R115–R130.