

A left 3-Engel element whose normal closure is not nilpotent

Noce, Marialaura; Tracey, Gareth; Traustason, Gunnar

Citation for published version (Harvard):

Noce, M, Tracey, G & Traustason, G 2020, 'A left 3-Engel element whose normal closure is not nilpotent', *Journal of Pure and Applied Algebra*, vol. 224, no. 3, pp. 1092-1101.

[Link to publication on Research at Birmingham portal](#)

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

A left 3-Engel element whose normal closure is not nilpotent

Marialaura Noce*, Gareth Tracey† and Gunnar Traustason‡

Abstract

We give an example of a locally nilpotent group G containing a left 3-Engel element x where $\langle x \rangle^G$ is not nilpotent.

2010 Mathematics Subject Classification: 20F45, 20F40

Keywords: Engel elements, Lie algebras, enveloping algebras.

1 Introduction

Let G be a group. An element $a \in G$ is a left Engel element in G , if for each $x \in G$ there exists a non-negative integer $n(x)$ such that

$$[[[x, a], a], \dots, a] = 1.$$

$\underbrace{\hspace{1.5cm}}_{n(x)}$

If $n(x)$ is bounded above by n then we say that a is a left n -Engel element in G . It is straightforward to see that any element of the Hirsch-Plotkin radical $HP(G)$ of G is a left Engel element and the converse is known to be true for some classes of groups, including solvable groups and finite groups (more generally groups satisfying the maximal condition on subgroups) [3,1]. The converse is however not true in general and this is the case even for

*University of Salerno - University of the Basque Country, mnoce@unisa.it

†University of Bath, g.m.tracey@bath.ac.uk

‡University of Bath, g.traustason@bath.ac.uk

bounded left Engel elements. In fact whereas one sees readily that a left 2-Engel element is always in the Hirsch-Plotkin radical this is still an open question for left 3-Engel elements. Recently there has been a breakthrough and in [6] it is shown that any left 3-Engel element of odd order is contained in $HP(G)$. From [10] one also knows that in order to generalise this to left 3-Engel elements of any finite order it suffices to deal with elements of order 2.

It was observed by William Burnside [2] that every element in a group of exponent 3 is a left 2-Engel element and so the fact that every left 2-Engel element lies in the Hirsch-Plotkin radical can be seen as the underlying reason why groups of exponent 3 are locally finite. For groups of 2-power exponent there is a close link with left Engel elements. If G is a group of exponent 2^n then it is not difficult to see that any element a in G of order 2 is a left $(n + 1)$ -Engel element of G (see the introduction of [11] for details). For sufficiently large n we know that the variety of groups of exponent 2^n is not locally finite [5,7]. As a result one can see (for example in [11]) that it follows that for sufficiently large n we do not have in general that a left n -Engel element is contained in the Hirsch-Plotkin radical. Using the fact that groups of exponent 4 are locally finite [9], one can also see that if all left 4-Engel elements of a group G of exponent 8 are in $HP(G)$ then G is locally finite.

Swapping the role of a and x in the definition of a left Engel element we get the notion of a right Engel element. Thus an element $a \in G$ is a right Engel element, if for each $x \in G$ there exists a non-negative integer $n(x)$ such that

$$[[[a, x], x], \dots, x] = 1.$$

$\underbrace{\hspace{1.5cm}}_{n(x)}$

If $n(x)$ is bounded above by n , we say that a is a right n -Engel element. By a classical result of Heineken [4] one knows that if a is a right n -Engel element in G then a^{-1} is a left $(n + 1)$ -Engel element.

In [8] M. Newell proved that if a is a right 3-Engel element in G then $a \in HP(G)$ and in fact he proved the stronger result that $\langle a \rangle^G$ is nilpotent of class at most 3. The natural question arises whether the analogous result holds for left 3-Engel elements. In this paper we show this is not the case by giving an example of a locally finite 2-group with a left 3-Engel element a such that $\langle a \rangle^G$ is not nilpotent.

2 An example of a left 3-Engel element x where $\langle x \rangle^G$ is not nilpotent

Our construction will be based on an example of a Lie algebra given in [12]. Let \mathbb{F} be the field of order 2 and consider a 4-dimensional vector space $V = \mathbb{F}x + \mathbb{F}u + \mathbb{F}v + \mathbb{F}w$ where

$$u \cdot v = u, \quad v \cdot w = w, \quad w \cdot u = v, \quad u \cdot x = 0, \quad v \cdot x = 0, \quad w \cdot x = u.$$

We then extend the product linearly on V . One can check that V is a Lie algebra with a trivial center and where $W = \mathbb{F}u + \mathbb{F}v + \mathbb{F}w$ is a simple ideal.

Let $E = \langle \text{ad}(x), \text{ad}(u), \text{ad}(v), \text{ad}(w) \rangle \leq \text{End}(V)$ be the associative enveloping algebra of V . Let

$$\begin{aligned} e_1 &= \text{ad}(w), & e_2 &= \text{ad}(w)^2, & e_3 &= \text{ad}(w)^3, & e_4 &= \text{ad}(v), \\ e_5 &= \text{ad}(v)\text{ad}(w), & e_6 &= \text{ad}(v)\text{ad}(w)^2, & e_7 &= \text{ad}(u), & e_8 &= \text{ad}(u)\text{ad}(w), \\ e_9 &= \text{ad}(u)\text{ad}(w)^2, & e_{10} &= \text{ad}(x)\text{ad}(v), & e_{11} &= \text{ad}(x)\text{ad}(w), & e_{12} &= \text{ad}(x)\text{ad}(w)^2. \end{aligned}$$

Lemma 2.1 *The associative enveloping algebra E is 12-dimensional with basis e_1, \dots, e_{12} .*

Proof We first show that E is spanned by products of the form

$$\text{ad}(x)^\epsilon \cdot \text{ad}(u)^r \cdot \text{ad}(v)^s \cdot \text{ad}(w)^t$$

where ϵ, r, s, t are non-negative integers. To see this we need to show that any product $\text{ad}(y_1) \cdots \text{ad}(y_m)$, with $y_1, \dots, y_m \in \{x, u, v, w\}$ can be written as a linear combination of elements of the required form. We use induction on m . This is obvious when $m = 1$. Now suppose $m \geq 2$ and that the statement is true for all shorter products. Suppose there are ϵ entries of x , r entries of u , s entries of v and t entries of w in the product $\text{ad}(y_1) \cdots \text{ad}(y_m)$. Using the fact that $\text{ad}(y_i)\text{ad}(y_j) = \text{ad}(y_j)\text{ad}(y_i) + \text{ad}(y_i y_j)$, we see that modulo shorter products we have

$$\text{ad}(y_1) \cdots \text{ad}(y_m) = \text{ad}(x)^\epsilon \text{ad}(u)^r \text{ad}(v)^s \text{ad}(w)^t.$$

Hence the statement is true for products of length m . This finishes the inductive proof of our claim.

From the fact that $\text{ad}(x)^2 = \text{ad}(x)\text{ad}(u) = 0$, $\text{ad}(v)^2 = \text{ad}(v)$, $\text{ad}(u)^2 = \text{ad}(x)\text{ad}(v)$ and $\text{ad}(u)\text{ad}(v) = \text{ad}(u) + \text{ad}(x)\text{ad}(w)$, we can assume that $0 \leq \epsilon, r, s \leq 1$ and that if $\text{ad}(u)$ is included then we can assume that neither $\text{ad}(x)$ nor $\text{ad}(v)$ is included. This together with $\text{ad}(x) = \text{ad}(x)\text{ad}(v)$ and $\text{ad}(w)^4 = \text{ad}(v)\text{ad}(w)^3 = \text{ad}(u)\text{ad}(w)^3 = \text{ad}(x)\text{ad}(w)^3 = 0$ shows that E is generated by e_1, \dots, e_{12} . It remains to see that these elements are linearly independent. Suppose $\alpha_1 e_1 + \dots + \alpha_{12} e_{12} = 0$ for some $\alpha_1, \dots, \alpha_{12} \in \mathbb{F}$. Then

$$0 = x(\alpha_1 e_1 + \dots + \alpha_{12} e_{12}) = \alpha_1 u + \alpha_2 v + \alpha_3 w$$

gives that $\alpha_1 = \alpha_2 = \alpha_3 = 0$. Then

$$0 = u(\alpha_4 e_4 + \dots + \alpha_{12} e_{12}) = \alpha_4 u + \alpha_5 v + \alpha_6 w$$

implies that $\alpha_4 = \alpha_5 = \alpha_6 = 0$. Likewise

$$0 = v(\alpha_7 e_7 + \dots + \alpha_{12} e_{12}) = \alpha_7 u + \alpha_8 v + \alpha_9 w$$

giving $\alpha_7 = \alpha_8 = \alpha_9 = 0$. Finally

$$0 = w(\alpha_{10} e_{10} + \alpha_{12} e_{11} + \alpha_{12} e_{12}) = \alpha_{10} u + \alpha_{11} v + \alpha_{12} w$$

and thus $\alpha_{10} = \alpha_{11} = \alpha_{12} = 0$. \square

We use this example to construct a certain locally nilpotent Lie algebra over \mathbb{F} of countably infinite dimension. For ease of notation it will be useful to introduce the following modified union of subsets of \mathbb{N} . We let

$$A \sqcup B = \begin{cases} A \cup B & (\text{if } A \cap B = \emptyset) \\ \emptyset & (\text{otherwise}) \end{cases}$$

For each non-empty subset A of \mathbb{N} we let W_A be a copy of the vector space W . That is $W_A = \{z_A : z \in W\}$ with addition $z_A + t_A = (z + t)_A$. We then take the direct sum of these

$$W^* = \bigoplus_{\emptyset \neq A \subseteq \mathbb{N}} W_A$$

that we turn into a Lie algebra with multiplication

$$z_A \cdot t_B = (zt)_{A \sqcup B}$$

when $z_A \in W_A$ and $t_B \in W_B$ that is then extended linearly on W^* . The interpretation here is that $z_\emptyset = 0$. Finally we extend this to a semidirect product with $\mathbb{F}x$

$$V^* = W^* \oplus \mathbb{F}x$$

induced from the action $z_A \cdot x = (zx)_A$.

Notice that V^* has basis

$$\{x\} \cup \{u_A, v_A, w_A : \emptyset \neq A \subseteq \mathbb{N}\}$$

and that

$$\begin{aligned} u_A \cdot u_B &= v_A \cdot v_B = w_A \cdot w_B = 0, \\ u_A \cdot x &= 0, \quad v_A \cdot x = 0, \quad w_A \cdot x = u_A \end{aligned}$$

and

$$u_A \cdot v_B = u_{A \sqcup B}, \quad v_A \cdot w_B = w_{A \sqcup B}, \quad w_A \cdot u_B = v_{A \sqcup B}.$$

Notice that any finitely generated subalgebra of V^* is contained in some $S = \langle x, u_{A_1}, \dots, u_{A_r}, v_{B_1}, \dots, v_{B_s}, w_{C_1}, \dots, w_{C_t} \rangle$. From the fact that $zxx = 0$ for all $z \in V^*$ it follows that S is nilpotent of class at most $2(r + s + t)$. Hence V^* is locally nilpotent. The next aim is to find a group $G \leq \text{GL}(V^*)$ containing $1 + \text{ad}(x)$ where $1 + \text{ad}(x)$ is a left 3-Engel element in G but where $\langle 1 + \text{ad}(x) \rangle^G$ is not nilpotent. The next lemma is a preparation for this.

Lemma 2.2 *The adjoint linear operator $\text{ad}(x)$ on V^* satisfies:*

- (a) $\text{ad}(x)^2 = 0$.
- (b) $\text{ad}(x)\text{ad}(y)\text{ad}(x) = 0$ for all $y \in V^*$.

Proof (a) Follows from the fact that $x \cdot x = u_A \cdot x = v_A \cdot x = 0$ and $(w_A \cdot x) \cdot x = u_A \cdot x = 0$.

(b) Follows from $w_A \cdot x \cdot u_B = u_A \cdot u_B = 0$, $w_A \cdot x \cdot v_B \cdot x = u_A \cdot v_B \cdot x = u_{A \sqcup B} \cdot x = 0$ and $w_A \cdot x \cdot w_B \cdot x = u_A \cdot w_B \cdot x = v_{A \sqcup B} \cdot x = 0$. \square

Let y be any of the generators x, u_A, v_A, w_A . As $\text{ad}(y)^2 = 0$ it follows that

$$(1 + \text{ad}(y))^2 = 1 + 2\text{ad}(y) + \text{ad}(y)^2 = 1.$$

Thus $1 + \text{ad}(y)$ is an involution in $\text{GL}(V^*)$.

Remark. Notice that for any $A, B \subseteq \mathbb{N}$, the pairs $(\text{ad}(u_A), \text{ad}(u_B))$, $(\text{ad}(v_A), \text{ad}(v_B))$ and $(\text{ad}(w_A), \text{ad}(w_B))$ consist of elements that commute. Thus the subgroups

$$\mathcal{U} = \langle 1 + \text{ad}(u_A) : A \subseteq \mathbb{N} \rangle, \mathcal{V} = \langle 1 + \text{ad}(v_A) : A \subseteq \mathbb{N} \rangle, \mathcal{W} = \langle 1 + \text{ad}(w_A) : A \subseteq \mathbb{N} \rangle$$

are elementary abelian of countably infinite rank. We will be working with the group $G = \langle 1 + \text{ad}(x), \mathcal{U}, \mathcal{V}, \mathcal{W} \rangle$.

Lemma 2.3 *The following commutator relations hold in G :*

- (a) $[1 + \text{ad}(u_A), 1 + \text{ad}(v_B)] = 1 + \text{ad}(u_{A \sqcup B})$.
- (b) $[1 + \text{ad}(v_A), 1 + \text{ad}(w_B)] = 1 + \text{ad}(w_{A \sqcup B})$.
- (c) $[1 + \text{ad}(w_A), 1 + \text{ad}(u_B)] = 1 + \text{ad}(v_{A \sqcup B})$.
- (d) $[1 + \text{ad}(u_A), 1 + \text{ad}(x)] = 1$.
- (e) $[1 + \text{ad}(v_A), 1 + \text{ad}(x)] = 1$.
- (f) $[1 + \text{ad}(w_A), 1 + \text{ad}(x)] = 1 + \text{ad}(u_A)$.

Proof (a) We have

$$\begin{aligned} [1 + \text{ad}(u_A), 1 + \text{ad}(v_B)] &= (1 + \text{ad}(u_A)) \cdot (1 + \text{ad}(v_B)) \cdot (1 + \text{ad}(u_A)) \cdot (1 + \text{ad}(v_B)) \\ &= 1 + \text{ad}(u_A)\text{ad}(v_B) + \text{ad}(v_B)\text{ad}(u_A) \\ &= 1 + \text{ad}(u_A v_B) \\ &= 1 + \text{ad}(u_{A \sqcup B}). \end{aligned}$$

(b) and (c) are proved similarly. For (f) we have

$$\begin{aligned} [1 + \text{ad}(w_A), 1 + \text{ad}(x)] &= (1 + \text{ad}(w_A)) \cdot (1 + \text{ad}(x)) \cdot (1 + \text{ad}(w_A)) \cdot (1 + \text{ad}(x)) \\ &= 1 + \text{ad}(w_A)\text{ad}(x) + \text{ad}(x)\text{ad}(w_A) + \text{ad}(x)\text{ad}(w_A)\text{ad}(x) \\ &= 1 + \text{ad}(w_A \cdot x) \\ &= 1 + \text{ad}(u_A). \end{aligned}$$

Here in the 2nd last equality, we have used Lemma 2.2. Parts (d) and (e) are proved similarly. \square

Remark. Notice that as V^* is locally nilpotent, it follows from Lemma 2.3 that G is locally nilpotent. Next proposition clarifies further the structure of G .

Proposition 2.4 *We have $G = \langle 1 + \text{ad}(x) \rangle \mathcal{U} \mathcal{V} \mathcal{W}$. Furthermore every element $g \in G$ has a unique expression $g = (1 + \text{ad}(x))^\epsilon r s t$ with $\epsilon \in \{0, 1\}$, $r \in \mathcal{U}$, $s \in \mathcal{V}$ and $t \in \mathcal{W}$.*

Proof We first deal with the existence of such a decomposition. Suppose

$$g = g_0(1 + \text{ad}(x))g_2 \cdots (1 + \text{ad}(x))g_n$$

where g_0, \dots, g_n are products of elements of the form $1 + \text{ad}(u_A)$, $1 + \text{ad}(v_A)$ and $1 + \text{ad}(w_A)$. From Lemma 2.3 we know that $(1 + \text{ad}(w_A))(1 + \text{ad}(x)) = (1 + \text{ad}(x))(1 + \text{ad}(w_A))(1 + \text{ad}(u_A))$ and $1 + \text{ad}(x)$ commutes with all products of the form $1 + \text{ad}(u_A)$ and $1 + \text{ad}(v_A)$. We can thus collect the $(1 + \text{ad}(x))$'s to the left, starting with the leftmost occurrence. This may introduce more elements of the form $(1 + \text{ad}(u_A))$ but no new $1 + \text{ad}(x)$. We thus see that

$$g = (1 + \text{ad}(x))^n g_1 \cdots g_m$$

where g_i is of the form $1 + \text{ad}(u_A)$, $1 + \text{ad}(v_A)$ or $1 + \text{ad}(w_A)$. This reduces our problem to the case when $g \in \langle \mathcal{U}, \mathcal{V}, \mathcal{W} \rangle$. Suppose

$$g = g_0(1 + \text{ad}(u_{A_1}))g_1 \cdots (1 + \text{ad}(u_{A_n}))g_n$$

where g_0, \dots, g_n are products of elements of the form $1 + \text{ad}(v_A)$ and $1 + \text{ad}(w_A)$. Suppose that the elements occurring in these products are $1 + \text{ad}(v_{A_{n+1}}), \dots, 1 + \text{ad}(v_{A_{n+l}}), 1 + \text{ad}(w_{A_{n+l+1}}), \dots, 1 + \text{ad}(w_{A_m})$.

Using Lemma 2.3 we know that $(1 + \text{ad}(v_B))(1 + \text{ad}(u_A)) = (1 + \text{ad}(u_A))(1 + \text{ad}(v_B))(1 + \text{ad}(u_{A \sqcup B}))$ and that $(1 + \text{ad}(w_B))(1 + \text{ad}(u_A)) = (1 + \text{ad}(u_A))(1 + \text{ad}(w_B))(1 + \text{ad}(v_{A \sqcup B}))$. We can thus collect $1 + \text{ad}(u_{A_1}), \dots, 1 + \text{ad}(u_{A_n})$ to the left. In doing so we may introduce new terms of the form $1 + \text{ad}(u_A)$, with A of the form $A_{i_1} \sqcup \cdots \sqcup A_{i_s}$, and $s \geq 2$. This shows that

$$g = (1 + \text{ad}(u_{A_1})) \cdots (1 + \text{ad}(u_{A_n}))g_1 \cdots g_m$$

where each g_j is of the form $1 + \text{ad}(v_B)$, $1 + \text{ad}(w_B)$ or $1 + \text{ad}(u_A)$, and A is a modified union of at least 2 sets from $\{A_1, \dots, A_m\}$. We can repeat this procedure, collecting all the new $(1 + \text{ad}(u_A))$ s. In doing so, we possibly introduce some new such elements but these will then be with an A that is a modified union of at least 3 sets from $\{A_1, \dots, A_m\}$. Continuing like this the procedure will end after at most m steps as every modified union of $m + 1$ sets from $\{A_1, \dots, A_m\}$ will be trivial. We have thus seen that $g = rh$ with $r \in \mathcal{U}$ and $h \in \langle \mathcal{V}, \mathcal{W} \rangle$. We are now only left with the situation when $g \in \langle \mathcal{V}, \mathcal{W} \rangle$. Suppose

$$g = g_0(1 + \text{ad}(v_{A_1}))g_1 \cdots (1 + \text{ad}(v_{A_n}))g_n$$

where g_0, g_1, \dots, g_n are of the form $1 + \text{ad}(w_A)$. As $(1 + \text{ad}(w_B))(1 + \text{ad}(v_A)) = (1 + \text{ad}(v_A))(1 + \text{ad}(w_B))(1 + \text{ad}(w_{A \sqcup B}))$, we can now collect $1 + \text{ad}(v_{A_1}), \dots, 1 + \text{ad}(v_{A_n})$ to the left and in doing so, all the new terms introduced will be of the form $1 + \text{ad}(w_A)$. Thus $g = st$ with $s \in \mathcal{V}$ and $t \in \mathcal{W}$. This completes the existence part. We now want to show that such a decomposition is unique. Suppose

$$\begin{aligned} (1 + \eta \text{ad}(x)) &= (1 + \tau \text{ad}(x)) \\ (1 + \alpha_1 \text{ad}(u_{A_1})) \cdots (1 + \alpha_r \text{ad}(u_{A_r})) &= (1 + \beta_1 \text{ad}(u_{A_1})) \cdots (1 + \beta_r \text{ad}(u_{A_r})) \\ (1 + \gamma_1 \text{ad}(v_{B_1})) \cdots (1 + \gamma_s \text{ad}(v_{B_s})) &= (1 + \delta_1 \text{ad}(v_{B_1})) \cdots (1 + \delta_s \text{ad}(v_{B_s})) \\ (1 + \epsilon_1 \text{ad}(w_{C_1})) \cdots (1 + \epsilon_t \text{ad}(w_{C_t})) &= (1 + \nu_1 \text{ad}(w_{C_1})) \cdots (1 + \nu_t \text{ad}(w_{C_t})). \end{aligned}$$

Applying both sides to $w_{\mathbb{N}}$ we get

$$w_{\mathbb{N}} + \eta u_{\mathbb{N}} = w_{\mathbb{N}} + \tau u_{\mathbb{N}}$$

from which we get $\eta = \tau$. Applying both sides to x we get

$$\begin{aligned} x + \epsilon_1 u_{C_1} + \cdots + \epsilon_r u_{C_t} &= x + \nu_1 u_{C_1} + \cdots + \nu_r u_{C_t} \\ \epsilon_1 \epsilon_2 v_{C_1 \sqcup C_2} + \cdots + \epsilon_{t-1} \epsilon_t v_{C_{t-1} \sqcup C_t} &= \nu_1 \nu_2 v_{C_1 \sqcup C_2} + \cdots + \nu_{t-1} \nu_t v_{C_{t-1} \sqcup C_t} \\ \epsilon_1 \epsilon_2 \epsilon_3 w_{C_1 \sqcup C_2 \sqcup C_3} + \cdots &= \nu_1 \nu_2 \nu_3 w_{C_1 \sqcup C_2 \sqcup C_3} + \cdots \\ + \epsilon_{t-2} \epsilon_{t-1} \epsilon_t w_{C_{t-2} \sqcup C_{t-1} \sqcup C_t} &= + \nu_{t-2} \nu_{t-1} \nu_t w_{C_{t-2} \sqcup C_{t-1} \sqcup C_t} \end{aligned}$$

from which we see that $\epsilon_1 = \nu_1, \dots, \epsilon_t = \nu_t$. Thus

$$\begin{aligned} (1 + \alpha_1 \text{ad}(u_{A_1})) \cdots (1 + \alpha_r \text{ad}(u_{A_r})) &= (1 + \beta_1 \text{ad}(u_{A_1})) \cdots (1 + \beta_r \text{ad}(u_{A_r})) \\ (1 + \gamma_1 \text{ad}(v_{B_1})) \cdots (1 + \gamma_s \text{ad}(v_{B_s})) &= (1 + \delta_1 \text{ad}(v_{B_1})) \cdots (1 + \delta_s \text{ad}(v_{B_s})). \end{aligned}$$

We can assume that $A_j \not\subseteq A_i$ and $B_j \not\subseteq B_i$ when $i < j$. Applying both sides to $u_{\mathbb{N} \setminus B_1}$ gives

$$u_{\mathbb{N} \setminus B_1} + \gamma_1 u_{\mathbb{N}} = u_{\mathbb{N} \setminus B_1} + \delta_1 u_{\mathbb{N}}$$

from which we see that $\gamma_1 = \delta_1$. Cancelling on both sides by $1 + \gamma_1 \text{ad}(v_{B_1})$ and then applying both sides to $u_{\mathbb{N} \setminus B_2}$ likewise gives $\gamma_2 = \delta_2$. Continuing in this manner gives $\gamma_1 = \delta_1, \dots, \gamma_s = \delta_s$. We then have

$$(1 + \alpha_1 \text{ad}(u_{A_1})) \cdots (1 + \alpha_r \text{ad}(u_{A_r})) = (1 + \beta_1 \text{ad}(u_{A_1})) \cdots (1 + \beta_r \text{ad}(u_{A_r})).$$

A similar argument as before, applying both sides to $v_{\mathbb{N} \setminus A_1}, v_{\mathbb{B} \setminus A_2}, \dots$ gives likewise $\alpha_1 = \beta_1, \dots, \alpha_r = \beta_r$. This finishes the proof. \square

We are now ready to prove the main result of this paper.

Theorem 2.5 *The element $1 + \text{ad}(x)$ is a left 3-Engel element in G . However $\langle 1 + \text{ad}(x) \rangle^G$ is not nilpotent.*

Proof Let $g = h(1 + \text{ad}(w_{A_1})) \cdots (1 + \text{ad}(w_{A_n}))$ be an arbitrary element in G where $h \in \langle (1 + \text{ad}(x)) \rangle \mathcal{UV}$. We want to show that

$$[(1 + \text{ad}(x))^g, {}_2(1 + \text{ad}(x))] = 1.$$

Notice first that if $y \in V$ then

$$\begin{aligned} (1 + \text{ad}(y))^{1 + \text{ad}(w_A)} &= (1 + \text{ad}(w_A))(1 + \text{ad}(y))(1 + \text{ad}(w_A)) \\ &= 1 + \text{ad}(y) + \text{ad}(yw_A). \end{aligned}$$

Notice that $(1 + \text{ad}(x))^g = (1 + \text{ad}(x))^{(1 + \text{ad}(w_{A_1})) \cdots (1 + \text{ad}(w_{A_n}))}$ and straightforward induction show that

$$(1 + \text{ad}(x))^g = 1 + \text{ad}(y)$$

where

$$y = x + \sum_{1 \leq i \leq n} u_{A_i} + \sum_{1 \leq i < j \leq n} v_{A_i \sqcup A_j} + \sum_{1 \leq i < j < k \leq n} w_{A_i \sqcup A_j \sqcup A_k}.$$

Since $\text{ad}(x)\text{ad}(y)\text{ad}(x) = 0$, the commutator of $(1 + \text{ad}(x))^g$ with $(1 + \text{ad}(x))$ is

$$\begin{aligned} (1 + \text{ad}(y))(1 + \text{ad}(x))(1 + \text{ad}(y))(1 + \text{ad}(x)) &= 1 + \text{ad}(y)\text{ad}(x) + \text{ad}(x)\text{ad}(y) \\ &\quad + \text{ad}(y)\text{ad}(x)\text{ad}(y). \end{aligned}$$

Then

$$\begin{aligned} [(1 + \text{ad}(x))^g, {}_2(1 + \text{ad}(x))] &= ((1 + \text{ad}(y))(1 + \text{ad}(x)))^4 \\ &= ((1 + \text{ad}(y)\text{ad}(x) + \text{ad}(x)\text{ad}(y) \\ &\quad + \text{ad}(y)\text{ad}(x)\text{ad}(y))^2 = 1 \end{aligned}$$

using again the fact that $\text{ad}(x)\text{ad}(y)\text{ad}(x) = 0$.

Then the normal closure of $1 + \text{ad}(x)$ in G is though not nilpotent as for $A_i = \{i\}$ we have

$$\begin{aligned} [1 + \text{ad}(w_{A_1}), 1 + \text{ad}(x), 1 + \text{ad}(w_{A_2}), 1 + \text{ad}(w_{A_3}), \dots \\ \dots, 1 + \text{ad}(x), 1 + \text{ad}(w_{A_{2n}}), 1 + \text{ad}(w_{A_{2n+1}})] = 1 + \text{ad}(w_A), \end{aligned}$$

where $A = A_1 \sqcup \dots \sqcup A_{2n+1} = \{1, 2, \dots, 2n+1\}$. \square

Our next aim is to show however that if we take any r conjugates $(1 + \text{ad}(x))^{g_1}, \dots, (1 + \text{ad}(x))^{g_r}$ of $(1 + \text{ad}(x))$ in G , they generate a nilpotent subgroup of r -bounded class that grows linearly with r .

We first work in a more general setting. For each $e \in E$ and $\emptyset \neq A \subseteq \mathbb{N}$, let $e(A) \in \text{End}(V^*)$ where

$$u_B e(A) = (ue)_{B \sqcup A}.$$

Then let $E^* = \langle \text{ad}(x), e(A) : e \in E \text{ and } \emptyset \neq A \subseteq \mathbb{N} \rangle$. As V^* is locally nilpotent, one sees readily that the elements of E^* are nilpotent and thus $1 + E^*$ is a subgroup of $\text{End}(V^*)$. We are going to see that $1 + E^*$ is of finite exponent.

Remark Notice that $\text{ad}(u_{\mathbb{N}}) = \text{ad}(v_{\mathbb{N}}) = 0$.

Lemma 2.6 *The elements $\text{ad}(w_{\mathbb{N}})$ and $\{e_i(A) : 1 \leq i \leq 12, \emptyset \neq A \subset \mathbb{N}\} \cup \{e_1(\mathbb{N}), e_2(\mathbb{N}), e_3(\mathbb{N})\}$ form a basis for E^* .*

Proof One sees that these elements span E^* as a vector space in a similar way as in the proof of Lemma 2.1. We thus skip over the details and only show that these elements are linearly independent. Suppose

$$\epsilon \text{ad}(x) + \sum_{i=1}^{12} \sum_A \epsilon_A^i e_i(A) = 0,$$

where only finitely many of the coefficients ϵ, ϵ_A^i are non-zero. Denote the left hand side by T . Then

$$0 = xT = \sum_A \epsilon_A^1 u_A + \sum_A \epsilon_A^2 v_A + \sum_A \epsilon_A^3 w_A$$

implying that $\epsilon_A^i = 0$ for all A and $i = 1, 2, 3$. Then

$$0 = u_{\mathbb{N} \setminus A} T = \sum_{B \subseteq A} \epsilon_B^4 u_{(\mathbb{N} \setminus A) \sqcup B} + \sum_{B \subseteq A} \epsilon_B^5 v_{(\mathbb{N} \setminus A) \sqcup B} + \sum_{B \subseteq A} \epsilon_B^6 w_{(\mathbb{N} \setminus A) \sqcup B}.$$

In particular $\epsilon_A^i = 0$ for all $A \neq \mathbb{N}$ and $i = 4, 5, 6$. We continue in a similar way. Next

$$0 = v_{\mathbb{N} \setminus A} T = \sum_{B \subseteq A} \epsilon_B^7 u_{(\mathbb{N} \setminus A) \sqcup B} + \sum_{B \subseteq A} \epsilon_B^8 v_{(\mathbb{N} \setminus A) \sqcup B} + \sum_{B \subseteq A} \epsilon_B^9 w_{(\mathbb{N} \setminus A) \sqcup B}$$

that shows that $\epsilon_A^i = 0$ for all A and $i = 7, 8, 9$. Finally

$$0 = w_{\mathbb{N}}T = \epsilon u_{\mathbb{N}}$$

giving $\epsilon = 0$ and

$$0 = w_{\mathbb{N} \setminus A}T = \sum_{B \subseteq A} \epsilon_B^{10} u_{(\mathbb{N} \setminus A) \sqcup B} + \sum_{B \subseteq A} \epsilon_B^{11} v_{(\mathbb{N} \setminus A) \sqcup B} + \sum_{B \subseteq A} \epsilon_B^{12} w_{(\mathbb{N} \setminus A) \sqcup B}$$

and $\epsilon_A^i = 0$ for all A and $i = 10, 11, 12$. This finishes the proof. \square

Corollary 2.7 *We have $(1 + E^*)^{32} = 1$.*

Proof Let \bar{E} be the subalgebra of E^* generated by all $e_i(A)$ where $1 \leq i \leq 12$ and $\emptyset \neq A \subseteq \mathbb{N}$. Let $f = \text{ad}(x) + e \in E$ where $e \in \bar{E}$. Then $f^2 = \text{ad}(x)^2 + e^2 + (\text{ead}(x) + \text{ad}(x)e) = e^2 + (\text{ead}(x) - \text{ad}(x)e)$. It is straightforward to see that $\text{ead}(x) - \text{ad}(x)e \in \bar{E}$ (\bar{E} is an ideal in the Lie algebra E^*) and thus $f^2 \in \bar{E}$. It thus suffices to show that $\bar{E}^{16} = 0$, as then it will follow that $(E^*)^{32} = 0$ and therefore $(1 + e)^{32} = 1 + e^{32} = 1$ for all $e \in E^*$.

Let $e = y_1 + \dots + y_m$ be any element in \bar{E} where y_1, \dots, y_m belong to the basis $\{e_i(A) : 1 \leq i \leq 12, \emptyset \neq A \subset \mathbb{N}\} \cup \{e_1(\mathbb{N}), e_2(\mathbb{N}), e_3(\mathbb{N})\}$ for \bar{E} given in Lemma 2.6. As any product with a repeated term is 0 we see that e^{16} is a sum of terms of the form

$$\sum_{\sigma \in S_{16}} f_{\sigma(1)} \cdots f_{\sigma(16)} \tag{1}$$

with $f_1, \dots, f_{16} \in \{y_1, \dots, y_m\}$. As $16 > 12$ some two of f_1, \dots, f_{16} must be of the same type. Without loss of generality we can assume that these are $f_{15} = e_i(A)$ and $f_{16} = e_i(B)$. Notice that the sum (1) splits naturally into $16!/2$ sums of pairs

$$\begin{aligned} \sum_{\sigma \in S_{16}} f_{\sigma(1)} \cdots f_{\sigma(16)} &= \sum_{\sigma \in S_{14}} (e_i(A)e_i(B) + e_i(B)e_i(A)) f_{\sigma(1)} \cdots f_{\sigma(14)} + \cdots \\ &\quad + \sum_{\sigma \in S_{14}} f_{\sigma(1)} \cdots f_{\sigma(14)} (e_i(A)e_i(B) + e_i(B)e_i(A)), \end{aligned}$$

one for each of the $\binom{16}{2}$ positions of the pair $(e_i(A), e_i(B))$ within the product. But for each such choice of positions the two elements in the pair have the same value and as the characteristic is 2, the sum of each pair is 0. Thus the sum in (1) is zero and we have shown that $\bar{E}^{16} = 0$. \square

Proposition 2.8 *Any r -generator subgroup of $1 + E^*$ is nilpotent of r -bounded class.*

Proof From Corollary 2.7 we know that $1 + E^*$ is of bounded exponent. The result thus follows from Zel'manov's solution to the Restricted Burnside Problem. \square

Despite the fact that the normal closure of $1 + \text{ad}(x)$ in G is not nilpotent, it turns out that the nilpotency class of the subgroup generated by any k conjugates grows linearly with respect to k . In order to see this we first introduce some more notation. Let A_1, A_2, \dots, A_k be any k subsets of \mathbb{N} . For each k -tuple $(i1, i2, \dots, ir)$ of non-negative integers and each $e \in E$ we let

$$e^{(i1, \dots, ir)} = \sum_{\substack{B_1 \subseteq A_1 \\ |B_1| = i1}} \dots \sum_{\substack{B_k \subseteq A_k \\ |B_k| = ik}} e(B_1 \sqcup B_2 \sqcup \dots \sqcup B_k).$$

Notice that

$$e^{(i1, \dots, ir)} f^{(j1, \dots, jr)} = \binom{i1 + j1}{i1} \dots \binom{ir + jr}{ir} (ef)^{(i1+j1, \dots, ir+jr)}.$$

Now notice that $\binom{3+i}{3}$ is even for $i = 1, 2, 3$ and the same is true for $\binom{2+2}{2}$ and $\binom{1+1}{1}$. However $\binom{2+1}{2}$ is odd. From this it follows that

$$Q = \langle \text{ad}(x), e^{(i1, \dots, ir)} : e \in E, 0 \leq i1, \dots, ir \leq 3, i1 + \dots + ir \geq 1 \rangle$$

is a subalgebra of E^* . A nonzero product in Q can have at most $2r$ elements of the form $e^{(i1, \dots, ir)}$ and as $\text{ad}(x)^2 = 0$ we could then have at most $1 + 2r$ occurrences of $\text{ad}(x)$ in a non-zero product. Thus $Q^{4r+2} = 0$.

Now take some r conjugates of $(1 + \text{ad}(x))$ in G . Recall that each conjugate is of the form $(1 + \text{ad}(x))^{(1+\text{ad}(w_{C_1}) \dots (1+\text{ad}(w_{C_j}))}$. For ease of notation we will assume that each C_k is a singleton set. The following argument also works for the more general case. Let

$$A_1 = \{1, \dots, k_1\}, A_2 = \{k_1 + 1, \dots, k_2\}, \dots, A_r = \{k_{r-1} + 1, \dots, k_r\}.$$

Then we have seen (see the proof of Theorem 2.5) that

$$\begin{aligned} (1 + \text{ad}(x))^{(1+\text{ad}(w_1)) \dots (1+\text{ad}(w_{k_1}))} &= 1 + \text{ad}(x) + e_7^{(1,0, \dots, 0)} + e_4^{(2,0, \dots, 0)} + e_1^{(3,0, \dots, 0)} \\ &\vdots \\ (1 + \text{ad}(x))^{(1+\text{ad}(w_{k_{r-1}+1})) \dots (1+\text{ad}(w_{k_r}))} &= 1 + \text{ad}(x) + e_7^{(0, \dots, 0, 1)} + e_4^{(0, \dots, 0, 2)} + e_1^{(0, \dots, 0, 3)}. \end{aligned}$$

In other words the r conjugates are all in $1 + Q$. Hence if H is the subgroup of $\mathrm{GL}(V^*)$ generated by the r conjugates then

$$\gamma_{4r+2}(H) \leq \gamma_{4r+2}(1 + Q) \leq 1 + Q^{4r+2} = 1.$$

We have thus proved the following.

Proposition 2.9 *Let $(1 + \mathrm{ad}(x))^{g_1}, \dots, (1 + \mathrm{ad}(x))^{g_r}$ be any r conjugates of $1 + \mathrm{ad}(x)$ in G . Then the group generated by these conjugates is nilpotent of class at most $4r + 2$.*

Acknowledgments

The first author is partially supported by the “National Group for Algebraic and Geometric Structures, and their Applications” (GNSAGA - INdAM). Also, she would like to thank the Department of Mathematics at the University of Bath for its excellent hospitality while this research was conducted.

The second and the third author acknowledge the EPSRC (grant number 1652316) for their support.

References

- [1] R. Baer, Engelsche Elemente Noetherscher Gruppen, *Math. Ann.* **133** (1957), 256-270.
- [2] W. Burnside, On an unsettled question in the theory of discontinuous groups, *Quart. J. Pure Appl. Math.* **37** (1901), 230-238.
- [3] K. W. Gruenberg, The Engel elements of a soluble group, *Illinois J. Math.* **3** (1959), 151-169.
- [4] H. Heineken, Eine Bemerkung über engelshe Elemente, *Arch. Math.* **12** (1960), 176-178.
- [5] S. V. Ivanov, The free Burnside groups of sufficiently large exponents, *Int. J. Algebra and Comp.* **4** (1994), 1-308.
- [6] E. Jabara and G. Traustason, Left 3-Engel elements of odd order in groups, *Proc. Am. Soc.*, to appear.

- [7] I. G. Lysenok, Infinite Burnside groups of even period, *Izv. Math.* **60** (1996), 453-654.
- [8] M. Newell, On right-Engel elements of length three, Proc. Royal Irish Ac. **96A**, No.1 (1996), 17-24.
- [9] I. N. Sanov, Solution of Burnside's Problem for Exponent Four, Leningrad State Univ., *Ann. Maths. Ser.* **10** (1940), 166-170.
- [10] G. Tracey and G. Traustason, Left 3-Engel elements in groups of exponent 60, *Int. J. Algebra and Comp.* **28** No. 4 (2018), 673-695.
- [11] G. Traustason, Left 3-Engel elements in groups of exponent 5, *J. Algebra*, **414** (2014), 41-71.
- [12] G. Traustason, Engel Lie-algebras, *Quart. J. Math. Oxford. Ser. (2)*, **44** (1993), 355-384.