UNIVERSITY^{OF} BIRMINGHAM University of Birmingham Research at Birmingham

Engel elements in branch groups

Fernandez-Alcober, Gustavo; Noce, Marialaura; Tracey, Gareth

Citation for published version (Harvard):

Fernandez-Alcober, G, Noce, M & Tracey, G 2020, 'Engel elements in branch groups', *Journal of Algebra*, vol. 554, pp. 54-77.

Link to publication on Research at Birmingham portal

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

•Users may freely distribute the URL that is used to identify this publication.

Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)

•Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

ENGEL ELEMENTS IN WEAKLY BRANCH GROUPS

GUSTAVO A. FERNÁNDEZ-ALCOBER, MARIALAURA NOCE, AND GARETH M. TRACEY

ABSTRACT. We study properties of Engel elements in weakly branch groups, lying in the group of automorphisms of a spherically homogeneous rooted tree. More precisely, we prove that the set of bounded left Engel elements is always trivial in weakly branch groups. In the case of branch groups, the existence of non-trivial left Engel elements implies that these are all p-elements and that the group is virtually a p-group (and so periodic) for some prime p. We also show that the set of right Engel elements of a weakly branch group is trivial under a relatively mild condition. Also, we apply these results to well-known families of weakly branch groups, like the multi-GGS groups.

1. INTRODUCTION

A rapidly developing area of group theory studies the properties of branch groups, a special kind of groups acting on spherically homogeneous rooted trees. These groups, which were first defined by Grigorchuk at the Groups St Andrews conference in Bath in 1997, are generalizations of the famous p-groups constructed by Grigorchuk himself [9], and by Gupta and Sidki [13]. Despite their relatively recent introduction, branch groups have appeared in the literature in the past, without being explicitly defined. For instance, the class of branch groups contains one of the three classes of groups in John Wilson's famous characterisation of just infinite groups [18]. This is one of the primary reasons for their study. Another important motivation for studying branch groups comes from the remarkable properties that some of these groups can possess, like intermediate growth, amenability, the congruence subgroup property, or providing a negative answer to the General Burnside Problem. In this setting, one can also consider the larger family of *weakly branch groups*, which preserve many of the most interesting features enjoyed by branch groups. We refer the reader to Section 2 for a quick introduction to these classes of groups.

With this motivation in mind, the purpose of this paper is to investigate Engel elements in weakly branch groups. Given two elements g and x in a group G, we define [g, n x] for all $n \in \mathbb{N} \cup \{0\}$ by means of [g, 0 x] = g and, for $n \ge 1$,

 $[g_{,n} x] = [[g_{,n-1} x], x] = [[\dots [[g, x], x], \dots], x],$ (where x appears n times).

²⁰¹⁰ Mathematics Subject Classification. 20E08, 20F45.

Key words and phrases. Branch groups, Engel elements.

The first two authors are supported by the Spanish Government, grant MTM2017-86802-P, partly with FEDER funds. The first author is also supported by the Basque Government, grant IT974-16. The second author is partially supported by the "National Group for Algebraic and Geometric Structures, and their Applications" (GNSAGA - INdAM). The third author acknowledges the EPSRC (grant number 1652316) for their support.

Engel conditions in group theory have to do with the triviality of these iterated left normed commutators. If $[g,_n x] = 1$ for some $n \in \mathbb{N}$, we say that x is Engel on g, and the smallest such n is the Engel degree of x on g. If x is Engel on all elements $g \in G$, we say that x is a left Engel element of G. Observe that the Engel degree of x can vary as g runs over G, and in principle could be unbounded. If there is a bound for the Engel degrees of x, i.e. if there exists $n \in \mathbb{N}$ such that $[g,_n x] = 1$ for all $g \in G$, we say that x is a bounded left Engel element of G. We denote by L(G) and $\overline{L}(G)$ the sets of left Engel elements and bounded left Engel elements of G, respectively. On the other hand, if $g \in G$ is such that every $x \in G$ is Engel on g, we say that g is a right Engel element of G. Bounded right Engel elements are defined in an obvious way. We write R(G) and $\overline{R}(G)$ for the sets of right Engel elements and bounded right Engel elements of G. Observe that $R(G) \subseteq L(G)^{-1}$ [17, 12.3.1], and that obviously $\overline{L}(G) \subseteq L(G)$ and $\overline{R}(G) \subseteq R(G)$. In particular, if L(G) = 1 then all four Engel sets are trivial.

We say that G is an Engel group if L(G) = G (or equivalently R(G) = G). On the other hand, if the identity $[g, x, .^n, x] = 1$ holds for all $g, x \in G$, i.e. if every $x \in G$ is a bounded left Engel element with a common bound for all $g \in G$ (or equivalently with the right Engel condition), then G is said to be an *n*-Engel group.

In every group G, the sets L(G), $\overline{L}(G)$, R(G), and $\overline{R}(G)$ contain some distinguished subgroups, namely the Hirsch-Plotkin radical, the Baer radical, the hypercenter and the ω -center, respectively. In his book A Course in the Theory of Groups, D.J.S. Robinson considers it one of the major goals of Engel theory to find conditions which will guarantee that these four sets of Engel elements coincide with the corresponding subgroups [17, Section 12.3]. For example, Baer proved that this is the case if G satisfies the maximal condition (see [3, Satz L'] or [17, 12.3.7]); in particular, L(G) coincides with the Fitting subgroup if G is finite. However, these equalities do not hold in general. It is then natural to ask whether L(G), $\overline{L}(G)$, R(G), and $\overline{R}(G)$ are always subgroups of G, and it was not until recently that the first counterexamples were found. It is here where branch groups come into play in Engel theory.

Let \mathfrak{G} be the first Grigorchuk group. This is a branch group acting on the binary tree, introduced by Grigorchuk [9] in 1980. In 2006 Bludov announced [6] that the wreath product $\mathfrak{G} \wr D_8$, with the natural action of D_8 on 4 points, can be generated by Engel elements but is not an Engel group. In particular, $L(\mathfrak{G})$ is not a subgroup. This example was never published, but ten years later, Bartholdi [4] showed that

$$\mathcal{L}(\mathfrak{G}) = \{ x \in \mathfrak{G} \mid x^2 = 1 \}$$

and, as a consequence, that $L(\mathfrak{G})$ is not a subgroup. To date, it is still an open question whether $\overline{L}(G)$, R(G) and $\overline{R}(G)$ are always subgroups.

The other major question in Engel theory is whether Engel groups are locally nilpotent. The answer is negative in general, the main example being the socalled Golod-Shafarevich groups. These groups also provide a negative answer to the General Burnside Problem (GBP), which can be equivalently formulated as the question of whether periodic groups are locally finite. This resemblance between Engel and Burnside problems, together with the fact that many groups answering GBP in the negative are branch groups, and Bartholdi's result on left Engel elements of the Grigorchuk group, makes it natural to study the behaviour of the Engel sets L(G), $\overline{L}(G)$, R(G) and $\overline{R}(G)$ in (weakly) branch groups. This is the specific goal that we are addressing in this paper.

Before proceeding to state our main theorems, let us mention some results in the literature regarding Engel elements in groups of automorphisms of spherically homogeneous rooted trees. In the aforementioned paper, Bartholdi also proved that, if G is the Gupta-Sidki 3-group, then L(G) = 1. On the other hand, in [7], Garreta and the first two authors proved that again L(G) = 1 if G is any fractal subgroup of a Sylow pro-p subgroup of the group of automorphisms of the p-adic tree satisfying the condition $|G': \operatorname{st}_G(1)'| = \infty$, and in particular if Gis non-abelian and has torsion-free abelianization. Also, Tortora and the second author showed in [16] that $\overline{L}(\mathfrak{G}) = \mathbb{R}(\mathfrak{G}) = 1$ for the Grigorchuk group \mathfrak{G} . As we next see, the situation in these classes of groups generalises to a great extent to weakly branch groups, which have a tendency to have trivial Engel sets.

Our first main result reads as follows.

Theorem A. Let G be a weakly branch group. Then the following hold:

- (i) $\overline{\mathrm{L}}(G) = 1$.
- (ii) If the set of finite order elements of L(G) is non-trivial then it is a p-set for some prime p, and the rigid stabilizer rst_G(n) is a p-group for some n ≥ 1.

Thus even if weakly branch groups provide examples in which L(G) is not a subgroup, they cannot be used to obtain similar examples for $\overline{L}(G)$. This result can be interpreted in a similar vein to the fact that weakly branch groups, being residually finite, cannot provide examples of finitely generated infinite groups of finite exponent: the "problem" in both cases is boundedness. On the other hand, part (ii) of Theorem A raises the following question.

Question 1. Can a weakly branch group G contain left Engel elements of infinite order? If the answer is negative, then L(G) consists entirely of p-elements for some prime p.

If instead of weakly branch the group is actually branch, then we have the following stronger version of Theorem A.

Theorem B. Let G be a branch group. If $L(G) \neq 1$ then G is periodic and there exists a prime p such that:

- (i) L(G) consists of p-elements.
- (ii) G is virtually a p-group.

Compare the results in Theorem B with the situation in the Grigorchuk group. In that case, $L(\mathfrak{G})$ consists of all elements of order 2 in \mathfrak{G} , and \mathfrak{G} is a 2-group. On the other hand, the prime p can be arbitrary in (i) and (ii): if \mathcal{F}_p is the group of p-finitary automorphisms of a p-adic tree then it is easy to see that $L(\mathcal{F}_p) = \mathcal{F}_p$, and this is a p-group. Observe however that, contrary to \mathfrak{G} , the group \mathcal{F}_p is not finitely generated.

Question 2. Are there any finitely generated (weakly) branch groups for which the set L(G) is non-trivial and consists of p-elements for an odd prime p?

In the following theorem we consider right Engel elements in weakly branch groups under a relatively mild condition.

Theorem C. Let G be a weakly branch group. If the rigid stabilizer $rst_G(n)$ is not an Engel group for any $n \in \mathbb{N}$, then R(G) = 1.

Question 3. Is R(G) = 1 for every finitely generated (weakly) branch group? By Theorem C, this seems closely linked to this other question: can a finitely generated (weakly) branch group be Engel?

Again, without finite generation, the group \mathcal{F}_p shows that the answer is negative in both cases. Regarding the last question, observe that weakly branch groups cannot satisfy a law (see [1, Corollary 1.4] or [14]) and so cannot be *n*-Engel for a fixed *n*. Thus we are asking whether finite generation makes it impossible for them to be Engel as well.

As an application of Theorems A, B, and C, we get the following corollary, which provides information about Engel elements in some specific families of weakly branch groups. The definition of these families is given either in Section 2 or right before the proof of the corresponding result.

Corollary D. Let \mathcal{T} be a spherically homogeneous rooted tree. Then the following hold:

- (i) If G is an infinitely iterated wreath product of finite transitive permutation groups of degree at least 2, then L(G) = 1. This applies in particular to the whole group of automorphisms of T, and also to its Sylow pro-p subgroups if T is a p-adic tree, where p is a prime.
- (ii) If F is the group of finitary automorphisms of T, and there are infinitely many levels in which the number of descendants is greater than 2, then L(F) = 1. If T is a p-adic tree and F_p is the group of p-finitary automorphisms of T, then L(F_p) = 1.
- (iii) If \mathcal{H} is the Hanoi Tower group then $L(\mathcal{H}) = 1$.
- (iv) If G is a multi-GGS groups then R(G) = 1, and if G is furthermore non-periodic then L(G) = 1.

We conclude this introduction by indicating how the paper is organised. In Section 2 we first give some generalities about groups of automorphisms of spherically homogeneous rooted trees, with special emphasis on weakly branch groups. Then we provide several results regarding orbits of such automorphisms that will be essential later on. Our approach to the study of Engel elements in weakly branch groups is through the reduction to wreath products. Section 3 is devoted to a careful analysis of the scenarios that will arise when we apply this kind of reduction. Finally, in Section 4 we prove Theorems A and B, regarding left Engel elements in weakly branch groups, and then in Section 5 we obtain Theorem C about right Engel elements. The proof of the applications given in Corollary D is split between these two sections.

Notation and terminology. If $f: X \to Y$ and $g: Y \to Z$ are two maps, we write fg for their composition instead of $g \circ f$. As usual, S_n stands for the symmetric group on n letters. We denote the direct product of groups G_1, \ldots, G_n by $\prod_{i=1}^n G_i$. Given a group G, an element $g \in G$ is a *p*-element, where p is a prime,

if its order is a power of p, and a subset X of G is a p-set if every element of X is a p-element. Also, if G is finite, F(G) denotes the Fitting subgroup of G.

2. Automorphisms of spherically homogeneous rooted trees

In this section, we first give some notation and general facts about groups of automorphisms of a spherically homogeneous rooted tree, and more specifically, about (weakly) branch groups. For further information on the topic one can see, for example, [5] or [10]. Then we provide some results regarding orbits of automorphisms of such trees that will be needed in the following sections.

Let $\overline{d} = \{d_n\}_{n=1}^{\infty}$ be an infinite sequence of integers greater than 1, and let $d = d_1$. We write $\mathcal{T}_{\overline{d}}$ to denote the *spherically homogeneous rooted tree* corresponding to \overline{d} . This is a rooted tree where all vertices at level n (i.e. at distance n from the root) have the same number d_{n+1} of immediate descendants. If \overline{d} takes the constant value d, we write \mathcal{T}_d for $\mathcal{T}_{\overline{d}}$, and we call it the *d*-adic *tree*. In order to ease notation, and unless it is strictly necessary to make the sequence \overline{d} explicit, all throughout the paper we will simply write \mathcal{T} to denote an arbitrary spherically homogeneous rooted tree. Also we will write $V(\mathcal{T})$ for the set of vertices of \mathcal{T} and, for every $n \in \mathbb{N}$, we let \mathcal{L}_n be the set of all vertices on the *n*th level of \mathcal{T} .



FIGURE 1. A spherically homogeneous rooted tree

Let Aut \mathcal{T} be the group of automorphisms of \mathcal{T} (i.e. bijective maps from $V(\mathcal{T})$ to itself that preserve the root and incidence) under the operation of composition. Every $f \in \operatorname{Aut} \mathcal{T}$ can be described by providing, at every vertex v of the tree, the permutation $f_{(v)}$ that indicates how f sends the descendants of v onto the descendants of f(v). This permutation is called the *label* of f at v, and if v lies at level n then $f_{(v)} \in S_{d_{n+1}}$. The collection of all labels of f constitutes the *portrait* of f, and there is a one-to-one correspondence between automorphisms of \mathcal{T} and portraits. An automorphism of \mathcal{T} is called *finitary* if it has finitely many non-trivial labels in its portrait. Finitary automorphisms form a locally finite subgroup \mathcal{F} of Aut \mathcal{T} . If \mathcal{T} is a p-adic tree for a prime p and we fix a p-cycle σ in S_p , the group \mathcal{F}_p of finitary automorphisms whose labels are all powers of σ constitute a subgroup of \mathcal{F} . We call this the *group of p-finitary automorphisms* of \mathcal{T} . (We give no reference to σ , since different

choices of the *p*-cycle give rise to isomorphic groups.) Observe that \mathcal{F}_p is locally a finite *p*-group.

We write $\operatorname{st}(v)$ for the stabilizer of $v \in V(\mathcal{T})$ in $\operatorname{Aut} \mathcal{T}$ and $\operatorname{st}(n)$ for the pointwise stabilizer of \mathcal{L}_n , i.e.

$$\operatorname{st}(n) = \bigcap_{v \in \mathcal{L}_n} \operatorname{st}(v).$$

The latter is a normal subgroup of finite index of Aut \mathcal{T} . The factor group Aut $\mathcal{T}/\operatorname{st}(n)$ is naturally isomorphic to the automorphism group of the finite tree consisting of all levels of \mathcal{T} up to (and including) the *n*th level. Then Aut \mathcal{T} is isomorphic to the inverse limit of these finite groups, and is so a profinite group. Also, we have

Aut
$$\mathcal{T} \cong \cdots (S_{d_n} \wr (\cdots (S_{d_3} \wr (S_{d_2} \wr S_{d_1}))))) \cdots)$$

where the iterated wreath product is permutational at every step. If we consider the *p*-adic tree \mathcal{T}_p , where *p* is a prime, and we consider a fixed *p*-cycle $\sigma \in S_p$, then the set Γ_p of all automorphisms of \mathcal{T} with labels in $\langle \sigma \rangle$ is a Sylow pro*p* subgroup of Aut \mathcal{T}_p . We say that Γ_p is a standard Sylow pro-*p* subgroup of Aut \mathcal{T}_p . Observe that

$$\Gamma_p \cong \cdots (C_p \wr (\cdots (C_p \wr (C_p \wr C_p)) \cdots)) \cdots$$

Let \mathcal{T}_v be the subtree hanging from the vertex v of the tree. We have $\mathcal{T}_u \cong \mathcal{T}_v$ for any two vertices u, v on the same level, and we denote by $\mathcal{T}_{\langle n \rangle}$ any tree isomorphic to a subtree with root in \mathcal{L}_n . If s denotes the shift operator that erases the first term of a sequence, then $\mathcal{T}_{\langle n \rangle}$ is isomorphic to the spherically homogeneous tree defined by the sequence $s^n(\overline{d})$.

Every $f \in \operatorname{Aut} \mathcal{T}$ naturally induces a bijection between \mathcal{T}_v and $\mathcal{T}_{f(v)}$ which, under the identification of these trees with $\mathcal{T}_{\langle n \rangle}$, defines an automorphism f_v of $\mathcal{T}_{\langle n \rangle}$. This is called the *section* of f at v. Sections satisfy the following rules, for all $f, g \in \operatorname{Aut} \mathcal{T}$ and $v \in V(\mathcal{T})$:

$$(fg)_v = f_v g_{f(v)}$$

and

$$(f^g)_{g(v)} = (g_v)^{-1} f_v g_{f(v)}.$$

As a consequence, if $f \in \operatorname{st}(v)$, we get

(2.1)
$$(f^g)_{q(v)} = (f_v)^{g_v}.$$

If f fixes the vertex v, then the section f_v is nothing but the restriction of f to \mathcal{T}_v . The assignment $f \mapsto f_v$ induces a homomorphism $\psi_v \colon \operatorname{st}(v) \to \operatorname{Aut} \mathcal{T}_{\langle n \rangle}$, and the map

$$\psi_n : \operatorname{st}(n) \longrightarrow \operatorname{Aut} \mathcal{T}_{\langle n \rangle} \times \overset{d_1 \dots d_n}{\cdots} \times \operatorname{Aut} \mathcal{T}_{\langle n \rangle}$$
$$f \longmapsto (f_v)_{v \in \mathcal{L}_n}.$$

is an isomorphism. If n = 1 we simply write ψ for ψ_1 . In the case of a *d*-adic tree, we get

$$\operatorname{st}(n) \cong \operatorname{Aut} \mathcal{T} \times \stackrel{d^n}{\cdots} \times \operatorname{Aut} \mathcal{T}.$$

Observe also that $\operatorname{Aut} \mathcal{T}$ splits over $\operatorname{st}(n)$ for every $n \ge 1$. One can take as a complement the subgroup

$$H_n = \{ f \in \operatorname{Aut} \mathcal{T} \mid f_v = 1 \text{ for all } v \in \mathcal{L}_n \}$$

= $\{ f \in \operatorname{Aut} \mathcal{T} \mid f_{(v)} = 1 \text{ for all } v \in \bigcup_{i \ge n} \mathcal{L}_i \}.$

The automorphisms in H_1 are called *rooted automorphisms* of \mathcal{T} . They act on \mathcal{T} by permuting rigidly the subtrees hanging from the root according to some permutation of S_d . Every automorphism $f \in \operatorname{Aut} \mathcal{T}$ can be uniquely written in the form gh, where $g \in \operatorname{st}(1)$ and h is rooted. If $\psi(g) = (g_1, \ldots, g_d)$ and h corresponds to a permutation $\sigma \in S_d$, we use the following shorthand notation to denote f:

$$f = (g_1, \ldots, g_d)\sigma$$

Now let G be a subgroup of Aut \mathcal{T} . We set $\operatorname{st}_G(n) = \operatorname{st}(n) \cap G$ for all $n \ge 1$. If v is a vertex of \mathcal{T} , the *rigid stabilizer* of v in G is defined as follows:

 $\operatorname{rst}_G(v) = \{g \in G \mid g(u) = u \text{ for all } u \text{ lying outside } \mathcal{T}_v\}.$

If V is a set of vertices, all lying on the same level of \mathcal{T} , we set

$$\operatorname{rst}_G(V) = \langle \operatorname{rst}_G(v) \mid v \in V \rangle,$$

the rigid stabilizer of V in G. It turns out that

$$\operatorname{rst}_G(V) = \prod_{v \in V} \operatorname{rst}_G(v).$$

We write $\operatorname{rst}_G(n)$ for the rigid stabilizer of \mathcal{L}_n , and call it the *nth rigid stabilizer* of G. It is the direct product of the rigid stabilizers of all vertices of \mathcal{L}_n , and it is the largest "geometrical" direct product inside $\operatorname{st}_G(n)$, in the sense that a subgroup H of $\operatorname{st}_G(n)$ satisfies

$$\psi_n(H) = \prod_{v \in \mathcal{L}_n} H_v$$

with $H_v \leq \operatorname{Aut} \mathcal{T}_v$ if and only if $H \leq \operatorname{rst}_G(n)$. Obviously, if G is the whole of $\operatorname{Aut} \mathcal{T}$ then the *n*th rigid stabilizer coincides with the *n*th level stabilizer. However, this is not usually the case for arbitrary subgroups of $\operatorname{Aut} \mathcal{T}$.

By (2.1), we have

(2.2)
$$\operatorname{rst}_G(v)^g = \operatorname{rst}_G(g(v))$$

for every $v \in V(\mathcal{T})$ and $g \in G$. Thus if G is spherically transitive, i.e. if G acts transitively on every \mathcal{L}_n , then each level rigid stabilizer $\operatorname{rst}_G(n)$ is a direct product of isomorphic subgroups for all $n \in \mathbb{N}$. We are now ready to introduce the class of groups that are the object of our study.

Definition 2.1. Let G be a spherically transitive subgroup of Aut \mathcal{T} . Then: (a) If $|G: \operatorname{rst}_G(n)| < \infty$ for all $n \in \mathbb{N}$, we say that G is a branch group. (b) If $\operatorname{rst}_G(n) \neq 1$ for all $n \in \mathbb{N}$, we say that G is a weakly branch group.

Notice that all rigid level stabilizers in a weakly branch group are infinite. Also, since spherically transitive groups are infinite, branch groups are obviously weakly branch.

After this quick introduction to groups of automorphisms of a spherically homogeneous rooted tree, we start developing the tools that we will use in the proof of Theorems A, B, and C. A key ingredient in our approach to Engel problems in weakly branch groups is the reduction of the action of an automorphism f from the whole tree to one or several "reduced trees" determined by some special orbits of f on $V(\mathcal{T})$. For this reason, we start by describing some properties of orbits of automorphisms of \mathcal{T} .

Definition 2.2. If $f \in \operatorname{Aut} \mathcal{T}$ and $v \in V(\mathcal{T})$, the *f*-orbit of v is the orbit of v under the action of $\langle f \rangle$ on $V(\mathcal{T})$, i.e. the set $\{f^i(v) \mid i \in \mathbb{Z}\}$. The *f*-orbit is *trivial* if it consists of only one vertex, that is, if f(v) = v.

In the statement of the following lemma, we consider the least common multiple of an unbounded family of positive integers to be infinity.

Lemma 2.3. Let $f \in \operatorname{Aut} \mathcal{T}$ and, for every vertex $v \in V(\mathcal{T})$, let \mathcal{O}_v be the f-orbit of v. Then the following hold:

- (i) If w is a descendant of v, then $|\mathcal{O}_v|$ divides $|\mathcal{O}_w|$.
- (ii) $|f| = \operatorname{lcm}(|\mathcal{O}_v| \mid v \in V(\mathcal{T})).$
- (iii) If |f| is finite then there exists a finite subset V of $V(\mathcal{T})$ satisfying that $|f| = \operatorname{lcm}(|\mathcal{O}_v| \mid v \in V)$ and that, whenever w is a descendant of a vertex $v \in V$, we have $|\mathcal{O}_w| = |\mathcal{O}_v|$. Also if f is non-trivial then all the orbits \mathcal{O}_v with $v \in V$ are non-trivial. Furthermore, V can be chosen to lie in \mathcal{L}_n for some n.

Proof. (i) This is obvious by the orbit-stabilizer theorem, since $st(w) \subseteq st(v)$.

(ii) Set $H = \langle f \rangle$. Then $|\mathcal{O}_v| = |H/\operatorname{st}_H(v)|$ for all $v \in V(\mathcal{T})$. The natural map φ from H to the cartesian product of finite groups $\prod_{v \in V(\mathcal{T})} H/\operatorname{st}_H(v)$ is injective, since the intersection of all vertex stabilizers is trivial. Consequently

 $|f| = |\varphi(f)| = \operatorname{lcm}(|f\operatorname{st}_{H}(v)| \mid v \in V(T)) = \operatorname{lcm}(|H/\operatorname{st}_{H}(v)| \mid v \in V(T)),$

which proves the result.

(iii) Let $L = \{ |\mathcal{O}_v| \mid v \in V(\mathcal{T}) \}$. If |f| is finite then, by (ii), it can be achieved as the least common multiple of a finite subset of L. Let k be the minimum cardinality of such a subset and let

$$\mathcal{S} = \{ S \subseteq L \mid |S| = k \text{ and } \operatorname{lcm}(S) = |f| \}.$$

Observe that \mathcal{S} is a finite set.

We introduce a relation \leq_{d} in S by letting $S \leq_{d} T$ if there exists a bijection $\alpha: S \to T$ such that $s \mid \alpha(s)$ for all $s \in S$. By (i), this models the situation when we pass from the orbits of a set of vertices to the orbits of a set of descendants of those vertices. We claim that \leq_{d} is an order relation in S. Obviously, only antisymmetry needs to be checked. Assume that $\alpha: S \to T$ and $\beta: T \to S$ are such that $s \mid \alpha(s)$ and $t \mid \beta(t)$ for all $s \in S$ and $t \in T$. Then s divides $\beta(\alpha(s))$ and, if they are not equal, we get $lcm(S \setminus \{s\}) = |f|$. This is contrary to the minimality condition imposed on k. Thus $\beta(\alpha(s)) = s$ and, since $s \mid \alpha(s)$ and $\alpha(s) \mid \beta(\alpha(s))$, we obtain that $\alpha(s) = s$ for all $s \in S$. We conclude that S = T, which proves antisymmetry of \leq_{d} .

Now choose S in S that is maximal with respect to the order \leq_d , and let $V = \{v_1, \ldots, v_k\} \subseteq V(\mathcal{T})$ be such that $S = \{|\mathcal{O}_{v_1}|, \ldots, |\mathcal{O}_{v_k}|\}$. Consider an arbitrary set of vertices $W = \{w_1, \ldots, w_k\}$, where each w_i is a descendant of v_i , and let $T = \{|\mathcal{O}_{w_1}|, \ldots, |\mathcal{O}_{w_k}|\}$. Then $S \leq_d T$ and, by the maximality of S, we

have S = T. This implies that $|\mathcal{O}_{w_i}| = |\mathcal{O}_{v_i}|$ for all $i = 1, \ldots, k$. Observe also that the minimality of k implies that, if f is non-trivial, no orbit \mathcal{O}_v with $v \in V$ is of length 1. Hence V satisfies the properties stated in (iii).

Finally, observe that also the set W satisfies the required properties. Thus by considering, for a suitable n, a subset of \mathcal{L}_n consisting of one descendant of each vertex in V, we may assume that $V \subseteq \mathcal{L}_n$.

Vertices and orbits as in part (iii) of the previous lemma will play a fundamental role in the rest of the paper, and it is convenient to introduce some terminology.

Definition 2.4. Let $f \in \operatorname{Aut} \mathcal{T}$ and let \mathcal{O} be an *f*-orbit. We say that \mathcal{O} is *totally splitting* if for every descendant w of a vertex $v \in \mathcal{O}$, the length of the *f*-orbit of w is equal to $|\mathcal{O}|$.

Equivalently, an f-orbit \mathcal{O} is totally splitting when the set of descendants of the vertices in \mathcal{O} at every level of the tree splits into the maximum possible number of f-orbits.

Definition 2.5. Let $f \in \operatorname{Aut} \mathcal{T}$ be an automorphism of finite order. If V is a finite set of vertices satisfying the conditions in (iii) of Lemma 2.3, all of them lying on the same level of \mathcal{T} , we say that V is a *fundamental system of vertices* for f.

Next we give a sufficient condition for two automorphisms of \mathcal{T} to generate a wreath product.

Lemma 2.6. Let $f \in \operatorname{Aut} \mathcal{T}$ be an automorphism of finite order m, and assume that the f-orbit of a vertex $v \in V(\mathcal{T})$ has length m. Then for every $g \in \operatorname{rst}(v)$, the subgroup $\langle g, f \rangle$ of $\operatorname{Aut} \mathcal{T}$ is isomorphic to the regular wreath product $\langle g \rangle \wr \langle f \rangle$.

Proof. Let \mathcal{O} be the *f*-orbit of *v*. Since $|\mathcal{O}| = |f|$, we have $\langle f \rangle \cap \operatorname{st}(v) = 1$. As a consequence, if *v* lies at level *n* of the tree, also $\langle f \rangle \cap \operatorname{st}(n) = 1$ and

(2.3)
$$\langle g, f \rangle = \langle f \rangle \langle g, g^f, \dots, g^{f^{m-1}} \rangle = \langle f \rangle \ltimes \langle g, g^f, \dots, g^{f^{m-1}} \rangle,$$

since $g \in \operatorname{rst}(v)$ implies that $\langle g, g^f, \dots, g^{f^{m-1}} \rangle \subseteq \operatorname{st}(n)$.

Now set $v_i = f^i(v)$ for all $i \in \mathbb{Z}$, so that $\mathcal{O} = \{v_0, v_1, \ldots, v_{m-1}\}$. Since $g \in \operatorname{rst}_G(v)$, from (2.2) we get $g^{f^i} \in \operatorname{rst}_G(v_i)$ for all $i = 0, \ldots, m-1$, and then

$$\langle g^{f^i} \rangle \cap \langle g, g^f, \dots, g^{f^{i-1}} \rangle \subseteq \operatorname{rst}_G(v_i) \cap \operatorname{rst}_G(\{v_1, \dots, v_{i-1}\}) = 1.$$

Also $[g^{f^i}, g^{f^j}] = 1$ for every $i, j \in \{0, \dots, m-1\}$. It follows that

$$\langle g, g^f, \dots, g^{f^{m-1}} \rangle = \langle g \rangle \times \langle g^f \rangle \times \dots \times \langle g^{f^{m-1}} \rangle,$$

= *a*, we conclude from (2.3) that $\langle a, f \rangle \cong \langle a \rangle \setminus \langle f \rangle.$

and since $g^{f^m} = g$, we conclude from (2.3) that $\langle g, f \rangle \cong \langle g \rangle \wr \langle f \rangle$.

The result in Lemma 2.6 raises the question of whether an automorphism $f \in \operatorname{Aut} \mathcal{T}$ of finite order m must have a regular orbit on $V(\mathcal{T})$, i.e. an orbit of length m. This is clearly the case if m is a prime power, by (ii) of Lemma 2.3, but it usually fails otherwise. Indeed, one can consider for example a rooted automorphism corresponding to a permutation whose order is strictly bigger than the lengths of its disjoint cycles. However, as we see in Lemma 2.9 below, it is always possible to derive a collection of automorphisms f_i from f, acting

not on \mathcal{T} but on some other rooted trees \mathcal{R}_i obtained from \mathcal{T} , and having the property that every f_i has a regular orbit on $V(\mathcal{R}_i)$. These automorphisms f_i will allow us to study Engel conditions regarding f by using Lemma 2.6.

As we will see, Lemma 2.9 is essentially a reformulation of (iii) of Lemma 2.3. Before proceeding we need to introduce the concept of reduced tree. Note that reduced trees are somehow related to the trees obtained by deletion of layers defined by Grigorchuk and Wilson in [12].

Definition 2.7. Let V be a subset of vertices of \mathcal{T} , all lying on the same level n. We define the *reduced tree* of \mathcal{T} at V, denoted by $\mathcal{R}(V)$, as the rooted tree consisting of the subtrees \mathcal{T}_v for $v \in V$, all connected to a common root. In other words, the set of vertices of $\mathcal{R}(V)$ is

$$\{\emptyset\} \cup \{vw \mid v \in V, \ w \in \mathcal{T}_{s^{n+1}(\overline{d})}\},\$$

where as before s denotes the shift operator on sequences.

For example, in the following figure, we consider the rooted automorphism f of the ternary tree \mathcal{T}_3 corresponding to the permutation (1 2 3) and we show in red the reduced tree at the orbit of the vertex 13:



FIGURE 2. An f-orbit and its corresponding reduced tree

Every $f \in \operatorname{Aut} \mathcal{T}$ such that f(V) = V induces by restriction an automorphism $f_V \in \operatorname{Aut} \mathcal{R}(V)$. Clearly, the map $\Phi_V : f \mapsto f_V$ is a homomorphism of groups. The effect of Φ_V is to focus on the action of f only on the subtrees \mathcal{T}_v with $v \in V$, so to speak. We will use reduced trees mainly in the case where V is an orbit of f.

Remark 2.8. If v is a vertex of the reduced tree $\mathcal{R}(V)$ and $f \in \operatorname{Aut} \mathcal{T}$ is such that f(V) = V, then the f_V -orbit of v coincides with the f-orbit of v as a vertex in $V(\mathcal{T})$. In particular, if \mathcal{O} is a totally splitting f-orbit and we consider the induced automorphism $x = \Phi_{\mathcal{O}}(f)$ of $\mathcal{R}(\mathcal{O})$, then (ii) of Lemma 2.3 implies that $|x| = |\mathcal{O}|$. In other words, \mathcal{O} is a regular orbit of x in $\mathcal{R}(\mathcal{O})$.

Given a subgroup G of Aut \mathcal{T} , we write G_V for the image of the setwise stabilizer of V in G under the homomorphism Φ_V . In other words,

$$G_V = \{ f_V \mid f \in G \text{ and } f(V) = V \}.$$

Then G_V is a subgroup of Aut $\mathcal{R}(V)$, and for every vertex $v \in V$ we have $\Phi_V(\operatorname{rst}_G(v)) \subseteq \operatorname{rst}_{G_V}(v)$ (the inclusion can be proper, since there can be automorphisms in G whose action is trivial on \mathcal{T}_w for every $w \neq v$ with $w \in V$, but non-trivial for some $w \notin V$).

On the other hand, if $f \in G$ stabilizes the set V and $x = \Phi_V(f)$ is the induced automorphism of $\mathcal{R}(V)$, then $f \in \mathcal{L}(G)$ or $f \in \overline{\mathcal{L}}(G)$ imply that $x \in \mathcal{L}(H)$ or $x \in \overline{\mathcal{L}}(H)$, respectively. In particular, by choosing V to be an f-orbit, this will allow us to transfer the analysis of a given Engel element in a subgroup of Aut \mathcal{T} to a more restricted situation where, for example, the Engel element acts transitively on the first level of the tree.

Actually the most convenient strategy is to reduce the tree to non-trivial totally splitting f-orbits, since the induced automorphisms will then have regular orbits. More precisely, we will rely on the following lemma, which is basically a rephrasing of part of Lemma 2.3 in the language of reduced trees.

Lemma 2.9. Let $f \in \operatorname{Aut} \mathcal{T}$ be an automorphism of finite order m > 1 and let $\{v_1, \ldots, v_k\}$ be a fundamental system of vertices for f. For every $i = 1, \ldots, k$, let \mathcal{O}_i be the f-orbit of v_i , set $\mathcal{R}_i = \mathcal{R}(\mathcal{O}_i)$, and let f_i be the automorphism of \mathcal{R}_i induced by f. Then the following hold:

- (i) $lcm(|O_1|, ..., |O_k|) = m.$
- (ii) \mathcal{O}_i is a non-trivial totally splitting f-orbit for every $i = 1, \ldots, k$.
- (iii) $|f_i| = |\mathcal{O}_i|$ for every $i = 1, \dots, k$.

Proof. The first two items follow from (iii) of Lemma 2.3, and (iii) from Remark 2.8. \Box

3. Some properties of Engel elements in wreath products

In this section we prove several results regarding Engel elements in wreath products. These will provide the basis for the proof of the main theorems in this paper, which will be addressed in Sections 4 and 5.

We start by studying left Engel elements lying outside the base group of a regular wreath product of two cyclic groups. To this purpose, we rely on the paper [15] by Liebeck.

Lemma 3.1. Let $X = \langle x \rangle$ and $Y = \langle y \rangle$ be two non-trivial cyclic groups, where X is finite, and let $W = Y \wr X$ be the corresponding regular wreath product. If $x \in L(W)$ then X and Y are finite p-groups for some prime p. Furthermore, the Engel degree of x on g = (y, 1, ..., 1) is equal to

$$|x| + \frac{1}{p}(\log_p |y| - 1)(p - 1)|x|.$$

Proof. Let m be the order of x, and let p be an arbitrary prime divisor of m. Also, write d for the Engel degree of x on g = (y, 1, ..., 1).

First of all, suppose that Y is finite. Then W is finite and, by Baer's theorem mentioned in the introduction, x lies in the Fitting subgroup F(W). We claim that Y is then a p-group. To this purpose, assume that |Y| is divisible by a prime $q \neq p$, and let $Z = \langle z \rangle \neq 1$ be the subgroup of Y of order q. Consider the direct product Z^X inside the base group of W. Since Z^X is abelian and normal in W, it lies in F(W). Now Z^X is a q-group and $x_p = x^{m/p}$ is a p-element, and both lie in the nilpotent group F(W). It follows that x_p centralizes Z^X , which is clearly a contradiction, since x_p does not commute with (z, 1, ..., 1). This proves the claim, and since this property holds for every prime divisor of m, it also follows that X is a p-group. Observe that, since both X and Y are finite p-groups, the proof of Theorem 5.1 of [15] yields that

(3.1)
$$d = m + \frac{1}{p} (\log_p |y| - 1)(p - 1)m$$

in this case.

Now it is easy to see that Y cannot be infinite. For a contradiction, suppose that Y is infinite and consider a prime q different from p. Then the wreath product $W_q = (Y/Y^q) \wr X$ can be seen as a factor group of W, and so x is a left Engel element in W_q . Since $|Y/Y^q| = q$ and p divides |X|, we get a contradiction with the previous paragraph. \Box

Now we digress from Engel elements for a moment, but still working with wreath products of cyclic groups, in order to prove that rigid stabilizers of weakly branch groups are not only infinite, but have infinite exponent (Proposition 3.3 below).

Lemma 3.2. Let $X = \langle x \rangle$ and $Y = \langle y \rangle$ be two finite cyclic groups, where Y is non-trivial, and let $W = Y \wr X$ be the corresponding regular wreath product. If g = (y, 1, ..., 1) then |xg| > |x|.

Proof. Set m = |x| and let $n \in \{1, ..., m\}$ be arbitrary. Then

$$(xg)^n = x^n g_1^n g_2^{\binom{n}{2}} \dots g_{n-1}^{\binom{n}{n-1}} g_n$$

where $g_1 = g$ and $g_i = [g, x, \stackrel{i-1}{\ldots}, x]$ for every $i = 2, \ldots, n$. Now observe that each g_i is of the form

$$g_i = (*, \ldots, *, y, 1, \ldots, 1),$$

where we use * to denote unspecified powers of y, and y occupies the *i*th position. It follows that

$$(xg)^n = x^n(*, \dots, *, y, 1, \dots, 1),$$

where y appears at the nth position. In particular, $(xg)^n \neq 1$ for $1 \leq n \leq m$, and consequently |xg| > m, as desired.

Proposition 3.3. Let G be a weakly branch group. Then the exponent of $rst_G(n)$ is infinite for every $n \in \mathbb{N}$.

Proof. By way of contradiction, assume that $\operatorname{rst}_G(n)$ has finite exponent. Thus $\operatorname{rst}_G(n)$ is periodic and there is a bound for the orders of its elements. For every $k \geq n$, let π_k be the (finite) set of prime divisors of the orders of the elements of $\operatorname{rst}_G(k)$. Then $\{\pi_k\}_{k\geq n}$ is a decreasing sequence of non-empty finite sets and consequently their intersection is also non-empty. Let p be a prime in $\cap_{k>n} \pi_k$.

Consider a *p*-element $f \in \operatorname{rst}_G(n)$ of maximum order, say *m*. Since the order of *f* is the least common multiple of the orders of the components of $\psi_n(f)$, we may assume without loss of generality that $f \in \operatorname{rst}_G(u)$ for some vertex *u* of the *n*th level. By (ii) of Lemma 2.3, there is a vertex *v* in the tree \mathcal{T} such that the *f*-orbit of *v* has length *m*. Of course, *v* must be a descendant of *u*. Now the choice of *p* allows us to consider a non-trivial *p*-element *g* in $\operatorname{rst}_G(v)$. Set $H = \langle g, f \rangle$. By Lemma 2.6, we have $H \cong \langle g \rangle \wr \langle f \rangle$. In particular, H is a finite p-group. On the other hand, by Lemma 3.2, H contains an element of order greater than m. This contradicts the choice of m, since $H \subseteq \operatorname{rst}_G(n)$.

Now we continue with our analysis of Engel elements in some wreath products. Before proceeding, we introduce some further notation. If G is a group and $S \subseteq G$, we write $L_G(S)$ to denote the set of all $x \in G$ that are left Engel elements on every element of S, that is, such that for all $s \in S$ there exists n = n(s, x) such that [s, n x] = 1. We define the set $\overline{L}_G(S)$ in the obvious way, and if $x \in \overline{L}_G(S)$ then the Engel degree of x on S is the maximum of the Engel degrees of x on the elements of S.

Lemma 3.4. Let $W = Y \wr X$ be a regular wreath product of two non-trivial groups, where X is finite cyclic of order n, and let $\pi : W \to X$ be the natural projection. Assume that $D = D_1 \times \cdots \times D_n \neq 1$ is a subgroup of the base group of W, and that $w \in W$ is such that $\pi(w)$ is a generator of X. Then the following hold:

- (i) If $w \in \overline{L}_W(D)$ has Engel degree d on D then $d \ge n$.
- (ii) If $w \in L_W(D)$ then $C_D(w)$ is periodic.

Proof. Write $w = (y_1, \ldots, y_n)x$, where $y_i \in Y$ and x generates X. We may assume that x permutes the components of the base group according to the cycle $(1 \ 2 \ \ldots \ n)$.

(i) Without loss of generality, we may assume that $D_1 \neq 1$. Choose a non-trivial element $g = (y, 1, ..., 1) \in D$ and let $1 \leq i \leq n-1$. One can easily check by induction on *i* that

$$[g_{,i}w] = (y^{(-1)^i}, \dots, y^{y_1\dots y_i}, 1, \dots, 1),$$

where the last non-trivial component is in position i+1. It follows that $[g_{n-1}w] \neq 1$ and $d \geq n$.

(ii) By contradiction, assume that $h = (z_1, \ldots, z_n) \in C_D(w)$ is of infinite order. For notational convenience, set $z_0 = z_n$ and $y_0 = y_n$. Then from the condition $h = h^w$ we get $z_i = z_{i-1}^{y_{i-1}}$ for all $i = 1, \ldots, n$. Hence all components of h are conjugate and they are all of infinite order.

Now let $g = (z_1, 1, ..., 1) \in D$. For every $k \ge 0$, let us write $[g_{k}, w] = (z_{k,1}, \ldots, z_{k,n})$ and, as before, set $z_{k,0} = z_{k,n}$. We claim that the following hold for every $k \ge 0$:

- (a) $z_{k,i} \in \langle z_i \rangle$ for every $i = 1, \ldots, n$.
- (b) If we write $z_{k,i} = z_i^{m_{k,i}}$, then there exists $i \in \{1, \ldots, n\}$ such that $m_{k,i} \neq m_{k,i-1}$.

We argue by induction on k. The result is obvious for k = 0, so assume $k \ge 1$ and that the claim is true for values less than k. Since $[g_{,k} w] = [g_{,k-1} w]^{-1} [g_{,k-1} w]^w$, it follows that

$$z_{k,i} = z_{k-1,i}^{-1} z_{k-1,i-1}^{y_{i-1}} = z_i^{-m_{k-1,i}} (z_{i-1}^{y_{i-1}})^{m_{k-1,i-1}} = z_i^{m_{k-1,i-1}-m_{k-1,i}}$$

for all i = 1, ..., n. This proves (a) and, if (b) does not hold, then

$$m_{k-1,1} - m_{k-1,2} = m_{k-1,2} - m_{k-1,3} = \dots = m_{k-1,n-1} - m_{k-1,n} = m_{k-1,n} - m_{k-1,1}$$

Now the sum of the n-1 first terms in this chain of equalities is the same as n-1 times the last one, i.e.

$$m_{k-1,1} - m_{k-1,n} = (n-1)(m_{k-1,n} - m_{k-1,1}).$$

From this, it readily follows that

$$m_{k-1,1} = m_{k-1,2} = m_{k-1,3} = \dots = m_{k-1,n},$$

which is contrary to the induction hypothesis.

Finally, observe that (b) above implies that $m_{k,i}$ and $m_{k,i-1}$ cannot both be zero. Since z_i and z_{i-1} are of infinite order, we conclude that $[g_{,k} w] \neq 1$ for all $k \geq 1$ and consequently $w \notin L_W(D)$. This contradiction completes the proof.

4. Left Engel elements in weakly branch groups

At this point, we can start combining all the machinery developed in Sections 2 and 3 in order to prove the main results of this paper. In this section we consider left Engel elements. The following is an expanded version of Theorem A.

Theorem 4.1. Let G be a subgroup of $\operatorname{Aut} \mathcal{T}$ in which all rigid vertex stabilizers are non-trivial. Then:

(i) If f is a non-trivial left Engel element of finite order, and O is a non-trivial totally splitting f-orbit, then for some prime number p the length of O is a p-power and rst_G(O) is a p-subgroup.

If G is furthermore weakly branch, then:

- (ii) If the set of finite order elements of L(G) is non-trivial then it is a p-set for some prime p, and $rst_G(n)$ is a p-group for some $n \ge 1$.
- (iii) $\overline{\mathrm{L}}(G) = 1$.

Proof. (i) Denote the reduced tree $\mathcal{R}(\mathcal{O})$ by \mathcal{R} , and set $x = \Phi_{\mathcal{O}}(f)$ and $H = G_{\mathcal{O}}$. We observe that $|x| = |\mathcal{O}|$ by Remark 2.8. Consider now a vertex v in \mathcal{O} and an arbitrary element $g \in \operatorname{rst}_G(v)$, and set $y = \psi_v(g)$ (here v is considered as a vertex in \mathcal{T}). Then $h = \Phi_{\mathcal{O}}(g)$ lies in $\operatorname{rst}_H(v)$ and $\psi_v(h) = y$ (here v is considered as a vertex in \mathcal{R}). By Lemma 2.6, we have $\langle h, x \rangle \cong \langle h \rangle \wr \langle x \rangle$. Since $x \in L(H)$, Lemma 3.1 implies that both |y| and |x| are p-powers for some prime p. Thus |g| and $|\mathcal{O}|$ are p-powers. Since $g \in \operatorname{rst}_G(v)$ was arbitrary and f acts transitively on \mathcal{O} , we conclude that $\operatorname{rst}_G(\mathcal{O})$ is a p-group.

(ii) Let again $f \in L(G)$ be a non-trivial element of finite order. By applying Lemma 2.9 to f, we obtain non-trivial totally splitting f-orbits $\mathcal{O}_1, \ldots, \mathcal{O}_k$, all lying on the same level n of the tree, such that $|f| = \operatorname{lcm}(|\mathcal{O}_1|, \ldots, |\mathcal{O}_k|)$. Let us fix $i \in \{1, \ldots, k\}$. By (i), there exists a prime p (in principle, depending on i) such that $|\mathcal{O}_i|$ is a p-power and $\operatorname{rst}_G(\mathcal{O}_i)$ is a p-group. Since G acts now level transitively on \mathcal{T} , all rigid vertex stabilizers are isomorphic by (2.2). It follows that p is the same for all i and consequently $\operatorname{rst}_G(n)$ is a p-group. Also the length of all orbits $\mathcal{O}_1, \ldots, \mathcal{O}_k$ is a power of p and f is a p-element.

(iii) By contradiction, assume that $f \in \overline{\mathcal{L}}(G)$, $f \neq 1$. Let d be the Engel degree of f.

Assume first that f is of finite order. Let \mathcal{O} be a non-trivial totally splitting f-orbit. Define x and y as in the proof of (i), and recall that these are p-elements. By Lemma 3.1,

$$d \ge |x| + \frac{1}{p}(\log_p |y| - 1)(p - 1)|x|.$$

On the other hand, since the exponent of $rst_G(n)$ is not finite by Proposition 3.3, the order of y is unbounded. This is a contradiction.

Assume now that the order of f is infinite. By Lemma 2.3, there exists an f-orbit \mathcal{O} of length $\ell > d$. Let once again \mathcal{R} be the reduced tree $\mathcal{R}(\mathcal{O})$, and set $h = \Phi_{\mathcal{O}}(f)$ and $H = G_{\mathcal{O}}$. Then $h \in S\langle x \rangle$, where S is the first level stabilizer in Aut \mathcal{R} (i.e. the stabilizer of \mathcal{O}) and x is a rooted automorphism corresponding to a cycle of length ℓ . Observe that $S\langle x \rangle$ is isomorphic to a regular wreath product $W = Y \wr X$, where Y is the stabilizer in \mathcal{R} of a vertex in \mathcal{O} and $X = \langle x \rangle$ is cyclic of order ℓ . Under this isomorphism, h corresponds to an element w with $\pi(w) = x$. Also h lies in $\overline{L}_H(D)$ with Engel degree at most d, where $D = \Phi_{\mathcal{O}}(\operatorname{rst}_G(\mathcal{O}))$ corresponds to a non-trivial direct product inside the base group of W. Now, by applying (i) of Lemma 3.4, we get $d \geq \ell$, which is a contradiction. This completes the proof of (iii).

Now we proceed to prove Theorem B.

Theorem 4.2. Let G be a branch group. If $L(G) \neq 1$ then G is periodic and there exists a prime p such that:

- (i) L(G) consists of p-elements.
- (ii) G is virtually a p-group.

Proof. It suffices to show that L(G) does not contain any elements of infinite order. Indeed, since $L(G) \neq 1$, the theorem then follows immediately from (ii) of Theorem 4.1, by taking into account that $|G : \operatorname{rst}_G(n)|$ is always finite if G is a branch group.

Let us assume then that $f \in L(G)$ is of infinite order. Consider an f-orbit \mathcal{O} in $V(\mathcal{T})$ of length $\ell \geq 2$, and let n be the level of \mathcal{T} containing \mathcal{O} . Set $\mathcal{R} = \mathcal{R}(\mathcal{O}), h = \Phi_{\mathcal{O}}(f)$ and $H = G_{\mathcal{O}}$. Then for every vertex $v \neq \emptyset$ of \mathcal{R} we have $\Phi_{\mathcal{O}}(\operatorname{rst}_G(v)) \subseteq \operatorname{rst}_H(v)$, and consequently all rigid vertex stabilizers of H are non-trivial. Also $h \in L(H)$.

If h has finite order, then by (i) of Theorem 4.1, the rigid stabilizer in H of some vertex $v \neq \emptyset$ of \mathcal{R} is periodic. Consequently $\operatorname{rst}_G(v)$ is periodic, and by level transitivity of G, also $\operatorname{rst}_G(n)$ is periodic. Since $|G : \operatorname{rst}_G(n)|$ is finite, it follows that G itself is periodic, which is a contradiction.

Assume now that the order of h is infinite. As in the proof of (iii) of Theorem 4.1, h lies in $S\langle x \rangle$, where S is the first level stabilizer of Aut \mathcal{R} , and x is a rooted automorphism corresponding to a cycle of length ℓ . We can identify $S\langle x \rangle$ with the regular wreath product $W = Y \wr X$, where $X = \langle x \rangle$ is cyclic of order ℓ and h maps onto x. Then $h \in L_W(D)$, where

$$D = \Phi_{\mathcal{O}}(\operatorname{rst}_G(\mathcal{O})) = \Phi_{\mathcal{O}}(\operatorname{rst}_G(n))$$

corresponds to a non-trivial direct product inside the base group of W. By (ii) of Lemma 3.4, $C_D(h)$ is periodic. However, since G is branch we have $f^k \in \operatorname{rst}_G(n)$ for some $k \geq 1$ and then $h^k = \Phi_O(f^k) \in D$. It follows that $h^k \in C_D(h)$ is an element of infinite order, which is a contradiction. Now we can apply Theorems A and B to some distinguished subgroups of Aut \mathcal{T} , and obtain the part of Corollary D regarding left Engel elements. Before proceeding, we will introduce some of the groups that appear in the following result.

First of all, the Hanoi Tower group \mathcal{H} is the subgroup of Aut \mathcal{T}_3 generated by the three automorphisms a, b and c given by the following recursive formulas:

$$a = (1, 1, a)(1 \ 2),$$

 $b = (1, b, 1)(1 \ 3),$
 $c = (c, 1, 1)(2 \ 3).$

This group models the popular Hanoi Tower puzzle on 3 pegs.

On the other hand, given an odd prime p and a non-trivial subspace \mathbf{E} of \mathbb{F}_p^{p-1} , we define the multi-GGS group (GGS standing for Grigorchuk, Gupta, and Sidki) $G_{\mathbf{E}}$ as the following subgroup of Aut \mathcal{T}_p . The group $G_{\mathbf{E}}$ is generated by the rooted automorphism a of order p corresponding to the p-cycle $(1 \ 2 \ \ldots \ p)$, and by the elementary abelian p-subgroup B consisting of all automorphisms $b_{\mathbf{e}}$, with $\mathbf{e} = (e_1, \ldots, e_{p-1}) \in \mathbf{E}$, defined recursively via

(4.1)
$$b_{\mathbf{e}} = (a^{e_1}, \dots, a^{e_{p-1}}, b_{\mathbf{e}}).$$

If dim $\mathbf{E} = 1$ then $G_{\mathbf{E}}$ is simply called a GGS group. Multi-GGS groups are usually presented by giving a basis $(\mathbf{e}_1, \ldots, \mathbf{e}_r)$ of \mathbf{E} and defining $b_i \in B$ from \mathbf{e}_i as in (4.1) for each $i = 1, \ldots, r$, so that $G_{\mathbf{E}} = \langle a, b_1, \ldots, b_r \rangle$. We refer the reader to the paper [2] by Alexoudas, Klopsch, and Thillaisundaram for general facts about multi-GGS groups. Multi-GGS groups are infinite and provide a wealth of examples giving a negative answer to the General Burnside Problem. For instance, the famous Gupta-Sidki *p*-group is the GGS group $\langle a, b \rangle$ with *b* corresponding to the vector $\mathbf{e} = (1, -1, 0, \ldots, 0)$. In general, a multi-GGS group is periodic if and only if \mathbf{E} is contained in the hyperplane of \mathbb{F}_p^{p-1} given by the equation $e_1 + \cdots + e_{p-1} = 0$ [2, Theorem 3.2]. On the other hand, multi-GGS groups are known to be branch unless $\mathbf{E} = \langle (1, \ldots, 1) \rangle$ consists of constant vectors, in which case it is weakly branch [2, Proposition 3.7].

Corollary 4.3. In all the following groups, the only left Engel element is the identity:

- (i) Every infinitely iterated wreath product of finite transitive permutation groups of degree at least 2. In particular, Aut \mathcal{T} and Γ_p , for p a prime.
- (ii) The group \$\mathcal{F}\$ of all finitary automorphisms of \$\mathcal{T}\$, provided that the sequence \$\vec{d}\$ defining \$\mathcal{T}\$ contains infinitely many terms greater than 2.
- (iii) All non-periodic multi-GGS groups G_E, i.e. those with at least one vector e ∈ E having non-zero sum in F_p.
- (iv) The Hanoi Tower group \mathcal{H} .

Proof. (i) For every $n \in \mathbb{N}$, let K_n be a finite transitive permutation group of degree $d_n \geq 2$, and let W be the iterated wreath product of all these groups. Let \mathcal{T} be the spherically homogeneous rooted tree corresponding to the sequence $\overline{d} = \{d_n\}_{n \in \mathbb{N}}$. Then W is isomorphic to the subgroup K of Aut \mathcal{T} consisting of all automorphisms whose labels at level n are elements of K_{n+1} . Observe that K

is a branch group, since every K_n is transitive and obviously $rst_K(n) = st_K(n)$ in this case.

According to Theorem B, we only need to construct an element of infinite order in K to conclude that L(W) = 1. To this purpose, we choose an infinite sequence $\{k_n\}_{n \in \mathbb{N}}$ of non-trivial permutations $k_n \in K_n$, and an infinite sequence $\{v_n\}_{n \in \mathbb{N} \cup \{0\}}$ of vertices, where $v_n \in \mathcal{L}_n$ and $k_n(v_n) \neq v_n$. Also, let \mathcal{O}_n denote the orbit of v_n under $\langle k_n \rangle$ and set $\ell_n = |\mathcal{O}_n|$.

Now we define f to be the automorphism of \mathcal{T} having label k_{n+1} at vertex v_n for all $n \in \mathbb{N} \cup \{0\}$. We claim that the length of the f-orbit of v_n is $\ell_1 \ldots \ell_n$ for all $n \in \mathbb{N}$. Since $\ell_i \geq 2$ for every i, we conclude that f is of infinite order by using (ii) of Lemma 2.3.

We prove the claim by induction on n. The result is obvious for n = 1, since f behaves as k_1 on the first level of \mathcal{T} . Then f^{ℓ_1} fixes all vertices in the orbit \mathcal{O}_1 , and a simple calculation shows that on all those vertices the section of f^{ℓ_1} coincides with the section of f at v_1 , let us call it g. Since v_n lies at level n - 1 for g, by induction the length of the g-orbit of v_n is $\ell_2 \ldots \ell_n$. From this one can readily see that the f-orbit of v_1 has length $\ell_1 \ldots \ell_n$, as desired.

(ii) Obviously, \mathcal{F} is spherically transitive and $\operatorname{rst}_{\mathcal{F}}(n) = \operatorname{st}_{\mathcal{F}}(n)$ for all $n \in \mathbb{N}$. Thus \mathcal{F} is a branch group. In this case, all elements of \mathcal{F} are of finite order, but we still get $L(\mathcal{F}) = 1$ from Theorem B, because there is no prime p for which \mathcal{F} is virtually a p-group. Indeed, assume for a contradiction that N is a normal psubgroup of \mathcal{F} of finite index m. Under this assumption, if H is a q-subgroup of \mathcal{F} for a prime $q \neq p$, the order of H cannot exceed m. However, as we see in the next paragraph, the condition on the sequence \overline{d} implies that \mathcal{F} has 2-subgroups and 3-subgroups of arbitrarily high order, and we get a contradiction.

Consider the following subset of \mathbb{N} :

$$S = \{ n \in \mathbb{N} \mid d_n \ge 3 \}.$$

By hypothesis, S is infinite. For every $n \in S$, let H_n be the subgroup of \mathcal{F} consisting of all automorphisms with labels lying in $\langle (1 \ 2) \rangle$ for all vertices in \mathcal{L}_n and trivial labels elsewhere. Then the order of H_n is $2^{d_1...d_n}$, which tends to infinity as $n \to \infty$. We can define similarly a subgroup J_n of order $3^{d_1...d_n}$ for every $n \in S$, by using the 3-cycle (1 2 3). Thus we get 2-subgroups and 3-subgroups of \mathcal{F} of arbitrarily high order, as desired.

(iii) If $\mathbf{E} = \langle (1, \ldots, 1) \rangle$ then $L(G_{\mathbf{E}}) = 1$ by [7, Theorem 7]. Otherwise $G_{\mathbf{E}}$ is a branch group, and the result follows immediately from Theorem B and from the characterisation of periodic multi-GGS groups given above.

(iv) The Hanoi Tower group is known to be a branch group [11, Theorem 5.1]. Let us see that the element $ab = (b, 1, a)(1 \ 2 \ 3)$ is of infinite order. Assume, for a contradiction, that |ab| = k is finite. Observe that $k = 3\ell$ for some ℓ , since ab has order 3 modulo the first level stabilizer. But then

$$(ab)^{3\ell} = ((ba)^{\ell}, (ab)^{\ell}, (ab)^{\ell})$$

implies that $(ab)^{\ell} = 1$, which is a contradiction.

Corollary 4.4. Let p be a prime and let \mathcal{F}_p be the group of p-finitary automorphisms of Aut \mathcal{T}_p . Then the following hold:

(i) $L(\mathcal{F}_p) = \mathcal{F}_p$.

(ii)
$$\overline{\mathrm{L}}(\mathcal{F}_p) = 1$$
.

Proof. Since \mathcal{F}_p is locally a finite *p*-group, (i) is clear. On the other hand, since \mathcal{F}_p is spherically transitive and $\operatorname{rst}_{\mathcal{F}_p}(n) = \operatorname{st}_{\mathcal{F}_p}(n)$ for all n, (ii) follows directly from Theorem A.

5. RIGHT ENGEL ELEMENTS IN WEAKLY BRANCH GROUPS

In this final section, we prove Theorem C, regarding right Engel elements in weakly branch groups, and then we apply it to show that R(G) = 1 whenever G is a GGS group. Before proceeding, we need a straightforward lemma.

Lemma 5.1. Suppose that \mathcal{T} has d vertices in the first level, and consider $x, y \in \operatorname{Aut} \mathcal{T}$ such that:

- (i) y = az, where a is the rooted automorphism corresponding to the cycle $(1 \ 2 \ \dots \ d)$ and $z \in st(1)$ is given by $\psi(z) = (z_1, \dots, z_d)$.
- (ii) $x \in \operatorname{st}(1)$ is given by $\psi(x) = (x_1, \ldots, x_d)$.

Then, for all $k \geq 2$, we have

$$\psi([y_{k}x]) = ([(x_{d}^{-1})^{z_{1}}, k-1, x_{1}]^{x_{1}}, \dots, [(x_{d-1}^{-1})^{z_{d}}, k-1, x_{d}]^{x_{d}}).$$

Proof. We have

$$\psi([y,x]) = \psi((x^{-1})^y x) = \psi((x^{-1})^a)^{\psi(z)} \psi(x)$$

= $((x_d^{-1})^{z_1} x_1, (x_1^{-1})^{z_2} x_2, \dots, (x_{d-1}^{-1})^{z_d} x_d).$

Now the result follows immediately by observing that taking subsequent commutators with x is performed componentwise.

Now we are ready to prove Theorem C.

Theorem 5.2. Let G be a weakly branch group. If $rst_G(n)$ is not an Engel group for all $n \in \mathbb{N}$, then R(G) = 1.

Proof. Let $f \in G$, $f \neq 1$, and assume by way of contradiction that $f \in \mathcal{R}(G)$. Choose a non-trivial f-orbit $\mathcal{O} = \{v_1, \ldots, v_d\}$, and assume that f permutes cyclically the vertices v_i . Let $\mathcal{R} = \mathcal{R}_{\mathcal{O}}$, $H = G_{\mathcal{O}}$ and $y = \Phi_{\mathcal{O}}(f) \in \mathcal{R}(H)$. Then we can write y = az, where a is rooted in \mathcal{R} corresponding to the cycle $(1 \ 2 \ \ldots \ d)$ and z is in the first level stabilizer. Write $\psi(z) = (z_1, \ldots, z_d)$.

Let n be the level of \mathcal{T} where \mathcal{O} lies. Since $\Phi_{\mathcal{O}}(\operatorname{rst}_G(n)) \subseteq \operatorname{rst}_H(1)$ and $\operatorname{rst}_G(n)$ is not Engel by hypothesis, it follows that $\operatorname{rst}_H(1)$ is not an Engel group. If L is the first component of the direct product $\psi(\operatorname{rst}_H(1))$ then L is not an Engel group either, and we can choose $a, b \in L$ such that $[b, k a] \neq 1$ for all $k \geq 1$. Now consider $r_1, r_2 \in \operatorname{rst}_H(1)$ such that

 $\psi(r_1)=(a,1,\ldots,1) \qquad \text{and} \qquad \psi(r_2)=(b,1,\ldots,1),$ and define $x=r_1(r_2^{-1})^{y^{-1}},$ so that

$$\psi(x) = (a, 1, \dots, 1, (b^{-1})^{z_1^{-1}}).$$

By applying the formula in Lemma 5.1, we get

$$\psi([y_{k} x]) = ([b_{k-1} a]^{a}, *, \dots, *)$$

and consequently $[y_k x] \neq 1$ for all $k \geq 2$. This is a contradiction, since $y \in \mathbf{R}(H)$ and $x \in H$.

Theorem C can be applied to show that GGS groups have no non-trivial right Engel elements. We first need to prove the weaker result that they are not Engel groups.

Lemma 5.3. Let G be a GGS group. Then G is not an Engel group.

Proof. We show that there is a power of b that is not a left Engel element of G. Let **e** be the defining vector of b. Consider any index $i \in \{1, \ldots, p-1\}$ such that $e_{p-i} \neq 0$ in \mathbb{F}_p , and choose $\lambda \in \mathbb{F}_p^{\times}$ such that $\lambda e_{p-i} = -i$. Then we have

(5.1)
$$\psi((b^{-\lambda})^{a^i}) = (*, \dots, *, a^i),$$

where we use * to denote unspecified elements of G.

Since $(b^{-\lambda})^{a^i} = [a^i, b^{\lambda}]b^{-\lambda}$, it follows that, for every $k \ge 2$,

$$\begin{split} [(b^{-\lambda})^{a^{i}}, b^{\lambda}, \overset{k-1}{\dots}, b^{\lambda}] &= [[a^{i}, b^{\lambda}]b^{-\lambda}, b^{\lambda}, \overset{k-1}{\dots}, b^{\lambda}] \\ &= [[a^{i}, b^{\lambda}, b^{\lambda}]^{b^{-\lambda}}, b^{\lambda}, \overset{k-2}{\dots}, b^{\lambda}] \\ &= [a^{i}, b^{\lambda}, \overset{k}{\dots}, b^{\lambda}]^{b^{-\lambda}}. \end{split}$$

By using (5.1), it follows that

(5.2) $\psi([a^{i}, b^{\lambda}, ..^{k}, .., b^{\lambda}]^{b^{-\lambda}}) = \psi([(b^{-\lambda})^{a^{i}}, b^{\lambda}, \overset{k-1}{\dots}, b^{\lambda}]) = (*, \dots, *, [a^{i}, b^{\lambda}, \overset{k-1}{\dots}, b^{\lambda}]).$

Now if b^{λ} is a left Engel element of G, choose the minimum $k \geq 1$ such that $[a^i, b^{\lambda}, .k^{\cdot}, b^{\lambda}] = 1$. Since a^i and b^{λ} do not commute, we have $k \geq 2$ and so $[a^i, b^{\lambda}, \frac{k-1}{2}, b^{\lambda}] \neq 1$. According to (5.2), this is a contradiction.

Corollary 5.4. Let G be a GGS group. Then R(G) = 1.

Proof. If the defining vector \mathbf{e} is constant, then L(G) = 1 by Theorem 7 of [7], and consequently also R(G) = 1. Thus in the remainder we assume that \mathbf{e} is not constant. By Lemmas 3.2 and 3.4 of [8], we know that G is regular branch over K, where $K = \gamma_3(G)$ if \mathbf{e} is symmetric and K = G' otherwise. Since $\operatorname{rst}_G(n)$ contains a copy of $K \times \stackrel{p^n}{\cdots} \times K$ for every $n \in \mathbb{N}$, if we prove that K is not Engel then Theorem C applies to conclude that R(G) = 1.

In order to show that K is not Engel, we are going to find a vertex v of the first level of the tree such that $\psi_v(K) = G$. Since G is not Engel by Lemma 5.3, it follows that K is not Engel either, as desired.

We consider separately the cases when \mathbf{e} is symmetric and non-symmetric. Assume first that \mathbf{e} is non-symmetric, so that K = G'. We have

$$\psi([b,a]) = (a^{-e_1}b, a^{e_1-e_2}, a^{e_2-e_3}, \dots, a^{e_{p-2}-e_{p-1}}, b^{-1}a^{e_{p-1}}).$$

Since **e** is not constant, there exists $i \in \{1, \ldots, p-2\}$ such that $e_i \neq e_{i+1}$ in \mathbb{F}_p . If v is the vertex i + 1 on the first level of the tree, then

$$\psi_v([b,a]) = a^{e_i - e_{i+1}}$$
 and $\psi_v([b,a]^{a^i}) = a^{-e_1}b.$

Since the subgroup $\langle a^{e_i-e_{i+1}}, a^{-e_1}b \rangle$ coincides with G, we get the desired equality $\psi_v(G') = G$.

Now let **e** be symmetric, i.e. such that $e_i = e_{p-i}$ for all $i = 1, \ldots, p-1$. Since **e** is not constant, this implies that $p \ge 5$. We have

$$\psi([b, a, a]) = \psi([b, a]^{-1})\psi([b, a]^{a})$$

= $(b^{-1}a^{e_{1}}b^{-1}a^{e_{p-1}}, a^{-2e_{1}+e_{2}}b, a^{e_{1}-2e_{2}+e_{3}}, \dots,$
 $a^{e_{p-3}-2e_{p-2}+e_{p-1}}, a^{-e_{p-1}}ba^{e_{p-2}-e_{p-1}}).$

If $e_i - 2e_{i+1} + e_{i+2} \neq 0$ for some $i \in \{1, \ldots, p-3\}$, we have a non-trivial power of a in one of the components of $\psi([b, a, a])$ and we can argue as above to prove that $\psi_v(\gamma_3(G)) = G$ for a vertex v in the first level. On the other hand, if $e_i - 2e_{i+1} + e_{i+2} = 0$ for all $i = 1, \ldots, p-3$, then

$$e_{3} = 2e_{2} - e_{1},$$

$$e_{4} = 2e_{3} - e_{2} = 3e_{2} - 2e_{1},$$

$$\vdots$$

$$e_{p-1} = 2e_{p-2} - e_{p-3} = (p-2)e_{2} - (p-3)e_{1}$$

Since $e_{p-1} = e_1$, the last equation implies that $e_1 = e_2$, and then using all other equations, we get that all components e_i are equal to e_1 . Thus the vector **e** is constant, which is a contradiction.

Acknowledgements. The authors want to thank A. Tortora and G. Traustason for helpful discussions.

References

- M. Abért, Group laws and free subgroups in topological groups, Bulletin of the London Mathematical Society 37 (2005), no. 4, 525–534.
- [2] T. Alexoudas, B. Klopsch, and A. Thillaisundaram, Maximal subgroups of multi-edge spinal groups, Groups, Geometry, and Dynamics 10 (2016), 619–648.
- [3] R. Baer, Engelsche elemente noetherscher gruppen, Mathematische Annalen 133 (1957), 256-270.
- [4] L. Bartholdi, Algorithmic decidability of Engel's property for automaton groups, Computer Science – Theory and Applications, Lecture Notes in Computer Science, Volume 9691, Springer, 2016, pp. 29–40.
- [5] L. Bartholdi, R.I. Grigorchuk, and Z. Sunik, *Branch groups*, Handbook of Algebra, Volume 3, North-Holland, 2003, pp. 989–1112.
- [6] V.V. Bludov, An example of not Engel group generated by Engel elements, A Conference in Honor of Adalbert Bovdi's 70th Birthday (2006).
- [7] G.A. Fernández-Alcober, A. Garreta, and M. Noce, Engel elements in some fractal groups, Monatshefte für Mathematik 189 (2019), 651–660.
- [8] G.A. Fernández-Alcober and A. Zugadi-Reizabal, GGS-groups: order of congruence quotients and Hausdorff dimension, Transactions of the American Mathematical Society 366 (2013), 1993–2017.
- R.I. Grigorchuk, On Burnside's problem on periodic groups, Funktsionalyi Analiz i ego Prilozheniya 14 (1980), 53–54.
- [10] _____, Just infinite branch groups, New Horizons in pro-p Groups, Progress in Mathematics, Volume 184, Birkhäuser, 2000, pp. 121–179.
- [11] R.I. Grigorchuk and Z. Sunic, Self-similarity and branching in group theory, Groups St Andrews 2005, Volume 1, Cambridge University Press, 2007, pp. 36–95.
- [12] R.I. Grigorchuk and J.S. Wilson, The uniqueness of the actions of certain branch groups on rooted trees, Geometriae Dedicata 100 (2003), 103–116.
- [13] N. Gupta and S. Sidki, On the Burnside problem for periodic groups, Mathematische Zeitschrift 182 (1983), 385–388.

- [14] Yu.G. Leonov, On identities of groups of automorphisms of trees, Visnyk of Kyiv State University of T.G. Shevchenko 3 (1997), 37–44.
- [15] H. Liebeck, Concerning nilpotent wreath products, Mathematical Proceedings of the Cambridge Philosophical Society 58 (1962), 443–451.
- [16] M. Noce and A. Tortora, A note on Engel elements in the first Grigorchuk group, International Journal of Group Theory 8 (2019), 9–14.
- [17] D. Robinson, A Course in the Theory of Groups, Graduate Texts in Mathematics, Springer, 1996.
- [18] J.S. Wilson, Groups with every proper quotient finite, Mathematical Proceedings of the Cambridge Philosophical Society 69 (1971), 373–391.

Department of Mathematics, University of the Basque Country UPV/EHU, Bilbao, Spain

E-mail address: gustavo.fernandez@ehu.eus

DIPARTIMENTO DI MATEMATICA, UNIVERSITÀ DI SALERNO, ITALY; DEPARTMENT OF MATH-EMATICS, UNIVERSITY OF THE BASQUE COUNTRY UPV/EHU, BILBAO, SPAIN *E-mail address:* mnoce@unisa.it

DEPARTMENT OF MATHEMATICAL SCIENCES, UNIVERSITY OF BATH, BATH BA2 7AL, UNITED KINGDOM

E-mail address: G.M.Tracey@bath.ac.uk