

Understanding potential mechanisms of harm

Jasper, Alice; Sapey, Elizabeth; Thickett, David; Scott, Aaron

DOI:

[10.1152/AJPLUNG.00081.2021](https://doi.org/10.1152/AJPLUNG.00081.2021)

License:

Creative Commons: Attribution (CC BY)

Document Version

Publisher's PDF, also known as Version of record

Citation for published version (Harvard):

Jasper, A, Sapey, E, Thickett, D & Scott, A 2021, 'Understanding potential mechanisms of harm: the drivers of electronic cigarette-induced changes in alveolar macrophages, neutrophils and lung epithelial cells', *American journal of physiology. Lung cellular and molecular physiology*, vol. 321, no. 2, pp. L336-L348.
<https://doi.org/10.1152/AJPLUNG.00081.2021>

[Link to publication on Research at Birmingham portal](#)

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

REVIEW

Electronic Cigarettes: Not All Good News?

Understanding potential mechanisms of harm: the drivers of electronic cigarette-induced changes in alveolar macrophages, neutrophils, and lung epithelial cells

Alice E. Jasper, Elizabeth Sapey, David R. Thickett, and Aaron Scott

Birmingham Acute Care Research, Institute of Inflammation and Ageing, University of Birmingham, Birmingham, United Kingdom

Abstract

Electronic (e-) cigarettes are growing in popularity despite uncertainties regarding their long-term health implications. The link between cigarette smoking and initiation of chronic lung disease took decades to unpick so in vitro studies mimicking e-cigarette exposure aim to detect early indicators of harm. In response to e-cigarette exposure, alveolar macrophages adopt a proinflammatory phenotype of increased secretion of proinflammatory cytokines, reduction in phagocytosis, and efferocytosis and reactive oxygen species generation. These effects are largely driven by free radical exposure, changes in PI3K/Akt signaling pathways, nicotine-induced reduction in phagocytosis receptors, and impaired lipid homeostasis leading to a foam-like lipid-laden phenotype. Neutrophils exhibit disrupted chemotaxis and transmigration to chemokines, reduced phagocytosis and bacterial killing, and an increase in protease secretion without corresponding antiproteases in response to e-cigarette exposure. This is driven by an altered ability to respond and to polarize toward chemoattractants, an activation of the p38 MAPK signaling pathway and inability to assemble NADPH oxidase. E-cigarettes induce lung epithelial cells to display decreased ciliary beat frequency and ion channel conductance as well as changes in chemokine secretion and surface protein expression. Changes in gene expression, mitochondrial function, and signaling pathways have been demonstrated in lung epithelial cells to explain these changes. Many functional outputs of alveolar macrophages, neutrophils, and lung epithelial cells have not been fully explored in the context of e-cigarette exposure and the underlying driving mechanisms are poorly understood. This review discusses current evidence surrounding the effects of e-cigarettes on alveolar macrophages, neutrophils, and lung epithelial cells with particular focus on the cellular mechanisms of change.

alveolar macrophages; e-cigarettes; lung epithelial cells; mechanisms; neutrophils

INTRODUCTION

Electronic (e-) cigarettes, also known as electronic nicotine delivery systems (ENDS), are noncombustible nicotine delivery devices, which were developed in 2003 as an alternative to tobacco cigarettes. Despite origins as a smoking cessation tool, only 41% of e-cigarette users have quitting or harm reducing intentions (1). Furthermore, 80% of those who switch from cigarettes to e-cigarettes continue to use the device 1 yr after switching (2). Therefore, it is important that e-cigarettes are not considered only as short-term therapy. E-cigarettes serve to heat e-cigarette liquids (e-liquids), delivering a vapor to the lungs at subcombustion temperatures. E-liquids usually contain nicotine, a variety of flavoring compounds with over 7,700 currently described (3), humectants, propylene glycol (PG), and vegetable glycerin (VG). Production of sweet and fruity e-liquid flavors and

exposure to advertising have increased the uptake of e-cigarette use by nonsmoker adolescents who are consequently 3.5 times more likely to initiate cigarette smoking, supporting the gateway to smoking hypothesis (4). The e-cigarette market is now worth an estimated \$22.6 billion, supported by over 3.6 million e-cigarette users in the United Kingdom (1, 5–8) and over 40 million e-cigarette users worldwide (9).

The emergence of e-cigarette- or vaping-associated lung injury (EVALI) in the United States, acute severe respiratory distress with an absence of infection in otherwise healthy adults, has highlighted the potential dangers of e-cigarette use. Clinical presentation of EVALI includes dyspnea, fever, and tachycardia with bilateral ground glass opacities on computed tomography (CT). Of 81 patients with EVALI, 89% reported having used tetrahydrocannabinol (THC) and 9% used cannabidiol (CBD) e-liquids in the previous 90 days before symptom onset, raising further concerns around e-



Correspondence: A. Scott (A.Scott@bham.ac.uk).
Submitted 19 February 2021 / Revised 6 May 2021 / Accepted 12 May 2021



liquid components (10). Vitamin E acetate has been strongly linked with EVALI cases, particularly those where THC use has been reported (11). Vitamin E acetate is a supplement used to thicken some e-liquids and is considered safe for ingestion and skin application (12). However, upon inhalation, vitamin E acetate is delivered directly to the surfactant lining of the epithelium, interfering with the chemical properties and contributing to respiratory dysfunction in EVALI (11, 13, 14). The discovery of the dangers of vitamin E acetate raises further concern about e-liquid ingredients and the potential toxicity caused by vaporization. Since August 2019, over 2,800 cases of EVALI have been confirmed by the Centers for Disease Control and Prevention (CDC) with 68 confirmed deaths in the United States and at least 2 in the United Kingdom (15, 16).

Smoking cessation is a priority to reduce ill-health. Cigarette smoking is associated with initiating and worsening pathology in almost every chronic inflammatory lung disease (17) as well as causing a number of cancers, cardiovascular diseases, and strokes. Most notably, cigarette smoke is a causative factor in chronic obstructive pulmonary disease (COPD) (18). COPD is characterized by postbronchodilator forced expiratory volume in 1s (FEV_1)/forced vital capacity (FVC) ratio of less than 0.70, symptoms including shortness of breath, chronic cough, sputum production, or wheezing and a history of smoking cigarettes or exposure to other noxious stimuli (19). COPD is associated with chronic lung inflammation resulting in progressively worsening and irreversible airflow obstruction driven by small airways destruction, emphysema, and chronic bronchitis (20). Risk of development of smoking-associated COPD is heightened by α -1 antitrypsin deficiency, the genetic condition resulting in the absence or dysfunction of the antiprotease α -1 antitrypsin that normally protects the lungs from destruction by proteinases such as neutrophil elastase (NE) (21). Neutrophils are a central contributor to the chronic inflammation associated with COPD, as their recruitment, accumulation, and degranulation drives tissue damage, and their dysregulation in stable disease contributes to increased inflammation and impaired bacterial clearance (22). Pathophysiology is further driven by alveolar macrophages adopting a proinflammatory state, whereby secretion of inflammatory chemokines, interleukin (IL)-8 and monocyte chemoattractant protein-1 (MCP-1) recruits further immune cells to the small airways (23). Hyperplasia of airway epithelial cells contributes to tissue remodeling, which drives progression of airflow obstruction due to the stiffening of the airways and lung parenchyma (24). However, the link between these pathological drivers of COPD and smoking took decades to uncover, in part due to the slowly progressive nature of cigarette-associated lung damage, with often a lag of over three decades between smoking initiation and the development of characteristic airflow obstruction (25). This picture is further complicated by individual variance such as smoking behaviors, occupational and environment exposures, and genetic factors. This “slow-burning” disease is complex to model in cell, tissue, or animal-based models, where a short duration of smoke exposure is used. The same challenges are faced by researchers attempting to understand the signals of harm of e-cigarettes, as a similar disease progression may need to be considered.

To determine whether and what the long-term harm of e-cigarettes might be, models to mimic the exposure conditions in vitro and in vivo have been developed by researchers yet lack of standardization in experimental methods has resulted in differing experimental methods (26). E-cigarette vapor extract systems rely on vapor constituents quickly dissolving into media with high degrees of dilution and loss of some insoluble elements. E-cigarette vapor extract is best for direct comparison with cigarette smoke extract, which has been used in cigarette studies. E-cigarette vapor condensate is generated by vaporizing and condensing e-liquids, which is then diluted with media and used to treat cells. This represents the physiological behavior of e-cigarette vapor, vaporizing in the e-cigarette and condensing in the airway, more accurately than e-cigarette vapor extract treatment (27). Both e-cigarette vapor extract and condensate exposure models account for chemical changes induced by vaporization, preferable compared with unvaporized liquid treatment. The models mentioned thus far do not simulate direct vapor exposure; however, in the air-liquid interface model, cells are cultured on membranes and e-cigarette vapor is drawn over the apical surface of the cells. Each exposure model has advantages and limitations, and various methods have been used to quantify level of exposure including measurements of particulate deposition by optical density and nicotine quantification by mass spectrometry. In addition to variations in exposure model is the further complication of variations in devices, device settings, and e-liquids. The first devices to be invented, first-generation or cig-a-like devices, were disposable, non-refillable, and looked like traditional cigarettes. Later came the more advanced pen-like second-generation devices, with refillable tank and rechargeable battery. Second-generation devices rapidly overtook first generation due to the cost-effectiveness of a refillable and rechargeable device. However, development of larger volume tanks and high-power batteries formed the basis of third-generation devices. Most recently, the fourth generation of e-cigarettes allows user modification of temperature, power output, and airflow settings. In addition, these devices have larger tanks, longer battery life, and more customizable features than ever. Studies have shown that higher battery outputs lead to increased carbonyl levels in vapor (28, 29). The ability to customize settings, along with the huge variety of e-liquids—from flavors to nicotine concentrations—allows the user to adapt their device to suit their preferred vaping habits. However, this poses its own challenges in assessing effects of vaping on health as the combination of device, settings, and e-liquid differ from user to user.

Given the primary cells implicated in chronic smoking-related lung disease are lung epithelial cells, alveolar macrophages, and neutrophils, this review will assess evidence of cellular dysfunction for these cell types in response to e-cigarette exposure, summarized in Fig. 1.

■ ALVEOLAR MACROPHAGES

Alveolar macrophages are specialized resident innate immune cells residing in the lung alveoli and proximal airways. Existing in alveolar spaces, alveolar macrophages constantly encounter and clear an array of inhaled toxins,




Figure 1. A summary of the effects of e-cigarette vapor exposure on lung homeostasis driven by alterations in important lung cells. In response to e-cigarette exposure, functional changes in a range of cell type and subsequent dysfunctional interactions lead to changes characteristic of oxidative stress: 1) bacterial colonization is driven by impaired phagocytosis and bacterial killing by alveolar macrophages and neutrophils, exacerbated by evidence of increased virulence by invading pathogens; 2) impaired lipid metabolism by alveolar macrophages and alveolar type II epithelial cells leads to damaged surfactant and the formation of lipid-laden macrophages that exhibit abnormal processing of both endogenous and exogenous lipids; 3) tissue damage to the interstitium and epithelial cell layer is caused by recruitment of neutrophils and their subsequent degranulation of proteases such as matrix metalloproteases (MMPs) and neutrophil elastase (NE) without a corresponding increase in antiproteases such as $\alpha 1$ antitrypsin, leading to degradation of the extracellular matrix; 4) increased cytokine release by alveolar macrophages and epithelial cells recruits proinflammatory neutrophils and monocytes to the airspaces, causing tissue destruction as they transigrate; 5) mucociliary disruption causes accumulation of mucus due to reduced ciliogenesis and ciliary beat frequency; 6) persistent inflammation driven by reduced efferocytosis by alveolar macrophages, subsequent accumulation of apoptotic cells and neutrophil NETosis. These substantial changes are driven by functional changes in neutrophils, alveolar macrophages, and epithelial cells as a result of e-cigarette exposure. As a result, the alveolar spaces become inflamed and changes, some irreversible, occur, which can lead to the development of a range of respiratory diseases including bronchiectasis, chronic obstructive pulmonary disease, and emphysema. IL-8, interleukin 8; NET, neutrophil extracellular trap.

microbes, and antigens, assessing necessity for inflammatory responses and maintaining tissue homeostasis (30). Disruptions in the function of alveolar macrophages can contribute to the pathogenesis of lung disease such as COPD and enhance susceptibility to infection such as pneumonia (31, 32). Cigarette smoke exposure causes a proinflammatory alveolar macrophage phenotype as well as aberrant autophagy, impaired phagocytosis, and efferocytosis-driving inflammation in the lungs and reducing the clearance of infections and cell debris (33–35).

Alveolar Macrophages and E-cigarettes

Table 1 details the experimental methods and downstream effects of e-cigarette exposure to primary alveolar macrophages or cell line-derived macrophages. In an e-cigarette vapor condensate model, significant cell toxicities have been shown, which are independent of, but worsened by, the presence of nicotine; yet in an e-cigarette vapor extract model there was no significant cytotoxicity (36, 37). This may be because e-cigarette vapor condensate captures more of the

Table 1. A summary of the effects of e-cigarette vapor exposure on alveolar macrophage effector functions

Effector Function	E-Cigarette Model	ECL Nicotine Concentration	Exposure Model	Exposure Duration	Cell Type	Results	Reference
Viability	Second generation (Kanger Ltd)	36 mg/mL and 0 mg/mL	ECVC (0.8% concentration)	24 h	Human alveolar macrophages	36 mg/mL nicotine: 40.87%, 0 mg/mL nicotine: 77.87% live cells vs. 92.5% live cells in UTC	(36)
	QHIT e-cigarette (Puff, Moncalleri, Italy) 3.7 V	8 mg/g and 0 mg/g	100% ECVE (40 × 60 mL puff, 3 s, 30 s break)	0, 3, 6 days	THP-1 cell line	No change in toxicity with or without nicotine	(37)
Reactive oxygen species	Second generation (Kanger Ltd)	36 mg/mL	ECVC (0.5% concentration)	4 h	Human alveolar macrophages	50-fold increase in ROS production vs. UTC	(36)
Phagocytosis/antimicrobial activity	Vape pen (Green smart living/Njoy/Xtreme Vaping)	24 mg/mL and 8 mg/mL	ECVC [0.2–0.26 mg/mL nicotine (1.2–1.6 mM)]	1 h	Human alveolar macrophages	24 mg/mL nicotine: 535%, 8 mg/mL nicotine: 395% increase in MRSA population size vs. UTC	(38)
	LAVABOX DNA 200 Box Mod (Volcano e-cigs)	0	ECVC (55% PG/45% VG) (1% concentration)	1 h	Human alveolar macrophages	Significant reduction in <i>S. aureus</i> phagocytosis vs. UTC	(39)
	Second generation (Kanger Ltd)	36 mg/mL and 0 mg/mL	ECVC (0.5% concentration)	6 h	Human alveolar macrophages	36 mg/mL nicotine: 30% reduction, 0 mg/mL nicotine: 50% reduction in <i>E. coli</i> phagocytosis vs. UTC	(36)
	QHIT e-cigarette (Puff, Moncalleri, Italy) 3.7 V	0	100% ECVE (40 × 60 mL puff, 3 s, 30 s break)	0, 3, 6 days	THP-1 cell line	Reduced phagocytosis of <i>M. tuberculosis</i> at all-time points	(37)
Efferocytosis	EVOD-2 3.7 V, 1.5 Ω	18 mg/mL and 0 mg/mL	100% ECVE (50 × 3 s puffs, 5 s break)	24 h	THP-1 cell line	18 mg/mL nicotine: Significant reduction in efferocytosis compared with controls 0 mg/mL nicotine: No effect on efferocytosis	(40)
Cytokine/protease release	Second generation (Kanger Ltd)	26 mg/mL	ECVC (0.5% concentration)	24 h	Human alveolar macrophages	Increased secretion of IL-6, TNF-α, IL-8, MCP-1, and MMP-9 vs. UTC	(36)
	EVOD-2 3.7 V, 1.5 Ω	18 mg/mL	ECVE (50 × 3 s puffs, 5 s break)	24 h	THP-1 cell line	No change in IL-8, IL-1β, MIP-1β, or MCP-1 secretion but a reduction production of IL-6, MIP-1α, and TNFα vs. UTC	(41)
	None	1.8% or 0	1% PG/VG (55:45 v/v) media	Overnight	THP-1 cell line	1.8% nicotine: Increased secretion and activity of MMP-2 and MMP-9 vs. UTC 0 nicotine: No change in MMP-2 or MMP-9 vs. UTC	(42)
	QHIT e-cigarette (Puff, Moncalleri, Italy) 3.7 V	8 mg/g and 0 mg/g	100% ECVE (40 × 60 mL puff, 3 s, 30 s break)	3 days	THP-1 cell line	IL-8 was the highest produced cytokine and IL-1β was highly produced	(37)

Studies investigating the effect of e-cigarettes on alveolar macrophage viability, reactive oxygen species generation, phagocytosis, efferocytosis, and cytokine release are included. The exposure design and cell type have been included for comparison between findings. Only experiments using the humectant bases with or without nicotine in the absence of any flavorings have been included to aid understanding of the fundamental effects of e-cigarette exposure on alveolar macrophages. *E. coli*, *Escherichia coli*; ECVC, e-cigarette vapor condensate; ECVE, e-cigarette vapor extract; IL, interleukin; MCP, monocyte chemoattractant protein; MIP, macrophage inflammatory protein; MMP, matrix metalloproteinase; MRSA, methicillin-resistant *Staphylococcus aureus*; *M. tuberculosis*, *Mycobacterium tuberculosis*; PG, propylene glycol; ROS, reactive oxygen species; *S. aureus*, *Staphylococcus aureus*; TNF, tumor necrosis factor; UTC, untreated control; VG, vegetable glycerin.

vapor constituents, as it does not rely upon dissolving. In a range of studies examining cytokine secretion from both THP-1 cell lines and bronchoalveolar lavage (BAL)-derived macrophages, proinflammatory cytokines, most notably the robust recruiter of neutrophils IL-8, are upregulated in response to e-cigarette vapor condensate, extract, and supplemented media treatments (36, 37, 42). However, contradictory findings have been reported for other cytokines, such as one study showing an upregulation of tumor necrosis factor (TNF)- α after alveolar macrophage exposure to 50% PG: 50% VG condensate whereas another study showed a downregulation after 70% PG: 30% VG extract exposure in THP-1 cells (36, 41). This form of comparison makes it difficult to deduce whether varying results are due to different exposure models, cell types, or base humectants.

Conversely, effects on antimicrobial activity appear to be independent of nicotine content. Alveolar macrophages exposed to e-cigarettes exhibited reduced antimicrobial activity to gram-positive and gram-negative bacteria regardless of nicotine content and this is exacerbated by the increased virulence and growth rates of infecting pathogens (36–39). E-liquids have been shown to upregulate Ca^{2+} signaling in THP-1 cells and have been implicated in many cellular functions, including phagocytosis, so may represent a driving mechanism of impaired phagocytosis (43, 44). Furthermore, alveolar macrophages exposed to nicotine-containing e-cigarette vapor extract exhibited significant reduction in efferocytosis of apoptotic epithelial cells (40). Inability to clear cellular debris and dying cells causes persistent inflammation of the airway.

In vitro models and even cohort studies carried out to date model the short-term effects of vaping. Murine studies have been used to further examine the chronic effects. Using a chronic vapor exposure model, Maddison et al. (45) recapitulated the lipid-laden phenotype of alveolar macrophages seen in EVALI (46).

Mechanisms

Despite several decades of research, the mechanism(s) of action of cigarette smoke-mediated harm related to macrophage function are still not completely understood. Among the 5,000 different chemicals present in cigarette smoke (47, 48), the presence of many of the same harmful constituents that drive respiratory disease mechanisms have been found in e-cigarette vapor including carbonyl compounds (such as acrolein and formaldehyde), volatile organic compounds, and tobacco-specific nitrosamines (49). Recent exploratory evidence examining the effects of e-cigarettes on alveolar macrophages can broadly define potential mechanisms as 1) nicotine independent and driven by pathways activated by free radicals, aldehydes, and the PI3K pathway; 2) nicotine dependent and driven by cellular changes as a result of nicotine binding cell surface receptors; and 3) lipid metabolism driven by the altered ability to process lipids.

Nicotine independent.

Although most vapers are ex-smokers, an emerging cohort of previously never-smoker vapers appears to have much greater prevalence of vaping without nicotine. Although flavorings differ, the humectant base components PG and VG

are common to all e-liquids. For this reason, it is important to assess the nicotine-independent effects of e-liquids. Phagocytosis is dysregulated by e-cigarette vapor condensate treatment independent of nicotine, and partially restored by antioxidant/antialdehyde, N-acetyl cysteine (NAC) cotreatment, suggesting a free radical/aldehyde driven mechanism (36). THP-1 macrophages exhibit upregulated reactive oxygen species (ROS) production in response to nicotine-free e-cigarette vapor condensate exposure (36). The PI3K/Akt pathway is known to mediate ROS production (50), and work has demonstrated pan-PI3K inhibition and specific PI3K α inhibition improved viability and partially restored phagocytosis after nicotine-free e-cigarette vapor condensate challenge (36, 51). This work demonstrates nicotine is not the sole active component of e-cigarette vapor, yet needs further exploration.

Nicotine dependent.

Nicotine concentrations in e-liquids differ significantly between vapers, with users titrating nicotine intake to meet their addiction needs (52). Reduced efferocytosis of apoptotic epithelial cells has been reported (40), driven by a reduction in apoptotic cell receptor (CD36 and CD44) expression on the surface of alveolar macrophages following exposure to humectant bases PG and VG as well as nicotine. However, this treatment alone did not translate into functional reduction in efferocytosis in the absence of nicotine (40). Similarly, alveolar macrophages exhibit reduced expression of phagocytic receptors SR-A1 and TLR2 in response to nicotine-containing e-cigarette vapor extract, contributing to the reduction in phagocytosis (41). Release of proteinases is driven by an increase in intracellular Ca^{2+} levels in response to the opening of the ligand-gated nicotine receptors [nicotinic acetylcholine receptor (nAChRs)]. Macrophages express four types of nAChRs: CHRN1, CHRNA5, CHRNA6, and CHRNA7 and their activation by nicotine increases intracellular Ca^{2+} levels and NF- κ B signaling (42, 53).

Lipid metabolism.

Macrophages contribute to lipid metabolism and homeostasis in a range of tissues including the liver, spleen, and lungs, as they clear lipids from cell debris and excess cholesterol (54). In the lungs, alveolar macrophages catabolize surfactant and along with alveolar type II cells regulate pulmonary surfactant reducing surface tension and contributing to immune host defense against invading pathogens (55). In patients with EVALI, particularly those who use THC products, lipid-laden macrophages that stain heavily with oil red O stain have been reported in histological analyses of bronchoalveolar lavage (BAL) samples (10, 56). The derivation of these unique foam-like cells in EVALI is uncertain due to the presence in young, otherwise healthy individuals with an absence of hyperlipidemia (57). In addition, exogenous lipid exposure from oil-based e-cigarette vapor and persistent inflammation driving formation of lipid-laden macrophages, further evidence has shown that disruption of lipid metabolism causes lipid accumulation from endogenous sources in a nicotine-independent mechanism (45, 58). Although lipid-laden macrophages are not unique to patients with EVALI, and have been shown to be present in nonvapers and smokers (59–61), they contribute to the pathogenesis of exogenous

lipoid pneumonia, as in the presentation of EVALI (62). These lipid-laden macrophages have impaired function, particularly phagocytosis (54, 63), so may explain some of the functional implications of e-cigarette exposure reported. Finally, in vivo, lamellar bodies that regulate surfactant catabolism appeared poorly organized and irregular, suggesting changes in surfactant processing had occurred (45). Impaired surfactant homeostasis by macrophages causes excess lipid deposition in the airway and accumulation of intracellular lipids (64).

NEUTROPHILS

Neutrophils are essential innate immune cells, constituting 70% of circulating leukocytes and are rapidly recruited to sites of infection or injury in tissues (65). Neutrophils play an important role in tissue homeostasis including inflammation initiation and resolution. Neutrophils have an arsenal of killing mechanisms, which can be deployed in response to infection, phagocytosis, degranulation and neutrophil extracellular traps (NET)osis (66).

Neutrophil dysfunction has been reported in a range of disease states including COPD, sepsis, and inflammaging (67–70). Cigarette smoking upregulates neutrophil numbers in BAL, hinders normal neutrophil function, and promotes a proinflammatory phenotype (71–73), leading to persistent inflammation and increased tissue damage, which can cause early changes in the lung associated with progression to COPD. In health, there is an obligate area of tissue damage following neutrophil degranulation; however, disturbances of the proteinase-antiproteinase balance lead to uncontrolled promotion of tissue destructive proteinases including NE without the corresponding inhibition by antiproteinases such as α -1 antitrypsin (74, 75).

Neutrophils and E-cigarettes

Murine models initially showed unchanged neutrophil numbers in BAL after e-cigarette exposure (76); however, further delineation of humectants, nicotine, and flavorants found significant infiltration of neutrophils into the lung after exposure to 12 mg/mL nicotine flavorless e-liquid (50/50 PG/VG) (77). These studies used whole body exposure methods, which may have diluted out the impact on lung tissue. This finding supports the recently reported EVALI cases, cytopathological findings revealed neutrophil infiltration into the airways (58% of BAL cells were neutrophils) (10). However, data from cohort studies have reported no difference in neutrophil numbers in either BAL or induced sputum samples of e-cigarette users compared with nonsmokers and smokers (42, 78). Despite mixed findings on lung infiltration of neutrophils as a result of e-cigarette use, neutrophilic granule enzymes including NE and matrix metalloproteinases (MMPs) are consistently elevated (42, 78, 79) allows unopposed tissue destruction by proteinases and increases the potential for bystander tissue damage upon neutrophil recruitment to the airways.

E-cigarette vapor extract has not been reported to be cytotoxic to neutrophils; however, markers of neutrophil activation are elevated after exposure, including morphological shape

change and activation marker expression (CD11b and CD66b) (79). Much of the literature on neutrophil function in response to e-cigarette vapor exposure is difficult to interpret, as authors neglect to report concomitant effects on neutrophil viability in these experiments.

Previous work has highlighted the ease in which primed neutrophils degranulate and subsequently easily get caught in the lung (80, 81). Neutrophils exhibit a diminished ability to produce ROS, which coincides with the reduced ability to migrate through a membrane toward a chemoattractant after exposure to e-cigarettes (82). Phagocytosis of *Escherichia coli* and *Staphylococcus aureus* by circulating neutrophils is significantly reduced by e-cigarette exposure independently of nicotine; indicating that e-cigarette use may limit the control and elimination of bacterial infections, even within the blood (39, 82). The effect of e-cigarette exposure on NETosis varies depending on the exposure model and NET inducer investigated, with mixed reports of increasing suppression and sensitivity to NET production. In a report examining neutrophils from vapor's BAL, neutrophils were more sensitive to NET-stimulating Phorbol 12-myristate 13-acetate (PMA) than controls (78). Table 2 summarizes the key studies investigating neutrophil responses to e-cigarette exposure, including the device type, exposure model, and functional findings.

Mechanisms

Few studies have investigated the mechanisms of e-cigarette effects on neutrophils and those that have are largely driven by speculative hypotheses. The mechanism for dysregulated chemotaxis remains unclear. One explanation is that chemotaxis is dysregulated by e-cigarette-mediated changes in membrane fluidity and inability to polarize f-actin polymerization in response to a chemotactic agent [*N*-formylmethionine-leucyl-phenylalanine (fMLP)] (82), yet the cause of this remains unclear. NETosis relies upon the activation of protein kinase C (PKC) activation (83); however, the impact of e-cigarette exposure on PKC activation in neutrophils or an alternative pathway to NETosis has not been investigated. Evidence from cigarette smoke extract and nicotine treatments on neutrophils shows upregulation of p38 mitogen-activated protein kinase (MAPK), extracellular signal-regulated kinase (ERK) and nuclear factor (NF)- κ B pathways (84). In response to high-dose e-cigarette vapor extract treatment, neutrophils exhibit p38 MAPK activation but no change in ERK or NF- κ B pathway activation (79). Further investigation revealed secretion of MMP-9 by e-cigarette-exposed neutrophils is driven by p38 MAPK activation and can be reduced by chemical inhibition of this pathway (79). This indicates that there are both differential and overlapping mechanisms of e-cigarette damage compared with cigarette damage, which need further investigating to understand the possible impacts of chronic vaping. The mechanism of e-cigarette driven impaired neutrophil phagocytosis remains unknown. There may be receptor sensitivity/cleavage that prevent pathogen recognition, cytoskeletal rearrangement issues that prevent the internalization of the microbe, issues with assembly of NADPH oxidase preventing bacterial killing or another cause of this dysfunction, requiring more robust studies to confirm the driving mechanism of these effects. However, baseline ROS generation from neutrophils treated with e-

Table 2. A summary of the effects of e-cigarette vapor exposure on neutrophil effector functions

Effector Function	E-Cigarette Model	ECL Nicotine Concentration	Exposure Model	Exposure Duration	Cell Type	Results	Reference
Viability	VIP 1100 mAh battery with V5/CE5 Clearomizer	24 mg	ECVE (optical density values: 0.001, 0.003, 0.01, 0.03, 0.1)	6 h	Isolated peripheral blood neutrophils of never smokers	No effect on neutrophil apoptosis or necrosis at any concentrations but significant morphological changes and increased cell activation markers (CD11b and CD66b)	(79)
Reactive oxygen species	Kanger Miniprotank glassomizers attached to Kanger eVOD variable voltage 1,000 mAh battery	24 mg/mL	ECVE (100%, 75%, 50%, 25% concentrations)	20 min	Isolated peripheral blood neutrophils	Baseline ROS production significantly reduced by 100%, 75%, and 50% ECVE vs. UTC PMA-induced ROS unaffected	(82)
Phagocytosis/antimicrobial activity	LAVABOX DNA 200 Box Mod (Volcano e-cigs) and SMOK TFV4 mini tank with sub-ohm coil	0	ECL (PG/VG vehicle control) (1%, 0.5%, or 0.25% concentration)	30 min	Isolated peripheral blood neutrophils	Significantly reduced phagocytosis of <i>S. aureus</i> vs. UTC	(39)
	Kanger Miniprotank glassomizers attached to Kanger eVOD variable voltage 1,000 mAh battery	24 mg/mL	ECVE (100%, 75%, 50%, 25% concentrations)	30 min	Isolated peripheral blood neutrophils	40% reduction in phagocytosis of <i>E. coli</i> and <i>S. aureus</i> when treated with 100% ECVE vs. UTC	(82)
Cytokine/proteinase release	VIP 1100 mAh battery with V5/CE5 clearomizer	24 mg	ECVE (optical density values: 0.001, 0.003, 0.01, 0.03, 1)	6 h	Isolated peripheral blood neutrophils of never smokers	Significantly upregulated MMP-9 secretion and activity at 0.003 and 0.01 OD concentrations vs. UTC. Increase in NE secretion at 0.003 OD but decreased secretion at 0.03 and 0.1 OD vs. UTC	(79)
	N/A	N/A	N/A	N/A	BAL proteinase levels	NE, MMP-2, and MMP-9 secretion and activity were significantly elevated in BAL of vapers compared with non-smokers Antiproteinases (A1AT, CLPI, and TIMP-1/2) unchanged in BAL of vapers compared with nonsmokers	(42)
	None	18 mg/mL	PG/VG (55:45 v/v)	4 h	Isolated peripheral blood neutrophils	Significantly increased NE secretion vs. UTC	(42)
Migration	Kanger Miniprotank glassomizers attached to Kanger eVOD variable voltage 1,000 mAh battery	24 mg/mL	ECVE (100%, 75%, 50%, 25% concentrations)	20 min	Isolated peripheral blood neutrophils	ECVE (50%, 75%, and 100%) significantly reduced migration of neutrophils to fMLP vs. UTC	(82)
NETosis	N/A	N/A	N/A	N/A	Isolated peripheral blood neutrophils from smokers, e-cigarette users and healthy individuals	Significantly more sensitive to NET stimuli than smokers or healthy individuals	(78)
	Kanger Miniprotank glassomizers attached to Kanger eVOD variable voltage 1,000 mAh battery	24 mg/mL and 0 mg/mL	ECVE (100%, 75%, 50%, 25% concentrations)	20 min	Isolated peripheral blood neutrophils	24 mg/mL: No change in spontaneous or nigericin-induced NET production vs. UTC, but increasing suppression of PMA-induced NETS with increasing ECVE concentration 0 mg/mL: Significantly suppressed PMA-induced NET production vs. UTC	(82)

Studies investigating the effect of e-cigarettes on neutrophil viability, reactive oxygen species generation, phagocytosis, cytokine release, migration, and NETosis are included. The exposure design and cell type has been included for comparison of findings. Only experiments using the humectant bases with or without nicotine in the absence of any flavorings have been included to aid understanding of the fundamental effects of e-cigarette exposure on neutrophils. A1AT, alpha 1 antitrypsin; CD, cluster of differentiation; *E. coli*, Escherichia coli; ECVC, e-cigarette vapor condensate; ECVE, e-cigarette vapor extract; IL, interleukin; MMP, matrix metalloproteinase; NET, neutrophil extracellular trap; OD, optical density; PG, propylene glycol; PMA, phorbol myristate acetate; ROS, reactive oxygen species; *S. aureus*, *Staphylococcus aureus*; SLPI, secretory leukocyte protease inhibitor; TIMP, tissue inhibitor of metalloproteinase; TNF, tumor necrosis factor; UTC, untreated control; VG, vegetable glycerin.

cigarette vapor extract may signify an inability to rapidly assemble NADPH oxidase, which has implications for bacterial killing (85). Neutrophils express nicotine receptor variants CHRN1, CHRNA5, and CHRNA6 (42, 86) and exhibit an increase in intracellular Ca^{2+} concentration in response to e-cigarette exposure without a corresponding upregulation of NF- κ B activity, as is seen with direct nicotine-only treatment (42, 79, 87). Therefore, the downstream pathways of this receptor and its consequences for neutrophil functions require further investigations. Many questions remain about the causes of neutrophil dysfunction in response to e-cigarette exposure, meaning robust *in vitro* studies are required.

EPITHELIAL CELLS

Once considered to function purely as a static barrier, the airway epithelium is now recognized for its diverse range of functions and its importance in orchestrating inflammation

and tissue remodeling (88). Consisting of a range of cell types including secretory, ciliated, and type I/II alveolar cells, the airway epithelium is a dynamic tissue required to protect the lung against invading pathogens (89). Airway epithelial cells are central to lung health and their dysfunction contributes to lung diseases including asthma, COPD, and cystic fibrosis (88–90). Cigarette smoke exposure increases epithelial layer permeability and reduces cell-cell contacts (91, 92). In addition, impairing the mucociliary escalator drives mucus build up in the airways by impairing differentiation and function of ciliated epithelial cells that contribute to the development of chronic bronchitis, a feature of COPD (93, 94).

Epithelial Cells and E-cigarettes

Experimental models have attempted to recapitulate lung conditions to investigate the functional impact of e-cigarettes on epithelial cells. Table 3 summarizes the key findings regarding epithelial cell viability, ciliary parameters,

Table 3. A summary of the effects of e-cigarette vapor exposure on epithelial cell effector functions

Effector Function	E-Cigarette Model	ECL Nicotine Concentration	Exposure Model	Exposure Duration	Cell Type	Results	Reference
Viability	Third-generation device	1.2%	ECVE	72 h	BEAS-2B	Significant toxicities at 5% and 10% concentrations vs. UTC	(95)
	EVOD-2 3.7 V, 1.5 Ω	18 mg/mL and 0 mg/mL	100% ECVE (50 \times 3 s puffs, 5 s break)	24 h	HBEC	No increase in apoptosis or necrosis	(96)
	VC-1 exposure system	36 mg/mL, 0 mg/mL, and 100 μ M (aerosolized, calculated final exposure: 100 nM)	Air-liquid interface	36 puffs (1 every 30 s)	NHBE from non-smoker donors	Viability was unaffected vs. UTC	(76)
	Voltage-variable ENDS device from Vision Spinner II 1,600 mAh	1.1%	Air-liquid interface	0–60 min	HBEC	No effect on cellular viability vs. UTC (except at 60-min exposure)	(97)
Ciliary parameters	VC-1 exposure system	36 mg/mL	Air-liquid interface	8 h	NHBE from non-smoker donors	Reduced ciliary beat frequency but no change in percentage of ciliated cells vs. UTC	(76)
	Information not available	2.4%	Air-liquid interface	20 puffs	NHBE from non-smoker donors	Reduced ciliary beat frequency	(98)
Ion channel function	Voltage-variable ENDS device from Vision Spinner II 1,600 mAh	1.1%	Air-liquid interface	0–60 min	HBEC	Dose-dependent inhibition of chloride ion transfer vs. UTC	(97)
Cytokine release	Third-generation device	1.2%	ECVE	72 h	BEAS-2B	Significantly increased CXCL8, collagen I, and fibronectin secretion vs. UTC	(95)
	N/A	Nicotine-supplemented media (0, 1 or 10 μ M)	Air-liquid interface	5 days	NHBE from non-smoker donors	Elevated secretion of IL-6, IL-8, and a trend toward an increase in MCP-1 secretion vs. UTC	(76)
	Steamo Nova2 e-cigarette. 2.2 Ω , 2.6 V	18 mg/mL (calculated final exposure: 2.4 mg)	Air-liquid interface	25 min (3 s puff every 29 s)	Calu-3 cell line	Significant increase in IL-8 secretion 24 h after exposure	(99)
	EVOD-2 3.7 V, 1.5 Ω	18 mg/mL	100% ECVE (50 \times 3 s puffs, 5 s break)	24 h	HBEC	Significant reduction in TNF- α , MIP-1 β , and IP-10 secretion; however, no change in IL-6, IL-8, or MIP-1 α	(96)
Protein expression	SIGLEI 150 W with TFV4 top refill tank	0 (PG/VG 55/45)	Air-liquid interface	36 puffs (measured 24 h later)	Primary HBEC	3–4 fold increase in MUC5AC levels and significant increase in CYP1B1	(100)

Studies investigating the effect of e-cigarettes on epithelial cell viability, ciliary parameters, ion channel function, cytokine release, and protein expression are included. The exposure model and cell type have been included for comparison of findings. Only experiments using the humectant bases with or without nicotine in the absence of any flavorings have been included for simplicity and to aid understanding of the fundamental effects of e-cigarette exposure on this important cell type. CYP1B1, cytochrome P450 family 1 subfamily B member 1; ECVC, e-cigarette vapor condensate; ECVE, e-cigarette vapor extract; HBEC, human bronchial epithelial cultures, IL, interleukin, MCP, methyl-accepting chemotaxis protein; MUC5A, mucin 5AC; NHBE, normal human bronchial epithelial; PG, propylene glycol; TNF, tumor necrosis factor; UTC, untreated control; VG, vegetable glycerin.

cytokine release, ion channel function, and protein expression after e-cigarette exposure.

High concentrations of e-cigarette vapor extract trigger epithelial cell toxicities comparable to cigarette smoke extract exposure but how these doses compare with user exposure is unclear, as no data regarding e-cigarette vapor extract characteristics were provided (95). Other findings have indicated no induction of apoptosis or necrosis by unflavored e-cigarette exposure (40, 76, 97). Dysfunctional cilia beat frequency and motility have been reported in human nasal epithelial cells (HNEC) and normal human bronchial epithelial (NHBE) cells exposed to e-cigarettes in a manner similar to that seen with cigarette smoke extract (76, 98). Although the authors describe a return to baseline function over time, they do not rule out the potential of lasting damage with chronic exposure (98). Beas-2B, NHBE and Calu-3 cells exposed to e-cigarette vapor exhibit an increase in IL8 secretion (76, 78, 95). Extracellular matrix (ECM) proteins, collagen 1 and fibronectin, are secreted in a concentration-dependent manner in Beas-2B and airway smooth muscle cells exposed to e-cigarette vapor extract, leading to formation of mesh-like fibers observed in e-cigarette vapor-exposed HNECs and contributes to tissue remodeling (95, 98).

Mechanisms

Signaling molecules have been implicated as contributors to the functional changes observed by e-cigarette exposure to epithelial cells. $\alpha 7$ nAChRs on the surface of epithelial cells upregulate the PKC and ERK signaling pathways after stimulation with nicotine (101, 102). These pathways are involved with inflammatory responses and controlled cell death in epithelial cells but focused investigations on these molecules and their contribution to dysfunction after e-cigarette exposure are lacking (103). The regulator of ciliogenesis, FOXJ1, is downregulated in nicotine-containing e-cigarette-exposed NHBE cells, suggesting a possible mechanism for the reduction in ciliary beat function (76). Cigarette smoke inhibits cystic fibrosis transmembrane regulation (CFTR) in a Ca^{2+} -dependent manner, which can be repressed by direct elevations in Ca^{2+} (104, 105). Independent of changes in gene expression, conductance of the CFTR channel is impaired by e-liquid in a similar manner to that caused by cigarette smoke extract (76, 97). This is an effect of vaporized but not unvaporized e-liquid and drives airway dehydration (97) indicating the change in chemical composition, which occurs during vaping is a key factor in this activity. Reactive carbonyl species generated during vaping (28, 106) must therefore be the key active agent in mediating this effect. In addition, ion channel conductance of the big potassium (BK) channel is reduced by e-cigarette vapor in airway epithelial cells due to a reduced expression in α subunit gene (KCNMA1) (76), which leads to impairment of fluid homeostasis and air surface liquid (107). E-cigarette-exposed Beas-2B cells exhibit mitochondrial uncoupling after exposure to e-cigarette vapor extract as well as increased glycolysis at high e-cigarette vapor extract concentrations, suggesting a pivotal role of dysfunctional mitochondria in the aberrant function of airway epithelial cells that lead to airway remodeling and scarring after persistent insult or injury such as that caused by e-cigarette usage (95, 108). The mechanism

driving cytokine release not been investigated; however, IL8 was not upregulated by nicotine-free vapor exposure in NHBE cells, indicating this may be a nicotine-dependent effect (76). These molecules identified as potential mechanisms of the effects seen on epithelial cells are promising but lack in-depth investigations to conclusively report the mechanism of damage.

SUMMARY

The current evidence surrounding the potential impact of e-cigarette exposure on alveolar macrophages, neutrophils, and lung epithelial cells is limited but building. To date, studies have been small and exposure durations short. There have been many variations in exposure model, e-cigarette device and e-liquid used, and there are still important gaps to be filled. However, there are consistent signals of short-term harm in alveolar macrophage, neutrophil, and epithelial dysfunction. These short-term effects are likely to have long-term implications with repeated exposure. For example, e-cigarette-driven decline in efferocytosis by alveolar macrophages leads to reduced clearance of apoptotic neutrophils, driving neutrophil-induced inflammation and secondary necrosis, which has been shown to contribute to the pathophysiology of COPD (109, 110). In addition, impaired neutrophil chemotaxis and susceptibility to NET formation, as caused by e-cigarette exposure, has also been demonstrated in neutrophils COPD patients (22, 111). Finally, increased IL-8 secretion from lung epithelial cells has been associated with worse obstruction in COPD, with a multifactorial impact on disease pathophysiology (112). Demonstrating the relationship between cellular dysfunction reported in e-cigarette studies and those driving pathophysiology in just one chronic inflammatory lung disease with significant quality of life burdens highlights potential for chronic disease development with persistent e-cigarette exposure. Extensive studies investigating the contribution of nicotine receptors, signaling pathways, transcriptional/epigenetic alterations, and metabolic changes across all cell types are needed. Although large longitudinal studies of vapers will be required to fully determine the effects of chronic vaping, these studies will take years, by which time e-cigarettes may be embedded within society. Better in vitro studies may give us crucial insight into how e-cigarettes affect key cell types and importantly how these mechanisms differ from that of cigarette smoking. Until these processes are understood, we cannot reliably inform policy makers, healthcare professionals, or the public on the safety of e-cigarettes.

GRANTS

This work was supported in part by Health Data Research-UK PIONEER 2019 (to E. Sapey), British Lung Foundation Grant PPRG16-12 (to D. R. Thickett and A. Scott), National Institute for Health Research Health Technology Assessment Programme Grant NIHR129593 (to E. Sapey, D. R. Thickett, and A. Scott), and United Kingdom Research and Innovation Medical Research Council Grant MR/S002782/1 (to E. Sapey, D. R. Thickett, and A. Scott).

DISCLOSURES

No conflicts of interest, financial or otherwise, are declared by the authors.

AUTHOR CONTRIBUTIONS

A.E.J. prepared figures; A.E.J. and A.S. drafted manuscript; A.E.J., E.S., and A.S. edited and revised manuscript; A.E.J., E.S., D.R.T., and A.S. approved final version of manuscript.

REFERENCES

- Goniewicz ML, Lingas EO, Hajek P. Patterns of electronic cigarette use and user beliefs about their safety and benefits: an internet survey. *Drug Alcohol Rev* 32: 133–140, 2013. doi:10.1111/j.1465-3362.2012.00512.x.
- Hajek P, Phillips-Waller A, Przulj D, Pesola F, Myers Smith K, Bisal N, Li J, Parrott S, Sasieni P, Dawkins L, Ross L, Goniewicz M, Wu Q, McRobbie HJ. A randomized trial of e-cigarettes versus nicotine-replacement therapy. *N Engl J Med* 380: 629–637, 2019. doi:10.1056/NEJMoa1808779.
- Zhu SH, Sun JY, Bonnevie E, Cummins SE, Gamst A, Yin L, Lee M. Four hundred and sixty brands of e-cigarettes and counting: implications for product regulation. *Tob Control* 23: iii3–iii9, 2014. doi:10.1136/tobaccocontrol-2014-051670.
- Soneji S, Barrington-Trimis JL, Wills TA, Leventhal AM, Unger JB, Gibson LA, Yang J, Primack BA, Andrews JA, Miech RA, Spindle TR, Dick DM, Eissenberg T, Hornik RC, Dang R, Sargent JD. Association between initial use of e-cigarettes and subsequent cigarette smoking among adolescents and young adults: a systematic review and meta-analysis. *JAMA Pediatr* 171: 788–797, 2017. doi:10.1001/jamapediatrics.2017.1488.
- Evans SE, Hoffman AC. Electronic cigarettes: abuse liability, topography and subjective effects. *Tob Control* 23, Suppl 2: ii23–ii29, 2014. doi:10.1136/tobaccocontrol-2013-051489.
- McQueen A, Tower S, Sumner W. Interviews with “vapers”: implications for future research with electronic cigarettes. *Nicotine Tob Res* 13: 860–867, 2011. doi:10.1093/ntr/ntr088.
- Zhu SH, Gamst A, Lee M, Cummins S, Yin L, Zoref L. The use and perception of electronic cigarettes and snus among the U.S. population. *PLoS One* 8: e79332, 2013. doi:10.1371/journal.pone.0079332.
- Health AoSa. Use of e-cigarettes (vaporisers) among adults in Great Britain (Online). <https://ash.org.uk/information-and-resources/factsheets/statistical/use-of-e-cigarettes-among-adults-in-great-britain-2020/> [2020 Oct 14].
- Euromonitor International. Smokeless tobacco and vapour products. <https://www.euromonitor.com/smokeless-tobacco-and-vapour-products> [2018 Sep].
- Layden JE, Ghinai I, Pray I, Kimball A, Layer M, Tenforde MW, Navon L, Hoots B, Salvatore PP, Elderbrook M, Haupt T, Kanne J, Patel MT, Saathoff-Huber L, King BA, Schier JG, Mikosz CA, Meiman J. Pulmonary illness related to e-cigarette use in Illinois and Wisconsin—final report. *N Engl J Med* 382: 903–916, 2020. doi:10.1056/NEJMoa1911614.
- Blount BC, Karwowski MP, Shields PG, Morel-Espinosa M, Valentin-Blasini L, Gardner M; Lung Injury Response Laboratory Working Group, et al. Vitamin E acetate in bronchoalveolar-lavage fluid associated with EVALI. *N Engl J Med* 382: 697–705, 2020. doi:10.1056/NEJMoa1916433.
- Reboul E. Vitamin E bioavailability: mechanisms of intestinal absorption in the spotlight. *Antioxidants (Basel)* 6: 95, 2017. doi:10.3390/antiox6040095.
- Massey JB, She HS, Pownall HJ. Interaction of vitamin E with saturated phospholipid bilayers. *Biochem Biophys Res Commun* 106: 842–847, 1982. doi:10.1016/0006-291X(82)91787-9.
- Zuo YY, Veldhuizen RA, Neumann AW, Petersen NO, Possmayer F. Current perspectives in pulmonary surfactant—inhibition, enhancement and evaluation. *Biochim Biophys Acta* 1778: 1947–1977, 2008. doi:10.1016/j.bbame.2008.03.021.
- Outbreak of Lung Injury Associated with E-cigarette Use, or Vaping Centers for Disease Control and Prevention. https://www.cdc.gov/tobacco/basic_information/e-cigarettes/severe-lung-disease.html [2020 Jul 1].
- Medicines and Healthcare products Regulatory Agency. Drug Safety Update 2020: E-cigarette use or vaping: reporting suspected adverse reactions, including lung injury (Online). <https://www.gov.uk/drug-safety-update/e-cigarette-use-or-vaping-reporting-suspected-adverse-reactions-including-lung-injury>.
- Centers for Disease Control and Prevention, National Center for Chronic Disease P, Health P, Office on S, and Health. Publications and reports of the surgeon general. In: *How Tobacco Smoke Causes Disease: The Biology and Behavioral Basis for Smoking-Attributable Disease: A Report of the Surgeon General*. Atlanta (GA): Centers for Disease Control and Prevention (US), 2010.
- Hill A, Gompertz S, Stockley R. Factors influencing airway inflammation in chronic obstructive pulmonary disease. *Thorax* 55: 970–977, 2000. doi:10.1136/thorax.55.11.970.
- Mirza S, Clay RD, Koslow MA, Scanlon PD. COPD guidelines: a review of the 2018 GOLD report. *Mayo Clin Proc* 93: 1488–1502, 2018. doi:10.1016/j.mayocp.2018.05.026.
- King PT. Inflammation in chronic obstructive pulmonary disease and its role in cardiovascular disease and lung cancer. *Clin Transl Med* 4: 68–68, 2015. doi:10.1186/s40169-015-0068-z.
- Senn O, Russi EW, Imboden M, Probst-Hensch NM. alpha1-Antitrypsin deficiency and lung disease: risk modification by occupational and environmental inhalants. *European Respiratory Journal* 26: 909–917, 2005. doi:10.1183/09031936.05.00021605.
- Stockley JA, Walton GM, Lord JM, Sapey E. Aberrant neutrophil functions in stable chronic obstructive pulmonary disease: the neutrophil as an immunotherapeutic target. *Int Immunopharmacol* 17: 1211–1217, 2013. doi:10.1016/j.intimp.2013.05.035.
- Barnes PJ. Cellular and molecular mechanisms of chronic obstructive pulmonary disease. *Clin Chest Med* 35: 71–86, 2014. doi:10.1016/j.ccm.2013.10.004.
- Wang Y, Xu J, Meng Y, Adcock IM, Yao X. Role of inflammatory cells in airway remodeling in COPD. *Int J Chron Obstruct Pulmon Dis* 13: 3341–3348, 2018. doi:10.2147/COPD.S176122.
- Lokke A, Lange P, Scharling H, Fabricius P, Vestbo J. Developing COPD: a 25 year follow up study of the general population. *Thorax* 61: 935–939, 2006. doi:10.1136/thx.2006.062802.
- National Academies of Sciences Engineering, and Medicine, Health and Medicine Division. ; Board on Population Health and Public Health Practice; Committee on the Review of the Health Effects of Electronic Nicotine Delivery Systems. *Public Health Consequences of E-Cigarettes*, edited by Eaton DL, Kwan LY, Stratton K. Washington (DC): National Academic Press, 2018. doi:10.17226/24952.
- Olmedo P, Navas-Acien A, Hess C, Jarmul S, Rule A. A direct method for e-cigarette aerosol sample collection. *Environ Res* 149: 151–156, 2016. doi:10.1016/j.envres.2016.05.008.
- Kosmider L, Sobczak A, Fik M, Knysak J, Zaciera M, Kurek J, Goniewicz ML. Carbonyl compounds in electronic cigarette vapors: effects of nicotine solvent and battery output voltage. *Nicotine Tob Res* 16: 1319–1326, 2014. doi:10.1093/ntr/ntu078.
- Bekki K, Uchiyama S, Ohta K, Inaba Y, Nakagome H, Kunugita N. Carbonyl compounds generated from electronic cigarettes. *Int J Environ Res Public Health* 11: 11192–11200, 2014. doi:10.3390/ijerph11111192.
- Allard B, Panariti A, Martin JG. Alveolar macrophages in the resolution of inflammation, tissue repair, and tolerance to infection. *Front Immunol* 9: 1777, 2018. doi:10.3389/fimmu.2018.01777.
- Vlahos R, Bozinovski S. Role of alveolar macrophages in chronic obstructive pulmonary disease. *Front Immunol* 5: 435, 2014. doi:10.3389/fimmu.2014.00435.
- Aberdein JD, Cole J, Bewley MA, Marriott HM, Dockrell DH. Alveolar macrophages in pulmonary host defence the unrecognized role of apoptosis as a mechanism of intracellular bacterial killing. *Clin Exp Immunol* 174: 193–202, 2013. doi:10.1111/cei.12170.
- Monick MM, Powers LS, Walters K, Lohan N, Zhang M, Gerke A, Hansdottir S, Hunninghake GW. Identification of an autophagy defect in smokers' alveolar macrophages. *J Immunol* 185: 5425–5435, 2010. doi:10.4049/jimmunol.1001603.
- Hodge S, Matthews G, Mukaro V, Ahern J, Shivam A, Hodge G, Holmes M, Jersmann H, Reynolds PN. Cigarette smoke-induced changes to alveolar macrophage phenotype and function are improved by treatment with procysteine. *Am J Respir Cell Mol Biol* 44: 673–681, 2011. doi:10.1165/rcmb.2009-0459OC.
- Hodge S, Hodge G, Ahern J, Jersmann H, Holmes M, Reynolds PN. Smoking alters alveolar macrophage recognition and phagocytic ability: implications in chronic obstructive pulmonary disease. *Am J*

- Respir Cell Mol Biol* 37: 748–755, 2007. doi:10.1165/rcmb.2007-0025OC.
36. Scott A, Lugg ST, Aldridge K, Lewis KE, Bowden A, Mahida RY, Grudzinska FS, Dosanjh D, Parekh D, Foronjy R, Sapey E, Naidu B, Thickett DR. Pro-inflammatory effects of e-cigarette vapour condensate on human alveolar macrophages. *Thorax* 73: 1161–1169, 2018. doi:10.1136/thoraxjnl-2018-211663.
 37. Gómez A-C, Rodríguez-Fernández P, Villar-Hernández R, Gibert I, Muriel-Moreno B, Lacoma A, Prat-Aymerich C, Domínguez J. E-cigarettes: effects in phagocytosis and cytokines response against *Mycobacterium tuberculosis*. *PLoS One* 15: e0228919, 2020. doi:10.1371/journal.pone.0228919.
 38. Hwang JH, Lyes M, Sladewski K, Enany S, McEachern E, Mathew DP, Das S, Moshensky A, Bapat S, Pride DT, Ongkeko WM, Crotty Alexander LE. Electronic cigarette inhalation alters innate immunity and airway cytokines while increasing the virulence of colonizing bacteria. *J Mol Med (Berl)* 94: 667–679, 2016. doi:10.1007/s00109-016-1378-3.
 39. Clapp PW, Pawlak EA, Lackey JT, Keating JE, Reeber SL, Glish GL, Jaspers I. Flavored e-cigarette liquids and cinnamaldehyde impair respiratory innate immune cell function. *Am J Physiol Lung Cell Mol Physiol* 313: L278–L292, 2017. doi:10.1152/ajplung.00452.2016.
 40. Ween MP, Hamon R, Macowan MG, Thredgold L, Reynolds PN, Hodge SJ. Effects of E-cigarette E-liquid components on bronchial epithelial cells: demonstration of dysfunctional efferocytosis. *Respirology* 25: 620–628, 2020. doi:10.1111/resp.13696.
 41. Ween MP, Whittall JJ, Hamon R, Reynolds PN, Hodge SJ. Phagocytosis and inflammation: exploring the effects of the components of E-cigarette vapor on macrophages. *Physiol Rep* 5: e13370, 2017. doi:10.14814/phy2.13370.
 42. Ghosh A, Coakley RD, Ghio AJ, Muhlebach MS, Esther CR Jr, Alexis NE, Tarran R. Chronic E-cigarette use increases neutrophil elastase and matrix metalloproteinase levels in the lung. *Am J Respir Crit Care Med* 200: 1392–1401, 2019. doi:10.1164/rccm.201903-0615OC.
 43. Bootman MD, Bultynck G. Fundamentals of cellular calcium signaling: a primer. *Cold Spring Harb Perspect Biol* 12: a038802, 2020. doi:10.1101/cshperspect.a038802.
 44. Ghosh A, Beyazcicek O, Davis ES, Onyenwoke RU, Tarran R. Cellular effects of nicotine salt-containing e-liquids. *J Appl Toxicol* 41: 493–505, 2021. doi:10.1002/jat.4060.
 45. Madison MC, Landers CT, Gu B-H, Chang C-Y, Tung H-Y, You R, Hong MJ, Baghaei N, Song L-Z, Porter P, Putluri N, Salas R, Gilbert BE, Levental I, Campen MJ, Corry DB, Kheradmand F. Electronic cigarettes disrupt lung lipid homeostasis and innate immunity independent of nicotine. *J Clin Invest* 129: 4290–4304, 2019. doi:10.1172/JCI128531.
 46. Butt YM, Smith ML, Tazelaar HD, Vaszar LT, Swanson KL, Cecchini MJ, Boland JM, Bois MC, Boyum JH, Froemming AT, Khoo A, Mira-Avendano I, Patel A, Larsen BT. Pathology of vaping-associated lung injury. *N Engl J Med* 381: 1780–1781, 2019. doi:10.1056/NEJMc1913069.
 47. Tobacco Smoke Components. *Beiträge zur Tabakforschung International/Contributions to Tobacco Research* 17: 61–66, 1997.
 48. Thielen A, Klus H, Müller L. Tobacco smoke: unraveling a controversial subject. *Exp Toxicol Pathol* 60: 141–156, 2008. doi:10.1016/j.etp.2008.01.014.
 49. Goniewicz ML, Knysak J, Gawron M, Kosmider L, Sobczak A, Kurek J, Prokopowicz A, Jablonska-Czapla M, Rosik-Dulewska C, Havel C, Jacob P 3rd, Benowitz N. Levels of selected carcinogens and toxicants in vapour from electronic cigarettes. *Tob Control* 23: 133–139, 2014. doi:10.1136/tobaccocontrol-2012-050859.
 50. Koundouros N, Poulogiannis G. Phosphoinositide 3-kinase/Akt signaling and redox metabolism in cancer. *Front Oncol* 8: 160, 2018. doi:10.3389/fonc.2018.00160.
 51. Baker JR, Vuppusetty C, Colley T, Papaioannou AI, Fenwick P, Donnelly L, Ito K, Barnes PJ. Oxidative stress dependent microRNA-34a activation via PI3K α reduces the expression of sir-tuin-1 and sir-tuin-6 in epithelial cells. *Sci Rep* 6: 35871, 2016. doi:10.1038/srep35871.
 52. Dawkins LE, Kimber CF, Doig M, Feyerabend C, Corcoran O. Self-titration by experienced e-cigarette users: blood nicotine delivery and subjective effects. *Psychopharmacology (Berl)* 233: 2933–2941, 2016. doi:10.1007/s00213-016-4338-2.
 53. Bai X, Stitzel JA, Bai A, Zambrano CA, Phillips M, Marrack P, Chan ED. Nicotine impairs macrophage control of *Mycobacterium tuberculosis*. *Am J Respir Cell Mol Biol* 57: 324–333, 2017. doi:10.1165/rcmb.2016-0270OC.
 54. Remmerie A, Scott CL. Macrophages and lipid metabolism. *Cell Immunol* 330: 27–42, 2018. doi:10.1016/j.cellimm.2018.01.020.
 55. Whitsett JA, Wert SE, Weaver TE. Alveolar surfactant homeostasis and the pathogenesis of pulmonary disease. *Annu Rev Med* 61: 105–119, 2010. doi:10.1146/annurev.med.60.041807.123500.
 56. Mukhopadhyay S, Mehrad M, Dammert P, Arrossi AV, Sarda R, Brenner DS, Maldonado F, Choi H, Ghobrial M. Lung biopsy findings in severe pulmonary illness associated with e-cigarette use (vaping): a report of eight cases. *Am J Clin Pathol* 153: 30–39, 2019. doi:10.1093/ajcp/aqz182.
 57. Guerrini V, Gennaro ML. Foam cells: one size doesn't fit all. *Trends Immunol* 40: 1163–1179, 2019. doi:10.1016/j.it.2019.10.002.
 58. Singanayagam A, Snelgrove RJ. Less burn, more fat: electronic cigarettes and pulmonary lipid homeostasis. *J Clin Invest* 129: 4077–4079, 2019. doi:10.1172/JCI131336.
 59. Pambuccian SE. Testing for lipid-laden macrophages in bronchoalveolar lavage fluid to diagnose vaping-associated pulmonary injury. Are we there yet? *J Am Soc Cytopathol* 9: 1–8, 2020. doi:10.1016/j.jasc.2019.10.002.
 60. Ghosh A, Ahmad S, Coakley RD, Sassano MF, Alexis NE, Tarran R. Lipid-laden macrophages are not unique to patients with e-cigarette or vaping product use-associated lung injury. *Am J Respir Crit Care Med* 203: 1030–1033, 2021. doi:10.1164/rccm.202009-3507LE.
 61. Shields PG, Song MA, Freudenheim JL, Brasky TM, McElroy JP, Reisinger SA, Weng DY, Ren R, Eissenberg T, Wewers MD, Shilo K. Lipid laden macrophages and electronic cigarettes in healthy adults. *EBioMedicine* 60: 102982, 2020. doi:10.1016/j.ebiom.2020.102982.
 62. Sung S, Tazelaar HD, Crapanzano JP, Nassar A, Saqi A. Adult exogenous lipid pneumonia: a rare and underrecognized entity in cytology—a case series. *Cytojournal* 15: 17, 2018. doi:10.4103/cytojournal.cytojournal_29_17.
 63. Gibeon D, Zhu J, Sogbesan A, Banya W, Rossios C, Saito J, Rocha JP, Hull JH, Menzies-Gow AN, Bhavsar PK, Chung KF. Lipid-laden bronchoalveolar macrophages in asthma and chronic cough. *Respir Med* 108: 71–77, 2014. doi:10.1016/j.rmed.2013.10.005.
 64. Trapnell BC, Whitsett JA, Nakata K. Pulmonary alveolar proteinosis. *N Engl J Med* 349: 2527–2539, 2003. doi:10.1056/NEJMr023226.
 65. Dancy JT, Deubelbeiss KA, Harker LA, Finch CA. Neutrophil kinetics in man. *J Clin Invest* 58: 705–715, 1976. doi:10.1172/JCI108517.
 66. Kolaczowska E, Kubes P. Neutrophil recruitment and function in health and inflammation. *Nat Rev Immunol* 13: 159–175, 2013. doi:10.1038/nri3399.
 67. Patel JM, Sapey E, Parekh D, Scott A, Dosanjh D, Gao F, Thickett DR. Sepsis induces a dysregulated neutrophil phenotype that is associated with increased mortality. *Mediators Inflamm* 2018: 1–10, 2018. doi:10.1155/2018/4065362.
 68. Sapey E, Greenwood H, Walton G, Mann E, Love A, Aaronson N, Insall RH, Stockley RA, Lord JM. Phosphoinositide 3-kinase inhibition restores neutrophil accuracy in the elderly: toward targeted treatments for immunosenescence. *Blood* 123: 239–248, 2014. doi:10.1182/blood-2013-08-519520.
 69. Jasper A, McIver W, Sapey E, Walton G. Understanding the role of neutrophils in chronic inflammatory airway disease [version 1; peer review: 2 approved]. *F1000Res* 8: 557, 2019. doi:10.12688/f1000research.18411.1.
 70. Liew PX, Kubes P. The neutrophil's role during health and disease. *Physiol Rev* 99: 1223–1248, 2019. doi:10.1152/physrev.00012.2018.
 71. Karimi R, Tornling G, Grunewald J, Eklund A, Skold CM. Cell recovery in bronchoalveolar lavage fluid in smokers is dependent on cumulative smoking history. *PLoS One* 7: e34232, 2012. doi:10.1371/journal.pone.0034232.
 72. Xu Y, Li H, Bajrami B, Kwak H, Cao S, Liu P, Zhou J, Zhou Y, Zhu H, Ye K, Luo HR. Cigarette smoke (CS) and nicotine delay neutrophil spontaneous death via suppressing production of diphosphoinositol pentakisphosphate. *Proc Natl Acad Sci USA* 110: 7726–7731, 2013. doi:10.1073/pnas.1302906110.
 73. Guzik K, Skret J, Smagur J, Bzowska M, Gajkowska B, Scott DA, Potempa JS. Cigarette smoke-exposed neutrophils die

- unconventionally but are rapidly phagocytosed by macrophages. *Cell Death Dis* 2: e131, 2011. doi:10.1038/cddis.2011.13.
74. Gadek JE, Fells GA, Crystal RG. Cigarette smoking induces functional antiprotease deficiency in the lower respiratory tract of humans. *Science* 206: 1315–1316, 1979. doi:10.1126/science.316188.
 75. Stockley RA. Neutrophils and protease/antiprotease imbalance. *Am J Respir Crit Care Med* 160: S49–S52, 1999. doi:10.1164/ajrccm.160.supplement_1.13.
 76. Garcia-Arcos I, Geraghty P, Baumlín N, Campos M, Dabo AJ, Jundi B, Cummins N, Eden E, Grosche A, Salathe M, Foronjy R. Chronic electronic cigarette exposure in mice induces features of COPD in a nicotine-dependent manner. *Thorax* 71: 1119–1129, 2016. doi:10.1136/thoraxjnl-2015-208039.
 77. Chapman DG, Casey DT, Ather JL, Aliyeva M, Daphtary N, Lahue KG, van der Velden JL, Janssen-Heininger YMW, Irvin CG. The effect of flavored e-cigarettes on murine allergic airways disease. *Sci Rep* 9: 13671, 2019. doi:10.1038/s41598-019-50223-y.
 78. Reidel B, Radicioni G, Clapp PW, Ford AA, Abdelwahab S, Rebuli ME, Haridass P, Alexis NE, Jaspers I, Kesimer M. E-cigarette use causes a unique innate immune response in the lung, involving increased neutrophilic activation and altered mucin secretion. *Am J Respir Crit Care Med* 197: 492–501, 2018. doi:10.1164/rccm.201708-1590OC.
 79. Higham A, Rattray NJ, Dewhurst JA, Trivedi DK, Fowler SJ, Goodacre R, Singh D. Electronic cigarette exposure triggers neutrophil inflammatory responses. *Respir Res* 17: 56, 2016. doi:10.1186/s12931-016-0368-x.
 80. Summers C, Singh NR, White JF, Mackenzie IM, Johnston A, Solanki C, Balan KK, Peters AM, Chilvers ER. Pulmonary retention of primed neutrophils: a novel protective host response, which is impaired in the acute respiratory distress syndrome. *Thorax* 69: 623–629, 2014. doi:10.1136/thoraxjnl-2013-204742.
 81. Condliffe AM, Kitchen E, Chilvers ER. Neutrophil priming: pathophysiological consequences and underlying mechanisms. *Clin Sci (Lond)* 94: 461–471, 1998. doi:10.1042/cs0940461.
 82. Corriden R, Moshensky A, Bojanowski CM, Meier A, Chien J, Nelson RK, Crotty Alexander LE. E-cigarette use increases susceptibility to bacterial infection by impairment of human neutrophil chemotaxis, phagocytosis, and NET formation. *Am J Physiol Cell Physiol* 318: C205–C214, 2020. doi:10.1152/ajpcell.00045.2019.
 83. Gray RD, Lucas CD, MacKellar A, Li F, Hiersemenzel K, Haslett C, Davidson DJ, Rossi AG. Activation of conventional protein kinase C (PKC) is critical in the generation of human neutrophil extracellular traps. *J Inflamm* 10: 12, 2013. doi:10.1186/1476-9255-10-12.
 84. Koethe SM, Kuhnmuensch JR, Becker CG. Neutrophil priming by cigarette smoke condensate and a tobacco anti-idiotypic antibody. *Am J Pathol* 157: 1735–1743, 2000. doi:10.1016/S0002-9440(10)64810-9.
 85. Segal AW. How neutrophils kill microbes. *Annu Rev Immunol* 23: 197–223, 2005. doi:10.1146/annurev.immunol.23.021704.115653.
 86. Zia S, Ndoye A, Nguyen VT, Sa G. Nicotine enhances expression of the alpha 3, alpha 4, alpha 5, and alpha 7 nicotinic receptors modulating calcium metabolism and regulating adhesion and motility of respiratory epithelial cells. *Res Commun Mol Pathol Pharmacol* 97: 243–262, 1997.
 87. Iho S, Tanaka Y, Takauji R, Kobayashi C, Muramatsu I, Iwasaki H, Nakamura K, Sasaki Y, Nakao K, Takahashi T. Nicotine induces human neutrophils to produce IL-8 through the generation of peroxynitrite and subsequent activation of NF- κ B. *J Leukoc Biol* 74: 942–951, 2003. doi:10.1189/jlb.1202626.
 88. Hiemstra PS, McCray PB Jr, Bals R. The innate immune function of airway epithelial cells in inflammatory lung disease. *Eur Respir J* 45: 1150–1162, 2015. doi:10.1183/09031936.00141514.
 89. Crystal RG, Randell SH, Engelhardt JF, Voynow J, Sunday ME. Airway epithelial cells: current concepts and challenges. *Proc Am Thorac Soc* 5: 772–777, 2008. doi:10.1513/pats.200805-041HR.
 90. Puchelle E, Zahm JM, Tournier JM, Coraux C. Airway epithelial repair, regeneration, and remodeling after injury in chronic obstructive pulmonary disease. *Proc Am Thorac Soc* 3: 726–733, 2006. doi:10.1513/pats.200605-126SF.
 91. Nishida K, Brune KA, Putcha N, Mandke P, O'Neal WK, Shade D, Srivastava V, Wang M, Lam H, An SS, Drummond MB, Hansel NN, Robinson DN, Sidhaye VK. Cigarette smoke disrupts monolayer integrity by altering epithelial cell-cell adhesion and cortical tension. *Am J Physiol Lung Cell Mol Physiol* 313: L581–L591, 2017. doi:10.1152/ajplung.00074.2017.
 92. Heijink IH, Brandenburg SM, Postma DS, van Oosterhout AJM. Cigarette smoke impairs airway epithelial barrier function and cell-cell contact recovery. *Eur Respir J* 39: 419–428, 2012. doi:10.1183/09031936.00193810.
 93. Amatngalim GD, Vieira RP, Meiners S, Bartel S. Novel insights into the effects of cigarette smoke on the airway epithelial surface-lesions learned at the European Respiratory Society International Congress 2018 in Paris. *J Thorac Dis* 10: S2977–S2982, 2018. doi:10.21037/jtd.2018.08.17.
 94. Smaldone GC, Foster WM, O'Riordan TG, Messina MS, Perry RJ, Langenback EG. Regional impairment of mucociliary clearance in chronic obstructive pulmonary disease. *Chest* 103: 1390–1396, 1993. doi:10.1378/chest.103.5.1390.
 95. Sohal SS, Eapen MS, Naidu VGM, Sharma P. IQOS exposure impairs human airway cell homeostasis: direct comparison with traditional cigarette and e-cigarette. *ERJ Open Res* 5: 00159–2018, 2019. doi:10.1183/23120541.00159-2018.
 96. Ween MP, Hamon R, Macowan MG, Thredgold L, Reynolds PN, Hodge SJ. Effects of E-cigarette E-liquid components on bronchial epithelial cells: demonstration of dysfunctional efferocytosis. *Respirology* 25: 620–628, 2020. doi:10.1111/resp.13696.
 97. Lin VY, Fain MD, Jackson PL, Berryhill TF, Wilson LS, Mazur M, Barnes SJ, Blalock JE, Raju SV, Rowe SM. Vaporized e-cigarette liquids induce ion transport dysfunction in airway epithelia. *Am J Respir Cell Mol Biol* 61: 162–173, 2019. doi:10.1165/rcmb.2017-0432OC.
 98. Carson JL, Zhou L, Brighton L, Mills KH, Zhou H, Jaspers I, Hazucha M. Temporal structure/function variation in cultured differentiated human nasal epithelium associated with acute single exposure to tobacco smoke or E-cigarette vapor. *Inhal Toxicol* 29: 137–144, 2017. doi:10.1080/08958378.2017.1318985.
 99. Herr C, Tsitouras K, Niederstraßer J, Backes C, Beisswenger C, Dong L, Guillot L, Keller A, Bals R. Cigarette smoke and electronic cigarettes differentially activate bronchial epithelial cells. *Respir Res* 21: 67, 2020. doi:10.1186/s12931-020-1317-2.
 100. Ghosh A, Coakley RC, Mascenik T, Rowell TR, Davis ES, Rogers K, Webster MJ, Dang H, Herring LE, Sassano MF, Livraghi-Butrico A, Van Buren SK, Graves LM, Herman MA, Randell SH, Alexis NE, Tarran R. Chronic E-cigarette exposure alters the human bronchial epithelial proteome. *Am J Respir Crit Care Med* 198: 67–76, 2018. doi:10.1164/rccm.201710-2033OC.
 101. Kim H, Kim SR, Je J, Jeong K, Kim S, Kim HJ, Chang KC, Park SW. The proximal tubular $\alpha 7$ nicotinic acetylcholine receptor attenuates ischemic acute kidney injury through Akt/PKC signaling-mediated HO-1 induction. *Exp Mol Med* 50: 1–17, 2018. doi:10.1038/s12276-018-0061-x.
 102. Zhang C, Ding X-P, Zhao Q-N, Yang X-J, An S-M, Wang H, Xu L, Zhu L, Chen H-Z. Role of $\alpha 7$ -nicotinic acetylcholine receptor in nicotine-induced invasion and epithelial-to-mesenchymal transition in human non-small cell lung cancer cells. *Oncotarget* 7: 59199–59208, 2016. doi:10.18632/oncotarget.10498.
 103. Kim H, Zamel R, Bai XH, Liu M. PKC activation induces inflammatory response and cell death in human bronchial epithelial cells. *PLoS One* 8: e64182, 2013. doi:10.1371/journal.pone.0064182.
 104. Rasmussen JE, Sheridan JT, Polk W, Davies CM, Tarran R. Cigarette smoke-induced Ca^{2+} release leads to cystic fibrosis transmembrane conductance regulator (CFTR) dysfunction. *J Biol Chem* 289: 7671–7681, 2014. doi:10.1074/jbc.M113.545137.
 105. Patel W, Moore PJ, Sassano MF, Lopes-Pacheco M, Aleksandrov AA, Amaral MD, Tarran R, Gray MA. Increases in cytosolic Ca^{2+} induce dynamin- and calcineurin-dependent internalisation of CFTR. *Cell Mol Life Sci* 76: 977–994, 2019. doi:10.1007/s00018-018-2989-3.
 106. Herrington JS, Myers C. Electronic cigarette solutions and resultant aerosol profiles. *J Chromatogr A* 1418: 192–199, 2015. doi:10.1016/j.chroma.2015.09.034.
 107. Bartoszewski R, Matalon S, Collawn JF. Ion channels of the lung and their role in disease pathogenesis. *Am J Physiol Lung Cell Mol Physiol* 313: L859–L872, 2017. doi:10.1152/ajplung.00285.2017.
 108. Prakash YS, Pabelick CM, Sieck GC. Mitochondrial dysfunction in airway disease. *Chest* 152: 618–626, 2017. doi:10.1016/j.chest.2017.03.020.

109. **Hoenderdos K, Condcliffe A.** The neutrophil in chronic obstructive pulmonary disease. *Am J Respir Cell Mol Biol* 48: 531–539, 2013. doi:[10.1165/rcmb.2012-0492TR](https://doi.org/10.1165/rcmb.2012-0492TR).
110. **Hikichi M, Mizumura K, Maruoka S, Gon Y.** Pathogenesis of chronic obstructive pulmonary disease (COPD) induced by cigarette smoke. *J Thorac Dis* 11: S2129–S2140, 2019. doi:[10.21037/jtd.2019.10.43](https://doi.org/10.21037/jtd.2019.10.43).
111. **Sapey E, Stockley JA, Greenwood H, Ahmad A, Bayley D, Lord JM, Insall RH, Stockley RA.** Behavioral and structural differences in migrating peripheral neutrophils from patients with chronic obstructive pulmonary disease. *Am J Respir Crit Care Med* 183: 1176–1186, 2011. doi:[10.1164/rccm.201008-1285OC](https://doi.org/10.1164/rccm.201008-1285OC).
112. **Damia Ade D, Gimeno JC, Ferrer MJ, Fabregas ML, Folch PA, Paya JM.** A study of the effect of proinflammatory cytokines on the epithelial cells of smokers, with or without COPD. *Arch Bronconeumol* 47: 447–453, 2011. doi:[10.1016/j.arbres.2011.04.007](https://doi.org/10.1016/j.arbres.2011.04.007).