

‘Snacktivity™’ to increase physical activity

Snacktivity Study Team

DOI:

[10.1016/j.ypmed.2021.106851](https://doi.org/10.1016/j.ypmed.2021.106851)

License:

Creative Commons: Attribution-NonCommercial-NoDerivs (CC BY-NC-ND)

Document Version

Peer reviewed version

Citation for published version (Harvard):

Snacktivity Study Team 2021, 'Snacktivity™' to increase physical activity: time to try something different?', *Preventive Medicine*, vol. 153, 106851. <https://doi.org/10.1016/j.ypmed.2021.106851>

[Link to publication on Research at Birmingham portal](#)

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

TITLE PAGE

Title

‘Snacktivity™ ’ to increase physical activity: Time to try something different?

Authors Names:

James P Sanders PhD^{1,2}, Stuart JH Biddle PhD ⁵, Kajal Gokal PhD ^{1,2}, Lauren B Sherar PhD ^{1,2}, Magdalena Skrybant PhD ^{3,4}, Helen M Parretti PhD⁶, Natalie Ives PhD ⁷, Thomas Yates PhD ^{8,9}, Nanette Mutrie PhD ¹⁰ & Amanda J Daley PhD ^{1,2} on behalf of the Snacktivity Study Team*

Affiliations

¹ National Centre for Sport and Exercise Medicine, School of Sport, Exercise and Health Sciences, Loughborough University, Loughborough, Leicestershire, UK

² Centre for Lifestyle Medicine and Behaviour, School of Sport, Exercise and Health Sciences, Loughborough University, Loughborough, Leicestershire, UK

³ Institute of Applied Health Research, University of Birmingham, Edgbaston, Birmingham, UK

⁴ NIHR Applied Research Collaboration (ARC) West Midlands, University of Warwick, Coventry, UK

⁵ Physically Active Lifestyles Research Group (USQ-PALs), Centre for Health Research, University of Southern Queensland, Springfield, QLD 4300, Australia

⁶ Norwich Medical School, University of East Anglia, Norwich, UK

⁷ Birmingham Clinical Trials Unit, Institute of Applied Health Research, University of Birmingham, Edgbaston, Birmingham, UK

⁸ Diabetes Research Centre, University of Leicester, Leicester, UK

⁹ NIHR Leicester Biomedical Research Centre, Leicester, UK

¹⁰ Physical Activity for Health Research Centre, Institute for Sport, Physical Education and Health Sciences, University of Edinburgh, Edinburgh, UK

Corresponding Author

Professor Amanda Daley

The Centre for Lifestyle Medicine and Behaviour (CLiMB)

National Centre for Sport and Exercise Medicine

School of Sport, Exercise and Health Sciences,

31 Loughborough University
32 Loughborough
33 Leicestershire, LE11 3TU, UK
34 a.daley@lboro.ac.uk
35
36 Abstract Word Count: 250
37 Main Text Word Count: 1679
38
39 .

Abstract

Evidence demonstrates that participation in regular physical activity (PA) reduces the risk of morbidity and mortality. However, current PA guidelines are focused on weekly accumulation of 150 minutes of moderate intensity PA as a threshold. Although recent developments of this guidance have discussed the merits of short bouts of physical activity, guidance that sets large behavioural goals for PA has not been successful in supporting the public to become sufficiently physically active and a ‘one-size fits all’ approach to PA guidelines may not be optimal. A complementary ‘whole day’ approach to PA promotion (i.e. incorporating PA throughout the day) that could motivate the population to be more physically active, is a concept we have called ‘Snacktivity™’. The Snacktivity™ approach promotes small or ‘bite’ size bouts (e.g. 2-5 minutes) of PA accumulated throughout the whole day. Snacktivity™ is consistent with the small change approach which suggest that behaviour change and habit formation are best achieved through gradual building of task self-efficacy, celebrating small successes. Snacktivity™ also offers opportunities to “piggyback” on to existing behaviours/habits, using them as prompts for Snacktivity™. Moreover, small behaviour changes are easier to initiate and maintain than larger ones. A plethora of evidence supports the hypothesis that Snacktivity may be a more acceptable and effective way to help the public reach, or exceed current PA guidelines. This paper outlines the evidence to support the Snacktivity™ approach and the mechanisms by which it may increase population levels of physical activity. Future research directions for Snacktivity™ are also outlined.

1 Introduction

2 Despite unequivocal evidence that physical activity (PA) reduces morbidity and mortality, many adults do not
 3 meet the accepted PA guidelines to perform at least 150-minutes of moderate-intensity PA/week¹. This
 4 guidance has historically been promoted as 30 minutes of moderate-to-vigorous intensity physical activity PA
 5 (MVPA) on at least five days/week. Furthermore, adults are also advised to undertake PA that improves muscle
 6 strength on at least two days per week. However, despite PA guidance having existed for many years,
 7 population-levels of PA remain low. Based on accelerometer-measured PA, and with the removal from the
 8 guidelines of the necessity to accumulate PA in bouts lasting 10 mins or more, approximately 45-95% of the
 9 population are meeting PA guidelines depending on the approach to assessment and analysis²⁻⁴ however, this
 10 still means that approximately half of the population may be inactive. Of particular concern are data suggesting
 11 approximately 1-16% of adults participate in strength-based PA each/week⁵⁻⁷. Collectively, this suggests a
 12 need to consider more innovative, and translational guidance messaging to encourage the population to regularly
 13 engage in PA. Guidelines themselves do not change health behaviour, it is having the means and motivation to
 14 achieve them that matters.

15
 16 One of the most critical obstacles to meeting PA guidance is it requires inactive populations to make significant
 17 lifestyle changes to achieve at least 150 MVPA minutes/week. Previous PA interventions have only had modest
 18 effects on initiation of PA behaviour, and we know very little about successful behavioural maintenance⁸. There
 19 is also concern about the amount of time the public spend in sedentary behaviours, with adults spending
 20 approximately 60-70% of waking hours sedentary⁹. For inactive adults, high levels of sedentary time have been
 21 associated with disease and all-cause mortality¹⁰. These data are of concern and there is no reason to assume this
 22 situation will improve unless effective strategies are put in place to address the problem. It is clear that guidance
 23 setting large goals for PA has not been successful in supporting those who are inactive, to become sufficiently
 24 physically active and current approaches to PA guidelines may therefore not be optimal. A shift in emphasis in
 25 facilitating PA behaviour is required.

27 Every minute counts: Snacktivity™ to promote physical activity

28 Updated guidelines from health agencies in 2019/20, has removed the need to complete PA in bouts of 10 mins
 29 or more^{1,11,12}. Although revised guidance now recognises the importance of making small changes to PA
 30 behaviour and that any PA is better than none^{1,11,12}, guidance still focuses on the public needing to achieve a
 31 behavioural goal of at least 150 MVPA minutes/week, which can be a daunting task for inactive populations^{1,12}.
 32 Additional or complementary strategies are needed to assist the public in becoming more physically active. A
 33 complimentary ‘whole day’ approach to PA promotion that seeks to motivate and support individuals to be more
 34 physically active throughout the day, is a concept we refer to as ‘Snacktivity™’. Rather than broadly
 35 encouraging at least 150 MVPA minutes/week, Snacktivity™ focuses on promoting small (e.g., 2-5 minutes),
 36 but frequent, bouts of MVPA throughout each day, to accumulate at least 150 MVPA minutes/week. For
 37 example six ‘activity snacks’ lasting five minutes/day would be required to meet the PA recommendations.
 38 Examples of Snacktivity™ include walk-talk conversations, using stairs rather than the lift/elevator, pacing
 39 whilst using the telephone, or squats while brushing teeth (**Error! Reference source not found.**)).



Figure 1 - Snacktivity ideas

What is the evidence that Snacktivity™ can improve health?

Evidence demonstrates an inverse dose-response relationship between PA and all-cause mortality. This means that for inactive people, any increase in PA is beneficial for health¹³. The relationship is also characterised by a steep early slope meaning the greatest gains in health are experienced with inactive people doing a little more activity per week (e.g. 2-3 MET/hrs per/week), rather than by encouraging those who are already physically active to do marginally more¹⁴. Improved cardio-metabolic health and aerobic fitness have been reported following brief bouts of PA^{15,16}, this association remained consistent across subjective and device based measures, with studies reporting no difference in cardiovascular fitness improvement between accumulated and continuous bouts of PA of the same total duration¹⁷. Furthermore, whilst evidence is limited, some research suggests that short bouts of PA can also reduce psychological parameters including stress, depressive symptoms, and improve self-esteem in adults^{18,19}, with recent data suggesting similar improvements in positive mood state and decrease negative mood state²⁰. This suggests that achieving short(er) bouts of Snacktivity™ throughout the day may achieve similar health benefits to long(er) bouts. Moreover, whilst the idea that small bouts of PA may improved health outcomes is not new, as investigated in experimental studies^{17,21} it is not a message that has been prominent in public guidance, in part, due to a lack of high quality, “real world” evidence.

How might Snacktivity™ work to improve population health?

A common barrier to PA is a perceived lack of time. For inactive adults, long(er) bouts of MVPA may seem difficult to achieve and it may be that many people believe that achieving 150 MVPA minutes/week requires too much cognitive effort, planning and physical exertion to be worthwhile. In contrast, Snacktivity™ may be perceived as more achievable because each ‘snack’ requires a small time commitment, and it involves less planning and effort. Snacktivity™ does not require skills, equipment, or a change of clothing, little or no preparation, can be performed in most settings, and easily incorporated into daily life, addressing both convenience and health inequalities²², and allowing for greater population reach. Simple actions may become more habitual than complex ones. This suggests that the integration of Snacktivity™ into usual routines may be a more feasible and appealing approach to sustaining PA behaviour than trying to achieve larger changes²³. One way this may be facilitated it through the use of action planning, to plan the when, where, and how to complete an activity snack within the day (e.g in the morning when I am brushing my teeth I will perform squats, or when I collect the children from school I will walk). This will help reduce the cognitive load of Snacktivity™, whilst utilising an effective behaviour change technique. Moreover, small changes are easier to initiate, and maintain, than large changes which is consistent with the small change approach to behaviour change.²⁴

How people feel about PA is an important predictor of whether they continue to engage with a a behaviour. Snacktivity™ may help to develop confidence among those who awho are inactive by encouraging them to ‘start small’. Psychological theory acknowledges that achieving small changes is important for individuals’ task and self-regulatory self-efficacy and habit formation²⁵. If individuals complete activity snacks, this should increase their self-efficacy for engagement, making them more likely to continue. Snacktivity™ might then be the gateway for more sustained participation in physical activity. Snacktivity™ may be particularly appropriate

for specific populations, such as the elderly and people with chronic diseases/disabilities, who may find it difficult to engage in PA.

An important component of PA guidance is that adults should undertake muscle-strengthening activity on at least two days/week. It is important that people perform strength-based activities, given its association with the risk of falls, fractures and osteoporosis¹². Snacktivity™ provides an opportunity to promote this message since many muscle-strengthening activities lend themselves to Snacktivity™ (e.g. When I am boiling the kettle, I will perform 20 squats) as they are traditionally shorter, stationery, and do not necessarily need special equipment or clothing.

A further benefit of Snacktivity™ is that it encourages PA while simultaneously breaking up prolonged periods of sedentary behaviour throughout the day. Snacktivity™ may therefore provide two health outcomes in a ‘buy one, get one free’ scenario, increasing the probability of cost-effectiveness. To support this hypothesis, experimental trials have shown that breaking up sedentary behaviour with periods of MVPA provides favourable changes in individuals’ cardio-metabolic risk²⁶.

Issues to consider in promoting Snacktivity™

While there may be advantages to Snacktivity™, there are also some issues to consider. Snacktivity™ may be disruptive to the day and easily forgotten. It might be difficult for the public to achieve MVPA in ‘bite sizes’, or difficult to think of ways in which to implement Snacktivity™ into daily life. Indeed, incorporating activity snacks into the home or work life, may require a change in social norms (e.g. making it socially acceptable to leave one’s desk to perform an activity snack).

Simply giving people information does not lead to sustained behaviour change. Additional strategies to encourage the public to engage in Snacktivity™ will be required, and a wide range of technologies are now available to facilitate this process. Consideration needs to be given to whether an accumulated Snacktivity™ approach means the public to consider PA too many times in a day/week, and whether this then requires too much cognitive energy to enact. Given this, Snacktivity™ may not be flexible enough and/or convenient to the public. Whilst all PA is important for health, greater intensity PA provides more benefit for the same amount of time, particularly for non-communicable diseases¹. However, Snacktivity™ may encourage participation in predominately light-intensity PA, without progressing towards sufficient MVPA leading to Snacktivity™ having a smaller impact on health. Furthermore, although Snacktivity™ encourages participants to break up sedentary behaviour with MVPA, this relies on activating two motivational systems, which may limit engagement from the public.

Finally, short bouts might be useful in increasing PA, no randomised controlled trial has directly tested if Snacktivity™ derives the same health benefits as current PA guidelines; we are currently gathering such evidence (<https://fundingawards.nihr.ac.uk/award/RP-PG-0618-20008>).

Conclusion

1 Given the lack of success in encouraging inactive populations to achieve large(r) bouts of PA. Snacktivity™
2 may be a complementary public health message that offers a method of implementing this guidance. It is not
3 suggested that current PA guidance should be abandoned, but current approaches may not be ideal.
4 Snacktivity™ should be achievable by most of the population and therefore addresses health inequalities
5 making it accessible to all of those who might benefit. Snacking is a common behaviour and for the first time,
6 the public could be encouraged to snack as much as they like, just not with unhealthy foods, but with
7 Snacktivity™.

Author contributions

AD developed the original Snacktivity idea and JS conceived the idea for this report. AD and JS wrote the initial draft for this report and SJHB, LS, KG, MS, HMP, NI, TY, NM and Snacktivity Investigators contributed at a later stage.

*Snacktivity study team includes Stuart Biddle, Amanda J Daley, Dale Esliger, Charlotte Edwardson, Emma Frew, Kajal Gokal, Colin Greaves Sheila M Greenfield Natalie Ives, Kate Jolly, Ralph Maddison, Nanette Mutrie, Helen M Parretti, James Sanders, Lauren Sherar, Magdalena Skrybant, and Tom Yates.

Conflicts of interest/Competing interests (include appropriate disclosures)

This work was supported by the National Institute for Health Research (NIHR). AJD is supported by a National Institute for Health Research (NIHR) Research Professorship award. This publication presents independent research funded by the NIHR. The views expressed are those of the author(s) and not necessarily those of the NHS, the NIHR or the Department of Health and Social Care.

Ethical Compliance

No ethical approval was sought as this is a commentary and no data was collected or analysed by the authorship team in the production of this article.

Financial Disclosures

No financial disclosures were reported by the authors of this paper

REFERENCES

1. UK Chief Medical Officer. UK Chief Medical Officers' Physical Activity Guidelines. Department of Health and Social Care. <https://www.gov.uk/government/publications/physical-activity-guidelines-uk-chief-medical-officers-report>. Published 2019. Accessed December 4, 2020.
2. Rowlands A V., Sherar LB, Fairclough SJ, et al. A data-driven, meaningful, easy to interpret, standardised accelerometer outcome variable for global surveillance. *J Sci Med Sport*. 2019;22(10):1132-1138. doi:10.1016/j.jsams.2019.06.016
3. Taylor A, Taylor RS, Ingram W, et al. Randomised controlled trial of an augmented exercise referral scheme using web-based behavioural support for inactive adults with chronic health conditions: the e-coachER trial. *Br J Sports Med*. 2021;55(8):444-450. doi:10.1136/BJSPORTS-2020-103121
4. Zenko Z, Willis EA, White DA. Proportion of Adults Meeting the 2018 Physical Activity Guidelines for Americans According to Accelerometers. *Front Public Heal*. 2019;0(JUN):135. doi:10.3389/FPUBH.2019.00135
5. NHS. Health Survey for England 2016 Physical activity in adults. NHS Digital. <https://digital.nhs.uk/data-and-information/publications/statistical/health-survey-for-england/health-survey-for-england-2016>. Published 2017. Accessed November 21, 2020.
6. Dankel SJ, Loenneke JP, Loprinzi PD. Determining the Importance of Meeting Muscle-Strengthening Activity Guidelines: Is the Behavior or the Outcome of the Behavior (Strength) a More Important Determinant of All-Cause Mortality? *Mayo Clin Proc*. 2016;91(2):166-174. doi:10.1016/J.MAYOCP.2015.10.017
7. Kraschewski JL, Sciamanna CN, Ciccolo JT, et al. Is exercise used as medicine? Association of meeting strength training guidelines and functional limitations among older US adults. *Prev Med (Baltim)*. 2014;66:1-5. doi:10.1016/J.YPMED.2014.05.012
8. King AC, Whitt-Glover MC, Marquez DX, et al. Physical Activity Promotion: Highlights from the 2018 Physical Activity Guidelines Advisory Committee Systematic Review. *Med Sci Sports Exerc*. 2019;51(6):1340-1353. doi:10.1249/MSS.0000000000001945
9. Biswas A, Oh PI, Faulkner GE, et al. Sedentary time and its association with risk for disease incidence, mortality, and hospitalization in adults. *Ann Intern Med*. 2015;162(2):123. doi:10.7326/M14-1651
10. Ekelund U, Brown WJ, Steene-Johannessen J, et al. Do the associations of sedentary behaviour with cardiovascular disease mortality and cancer mortality differ by physical activity level? A systematic

- review and harmonised meta-analysis of data from 850 060 participants. *Br J Sports Med.* 2019;53(14):886-894. doi:10.1136/bjsports-2017-098963
11. Bull FC, Al-Ansari SS, Biddle S, et al. World Health Organization 2020 guidelines on physical activity and sedentary behaviour. *Br J Sports Med.* 2020;54(24):1451-1462.
 12. 2018 Physical Activity Guidelines Advisory Committee. *2018 Physical Activity Guidelines Advisory Committee Scientific Report.*; 2018. doi:10.1115/1.802878.ch1
 13. Arem H, Moore SC, Patel A, et al. Leisure time physical activity and mortality: A detailed pooled analysis of the dose-response relationship. *JAMA Intern Med.* 2015;175(6):959-967. doi:10.1001/jamainternmed.2015.0533
 14. Ekelund U, Tarp J, Steene-Johannessen J, et al. Dose-response associations between accelerometry measured physical activity and sedentary time and all cause mortality: Systematic review and harmonised meta-analysis. *BMJ.* 2019;366. doi:10.1136/bmj.l4570
 15. Hupin D, Roche F, Gremeaux V, et al. Even a low-dose of moderate-to-vigorous physical activity reduces mortality by 22% in adults aged ≥ 60 years: A systematic review and meta-analysis. *Br J Sports Med.* 2015;49(19):1262-1267. doi:10.1136/bjsports-2014-094306
 16. Jakicic JM, Kraus WE, Powell KE, et al. Association between Bout Duration of Physical Activity and Health: Systematic Review. *Med Sci Sports Exerc.* 2019;51(6):1213. doi:10.1249/MSS.0000000000001933
 17. Murphy MH, Lahart I, Carlin A, Murtagh E. The effects of continuous compared to accumulated exercise on health: A meta-analytic review. *Sport Med.* 2019;49:1585-1607. doi:10.1007/s40279-019-01145-2
 18. Teychenne M, White RL, Richards J, Schuch FB, Rosenbaum S, Bennie JA. Do we need physical activity guidelines for mental health: What does the evidence tell us? *Ment Health Phys Act.* 2020;18:100315. doi:10.1016/J.MHPA.2019.100315
 19. Barr-Anderson DJ, Auyoung M, Whitt-Glover MC, Glenn BA, Yancey AK. Integration of Short Bouts of Physical Activity Into Organizational Routine: A Systematic Review of the Literature. *Am J Prev Med.* 2011;40(1):76-93. doi:10.1016/J.AMEPRE.2010.09.033
 20. Stenling A, Moylan A, Fulton E, Machado L. Effects of a Brief Stair-Climbing Intervention on Cognitive Performance and Mood States in Healthy Young Adults. *Front Psychol.* 2019;0(OCT):2300. doi:10.3389/FPSYG.2019.02300

21. Boreham CAG, Kennedy RA, Murphy MH, Tully M, Wallace WFM, Young I. Training effects of short bouts of stair climbing on cardiorespiratory fitness, blood lipids, and homocysteine in sedentary young women. *Br J Sports Med.* 2005;39(9):590-593. doi:10.1136/bjsm.2002.001131
22. Reichert FF, Barros AJD, Domingues MR, Hallal PC. The role of perceived personal barriers to engagement in leisure-time physical activity. *Am J Public Health.* 2007;97(3):515-519. doi:10.2105/AJPH.2005.070144
23. Hill JO. Can a small-changes approach help address the obesity epidemic? A report of the joint task force of the american society for nutrition, institute of food technologists, and international food information council. *Am J Clin Nutr.* 2009;89(2):477-484. doi:10.3945/ajcn.2008.26566
24. Fogg B. *Tiny Habits: The Small Changes That Change Everything.* Houghton Mifflin Harcourt; 2019.
25. Gardner B, Rebar A., Lally P. Habit Interventions. In: *The Handbook of Behavior Change.* Cambridge: Cambridge University Press; 2020:599-616.
26. Frith E, Loprinzi PD. Accelerometer-assessed light-intensity physical activity and mortality among those with mobility limitations. *Disabil Health J.* 2018;11(2):298-300. doi:10.1016/j.dhjo.2017.08.007