# **UNIVERSITY** OF BIRMINGHAM University of Birmingham Research at Birmingham

# A new maximal subgroup of E<sub>8</sub> in characteristic 3

Craven, David; Stewart, David; Thomas, Adam

License: Creative Commons: Attribution-NonCommercial-NoDerivs (CC BY-NC-ND)

Document Version Peer reviewed version

Citation for published version (Harvard):

Craven, D, Stewart, D & Thomas, A 2022, 'A new maximal subgroup of E in characteristic 3', Proceedings of the American Mathematical Society, vol. 150, no. 4, pp. 1435–1448. <https://www.ams.org/journals/proc/2022-150-04/S0002-9939-2022-15759-1/home.html>

Link to publication on Research at Birmingham portal

#### **Publisher Rights Statement:**

First published in Proceedings of the American Mathematical Society in Volume 150, Number 4, April 2022, Pages 1435–1448, published by the American Mathematical Society. © 2022 American Mathematical Society

#### **General rights**

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

•Users may freely distribute the URL that is used to identify this publication.

•Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research. •User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)

•Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

#### Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

# A New Maximal Subgroup of $E_8$ in Characteristic 3

David A. Craven, David I. Stewart and Adam R. Thomas

### March 22, 2021

#### Abstract

We prove the existence and uniqueness of a new maximal subgroup of the algebraic group of type  $E_8$  in characteristic 3. This has type  $F_4$ , and was missing from previous lists of maximal subgroups produced by Seitz and Liebeck–Seitz. We also prove a result about the finite group  $H = {}^{3}D_4(2)$ , that if H embeds in  $E_8$  (in any characteristic p) and has two composition factors on the adjoint module then p = 3 and H lies in this new maximal  $F_4$  subgroup.

# 1 Introduction

The classification of the maximal subgroups of positive dimension of exceptional algebraic groups [13] is a cornerstone of group theory. In the course of understanding subgroups of the finite groups  $E_8(q)$  in [3], the first author ran into a configuration that should not occur according to the tables in [13].

We elicit a previously undiscovered maximal subgroup of type  $F_4$  of the algebraic group  $E_8$  over an algebraically closed field of characteristic 3. This discovery corrects the tables in [13], and the original source [17] on which it depends.

**Theorem 1.1.** Let **G** be a simple algebraic group of type  $E_8$  over an algebraically closed field of characteristic 3. Then **G** contains a unique conjugacy class of simple maximal subgroups of type  $F_4$ .

If **X** is in this class, then the restriction of the adjoint module  $L(E_8)$  to **X** is isomorphic to  $L_{\mathbf{X}}(1000) \oplus L_{\mathbf{X}}(0010)$ , where the first factor is the adjoint module for **X** of dimension 52 and the second is a simple module of dimension 196 for **X**.

The classification from [13] states that the maximal subgroups are maximal-rank or parabolic subgroups, or one of a short list of reductive subgroups that exist for all but a few small primes, together with  $G_2$ inside  $F_4$  for p = 7. This last case arises from a generic embedding of  $G_2$  in  $E_6$ , which falls into  $F_4$ on reduction modulo the prime p = 7 only. This new  $E_8$  subgroup is therefore the only example of a maximal subgroup that exists for a single prime, whose embedding cannot be explained using generic phenomena.

The structure of this note is as follows. The existence and uniqueness up to conjugacy of a maximal Lie subalgebra  $\mathfrak{f}_4 \subset \mathfrak{e}_8$  is first established. From this we are able to write down the root elements of  $\mathfrak{f}_4$ ; exponentiation gives expressions for the root groups of  $\mathbf{X}$  in terms of the root groups of  $E_8$ , providing an explicit construction of  $\mathbf{X}$ .

In the final section we determine various results providing extra details on this new class of maximal subgroups. For each unipotent class in  $\mathbf{X}$  we determine the corresponding unipotent class in  $E_8$  that contains it, and we do the same for nilpotent orbits of the corresponding Lie algebras. We also consider the maximal connected subgroups of  $\mathbf{X}$ . The maximal parabolic subgroups of  $\mathbf{X}$  will be contained in parabolic subgroups of  $E_8$  by the Borel–Tits Theorem. There are four classes of reductive maximal connected subgroups of  $\mathbf{X}$  when p = 3 with types  $B_4, A_1C_3, A_1G_2, A_2A_2$ . We show that all of these classes are contained in other maximal connected subgroups of  $E_8$  and we specify such an overgroup. Moreover, we determine that the first three classes are  $E_8$ -irreducible but the last class is not. (A subgroup is  $\mathbf{G}$ -irreducible if it is not contained in any proper parabolic subgroup of  $\mathbf{G}$ .)

In establishing the existence of  $\mathbf{X}$ , we prove the following extra result, of use in the project to classify maximal subgroups of the finite exceptional groups of Lie type.

**Proposition 1.2.** Let H be the group  ${}^{3}D_{4}(2)$ , let p be a prime, and suppose that H embeds in the algebraic group  $E_{8}$  in characteristic p. If the composition factors of the action of H on the adjoint module  $L(E_{8})$  have dimensions 52 and 196, then p = 3 and H is contained in a maximal subgroup  $\mathbf{X}$  of type  $F_{4}$ ; furthermore, H and  $\mathbf{X}$  stabilize the same subspaces of  $L(E_{8})$ .

# **2** From the Thompson group to $F_4$

One path to a construction of the  $F_4$  subgroup of  $E_8$  starts with the Thompson group, which contains a copy of  $H \cong {}^{3}D_4(2)$ , acting on  $L(E_8)$  with composition factors of dimensions 52 and 196. In fact, every H-invariant alternating bilinear form on the 248-dimensional module is invariant under a suitable copy of  $F_4 \leq \text{GL}_{248}(k)$ .

We cannot quite show this without a computer. Splitting  $L(E_8)$  up as the sum of 52 and 196 fragments the space of alternating forms into six components. For five of these six we can show that the *H*-invariant maps are  $F_4$ -invariant, but for the sixth we cannot do so without a computer. With a computer we can check that this sixth component is at least  $F_4(9)$ -invariant, and thus every subgroup *H* of  $E_8$  is contained in a copy of  $F_4(9)$ . But  $F_4(9)$  contains elements of order 6562, and thus there is an  $F_4$  subgroup of  $E_8$ containing it, via [12, Proposition 2].

We then show that this  $F_4$  subgroup is unique up to  $E_8$ -conjugacy, obtaining as a by-product that H is unique up to  $E_8$ -conjugacy.

We start with a copy J of the Thompson sporadic simple group. This has a 248-dimensional simple module M over  $\mathbb{C}$ , and it remains simple upon reduction modulo all primes. From [1, p.176], we see that there are elements of order 9 with Brauer character value 5 on M. The only integers that are the traces of semisimple elements of order 9 in  $E_8$  (on the adjoint module) are -1, 2, 8 and 29. Thus these elements cannot be semisimple, and in particular,  $p \mid 9$ . Thus we see that if J embeds in the algebraic group  $E_8$  in any characteristic p, then p = 3. It is a famous result that J does indeed embed in  $E_8(3)$ , and is unique up to conjugacy.

It is well known that J contains a subgroup H isomorphic to  ${}^{3}D_{4}(2)$ . From [1, p.90] and [6, pp.251–253], we see that in characteristic not 2, the restriction of M to H is the direct sum of a 52-dimensional simple module  $M_1$  and a 196-dimensional simple module  $M_2$ . However, all elements of order 9 in H act on  $M_1 \oplus M_2$  with trace 2, so we cannot use the previous method to show that H cannot embed in  $E_8$  in characteristic  $p \neq 2, 3$  acting on  $L(E_8)$  as  $M_1 \oplus M_2$ .

**Lemma 2.1.** Let p be an odd prime, let H denote the group  ${}^{3}D_{4}(2)$ , and suppose that H embeds in the algebraic group  $E_{8}$  in characteristic p, acting on the adjoint module with composition factors of dimensions 52 and 196. Then the submodule 52 carries the structure of a Lie algebra of type  $\mathfrak{f}_{4}$ , and in particular p = 3. Furthermore, such an  $\mathfrak{f}_{4}$  subalgebra of  $\mathfrak{e}_{8}$  does exist for p = 3.

*Proof.* Let  $M_1$  denote the 52-dimensional kH-submodule of the adjoint and  $M_2$  the 196-dimensional submodule. Note that  $|H| = 2^{12} \cdot 3^4 \cdot 7^2 \cdot 13$ , so either  $p \nmid |H|$  and we are essentially in characteristic 0, or p = 7, 13, or p = 3.

From [4], there is a unique conjugacy class of subgroups H in  $F_4$  for any odd characteristic p, acting irreducibly on the minimal and adjoint modules. In particular, we see from [4] that  $\operatorname{Hom}_{kH}(\Lambda^2(M_1), M_1)$ is 1-dimensional for all odd primes p. From an ordinary character calculation, we see that

$$\Lambda^2(\chi_{52}) = \chi_{52} + \chi_{1274},$$

where  $\chi_i$  is the irreducible character of degree *i*. For both 7 and 13, the reduction modulo *p* of  $\chi_{1274}$  is irreducible, hence the reduction modulo *p* is irreducible modulo *p* for all *p* > 3 (see [6, pp.252–253]). If *p* = 3 then  $\chi_{1274}$  has Brauer character constituents of degrees 52 and 1222 from [6, p.251]. Since

 $\operatorname{Hom}_{kH}(\Lambda^2(M_1), M_1) = k$ , we see that the exterior square is uniserial, with layers of dimensions 52, 1222 and 52. Moreover,

$$\operatorname{Hom}_{kH}(\Lambda^2(M_1), M_2) = 0.$$

Thus  $M_1$  forms an H-invariant subalgebra, which must be non-zero, since  $\mathfrak{e}_8$  contains no abelian subspace of dimension 52 by [5, Proposition 2.3]. Furthermore, as  $M_1$  is irreducible for H, the restriction of the Lie bracket to  $M_1$  furnishes it with the structure of a semisimple Lie algebra. Since  $\operatorname{Hom}_{kH}(\Lambda^2(M_1), M_1) = k$ , there is at most one isomorphism class of such, but as the algebraic k-group  $F_4$  does contain a subgroup isomorphic to H, acting as  $\chi_{52}$  on its adjoint module, it follows that  $M_1 \cong \mathfrak{f}_4$ . In particular, this means that the  $f_4$  Lie algebra must have a simple module of dimension 196, as it acts *H*-equivariantly on  $L(E_8)$ .

Such a simple module must be restricted: if not the p-closure  $L_p$  of the image L of  $\mathfrak{f}_4$  in  $\mathfrak{e}_8$  will contain a centre [19, 2.5.8(2)]; but L has no 1-dimensional submodules on M. By Curtis's theorem,  $M_2$  arises by differentiation of a restricted representation for the algebraic group  $F_4$ , whence p = 3, from the tables in [14].

The embedding and subalgebra do exist for p = 3 via the Thompson group, as seen above. 

We will prove that the  $f_4$ -subalgebra is the Lie algebra of an  $F_4$  algebraic subgroup of  $E_8$ . To do so, we will actually prove that every H-invariant alternating product on the kH-module  $M = M_1 \oplus M_2$  for p=3 is also  $F_4$ -invariant, for the unique  $F_4 \leq GL_{52}(k)$  containing H. To do so, we need to understand the space

$$\operatorname{Hom}_{kH}(\Lambda^2(M), M)$$

of alternating products on M. Using  $M = M_1 \oplus M_2$ , and the formula

$$\Lambda^2(A \oplus B) \cong \Lambda^2(A) \oplus \Lambda^2(B) \oplus A \otimes B$$

we split the space of products up into six components. The next result gives the dimensions of these components.

**Proposition 2.2.** We have

. . 9 . . . .

*Proof.* One may use a computer to check these with ease. Some may be checked easily by hand as well, using the ordinary character table and the 3-decomposition matrix for H. For example, using those two tables,  $M_1 \otimes M_2$  does not possess a composition factor  $M_1$ , and thus

$$\operatorname{Hom}_{kH}(M_1 \otimes M_2, M_1) = \operatorname{Hom}_{kH}(\Lambda^2(M_1), M_2) = 0.$$

The statement that  $\operatorname{Hom}_{kH}(\Lambda^2(M_1), M_1) = k$  appears in [4, Section 4.3.4] (where it is proved by computer). 

At least the existence of two of the three remaining maps is clear from the fact that H embeds in  $E_8(3)$ with representation  $M_1 \oplus M_2$ . If  $\operatorname{Hom}_{kH}(M_1 \otimes M_2, M_2) = 0$  then  $M_1$  would be an ideal of the Lie algebra, which is not possible. A character calculation shows that  $S^2(M_2)$  does not have a composition factor  $M_1$ , so

$$\operatorname{Hom}_{kH}(\Lambda^2(M_2), M_1) = \operatorname{Hom}_{kH}(M_1 \otimes M_2, M_2).$$

It is only  $\operatorname{Hom}_{kH}(\Lambda^2(M_2), M_2)$  that cannot easily be seen. Indeed, this space will cause us a problem later on.

We now prove that five of the six Hom-spaces extend to the algebraic group  $\mathbf{X} = F_4$ , with only  $\operatorname{Hom}_{kH}(\Lambda^2(M_2), M_2)$  missing. If one is happy to use a computer for all of this, one simply checks that all H-invariant maps are  $F_4(3)$ - and even  $F_4(9)$ -invariant, and thus one does not need to prove the next proposition.

**Proposition 2.3.** Let  $\mathbf{X}$  be an algebraic k-group of type  $F_4$ . We have

$$\operatorname{Hom}_{\mathbf{X}}(\Lambda^2(L(1000)), L(1000)) = k$$

 $\operatorname{Hom}_{\mathbf{X}}(\Lambda^{2}(L(1000)), L(0010)) = \operatorname{Hom}_{\mathbf{X}}(L(1000) \otimes L(0010), L(1000)) = 0, \text{ and} \\ \operatorname{Hom}_{\mathbf{X}}(\Lambda^{2}(L(0010)), L(1000)) = \operatorname{Hom}_{\mathbf{X}}(L(1000) \otimes L(0010), L(0010)) = k.$ 

*Proof.* Note that each of these spaces must have dimension at most the dimension of the corresponding space for H. This yields the two 0-dimensional spaces, and that the other spaces have dimension at most 1. The first statement holds because **X** is a non-abelian Lie algebra, and therefore the space is non-zero.

For the last statement, since  $S^2(M_2)$  has no composition factor isomorphic to  $M_1$ , certainly  $S^2(L(0010))$  has no composition factor isomorphic to L(1000). Thus the two Hom-spaces are isomorphic, so it remains to find a non-zero map in the latter space.

The composition factors of the kH-module  $M_1 \otimes M_2$  are of dimensions

25, 196, 196, 441, 1963, 2457, 2457, 2457.

The highest-weight module L(1010), which must appear as a composition factor in  $L(1000) \otimes L(0010)$ , has dimension 7371, and must restrict to kH to be the sum of the three (non-isomorphic) modules of dimension 2457. The rest of the composition factors, in total, have dimension 2821, so there must be a composition factor of dimension between 1963 and 2821. Consulting [14, Appendix A.50], we find exactly one such module: L(0011) of dimension 2404 = 1963 + 441. The remaining kH-modules, 25, 196 and 196, must be the other composition factors for  $F_4$ , because  $F_4$  has no simple modules of dimension 25 + 196, 196 + 196, or 25 + 196 + 196.

Thus the composition factors of  $L(1000) \otimes L(0010)$  have dimensions 25, 196, 196, 2404 and 7371. Since L(0010) is the unique module to appear more than once, and the tensor product is self-dual, L(0010) must be a submodule, and the maps in  $\operatorname{Hom}_{kH}(M_1 \otimes M_2, M_2)$  extend to **X**.

The last remaining Hom-space to check is  $\operatorname{Hom}_{\mathbf{X}}(\Lambda^2(L(0010)), L(0010))$ . This seems difficult to do by hand, and we resort to a computer. There are two ways to proceed. The first is to prove that that there is an  $F_4(3)$ -invariant map in the space (this takes a couple of minutes), and thus the H is contained in a copy of  $F_4(3)$  in  $E_8$ . We then apply [3, Proposition 6.8], which states that  $F_4(3)$  is contained in a positive-dimensional subgroup stabilizing the same subspaces of  $L(E_8)$ , which are  $M_1$  and  $M_2$ . This must be a copy of  $F_4$  (as it stabilizes an  $\mathfrak{f}_4$ -Lie subalgebra), and we are done. Alternatively, we prove the same statement for  $F_4(9)$  (which contains elements of order  $9^4 + 1 = 6562$ , this takes about half an hour on one of the first author's computers) and then apply [12, Proposition 2], which yields the same positive-dimensional subgroup.

In either case, we obtain the following.

**Proposition 2.4.** Let p = 3 and let H be a subgroup  ${}^{3}D_{4}(2)$  of  $E_{8}$ , acting on  $L(E_{8})$  with composition factors of dimensions 52 and 196. Then H is contained in a positive-dimensional subgroup of type  $F_{4}$ , stabilizing exactly the same subspaces of  $L(E_{8})$  that are stabilized by H.

It suffices to ascertain the uniqueness up to conjugacy of the hypothesized subgroup of type  $F_4$ , and as a by-product we also obtain uniqueness of H up to conjugacy.

Let t be an involution in  $\mathbf{X} = F_4$  with centralizer  $B_4$ . The trace of t on  $L(E_8)$  is -8, and so the centralizer of t in  $E_8$  is  $D_8$ , which acts with composition factors  $L(\lambda_2)$  and  $L(\lambda_7)$ , of dimensions 120 and 128 respectively. The restriction of the  $k\mathbf{X}$ -module L(1000) to  $B_4$  is  $L(0001) \oplus L(0100)$ , with dimensions 16 and 36 respectively. The restriction of L(0010) to  $B_4$  is  $L(0010) \oplus L(1001)$ , of dimensions 84 and 112 respectively. (This can be checked using weights or quickly on a computer for  $F_4(3)$ .) From [20, Table 60] we see that this  $B_4$  is subgroup  $E_8(\#45)$  and is unique up to conjugacy. But clearly there is a unique way to assemble the numbers 16, 36, 84 and 112 to make 52 and 196. Thus given any subgroup of type  $B_4$  there exists at most one  $F_4$  containing it, which must stabilize the submodule  $L(0001) \oplus L(0100)$ . Thus we obtain the result that  $\mathbf{X}$  is unique up to conjugacy.

This completes the proof of uniqueness of  $\mathbf{X}$ , and thus Theorem 1.1 is proved.

# **3** Presentation of $F_4$

For their application to future explicit computations, we give expressions for the root elements  $x_{\pm\beta_i}(t)$ for  $\beta_i$  a root of the maximal subgroup  $F_4$  as products of root elements of  $E_8$ ; see [2, §4.4] for notation. We also provide the elements  $h_{\beta_i}(t)$  written in terms of those of  $E_8$ . Note that since the coefficients  $t_j$ we exhibit are elements of GF(3), it follows that the subgroup  $\mathbf{Y}$  we produce is defined over GF(3). As the factors of  $x_{\beta_i}(t) = \prod_j x_{\alpha_j}(c_j t)$  commute we get easily that  $x_{-\beta_i}(t) = \prod_j x_{-\alpha_j}(c_j t)$ . For this reason we only list the root group elements for positive roots. Note also that the element  $e_{\beta_i}$  of a Chevalley basis for  $\mathfrak{h}$  is recovered from such an expression as  $\sum c_j e_{\alpha_j}$ .

```
x_{1000}(t) = x_{00010000}(t)x_{00000100}(t)
x_{0100}(t) = x_{00100000}(-t)x_{00000010}(t)
x_{0010}(t) = x_{10000000}(-t)x_{00000001}(t)x_{00011000}(-t)x_{00001100}(-t)
x_{0001}(t) = x_{11110000}(t)x_{01121000}(-t)x_{01111100}(-t)x_{01011110}(t)
x_{1100}(t) = x_{00110000}(-t)x_{00000110}(-t)
x_{0110}(t) = x_{10100000}(t)x_{00000011}(-t)x_{00111000}(-t)x_{00001110}(-t)
x_{0011}(t) = x_{11121000}(t)x_{1111100}(t)x_{01011111}(t)x_{01122100}(-t)
x_{1110}(t) = x_{10110000}(t)x_{00000111}(t)x_{00111100}(-t)x_{00011110}(t)
x_{0120}(t) = x_{10111000}(-t)x_{00001111}(-t)
x_{0111}(t) = x_{11221000}(t)x_{1111110}(t)x_{01111111}(t)x_{01122110}(-t)
x_{1120}(t) = x_{10111100}(-t)x_{00011111}(t)
x_{1111}(t) = x_{11221100}(t)x_{11121110}(-t)x_{01121111}(-t)x_{01122210}(t)
x_{0121}(t) = x_{1111111}(-t)x_{11222100}(-t)x_{11122110}(t)x_{01122111}(t)
x_{1220}(t) = x_{10111110}(t)x_{00111111}(-t)
x_{1121}(t) = x_{11121111}(t)x_{11232100}(t)x_{11122210}(-t)x_{01122211}(-t)
x_{0122}(t) = x_{12232111}(-t)x_{12233210}(t)
x_{1221}(t) = x_{11221111}(-t)x_{11232110}(-t)x_{11222210}(t)x_{01122221}(-t)
x_{1122}(t) = x_{12232211}(t)x_{12243210}(-t)
x_{1231}(t) = x_{11232111}(t)x_{11222211}(-t)x_{11122221}(-t)x_{11233210}(t)
x_{1222}(t) = x_{12232221}(t)x_{12343210}(t)
x_{1232}(t) = x_{22343210}(t)x_{12343211}(t)x_{12243221}(t)x_{1223321}(-t)
x_{1242}(t) = x_{22343211}(t)x_{12244321}(t)
x_{1342}(t) = x_{22343221}(-t)x_{12344321}(t)
x_{2342}(t) = x_{22343321}(t)x_{12354321}(-t)
h_{1000}(t) = h_{00010000}(t)h_{00000100}(t)
h_{0100}(t) = h_{00100000}(t)h_{00000010}(t)
```

 $h_{0010}(t) = h_{10000000}(t)h_{00010000}(t)h_{00001000}(t^2)h_{00000100}(t)h_{00000001}(t)$ 

 $h_{0001}(t) = h_{1000000}(t)h_{01000000}(t^4)h_{00100000}(t^3)h_{00010000}(t^5)h_{00001000}(t^3)h_{00000100}(t^2)h_{00000010}(t)$ 

**Remark 3.1.** We note that the 24-dimensional unipotent subgroup generated by the positive root elements of  $F_4$  is contained in the 120-dimensional unipotent subgroup generated by the positive roots of  $E_8$ . The given presentation has the slightly unfortunate property that the structure constants are not the same as those used in Magma, a standard choice. However, one can if one wishes rectify this by choosing a different base: let  $\tilde{x}_{\alpha_1}(t) = x_{-0100}(t)$ ,  $\tilde{x}_{\alpha_2}(t) = x_{-1242}(t)$ ,  $\tilde{x}_{\alpha_3}(t) = x_{1232}(t)$ ,  $\tilde{x}_{\alpha_4}(t) = x_{-0001}(t)$ . Then we generate the same maximal subgroup  $F_4$  but this time the structure constants do agree with those in Magma.

#### 3.1 How did we find this presentation?

We start with the 248-dimensional module M for  $H \cong {}^{3}D_{4}(2)$ . We use the Ryba space, as used in [4] (see also the description as the 'Lie product method' in [5]), to construct an explicit Lie product on M that turns M into a copy of  $\mathfrak{e}_{8}$ . This gives us explicit structure constants. The module M splits as the sum of 52- and 196-dimensional H-stable submodules. As explained above, the first of these subspaces is forced to be a subalgebra  $\mathfrak{h}$  of  $\mathfrak{e}_{8}$  isomorphic to  $\mathfrak{f}_{4}$  and so Magma could write down a basis for it in terms of a Chevalley basis of  $\mathfrak{e}_{8}$ . However, this process left us with basis elements for  $\mathfrak{h}$  with around 120 non-zero coefficients in terms of a basis of  $\mathfrak{e}_{8}$ .

We found that four of the basis vectors for  $\mathfrak{h}$  were toral and commuted with each other, thus spanning a maximal toral subalgebra  $\mathfrak{t}$ . We then searched for a **G**-conjugate of  $\mathfrak{h}$  such that the corresponding conjugate of  $\mathfrak{t}$  was contained in the standard toral subalgebra of  $\mathfrak{g}$ . To do this we used the inbuilt **InnerAutomorphism** function to construct the automorphisms of  $\mathfrak{g}$  corresponding to  $x_{\gamma}(\pm 1)$  for all roots  $\gamma$  in the root system of  $E_8$ .

Our strategy was to implement a naive hill climb for the first basis element  $t_1$  of t. Indeed, we searched through all 480 possible conjugating elements and selected the one that yielded the largest number of zero coefficients when expressing  $t_1^g$  in terms of the basis of  $\mathfrak{g}$ . We remembered the elements we used at each step. This meant that when we could no longer increase the number of zero coefficients we could trace our steps back and take the next best conjugating element and continue the process. This lead to a significant increase in the number of zero coefficients but nowhere near the 240 we needed.

We then slightly upgraded our hill climb algorithm to include using a random conjugating element at fixed intervals. Every 100 steps we chose a random conjugating element and used this, regardless of what it did to the number of zero coefficients. This method was not optimized; it could be that a better choice would have been every 5 steps, or 500 steps. But this hill climb was enough for us; it quickly led us to a conjugating element  $g_0$  which took  $t_1$  into the standard toral subalgebra of  $\mathfrak{g}$ . At this point, the remaining three basis elements  $t_2, t_3$  and  $t_4$  were not sent by  $g_0$  to something in the standard toral subalgebra of  $\mathfrak{g}$ , but they had significantly fewer non-zero coefficients. We then repeated the algorithm looking to increase the total number of zero coefficients in  $t_1^g, \ldots, t_4^g$  and this quickly converged, yielding a conjugating element  $g_1$  which sent  $t_1, \ldots, t_4$  to the four toral elements corresponding to the generators of the maximal torus given in the presentation. From the toral subalgebra of  $\mathfrak{h}^{g_1}$  it was then routine to take a Cartan decomposition and find the corresponding root elements.

It turned out that the root elements of  $\mathfrak{h}$  were expressed as a sum of commuting root elements of  $\mathfrak{e}_8$ ; in fact long root elements of  $\mathfrak{h}$  were of type  $2A_1$  in  $\mathfrak{e}_8$ , whereas short root elements were of type  $4A_1$ . For pairwise commuting root elements  $e_{\alpha_1}, \ldots, e_{\alpha_k}$  of  $\mathfrak{e}_8$ , the operators ad  $e_{\alpha_i}$  pairwise commute and so one has

$$\exp(t_1 \operatorname{ad} e_{\alpha_1} + \dots + t_j \operatorname{ad} e_{\alpha_k}) = \exp(t_1 e_{\alpha_1}) \dots \exp(t_k e_{\alpha_k}).$$

Thus if  $e_{\beta_i} = \sum t_j e_{\alpha_j} \in \mathfrak{h}$  is of this form, then evidently the left-hand side of the displayed equation normalizes  $\mathfrak{h}_i$  in the group  $\operatorname{GL}_{248}(k)$ , and the right-hand side belongs to  $E_8(k)$ , hence for  $t \in k$ , the elements  $x_{\beta_i}(t)$  generate a connected smooth subgroup  $\mathbf{Y}$  of  $E_8$  for which  $\mathfrak{h} \subseteq \operatorname{Lie}(\mathbf{Y})$ . But now maximality of  $\mathfrak{h}$  forces  $\operatorname{Lie}(\mathbf{Y}) = \mathfrak{h}$ . The only connected smooth affine k-group whose Lie algebra is a simple Lie algebra of type  $F_4$  is a group of type  $F_4$  itself and using the maximality of  $\mathfrak{f}_4$  we conclude that  $\mathbf{Y}$  must be a maximal connected subgroup.

# 4 Consequences

We extend the results of [10] to this new maximal subgroup, determining which unipotent classes of  $E_8$  meet the  $F_4$  non-trivially.

Class in $F_4$	Class in $E_8$
$A_1$	$2A_1$
$\tilde{A}_1$	$4A_1$
$A_1 + \tilde{A}_1$	$A_2 + 2A_1$
$A_2$	$2A_2$
$ ilde{A}_2$	$2A_2$
$A_2 + \tilde{A}_1$	$2A_2 + A_1$
$\tilde{A}_2 + A_1$	$2A_2 + 2A_1$
$B_2$	$2A_3$
$C_{3}(a_{1})$	$A_4 + 2A_1$
$F_4(a_3)$	$A_4 + A_2$
$B_3$	$A_6$
$C_3$	$D_{6}(a_{1})$
$F_4(a_2)$	$D_5 + A_2$
$F_4(a_1)$	$E_{8}(b_{6})$
$F_4$	$E_8(b_4)$

Table 4.1: Fusion of unipotent classes of maximal  $F_4$  into  $E_8$ . (Horizontal lines separate elements of different orders.)

**Proposition 4.1.** If u is a unipotent element of the maximal  $F_4$  subgroup of  $E_8$ , then the class of u in  $F_4$  and  $E_8$  is given in Table 4.1.

*Proof.* The proof is a fast computer check. Randomly generate elements u of orders 3, 9 and 27 in  $F_4(3)$  until we hit each class. (The class to which u belongs can be deduced from [8, Tables 3 and 4].) The Jordan blocks of the action of u on the sum of L(1000) and L(0010) are trivial to compute then. From [8] we obtain the class in  $E_8$  to which u belongs.

However, note that there is an error in [8], due to an error in [16], which leads to a single class having the wrong Jordan blocks in characteristic 3. This is corrected in [9], and it concerns exactly the class  $E_8(b_6)$  in the table. It has Jordan blocks  $9^{26}, 7, 3^2, 1$  on  $L(E_8)$ , not  $9^{25}, 8^2, 2^2, 1^3$  as stated in [8]. With this correction, the Jordan block structure of u on  $L(E_8)$  determines the unipotent class to which u belongs, and thus we are done.

For completeness we do the same thing for nilpotent orbits of  $f_4$ .

**Proposition 4.2.** If x is a nilpotent element of the maximal  $\mathfrak{f}_4$  subalgebra of  $\mathfrak{e}_8$ , then the class of x in  $\mathfrak{f}_4$  and  $\mathfrak{e}_8$  is given in Table 4.2.

*Proof.* Using the root elements constructed for  $\mathfrak{f}_4$  we find a set of orbit representatives for  $\mathfrak{f}_4 \subset \mathfrak{e}_8$  using [18, Appendix]. For each representative x we use Magma to calculate the Jordan block structure for the adjoint action of x and the normalizer of each term of the derived series of  $C_{\mathfrak{e}_8}(x)$ . Using [7, Proposition 1.5], we then find the  $\mathfrak{e}_8$  class of x.

**Proposition 4.3.** If M is a maximal connected reductive subgroup of the maximal  $F_4$  subgroup X of the algebraic group G of type  $E_8$ , then M is conjugate to one of the following four subgroups.

- (i)  $\mathbf{M}_1 = B_4 < D_8$  embedded via the spin module  $L_{B_4}(0001)$ . There are two such conjugacy classes which, when p is odd, are distinguished by whether or not they are contained in a maximal subgroup of type  $A_8$ . This is the subgroup which is not contained in such an  $A_8$ . It is **G**-irreducible and denoted  $E_8(\#45)$  in [20].
- (ii)  $\mathbf{M}_2 = A_1 C_3 < D_8$  embedded via  $L_{A_1}(2) \oplus L_{C_3}(010)$ . This subgroup is **G**-irreducible and denoted by  $E_8(\#774)$  in [20].
- (iii)  $\mathbf{M}_3 = A_1G_2 < A_1E_7$  embedded as follows:  $E_7$  has a maximal subgroup of type  $A_1G_2$  (when  $p \neq 2$ ). Therefore  $A_1E_7$  has a maximal subgroup  $A_1A_1G_2$  and  $M_3$  is embedded diagonally in this subgroup.

Class in $\mathfrak{f}_4$	Class in $\mathfrak{e}_8$
$A_1$	$2A_1$
$ ilde{A}_1$	$4A_1$
$A_1 + \tilde{A}_1$	$A_2 + 2A_1$
$A_2$	$2A_2$
$ ilde{A}_2$	$A_2 + 3A_1$
$A_2 + \tilde{A}_1$	$2A_2 + A_1$
$B_2$	$2A_3$
$\tilde{A}_2 + A_1$	$2A_2 + A_1$
$C_{3}(a_{1})$	$A_4 + 2A_1$
$F_4(a_3)$	$A_4 + A_2$
$B_3$	$A_6$
$C_3$	$D_5 + A_1$
$F_4(a_2)$	$E_{7}(a_{4})$
$F_4(a_1)$	$E_{6}(a_{1})$
$F_4$	$E_6$

Table 4.2: Fusion of nilpotent classes of maximal  $\mathfrak{f}_4$  into  $\mathfrak{e}_8$ .

One has to twist the embedding in the first  $A_1$  factor by the Frobenius morphism. This subgroup is again **G**-irreducible and denoted by  $E_8(\#967^{\{1,0\}})$  in [20].

(iv)  $\mathbf{M}_4 = A_2 A_2 < \bar{A}_2 E_6$  embedded as follows:  $E_6$  has a maximal subgroup  $A_2 G_2$ , and  $G_2$  has a maximal subgroup  $\tilde{A}_2$  generated by short root subgroups of the  $G_2$  when p = 3. Thus  $\bar{A}_2 E_6$  has a subgroup  $H = \bar{A}_2 A_2 \tilde{A}_2$  (denoted  $E_8(\#1012)$  in [20]). The first  $A_2$  factor of  $\mathbf{M}_4$  is the the second  $A_2$  factor of  $\mathbf{M}_4$  is diagonally embedded in the first and third factors of H (with no twisting by field or graph automorphisms). Note that  $\mathbf{M}_4$  is not  $\mathbf{G}$ -irreducible.

*Proof.* By [13, Corollary 2],  $F_4$  has four conjugacy classes of maximal connected subgroups in characteristic 3, which are indeed of types  $B_4$ ,  $A_1C_3$ ,  $A_1G_2$  and  $A_2A_2$ . The first two maximal subgroups are centralizers of involutions. It follows from the action of  $F_4$  on the Lie algebra of  $E_8$  that the centralizer in  $E_8$  of both of these involutions is  $D_8$ . Thus  $B_4$  and  $A_1C_3$  are contained in a maximal subgroup of type  $D_8$ . By [11, Table 8.1], there are only three  $E_8$ -conjugacy classes of  $B_4$  subgroups in  $D_8$ . Calculating the composition factors of the action of  $B_4 < F_4$  on the Lie algebra of  $E_8$  yields that it is indeed conjugate to  $\mathbf{M}_1$ , which is **G**-irreducible by [20, Theorem 1]. Calculating the composition factors of  $A_1C_3 < F_4$  on the Lie algebra of  $E_8$ , we find that it has no trivial composition factors. Therefore it must be **G**-irreducible (by [20, Corollary 3.8]) and we may use the classification of irreducible subgroups determined in [20]. In particular, the composition factors on the Lie algebra of  $E_8$  are enough to determine conjugacy and it follows that  $A_1C_3$  is conjugate to  $\mathbf{M}_2$ , as required.

Another calculation shows that  $A_1G_2 < F_4$  has no trivial composition factors on the Lie algebra of  $E_8$ . We can therefore use the same method as the previous case to deduce that it is conjugate to  $\mathbf{M}_3$ .

For  $\mathbf{AB} = A_2 A_2 < F_4$  we start by considering the first  $A_2$  factor  $\mathbf{A}$ , which we define to be the  $A_2$  subgroup generated by long root subgroups of  $F_4$ . This is the derived subgroup of a long  $A_2$ -Levi subgroup, and is thus a subgroup of  $B_4$ . Since  $B_4$  is conjugate to  $\mathbf{M}_1$ , we find that  $\mathbf{A}$  is contained in a maximal subgroup of type  $A_8$  and acts as  $L_{A_2}(10) \oplus L_{A_2}(01) \oplus L_{A_2}(00)^{\oplus 3}$  on the natural 9-dimensional module of  $A_8$ . Therefore,  $\mathbf{A}$  is a diagonal subgroup (without field or graph twists) of the derived subgroup of an  $A_2A_2$ -Levi subgroup of  $E_8$ . We now claim that the connected centralizer of  $\mathbf{A}$  in  $E_8$  is  $\overline{A}_2G_2$ . Indeed, we use [20, Theorem 1] to see that  $\mathbf{C} = \mathbf{A}\overline{A}_2G_2$  is  $E_8$ -irreducible (denoted by  $E_8(\#978)$ ) and the only reductive connected overgroups of  $\mathbf{C}$  are the maximal subgroups  $\overline{A}_2E_6$  and  $G_2F_4$ . Therefore  $\overline{A}_2G_2 \leq C_{E_8}(\mathbf{A})^{\circ}$ and it must be an equality since  $\mathbf{A}C_G(\mathbf{A})^{\circ}$  is  $E_8$ -irreducible since  $\mathbf{C}$  is.

Therefore, **AB** is contained in  $\bar{A}_2E_6$  with **B** contained in  $\bar{A}_2G_2$ . It is straightforward to list all  $A_2$  subgroups of  $\bar{A}_2G_2$ , noting that  $G_2$  has precisely two classes of  $A_2$  subgroups when the characteristic is 3. Computing the composition factors of **B** on the Lie algebra of  $E_8$  shows that **B** is conjugate to the

subgroup claimed and hence **AB** is conjugate to  $\mathbf{M}_4$ . The fact that  $\mathbf{M}_4$  is **G**-reducible follows from [20, Theorem 1].

# References

- John Conway, Robert Curtis, Simon Norton, Richard Parker, and Robert Wilson, Atlas of finite groups, Oxford University Press, Eynsham, 1985.
- [2] Roger W. Carter, Simple groups of Lie type, John Wiley & Sons Inc., 1989.
- [3] David A. Craven, On medium-rank Lie primitive and maximal subgroups of exceptional groups of Lie type, Mem. Amer. Math. Soc., to appear.
- [4] \_\_\_\_\_, The maximal subgroups of the exceptional groups  $F_4(q)$ ,  $E_6(q)$  and  ${}^2E_6(q)$  and related almost simple groups, preprint, 2020.
- [5] \_\_\_\_\_, The maximal subgroups of the exceptional groups  $E_7(q)$  and related almost simple groups, preprint, 2021.
- [6] Christoph Jansen, Klaus Lux, Richard Parker, and Robert Wilson, An atlas of Brauer characters, Oxford University Press, New York, 1995.
- [7] Mikko Korhonen, David Stewart and Adam Thomas, *Representatives for unipotent classes and nilpotent orbits*, preprint.
- [8] Ross Lawther, Jordan block sizes of unipotent elements in exceptional algebraic groups, Comm. Algebra 23 (1995), 4125–4156.
- [9] \_\_\_\_\_, Correction to: "Jordan block sizes of unipotent elements in exceptional algebraic groups", Comm. Algebra 26 (1998), 2709.
- [10] \_\_\_\_\_, Unipotent classes in maximal subgroups of exceptional algebraic groups, J. Algebra 322 (2009), 270–293.
- [11] Martin Liebeck and Gary Seitz, Reductive subgroups of exceptional algebraic groups, Mem. Amer. Math. Soc. 580 (1996), no. 580, vi+111.
- [12] \_\_\_\_\_, On the subgroup structure of exceptional groups of Lie type, Trans. Amer. Math. Soc. 350 (1998), 3409–3482.
- [13] \_\_\_\_\_, The maximal subgroups of positive dimension in exceptional algebraic groups, Mem. Amer. Math. Soc. 169 (2004).
- [14] Frank Lübeck, Small degree representations of finite Chevalley groups in defining characteristic, LMS J. Comput. Math. 4 (2001), 135–169.
- [15] Anatoly Mal'cev, Commutative subalgebras of semi-simple Lie algebras, Bull. Acad. Soc. URSS Ser. Math. 9 (1945), 291–300.
- [16] Kenzo Mizuno, The conjugate classes of unipotent elements of the Chevalley groups  $E_7$  and  $E_8$ , Tokyo J. Math. **3** (1980), 391–461.
- [17] Gary Seitz, Maximal subgroups of exceptional algebraic groups, Mem. Amer. Math. Soc. 90 (1991), no. 441, iv+197.
- [18] David Stewart, On the minimal modules for exceptional Lie algebras: Jordan blocks and stabilizers, LMS J. Comput. Math. 19 (2016), 235–258.
- [19] Helmut Strade and Rolf Farnsteiner, Modular Lie algebras and their representations, Monographs and Textbooks in Pure and Applied Mathematics, vol. 116, Marcel Dekker Inc., New York, 1988.
- [20] Adam Thomas, The irreducible subgroups of exceptional algebraic groups, Mem. Amer. Math. Soc. 268 (2021), no. 1307, vi+191.