

## Maximizing use of available population-based data on cardiometabolic diseases

Global Health & Population Project on Access to Care for Cardiometabolic Diseases (HPACC)

DOI:

[10.1016/S2213-8587\(21\)00328-4](https://doi.org/10.1016/S2213-8587(21)00328-4)

License:

Creative Commons: Attribution-NonCommercial-NoDerivs (CC BY-NC-ND)

*Document Version*

Peer reviewed version

*Citation for published version (Harvard):*

Global Health & Population Project on Access to Care for Cardiometabolic Diseases (HPACC) 2022, 'Maximizing use of available population-based data on cardiometabolic diseases', *The Lancet Diabetes and Endocrinology*. [https://doi.org/10.1016/S2213-8587\(21\)00328-4](https://doi.org/10.1016/S2213-8587(21)00328-4)

[Link to publication on Research at Birmingham portal](#)

### General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

### Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact [UBIRA@lists.bham.ac.uk](mailto:UBIRA@lists.bham.ac.uk) providing details and we will remove access to the work immediately and investigate.

# Maximizing Use of Available Population-Based Data on Cardiometabolic Diseases

David Flood<sup>1-3</sup>, David Guwatudde<sup>4</sup>, Albertino Damasceno<sup>5-7</sup>, Jennifer Manne-Goehler<sup>8,9\*</sup>, Justine I. Davies<sup>10-12\*</sup>, for the Global Health & Population Project on Access to Care for Cardiometabolic Diseases (HPACC)<sup>†</sup>

<sup>1</sup>Division of Hospital Medicine, Department of Medicine, University of Michigan, Ann Arbor, Michigan, USA; <sup>2</sup>Center for Indigenous Health Research, Wuqu' Kawoq, Tecpán, Guatemala; <sup>3</sup>Research Center for the Prevention of Chronic Diseases, Institute of Nutrition of Central America and Panama, Guatemala City, Guatemala; <sup>4</sup>Department of Epidemiology and Biostatistics, School of Public Health, Makerere University, Kampala, Uganda; <sup>5</sup>Department of Public and Forensic Health Sciences and Medical Education, Faculty of Medicine, University of Porto, Porto, Portugal; <sup>6</sup>EPIUnit, Institute of Public Health, University of Porto, Porto, Portugal; <sup>7</sup>Faculty of Medicine, Eduardo Mondlane University, Maputo, Mozambique; <sup>8</sup>Division of Infectious Diseases, Brigham and Women's Hospital, Harvard Medical School, Boston, MA, USA; <sup>9</sup>Medical Practice Evaluation Center, Massachusetts General Hospital, Harvard Medical School, Boston, MA, USA; <sup>10</sup>Institute for Applied Health Research, University of Birmingham, UK; <sup>11</sup>Centre for Global Surgery, Department of Global Health, Stellenbosch University, Cape Town, South Africa; <sup>12</sup>Medical Research Council/Wits University Rural Public Health and Health Transitions Research Unit, Faculty of Health Sciences, School of Public Health, University of the Witwatersrand, Johannesburg, South Africa

\*Joint senior authors

<sup>†</sup>Members listed at end of paper

26 Corresponding author:  
27 Justine I. Davies, MD  
28 Institute ~~for~~of Applied Health Research  
29 University of Birmingham  
30 Birmingham, United Kingdom  
31 B15 2TT  
32 [J.davies.6@bham.ac.uk](mailto:J.davies.6@bham.ac.uk)  
33  
34 Word count: 907947  
35 References: 10

36 The absolute worldwide burden of adult cardiometabolic diseases such as hypertension,  
37 diabetes, obesity, and dyslipidemia continues its relentless ascent. Scaling up ~~the~~ prevention,  
38 management, and control of cardiometabolic diseases is cost-effective but requires strong  
39 health systems.<sup>14</sup> Building ~~these strong health systems~~ requires data that are accurate, timely,  
40 and transparent, as we have previously argued in this journal.<sup>2</sup> In particular, data from high-  
41 quality population-based surveys are critical, as they reflect the spectrum of community-dwelling  
42 adults ~~in a particular geography~~, including those who are not reached by the health system.

44 ~~There~~Of late, ~~there~~ has been tremendous progress in making population-based survey data  
45 available for cardiometabolic diseases. Emblematic of this has been the release in 2018 of the  
46 World Health Organization (WHO) Noncommunicable Disease (NCD) Microdata Repository.<sup>3</sup>  
47 This hosts over 130 surveys conducted using the STEPwise approach to NCD surveillance  
48 (STEPS) methodology that are now available ~~after to users who submit~~ a brief application. Most  
49 STEPS surveys are conducted in low- and middle-income countries (LMICs) where a majority of  
50 the cardiometabolic disease burden occurs. Thus, this resource fills a critical gap in openly  
51 accessible population-based survey data on cardiometabolic risk factors and health care access  
52 in these settings.

54 Yet, there is more work to be done. The availability of population-based data, while necessary,  
55 is insufficient ~~by itself~~ to ensure their effective use to shape programs, strategies, and policies  
56 addressing cardiometabolic diseases. In this Comment, we highlight three other crucial actions  
57 ~~needed~~ to maximize the use of population data: harmonization, alignment with monitoring  
58 indicators to benchmark health system performance, and capacity-building initiatives to  
59 democratize data use [\(figure\)](#).

Our perspective is informed by our experience in the Global Health and Population Project on Access to Care for Cardiometabolic diseases (HPACC), an international research consortium with collaborators in more than 30 countries. ~~HPACC has created a dynamic repository of harmonized, nationally representative survey data currently representing 1.3 million individuals in more than 75 LMICs (including more than 50 STEPS surveys) to address questions of relevance to health system planning and evaluation for cardiometabolic diseases.~~

First, while population-based data ~~can and~~ should be used at the national level, these data also should be harmonized to maximize ~~its~~ use by international advocacy organizations, policymakers, and researchers. Harmonization refers to the process of bringing together distinct data sources into a single comparable format. Harmonized survey data are available in the area of maternal and child health,<sup>4</sup> but no such resource exists for cardiometabolic diseases. Such harmonized data allows for assessing health system effectiveness ~~and responsiveness~~, as our study of the state of hypertension care in 44 LMICs illustrates.<sup>55</sup> Harmonization ~~also~~ provides larger and more diverse samples, giving added power to study variations in cardiometabolic risk factors, including biological measures such as blood glucose and behavioral risk factors such as physical activity and diet. Understanding these variations is important, as it cannot be assumed that epidemiologic patterns of clinical relevance observed in well-studied high-income countries will be conserved in LMICs. Indeed, we have found that the association between diabetes and body mass index (BMI) is highly variable across world regions, implying that BMI thresholds generated using European or North American data cannot simply be applied ~~elsewhere in other world regions.~~<sup>6</sup> ~~Harmonization also allows for the construction of sophisticated clinical and policy models for the prevention, treatment, and control of cardiometabolic diseases to predict effects of change.~~<sup>1,76</sup> Importantly, to ensure that data are useful for cross-country comparisons, prior to data collection, time should be spent ensuring survey instruments and data collection are standardized and aligned with the highest priority global health metrics.

87

88 Second, population data on cardiometabolic diseases should be harnessed to benchmark and  
89 monitor health system performance. At present, these data are underutilized for this purpose.

90 Harmonized data from STEPS and ~~similar~~~~non-STEPS~~ surveys can reveal progress on monitoring indicators  
91 in the NCD Global Monitoring Framework<sup>7</sup> and inform new targets such as those proposed by  
92 the WHO Global Diabetes Compact, a recently established initiative to improve global diabetes  
93 care.<sup>88</sup> To show global variation in health system performance, harmonized data ideally should  
94 include not only LMICs but also high-income countries, though unfortunately data from high-  
95 income countries are currently less available.

96

97 Third, given limited research capacity in many LMICs, there is a need to build capacity to ensure  
98 the wide usability of population data on cardiometabolic diseases, ~~most especially~~ by those who have collected  
99 it. Local researchers—especially those in LMICs—who design and conduct surveys should be  
100 empowered to use harmonized data to answer their policy-relevant questions, conduct  
101 independent analyses, and publish in lead-author roles.<sup>9</sup> In addition to this being a step towards  
102 decolonialization of global health, these collaborators add critical contextual interpretation that  
103 may not be fully perceived or appreciated by those outside their settings.

104 maximizing use of available population data on cardiometabolic diseases, it is important to  
105 continue data-sharing efforts. Many STEPS and comparable ~~non-STEPS~~ household surveys remain  
106 unavailable, as are more than two dozen nationally representative health facility surveys  
107 conducted using the WHO Service Availability Readiness Assessment (SARA) methodology.<sup>10</sup>  
108 Additionally, many other data sources, for example, from subnational research studies, remain  
109 inaccessible. Finally, cardiometabolic disease epidemiology is rapidly evolving, but data are  
110 often historical. As ~~is done~~ for HIV, data collection for cardiometabolic diseases needs to be ongoing to  
111 assess temporal trends in disease prevalence and health system performance.

112

113 While we focus on maximizing use of available population data on cardiometabolic diseases, it  
114 is important to continue data-sharing efforts. Many STEPS and comparable ~~non-STEPS~~  
115 household surveys remain unavailable, as are more than two dozen nationally representative  
116 health facility surveys conducted using the WHO Service Availability Readiness Assessment  
117 (SARA) methodology.<sup>10</sup> Additionally, many other data sources, for example, from subnational  
118 research studies, remain inaccessible. Finally, cardiometabolic disease epidemiology is rapidly  
119 evolving, but data are often historical. As ~~is done~~ for HIV, data collection for cardiometabolic  
120 diseases needs to be ongoing to assess temporal trends in disease prevalence and health  
121 system performance.

122  
123 The staggering burden of cardiometabolic diseases brings ~~with it~~ an imperative to maximize the use of  
124 ~~these~~ data. Many ~~countries and individuals~~ LMICs ~~already~~ have invested substantial resources  
125 in producing these data, which are a global public good. However, while they are increasingly  
126 available, in practice ~~they~~ ~~these data~~ are still too sparse and underutilized given the toll these  
127 diseases are taking on people worldwide. We call on funders and international health  
128 organizations to invest in efforts to collect, harmonize and make available these data with an  
129 urgency befitting the magnitude of the global burden of cardiometabolic diseases.

## Declaration of interests

DF reports grant funding within the 3 years from a Pilot and Feasibility Grant funded by the Michigan Center for Diabetes Translational Research (NIH Grant P30-DK092926) and a grant from the Swinmurn Foundation to implement a sustainable diabetes clinic in Guatemala. DF also reports volunteer affiliations with Wuqu' Kawoq and GlucoSalud, outside the submitted work. During the course of this work, DF has received research fellowship funding from National Clinician Scholars Program at the University of Michigan Institute for Healthcare Policy & Innovation. TB reports grants from the Alexander von Humboldt Foundation, Wellcome Trust, and U.S. National Institutes of Health. RA reports contracts with Novo Nordisk, Union for International Cancer Control's (UICC), Novo Nordisk, Hoffman-La Roche, and Sloan Memorial Kettering Hospital, outside the submitted work. RA also reports payments or honorario from Merck and Hoffmann-La Roche, outside the submitted work. All other authors declare no competing interests.

## **References**

1. Basu S, Flood D, Geldsetzer P, et al. Estimated impact of increased diagnosis, treatment, and control of diabetes mellitus among low-and middle-income countries: A microsimulation model. *Lancet Global Health* 2021; (Accepted; in press).
2. Davies J, Yudkin JS, Atun R. Liberating data: the crucial weapon in the fight against NCDs. *Lancet Diabetes Endocrinol* 2016; 4(3): 197-8.
3. World Health Organization. NCD Microdata Repository. 2021. <https://extranet.who.int/ncdsmicrodata/index.php/catalog> (accessed July 19, 2021).
4. IPUMS. IPUMS Global Health. 2021. <https://globalhealth.ipums.org/> (accessed October 25, 2021).



- 156 5. Geldsetzer P, Manne-Goehler J, Marcus ME, et al. The state of hypertension care in 44  
157 low-income and middle-income countries: a cross-sectional study of nationally representative  
158 individual-level data from 1.1 million adults. *Lancet* 2019; 394(10199): 652-62.
- 159 6. Teufel F, Seiglie JA, Geldsetzer P, et al. Body-mass index and diabetes risk in 57 low-  
160 income and middle-income countries: a cross-sectional study of nationally representative,  
161 individual-level data in 685 616 adults. *Lancet* 2021; 398(10296): 238-48.
- 162 7. WHO. Noncommunicable Diseases Global Monitoring Framework: Indicator Definitions  
163 and Specifications. 2014. [https://www.who.int/nmh/ncd-](https://www.who.int/nmh/ncd-tools/indicators/GMF_Indicator_Definitions_Version_NOV2014.pdf)  
164 [tools/indicators/GMF\\_Indicator\\_Definitions\\_Version\\_NOV2014.pdf](https://www.who.int/nmh/ncd-tools/indicators/GMF_Indicator_Definitions_Version_NOV2014.pdf) (accessed September 29,  
165 2021).
- 166 8. Hunt D, Hemmingsen B, Matzke A, et al. The WHO Global Diabetes Compact: a new  
167 initiative to support people living with diabetes. *Lancet Diabetes Endocrinol* 2021; 9(6): 325-7.
- 168 9. Wang W, Assaf S, Pullum T, Kishor S. The Demographic and Health Surveys Faculty  
169 Fellows Program: Successes, Challenges, and Lessons Learned. *Glob Health Sci Pract* 2021;  
170 9(2): 390-8.
- 171 10. Mangipudi S, Leather A, Seedat A, Davies J. Oxygen availability in sub-Saharan African  
172 countries: a call for data to inform service delivery. *The Lancet Global Health* 2020; 8(9): e1123-  
173 e4.  
174  
175

## HPACC members

David Flood<sup>1-3</sup>, David Guwatudde<sup>4</sup>, Albertino Damasceno<sup>5-7</sup>, Krishna K. Aryal<sup>8</sup>, Rifat Atun<sup>9,10</sup>, Till W. Bärnighausen<sup>11-13</sup>, Brice Wilfried Bicaba<sup>14</sup>, Pascal Bovet<sup>15,16</sup>, Gary Brian<sup>17</sup>, Maria Dorobantu<sup>18</sup>, Farshad Farzadfar<sup>19</sup>, Gladwell Gathecha<sup>20</sup>, Pascal Geldsetzer<sup>11,21</sup>, Mongal Singh Gurung<sup>22</sup>, Corine Houehanou<sup>23</sup>, Nahla Hwalla<sup>24</sup>, Lindsay Jaacks<sup>25</sup>, Jutta Jorgensen<sup>26</sup>, Gibson Kagaruki<sup>27</sup>, Khem Karki<sup>28</sup>, Demetre Labadarios<sup>29</sup>, Nuno Lunet<sup>5,30</sup>, Maja E. Marcus<sup>31</sup>, Joao Martins<sup>32</sup>, Mary Mayige<sup>27</sup>, Omar Mwalim<sup>33,34</sup>, Kibachio Joseph Mwangi<sup>20,35</sup>, Bolormaa Norov<sup>36</sup>, Rebekka Rühle<sup>31</sup>, Sahar Saeedi Moghaddam<sup>37</sup>, Jacqueline A. Seiglie<sup>38</sup>, Abla M. Sibai<sup>39</sup>, Bahendeka Karaireho Silver<sup>40</sup>, Andrew Stokes<sup>41</sup>, Lela Sturua<sup>42</sup>, Adil Supiyev<sup>43</sup>, Michaela Theilmann<sup>11</sup>, Lindiwe Tsabedze<sup>44</sup>, Sebastian Vollmer<sup>31</sup>, Kehinde D. Whyte-Ilori<sup>45</sup>, Zhaxybay Zhumadilov<sup>46</sup>, Jennifer Manne-Goehler<sup>47,48</sup>, Justine I. Davies<sup>49-51</sup>

## Affiliations

<sup>1</sup>Division of Hospital Medicine, Department of Medicine, University of Michigan, Ann Arbor, Michigan, USA; <sup>2</sup>Center for Indigenous Health Research, Wuqu' Kawoq, Tecpán, Guatemala; <sup>3</sup>Research Center for the Prevention of Chronic Diseases, Institute of Nutrition of Central America and Panama, Guatemala City, Guatemala; <sup>4</sup>Department of Epidemiology and Biostatistics, School of Public Health, Makerere University, Kampala, Uganda; <sup>5</sup>Department of Public and Forensic Health Sciences and Medical Education, Faculty of Medicine, University of Porto, Porto, Portugal; <sup>6</sup>EPIUnit, Institute of Public Health, University of Porto, Porto, Portugal; <sup>7</sup>Faculty of Medicine, Eduardo Mondlane University, Maputo, Mozambique; <sup>8</sup>Monitoring Evaluation and Operational Research Project, Abt Associates, Kathmandu, Nepal; <sup>9</sup>Department of Global Health and Population, Harvard T.H. Chan School of Public Health, Harvard University, Boston, MA, USA; <sup>10</sup>Department of Global Health and Social Medicine, Harvard Medical School, Harvard University, Boston, MA, USA; <sup>11</sup>Heidelberg Institute of Global Health, Heidelberg University, Heidelberg, Germany; <sup>12</sup>Africa Health Research Institute, Somkhele, South Africa; <sup>13</sup>Harvard Center for Population and Development Studies, Cambridge, USA; <sup>14</sup>Institut National de Santé Publique, Burkina Faso; <sup>15</sup>Ministry of Health, Victoria, Seychelles; <sup>16</sup>Institute of Social and Preventive Medicine, Lausanne, Switzerland; <sup>17</sup>The Fred Hollows Foundation New Zealand; <sup>18</sup>University of Medicine and Pharmacy Carol Davila, Bucharest, Romania; <sup>19</sup>Non-Communicable Diseases Research Center, Endocrinology and Metabolism Population Sciences Institute, Tehran University of Medical Sciences, Tehran, Iran; <sup>20</sup>Division of Non-Communicable Diseases, Ministry of Health, Nairobi, Kenya; <sup>21</sup>Division of Primary Care and Population Health, Stanford University; <sup>22</sup>Health Research and Epidemiology Unit, Ministry of Health, Thimphu, Bhutan; <sup>23</sup>Laboratory of Epidemiology of Chronic and Neurological Diseases, Faculty of Health Sciences, University of Abomey-Calavi, Cotonou, Benin; <sup>24</sup>Faculty of Agricultural and Food Sciences, American University of Beirut, Beirut, Lebanon; <sup>25</sup>Global Academy of Agriculture and Food Security, The University of Edinburgh, Midlothian, United Kingdom; <sup>26</sup>Department of Public Health, University of Copenhagen, Copenhagen, Denmark; <sup>27</sup>National Institute for Medical Research, Dar es Salaam, Tanzania; <sup>28</sup>Department of Community Medicine and Public Health, Institute of Medicine, Tribhuvan University, Kathmandu, Nepal; <sup>29</sup>Faculty of Medicine and Health Sciences, Stellenbosch University, Stellenbosch, South Africa; <sup>30</sup>EPIUnit Institute of Public Health, University of Porto, Porto, Portugal; <sup>31</sup>Department of Economics and Centre for Modern Indian Studies, University of Göttingen, Göttingen, Germany; <sup>32</sup>Faculty of Medicine and Health Sciences, National University of East Timor, Dili, Timor-Leste; <sup>33</sup>Ministry of Health and Social Welfare, Elderly, Gender and Children, Zanzibar, Tanzania; <sup>34</sup>Bergen Centre for Ethics and Priority Setting (BCEPS), Department of Global Public Health and Primary Care, University of Bergen, Norway; <sup>35</sup>Faculté de Médecine, Université de Genève, Geneva, Switzerland; <sup>36</sup>National Center for Public Health, Ulaanbaatar, Mongolia;

<sup>37</sup>Endocrinology and Metabolism Research Center, Endocrinology and Metabolism Clinical Sciences Institute, Tehran University of Medical Sciences, Tehran, Iran; <sup>38</sup>Diabetes Unit, Massachusetts General Hospital, Boston, United States of America; <sup>39</sup>Epidemiology and Population Health Department, Faculty of Health Sciences American University of Beirut, Beirut, Lebanon; <sup>40</sup>Uganda Martyrs University | Saint Francis Hospital Nsambya, Kampala, Uganda; <sup>41</sup>Department of Global Health, Boston University School of Public Health, Boston, United States of America; <sup>42</sup>Non-Communicable Disease Department, National Center for Disease Control and Public Health, Tbilisi, Georgia; <sup>43</sup>Laboratory of Epidemiology and Public Health, Center for Life Sciences, Nazarbayev University, Astana, Kazakhstan; <sup>44</sup>Ministry of Health, Mbabane, Eswatini; <sup>45</sup>School of Medicine, University of Leeds, Leeds, West Yorkshire, United Kingdom; <sup>46</sup>Nazarbayev University School of Medicine, Nur-Sultan, Kazakhstan; <sup>47</sup>Division of Infectious Diseases, Brigham and Women's Hospital, Harvard Medical School, Boston, MA, USA; <sup>48</sup>Medical Practice Evaluation Center, Massachusetts General Hospital, Harvard Medical School, Boston, MA, USA; <sup>49</sup>Institute for Applied Health Research, University of Birmingham, UK; <sup>50</sup>Centre for Global Surgery, Department of Global Health, Stellenbosch University, Cape Town, South Africa; <sup>51</sup>Medical Research Council/Wits University Rural Public Health and Health Transitions Research Unit, Faculty of Health Sciences, School of Public Health, University of the Witwatersrand, Johannesburg, South Africa;"