

Determination of 4-(Methylnitrosamino)-1-(3-Pyridyl)-1-Butanone (NNK) arising from tobacco smoke in airborne particulate matter

Aquilina, Noel J.; Havel, Christopher M.; Harrison, Roy M.; Ho, Kin Fai; Benowitz, Neal L.; Jacob, Peyton

DOI:

[10.1016/j.envint.2021.106992](https://doi.org/10.1016/j.envint.2021.106992)

License:

Creative Commons: Attribution-NonCommercial-NoDerivs (CC BY-NC-ND)

Document Version

Publisher's PDF, also known as Version of record

Citation for published version (Harvard):

Aquilina, NJ, Havel, CM, Harrison, RM, Ho, KF, Benowitz, NL & Jacob, P 2022, 'Determination of 4-(Methylnitrosamino)-1-(3-Pyridyl)-1-Butanone (NNK) arising from tobacco smoke in airborne particulate matter', *Environment International*, vol. 158, 106992. <https://doi.org/10.1016/j.envint.2021.106992>

[Link to publication on Research at Birmingham portal](#)

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.



Determination of 4-(Methylnitrosamino)-1-(3-Pyridyl)-1-Butanone (NNK) arising from tobacco smoke in airborne particulate matter

Noel J. Aquilina^{a,b,*}, Christopher M. Havel^b, Roy M. Harrison^{c,1}, Kin-Fai Ho^d, Neal L. Benowitz^b, Peyton Jacob III^b

^a Department of Chemistry, University of Malta, Msida MSD 2080, Malta

^b Program in Clinical Pharmacology, Division of Cardiology, Department of Medicine, University of California, San Francisco, CA 94143, United States

^c Division of Environmental Health and Risk Management, School of Geography, Earth and Environmental Sciences, University of Birmingham, Birmingham B15 2TT, United Kingdom

^d The Jockey Club School of Public Health and Primary Care, The Chinese University of Hong Kong, Hong Kong, China

ARTICLE INFO

Handling Editor: Dr. Xavier Querol

Keywords:

Tobacco Smoke

Tobacco-Specific Nitrosamines

NNK

Particulate Matter

ABSTRACT

The most important tobacco-specific nitrosamine found in cigarette smoke and formed in ageing smoke after cigarettes are extinguished is 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone (NNK). It is formed from nitrosation of nicotine, under particular conditions both in indoor and outdoor environments. NNK has been classified as a potent lung carcinogen which is expected to be found primarily in the particle-phase and to be stable in particulate matter. In this study tests have been carried out to show that a bisulfate-treated filter is more efficient than an untreated filter to collect both nicotine and NNK, and that the latter is stable in outdoor particulate matter. To characterize NNK in the outdoor environment, airborne samples were collected from 11 cities in USA, UK, Hong Kong and Malta with characteristics varying from low to high population densities and from urban to suburban to rural, and with desert characteristics and distinct climates. It has been shown that airborne particle + gas phase nicotine and particle-phase NNK behave in a linearly correlated manner. A seasonal analysis was carried out on a subset of data available from five sites in California, where the load of NNK in PM₁₀ is driven by long range transport of the air masses passing over densely populated cities. In the winter season, the load of NNK in PM is higher than in summer in a statistically significant manner. The contamination of PM with NNK shows variability, but is observed at all sites. This paper highlights the potential risk of chronic exposure to NNK in particulate matter by the inhalation pathway.

1. Introduction

Nitrosamines are an important sub-group of organic nitrogen compounds that are found in several matrices (Farren et al., 2015) and are classified as extremely potent carcinogens by the International Agency for Research on Cancer (IARC) (Cogliano et al., 2011). They are challenging to characterise as they are found in minute quantities compared to inorganic components, hydrocarbons and oxygenated constituents of airborne particulate matter (PM) (Akyüz and Ata, 2013; Aragón et al., 2013; Chao et al., 2008; Farren et al., 2015; Häger and Niessner, 1996; Marano et al., 1982; Ramírez et al., 2012; Rounbehler et al., 1980). They are emitted directly to the atmosphere from several sources (Brown

et al., 2003) but are also formed through nitrosation or oxidation/nitrosation reactions of their precursor amines (Ge et al., 2011).

Tobacco use increases significantly the nitrosamine exposure through a group termed tobacco-specific nitrosamines (TSNAs), formed by the reaction of nicotine and other alkaloids with nitrous acid and other atmospheric oxidants (Hoffmann et al., 1984; Sleiman et al., 2010). The most important TSNA is 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone (NNK) as it is classified as a group 1 carcinogen by the IARC (Hoffmann et al., 1984; IARC, 2007), is on the list of the harmful and potentially harmful constituents in tobacco products and tobacco smoke, and it has been prioritized for regulatory purposes (Gunduz et al., 2016). In recent studies NNK has been emphasised as an

* Corresponding author.

E-mail address: noel.aquilina@um.edu.mt (N.J. Aquilina).

¹ Also at: Department of Environmental Sciences / Center of Excellence in Environmental Studies, King Abdulaziz University, PO Box 80203, Jeddah, 21589, Saudi Arabia.

<https://doi.org/10.1016/j.envint.2021.106992>

Received 27 September 2021; Received in revised form 9 November 2021; Accepted 16 November 2021

Available online 19 November 2021

0160-4120/© 2021 The Authors.

Published by Elsevier Ltd.

This is an open access article under the CC BY-NC-ND license

(<http://creativecommons.org/licenses/by-nc-nd/4.0/>).




Fig. 1. Schematic diagram for sample preparation, extraction, cleaning and analysis of Nicotine and NNK.

environmental contaminant of increasing health concern (Farren et al., 2015; Whitehead et al., 2015). Given its carcinogenic potency especially to induce lung cancer by metabolic activation (Hoffmann et al., 1994), the presence of NNK in PM can pose a potential population health risk (Apelberg et al., 2013; Farren et al., 2015; Hoffmann et al., 1991; Jacob et al., 2013; Kavvadias et al., 2009).

A higher population density and a higher tobacco smoking prevalence should result in higher airborne nicotine concentrations. Given that nicotine is subject to atmospheric reactivity, primarily with nitrous acid and ozone, it does not necessarily mean that the airborne NNK concentration will be higher. Whilst nicotine is found almost exclusively in the gas-phase (Alonso et al., 2010; Daisey, 1999), NNK is primarily found in the particle-phase (Jacob et al., 2013). In a recent study, it has been shown that PM contains a ubiquitous contribution of tobacco smoke (Aquilina et al., 2021). Although NNK tends to be very stable in the particle-phase, in chamber studies it was shown that NNK forms mainly indoors especially if nitrous acid is in high concentrations, but is reactive towards ozone and NO_x (Petrick et al., 2010; Petrick et al., 2011a; Petrick et al., 2011b; Sleiman et al., 2010) that would lead to its destruction. We are aware of only one study that has analysed NNK in airborne PM in London, and discussed the variability of NNK across winter and summer (Farren et al., 2015), but no studies have characterised the NNK in outdoor PM collected from various geographic locations under different atmospheric conditions. It is important to quantify the load of NNK in PM in view of the potential risk associated with the exposure to this potent carcinogen when exposed to outdoor PM, and in non-smoking indoor environments due to infiltration of PM from outdoors. For this purpose, specific tests have been carried out to check the stability of NNK in airborne PM. This is also the first study presenting an analysis of seasonal variability of NNK in terms of its load in PM, based on airborne samples collected in 11 sites, representing a range of geographic locations, from urban to rural to remote desert, in three continents.

2. Materials and methods

2.1. Reagents and standards

Water, methanol, dichloromethane (DCM), pentane, isopropyl alcohol (IPA), ethyl acetate (EtOAc), toluene and butanol (HPLC grade)

and sulfuric acid, hydrochloric acid, potassium carbonate, tetrasodium EDTA, acetone and ammonium formate (reagent grade) were purchased from Fisher Scientific, USA. Nicotine- d_4 used as an internal standard was synthesized and converted to the tartrate salt as described by (Jacob et al., 1988). Nicotine- d_0 base was purchased from Sigma Aldrich, USA, and converted to the tartrate salt based as previously described in (Jacob et al., 1988). Deuterated NNK (NNK- d_4) and NNK- d_0 were purchased from Toronto Research Chemicals, Canada. The Standard Reference Material (SRM) 1649b (Urban Dust) was purchased from the National Institute of Standards and Technology, USA.

2.2. Potential *in situ* formation and stability of NNK and collection efficiency of filters

Three tests were performed to characterise the nature of NNK in airborne PM samples.

Test 1: A test was run to check if any NNK collected on a quartz fibre filter is a sampling artefact due to *in situ* formation. A 100 mL bisulfate solution was prepared as described by (Leaderer and Hammond, 1991) and was spiked with 1 mL of 2 mg mL^{-1} nicotine- d_4 base in 0.02 N hydrochloric acid. The solution was placed in a DCM cleaned trough and two filters were allowed to soak in the solution for 30 min, then allowed to dry in a fume hood and stored in foil in a freezer. One control filter was left in the freezer and the other filter sampled air for 6 h at an average flowrate of 1.22 $\text{m}^3 \text{min}^{-1}$ at the San Francisco site where HONO concentrations are expected to be low, consistent with the typical North American urban background. Diurnally-averaged concentrations measured in Bakersfield, CA were below 0.5 ppb (VandenBoer et al., 2014). The two filters were extracted and analysed together for any deuterated NNK formed during sampling.

Test 2: If NNK does not degrade when exposed to normal atmospheric conditions, then the sum of the NNK collected on two consecutive 24-hour samples should add up to a co-located 48-hour sample. Several samples were run to check the stability of NNK.

Test 3: Although the use of a bisulfate-treated filter is used primarily used to capture the gas + particle phases of nicotine, the efficiency of the use of such filters in the collection of NNK needed to be evaluated. For this purpose, two identical calibrated Leckel low-volume samplers with $\text{PM}_{2.5}$ heads were used to collect several concurrent 24-hour samples at an average flowrate of 0.038 $\text{m}^3 \text{min}^{-1}$, one loaded with a bisulfate-

Table 1

Precision and accuracy for determination of NNK in analyte spiked filters used as QC samples.

Analyte	Added Amount (ngmL ⁻¹)	Measured Mean (ngmL ⁻¹)	Accuracy (% of Expected)	Precision (%RSD)
NNK	0 ^a	BLOQ	NA	NA
LOQ = 0.025, N = 44,	0 ^b	BLOQ	NA	NA
16 Analytical Runs	0 ^c	BLOQ	NA	NA
	1	1.11	111	10.8
	2	1.77	88	10.5
	5	5.51	110	2.7
	10	11.2	112	5.3

a. Analytical blanks; b. Travel blanks; c. Field blanks; N: Number of replicate samples; LOQ: Limit of Quantitation; BLOQ: Below Limit of Quantitation; RSD: Relative Standard Deviation; NA: Not Applicable.

treated filter and the second with an untreated one.

2.3. Outdoor atmospheric sampling

Airborne PM samples were collected at different times from eleven sites in four countries with different, characteristic microclimates. The sites were Bakersfield (BCA), Canebrake (CB), Sacramento (STS), Yosemite Village (YTD), Shasta Lake (SLM) in 2010 and San Francisco (SF) in 2016 in California, USA, Tseung Kwan O Industrial Estate (MA), Tseung Kwan O (TKO) and Tin Shui Wai (TSW) in Hong Kong, China in 2014–2015, Birmingham (BHAM), UK in 2016, and Msida (MSD), Malta in 2016 and 2018. The sites represent a variety of industrial, urban, suburban and rural settings. The sampling site information, sampling periods, instrumentation and filters used, PM fractions collected, calibration protocols and other details are discussed in (Aquilina et al., 2021). Whenever bisulfate treated filters were used, the samples were analysed for both nicotine and NNK.

2.4. Sample preparation

2.4.1. Working standards

The concentrations of the standards, which span the expected range were typically 0.025–100 ng mL⁻¹ for NNK and 4–1200 ng mL⁻¹ for nicotine. For quality assurance purposes, 1, 2, 5 and 10 ng mL⁻¹ NNK-d₀ spiked filters and 5, 10, 125, 370 and 1111 ng mL⁻¹ nicotine-d₀ were included with every 10 samples. Field and analytical blank filters were run every five samples.

2.4.2. Extraction and analyses of nicotine and NNK

Details of the filter extraction, extract purification and concentration, and parameters for the analyses of NNK by LC-MS/MS and Nicotine by GC-MS/MS are described in our previous paper (Aquilina et al., 2021) and (Whitehead et al., 2015) and shown as a schematic summary in Fig. 1. The same procedure was applied on 10 mg of SRM 1649b-Urban dust (NIST, 2017).

3. Results and discussion

3.1. Extraction of filters and analytical method validation

Performance of the analytical method including precision and accuracy, Limits of Quantitation (LOQs) and analysis of blanks that verify lack of interfering substances is summarized in Table 1. Precision and accuracy of the method were determined from the results of analysis of the spiked filters described above. LOQs were determined as the minimum quantity on the calibration curve that does not exceed an RSD of 15% for blank filters or blank filters spiked with analytes. For NNK the LOQ was 0.025 ngmL⁻¹. The details of the performance for the




Fig. 2. (a) NNK vs Nicotelline in Total Particulate Matter (TPM) in chamber study and (b) airborne NNK vs Nicotelline in Birmingham (BHAM-PM₁₀■), San Francisco (SF-TSP●) and Msida (MSD-PM₁₀◆) in 2016 and Msida (MSD-PM_{2.5}▲) in 2018 on treated filters.

analytical method for nicotine are described in our previous paper (Aquilina et al., 2021). As part of the analytical quality control protocol to document the precision of the analytical method, the amount of nicotine and NNK found in the SRM 1649b (NIST, 2017) analysed during this study are reported. These compounds are not listed in the NIST certificate of this SRM and therefore the results for precision only are reported. During the method development for the analysis of alkaloids in house dust, from a very limited number of samples (N = 2), the mean NNK level in the same SRM used in this study was 11.6 ng/g and for nicotine it was 66,000 ng/g (Whitehead et al., 2015). In this study, similar results for more samples (N = 9) and with higher confidence can be reported. The mean NNK was 6.8 ng/g (RSD 13.1%) and mean nicotine was 52,800 ng/g (RSD 8.4%). The validation data for the proposed method is considered suitable and is comparable to the previously reported method (Jacob et al., 2013).

3.2. Characterising the nature of NNK in airborne PM samples

Acidic conditions tend to favour nitrosation. In the NNK molecule, protonation of the pyridine nitrogen would protect it from oxidation at that site but perhaps not against reactions at the side chain, with the carbonyl group. This means that here we have to give a caveat. NNK can show an intrinsic and significant reactivity under normal atmospheric conditions.

Fig. 2(a) represents the relationship of the load of NNK in total particulate matter (TPM) as compared to the nicotelline (the tobacco-smoke derived particulate marker) load in TPM from a chamber study involving reference cigarettes smoked by a machine as described in (Schick et al., 2014; Whitlatch and Schick, 2019). Tobacco smoke was aged for 30 min, where upon emission, the smoke undergoes many




Fig. 3. NNK (in ng) on bisulfate-treated (T) and untreated (U) filters. (Solid line represents $Y = X$).

changes, loss of SVOCs from PM and further chemical reactions due to reactive species. The large intercept may be attributed to the above and to the fact that NNK is present in tobacco and formed by pyrosynthesis during the combustion.

Fig. 2(b) shows the relationship of airborne NNK and nicotine under normal atmospheric conditions. These samples were collected in this study on treated filters and in different PM fractions. Taking into account the caveat explained above, the relationship between NNK and nicotine in both chamber and atmospheric data is consistent. The slope of Fig. 2(b) is higher than that in Fig. 2(a) most probably indicating that NNK may have formed from nicotine under normal atmospheric conditions.

Test 1: Potential formation of NNK and stability: From the analysis of the pre-treated control and the sampled filters that were both spiked with nicotine- d_4 as described in Section 2.2, no deuterated NNK was detected, confirming that *in situ* formation of this compound on the filter during outdoor sampling did not occur at a significant rate. From the indoor scenario study by Sleiman et al., NNK forms rapidly (≈ 3 h) and then undergoes partial degradation by nitrous acid, to reach steady-state concentrations (Sleiman et al., 2010). A sampling volume of 440 m^3 was collected. This represents a substantial volume to collect sufficient PM to check if detected NNK is a sampling artefact.

Test 2: From the co-located sampling experiment ($N = 7$) described in Section 2.2 to compare the collection efficiency of a bisulfate-treated

filter (T) to an untreated filter (U), the average NNK mass obtained on a T filter was 0.88 ± 0.55 (in $\text{ng} \pm \text{SD}$) and 95% confidence interval (0.37–1.39 ng) as compared to the average NNK mass on a U filter which was 0.42 ± 0.29 and 95% confidence interval (0.14–0.69 ng). The ratio of NNK mass collected on T filters to that collected on U filters (T/U) is 2.35 ± 1.29 (mean \pm SD). The regression of the T to U filters data indicates a slope of 1.06 ($p < 0.186$) (see Fig. 3). For nicotine, the average nicotine mass obtained on a T filter was 1719 ± 1108 (in $\text{ng} \pm \text{SD}$) as compared to that on a U filter which was 291 ± 426 ng. The T/U ratio is 15.4 ± 18.8 , as expected due to, primarily, the volatility of the nicotine base. Given the small datasets, a Mann-Whitney U non-parametric test was run for NNK and nicotine mass on T and U filters. The T filters have the highest NNK and nicotine mass from the highest mean ranks and the test statistics indicate that for nicotine, there is a statistically significant difference between the treatments ($p = 0.006$) whilst for NNK there is no statistical difference between the U and the T filters ($p = 0.062$), yet T filters appear to be more efficient in collecting NNK as summarised in Table S1 in the Supporting Information (SI).

Test 3: When untreated filters were exposed to the same atmospheric conditions, the ratio of the average NNK mass on the filter after 48 h to the sum of two 24-hour consecutively sampled filters was 1.17 ± 0.25 ng. From the individual instances it could be noted that in some cases the 48-hour samples had more material than the individual 24-hour samples and in some cases the opposite happened. In principle, the degradation of NNK on the filter is minimal however under specific atmospheric conditions it could not be excluded that the 48-hour filter gave rise to further nitrosation of the collected nicotine or degradation of NNK in comparison to the 24-hour filters that were immediately removed from the sampler and stored away at -20°C .

3.3. Airborne nicotine versus NNK

Aquilina et al. have shown that PM has a ubiquitous percentage of secondhand smoke-derived particulates (0.03–0.08%) that is influenced by population density and also by tobacco smoke prevalence (Aquilina et al., 2021). In order to verify if the airborne NNK is sensitive to these characteristics, namely by comparing to airborne nicotine, samples using bisulfate-treated filters were collected from cities that have different urban fabric and climates as shown in Fig. 4. The figure also shows that NNK in PM is proportional to nicotine even if different PM size fractions were sampled. Msida (MSD) was sampled twice in 2016




Fig. 4. Airborne NNK vs nicotine (both in pgm^{-3}) in Birmingham (BHAM- $\text{PM}_{1.0}$ ■), San Francisco (SF-TSP-●) and Msida (MSD- $\text{PM}_{1.0}$ -◆) in 2016 and Msida (MSD- $\text{PM}_{2.5}$ -▲) in 2018.




Fig. 5. Airborne NNK vs Nicotelline (from untreated filters) in five California sites in 2010.

and 2018 across the colder months (Oct-Dec). The sampler in San Francisco (SF) (Aug-Sep) collected Total Suspended Particulates (TSP) where one would expect higher concentrations of NNK. This was the only site where sampling was done at a much higher elevation from the ground compared to the other locations. The samples in Birmingham (BHAM) were collected in the cold months (Oct-Nov) where NNK was quantified only in two out of fourteen samples. Samples below the LOQ were replaced by $0.5 \times \text{LOQ}$. The concentration of nicotine was also very low (median – 2280 pgm^{-3} and Interquartile Range – 3141 pgm^{-3}) compared to the other cities and thus most probably, loss of nicotine, resulted in NNK values to be below the LOQ. This tentative conclusion is supported by the study by Aquilina et al., where the airborne concentration of nicotelline, was also low in Birmingham, compared to the other cities (Fig. 2(b)). Furthermore, although NNK is degraded by photochemically-produced hydroxyl radicals (Pitts et al., 1978), it does not contain chromophores that absorb at wavelengths $> 290 \text{ nm}$ and therefore would not be expected to be susceptible to direct photolysis by sunlight (National Center for Biotechnology Information, 2021).

Further statistics associated with airborne nicotine and NNK are described in Table S2 in the SI.

3.4. Spatial and seasonal variability of NNK

Samples collected every week in 2010, in five cities in California (Fig. 6 (a)), were accompanied by the 24-hour average PM_{10} gravimetric concentrations. BCA and STS are areas with densely built environment with a population of < 0.5 million inhabitants. SLM and YTD are defined as suburban, whose population increases considerably in the summer periods. CB is a very small, low populated, rural desert village. The sites were exposed to annual mean winds $< 2 \text{ ms}^{-1}$ apart from YTD and CB that were characterised by mean winds typically $> 4 \text{ ms}^{-1}$ throughout the year (Aquilina et al., 2021).

In our previous paper (Aquilina et al., 2021), the untreated filters were also analysed for nicotelline. NNK was compared to nicotelline for each site, as shown in Fig. 5. With reference to Fig. 2, even for untreated filters, the relationship between NNK and the marker nicotelline is appreciably linear and compared to the chamber data certainly some loss of NNK occurs under normal atmospheric conditions. For the cities here studied, the claim that nicotine and NNK concentrations vary linearly (as also noticed in Fig. 4), is a characteristic depending both on population densities and smoking prevalence, is supported by air masses passing over the major and densely populated cities such as Sacramento, San Francisco, San Jose, Fresno and Oakland (Fig. 6(a)) that appear to contribute to higher NNK concentrations in the sites closer to these cities, unlike CB. Further details are provided in Figures S1-S5 in the SI.




Fig. 6. (a) Locations of sites in California within different air basins and (b) NNK (in pgm^{-3}) and PM_{10} concentrations (in μgm^{-3}) and NNK in PM_{10} ($\times 10 \text{ ng/g}$) for different sites in California in 2010.

The same filters were analysed for NNK in PM_{10} . Fig. 6(b) compares the variability of PM_{10} and NNK airborne concentrations and the load of NNK in PM_{10} by location over the entire 2010.

All sites show substantial variability in the gravimetric PM_{10} concentration, not unexpected in the different air basins within the state of California, but only YTD and even more BCA are represented by a higher median. The median NNK airborne concentrations tend to be similar in all cities ($2.24\text{--}3.33 \text{ pgm}^{-3}$) except for CB, where an order of magnitude less was recorded. This observation is not surprising given the very small population located south of the Sequoia National Forest and aside the Mojave Desert (see Table S2, for statistical details).

For the sites SLM, STS and YTD, the load of NNK in PM_{10} is similar whilst for CB it is the lowest given the PM_{10} concentration was similar to the other cities as confirmed in Fig. 6(b), Table S4 in the SI. For BCA, the load of NNK is lower than the three other cities because the median of the PM_{10} concentration was higher and appears that the highest concentrations are more of local origin (Figure S3 in SI). This is also




Fig. 7. Seasonal variability of NNK in PM_{10} ($\times 10$ ng/g) and PM_{10} concentrations (in $\mu g m^{-3}$) for different sites in California in 2010.




Fig. 8. (a) Airborne $PM_{2.5}$ (CHN, MLT) and PM_{10} (UK, USA) concentration (in $\mu g m^{-3}$) and (b) NNK in PM (ng/g) for different sites and countries.

reflected in the airborne concentration of NNK. YTD shows similar characteristics to BCA, but regional transport appears to be responsible from the NE and SE sectors for all analytes. For SLM, regional transport is also responsible for all analyte concentrations. At STS, high concentrations are more of local origin, but for NNK, the inference is less clear. These observations are confirmed in the conditional probability functions (CPF) and polar plots presented in [Figures S1-S5](#) in the SI.

A Kruskal–Wallis H test was run to evaluate whether absolute measured values of NNK and PM₁₀ concentrations and NNK in PM₁₀ differed across sites and seasons. The tests revealed, that in different seasons, insignificant differences ($p = 0.075$) in the airborne NNK concentrations were observed while significant differences ($p < 0.05$) were noticed for airborne PM₁₀ concentrations (JJA, SON > DJF, MAM) and NNK in PM₁₀ (, DJF, MAM > JJA, SON) (Table S3 in SI). However similar to the observations of Farren et al., NNK concentrations ranked higher in SON, DJF compared to MAM, JJA (Farren et al., 2015). The same test revealed significant differences ($p < 0.05$) for the three dependent variables across the five sites.

[Fig. 7](#) shows the seasonal variability of PM₁₀ concentration and load of NNK in PM₁₀ across the same five sites (further statistics are given in Table S3). In all the seasons, the median PM₁₀ concentration is highest in BCA and although it is a populated, busy city, long range transport might explain why the load of NNK in PM₁₀ is lower compared to the other sites. Similar to what was noted in [Fig. 6\(b\)](#), it can be confirmed that in CB, with the only exception of season DJF, although the median PM₁₀ concentration was similar to the other sites, the load of NNK was the lowest here, also confirmed in [Fig. 7](#).

3.5. NNK in different campaigns

[Fig. 8](#) represents the comparison in airborne PM and load of NNK in PM samples obtained from several countries. Using untreated filters, the PM gravimetric samples were collected from UK, Malta and Hong Kong. The sampling site details have been described in our previous paper (Aquilina et al., 2021). As these were relatively short air sampling campaigns, the variability in the concentrations cannot be discussed in direct comparison with the samples obtained in the California sites (USA) that were discussed in [Section 3.4](#). Nevertheless, as the UK, Hong Kong and Malta samples were collected in the period 2014–2018, they are still valuable samples to understand better the distribution of NNK in PM across different urban conglomerations and the fate of nicotine in outdoor conditions and hence the formation of NNK or otherwise.

The samples from Hong Kong show a higher median PM concentration than those from Malta and UK, whilst these two were more comparable to the USA levels ([Fig. 8\(a\)](#)). However [Fig. 8\(b\)](#) indicates that the NNK load represents a different perspective. In Hong Kong, it is understandable that the site in MA has a high median of PM, yet low load of NNK as it is a landfill taking construction waste, whilst TKO and TSW were urban sites located in residential/commercial areas. Apart from a substantial contribution to PM from traffic, it is reasonable to argue that densely populated cities contribute to a higher load of NNK in PM. Similar arguments can be advanced for the site MSD. Although the population in MSD is relatively small, it lies within the concentrated urban area of the whole island, thus the low median and high variability of the load of NNK. As for BHAM, the sampling site was in a suburban area of Birmingham and since very low PM concentrations and low nicotine concentration compared to the other sites were recorded, it is understandable that NNK was not detected, given that untreated filters appear to be less efficient to capture NNK, as discussed in [Section 3.3](#).

4. Conclusion

Following a previous publication indicating that PM is contaminated with tobacco-smoke derived particulates (Aquilina et al., 2021), this paper presents the first comprehensive study to demonstrate the ubiquitous presence of a tobacco-specific, potent lung carcinogen NNK in

airborne PM. The concentrations measured in this study fall well below those reported by (Farren et al., 2015) from the campaign in London, the only study looking for NNK in PM to date. This may be a consequence of different analytical methods and/or highly variable atmospheric concentrations influenced by local sources, possible atmospheric NNK formation and reactivity. Tests have shown that what was quantified is not a sampling artefact but a new addition to the knowledge of the chemical composition of PM. This information has been verified by using a pre-treated bisulfate filter, proved to be easy to prepare, yet innovative in the quest of sampling both gas- and particle-phase nicotine alongside NNK, that is a minor component of tobacco-derived PM. The use of a bisulfate-treated filter is more efficient in collecting NNK amidst the challenges posed by the reactivity of NNK as explained earlier. Using nicotelline as a suitable marker for the tobacco-smoke derived PM, it has been shown that NNK correlates very well with it when using treated filters. Although untreated filters are less efficient, [Fig. 5](#) shows that from five datasets, NNK still correlates appreciably well with the marker and hence the discussion on the NNK levels on untreated filters can be done with confidence. This is important for the use of untreated archived filters.

Analyses of samples gathered from four countries, all having distinct micro-climates, strengthens the claim that tobacco related pollutants increase with tobacco smoke prevalence and population density. Msida is in the middle of the most urbanized area in Malta where adult tobacco smoke prevalence is still relatively high (20%) (World Health Organization, 2019), hence the relatively higher nicotine and NNK levels. In a city like San Francisco, where tobacco smoke prevalence is much lower (12.3% in 2012) (SFTFPM.org, 2018) NNK was detected even on a high building. Birmingham in the UK, although the population is not small and adult smoking prevalence is 18.4% (in 2014) ((Local Government Association, 2018)), lower concentrations of nicotine and corresponding low levels of NNK were quantified only when using treated filters. In Hong Kong, the smoking prevalence is also low (11.4% in 2015) (Hong Kong Council on Smoking and Health, 2018) and although sampling was carried out in distinctly different areas, NNK levels were similar to other cities. From samples knowing the gravimetric PM concentration, the load of NNK could be calculated. This information has shown that while seasonality in PM is associated with several sources, topography and meteorology, the load of NNK is primarily associated with the passage of air masses over densely populated cities where most probably smoking is higher and in the colder months the load of NNK in PM is higher than in warmer months.

The main limitation of this study arises from relatively small datasets. Ideally more samples would be obtained from areas that have smaller populations and also from areas where smoking is known to be still predominant like Asia and Southern Europe in order to better quantify the level of contamination of airborne PM. Further research is necessary to evaluate the health implications of a chronic exposure, by inhalation, to the load of NNK in PM.

CRedit authorship contribution statement

Noel J. Aquilina: Conceptualization, Data curation, Formal analysis, Methodology, Writing – original draft. **Christopher M. Havel:** Formal analysis, Methodology. **Roy M. Harrison:** Data curation, Methodology, Writing – review & editing. **Kin-Fai Ho:** Data curation, Writing – review & editing. **Neal L. Benowitz:** Supervision, Writing – review & editing. **Peyton Jacob III:** Conceptualization, Writing – review & editing.

Declaration of Competing Interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Acknowledgements

The authors would like to thank the following persons in helping out to provide several filters for this study.

Lane Sharon and Brenda Saldana from the California Air Resources Board (CARB).

Polly Cheung from UCSF for helping out in the analytical work.

Maria Thompson from the University of Birmingham, UK.

Sara Fenech and Renato Camilleri from the Faculty of Science, University of Malta, Malta. Measurements in Malta were carried out using the Mobile Air Quality Laboratory within the University of Malta, procured through the European Regional Development Fund, Project ERDF-310 "Expanding the Physics and applied interdisciplinary research capabilities at the Faculty of Science". All funding bodies are acknowledged for their support to carry out this research.

Funding Sources

This work was supported by the California Tobacco-Related Disease Research Program Grant 23PT-0013 - Impacts of Thirdhand Smoke on Public Health (Phase-2), National Institute on Drug Abuse, P30 DA012393, the National Center for Research Resources, S10 RR026437, UCSF Bland Lane Center of Excellence on Secondhand Smoke.

Appendix A. Supplementary material

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.envint.2021.106992>.

References

- Akyüz, M., Ata, Ş., 2013. Seasonal variations of particle-associated nitrosamines by gas chromatography-mass spectrometry in the atmospheric environment of Zonguldak. *Turkey. Environ. Sci. Pollut. Res.* 20 (10), 7398–7412. <https://doi.org/10.1007/s11356-013-1758-y>.
- Apelberg, B.J., Hepp, L.M., Avila-Tang, E., Gundel, L., Hammond, S.K., Hovell, M.F., Hyland, A., Klepeis, N.E., Madsen, C.C., Navas-Acien, A., Repace, J., Samet, J.M., Breyse, P.N., 2013. Environmental monitoring of secondhand smoke exposure. *Tob. Control* 22 (3), 147–155. <https://doi.org/10.1136/tobaccocontrol-2011-050301>.
- Aquilina, N.J., Havel, C.M., Cheung, P., Harrison, R.M., Ho, K.-F., Benowitz, N.L., Jacob III, P., 2021. Ubiquitous atmospheric contamination by tobacco smoke: Nicotine and a new marker for tobacco smoke-derived particulate matter, nicotelline. *Environ. Int.* 150, 106417. <https://doi.org/10.1016/j.envint.2021.106417>.
- Aragón, M., Marcé, R.M., Borrull, F., 2013. Determination of N-nitrosamines and nicotine in air particulate matter samples by pressurised liquid extraction and gas chromatography-ion trap tandem mass spectrometry. *Talanta* 115, 896–901. <https://doi.org/10.1016/j.talanta.2013.07.010>.
- Brown, B.G., Borschke, A.J., Doolittle, D.J., 2003. An Analysis of the Role of Tobacco-Specific Nitrosamines in the Carcinogenicity of Tobacco Smoke. *Nonlinearity Biol. Toxicol. Med.*
- Chao, M.-R., Wang, C.-J., Wu, M.-T., Pan, C.-H., Kuo, C.-Y., Yang, H.-J., Chang, L.W., Hu, C.-W., 2008. Repeated Measurements of Urinary Methylated/Oxidative DNA Lesions, Acute Toxicity, and Mutagenicity in Coke Oven Workers. *Cancer Epidemiol. Biomarkers & Prev.* 17, 3381 LP – 3389. DOI: 10.1158/1055-9965.EPI-08-0721.
- Cogliano, V.J., Baan, R., Straif, K., Grosse, Y., Lauby-Secretan, B., El Ghissassi, F., Bouvard, V., Benbrahim-Tallaa, L., Guha, N., Freeman, C., Galichet, L., Wild, C.P., 2011. Preventable Exposures Associated With Human Cancers. *JNCI J. Natl. Cancer Inst.* 103 (24), 1827–1839. <https://doi.org/10.1093/jnci/djr483>.
- Daisey, J.M., 1999. Tracers for assessing exposure to environmental tobacco smoke: What are they tracing? *Environ. Health Perspect.* 107, 319–327. <https://doi.org/10.2307/3434424>.
- Farren, N.J., Ramirez, N., Lee, J.D., Finessi, E., Lewis, A.C., Hamilton, J.F., 2015. Estimated Exposure Risks from Carcinogenic Nitrosamines in Urban Airborne Particulate Matter. *Environ. Sci. Technol.* 49.
- Ge, X., Wexler, A.S., Clegg, S.L., 2011. Atmospheric amines - Part I. A review. *Atmos. Environ.* 45 (3), 524–546.
- Gunduz, I., Kondylis, A., Jaccard, G., Renaud, J.M., Hofer, R., Ruffieux, L., Gadani, F., 2016. Tobacco-specific N-nitrosamines NNN and NNK levels in cigarette brands between 2000 and 2014. *Regul. Toxicol. Pharmacol.* 76, 113–120. <https://doi.org/10.1016/j.yrtph.2016.01.012>.
- Häger, B., Niessner, R., 1996. Determination of N-nitrosomethylaniline and methylaniline in the gas phase. *Microchim. Acta* 122 (1–2), 35–44. <https://doi.org/10.1007/BF01252403>.
- Hoffmann, D., Brunemann, K.D., Prokopczyk, B., Djordjevic, M.V., 1994. Tobacco-specific N-nitrosamines and areca-derived N-nitrosamines: chemistry, biochemistry, carcinogenicity, and relevance to humans. *J. Toxicol. Environ. Heal.* 41 (1), 1–52.
- Hoffmann, D., Rivenson, A., Amin, S., Hecht, S.S., 1984. Dose-response study of the carcinogenicity of tobacco-specific N-nitrosamines in F344 rats. *J. Cancer Res. Clin. Oncol.* 108 (1), 81–86. <https://doi.org/10.1007/BF00390978>.
- Hoffmann, D., Rivenson, A., Chung, F.-L., Hecht, S.S., 1991. Nicotine-Derived N-Nitrosamines (TSNA) and their Relevance in Tobacco Carcinogenesis. *Crit. Rev. Toxicol.* 21 (4), 305–311. <https://doi.org/10.3109/10408449109017917>.
- Hong Kong Council on Smoking and Health, 2018. *Smoke Free Hong Kong - Annual Report 2017-2018*.
- IARC, 2007. Smokeless tobacco and some tobacco-specific N-nitrosamines. *IARC Monogr. Eval. Carcinog. risks to humans* 89, 1–592.
- Jacob, P., Benowitz, N.L., Shulgin, A.T., 1988. Synthesis of optically pure deuterium-labelled nicotine, nor nicotine and cotinine. *J. Label. Compd. Radiopharm.* 25 (10), 1117–1128. <https://doi.org/10.1002/jlcr.v25:1010.1002/jlcr.2580251009>.
- Jacob, P., Goniewicz, M.L., Havel, C.M., Schick, S.F., Benowitz, N.L., 2013. Nicotelline: A proposed biomarker and environmental tracer for particulate matter derived from tobacco smoke. *Chem. Res. Toxicol.* 26 (11), 1615–1631. <https://doi.org/10.1021/tx400094y>.
- Kavvadias, D., Scherer, G., Urban, M., Cheung, F., Errington, G., Shepperd, J., McEwan, M., 2009. Simultaneous determination of four tobacco-specific N-nitrosamines (TSNA) in human urine. *J. Chromatogr. B* 877 (11–12), 1185–1192. <https://doi.org/10.1016/j.jchromb.2009.03.009>.
- Leaderer, B.P., Hammond, S.K., 1991. Evaluation of Vapor-Phase Nicotine and Respirable Suspended Particle Mass as Markers for Environmental Tobacco Smoke. *Environ. Sci. Technol.* 25 (4), 770–777. <https://doi.org/10.1021/es00016a023>.
- Local Government Association, 2018. *Smoking Prevalence in adults - current smokers (IHS) in Birmingham* [WWW Document]. URL http://ginform.local.gov.uk/reports/Agastandard?mod-metric=3527&mod-area=E08000025&mod-group=AllRegions_England&mod-type=namedComparisonGroup (accessed 2.8.18).
- Marano, R.S., Updegrove, W.S., Machen, R.C., 1982. Determination of trace levels of nitrosamines in air by gas chromatography/low-resolution mass spectrometry. *Anal. Chem.* 54 (12), 1947–1951. <https://doi.org/10.1021/ac00249a008>.
- National Center for Biotechnology Information, 2021. *PubChem Annotation Record for 4-(N-Nitrosomethylamino)-1-(3-pyridyl)-1-butanone*, Source: Hazardous Substances Data Bank (HSDB) [WWW Document]. URL <https://pubchem.ncbi.nlm.nih.gov/substance/hsdb/7771>.
- NIST, 2017, 2017–. (Accessed 17 November 2021).
- Petrick, L., Destailats, H., Zouev, L., Sabach, S., Dubowski, Y., 2010. Sorption, desorption, and surface oxidative fate of nicotine. *Phys. Chem. Chem. Phys.* 12, 10356–10364. <https://doi.org/10.1039/c002643c>.
- Petrick, L.M., Sleiman, M., Dubowski, Y., Gundel, L.A., Destailats, H., 2011a. Tobacco smoke aging in the presence of ozone: A room-sized chamber study. *Atmos. Environ.* 45 (28), 4959–4965. <https://doi.org/10.1016/j.atmosenv.2011.05.076>.
- Petrick, L.M., Svidovsky, A., Dubowski, Y., 2011b. Thirdhand smoke: Heterogeneous oxidation of nicotine and secondary aerosol formation in the indoor environment. *Environ. Sci. Technol.* 45 (1), 328–333.
- Pitts, J.N., Grosjean, D., Van Cauwenbergh, K., Schmid, J.P., Fitz, D.R., 1978. Photooxidation of Aliphatic Amines Under Simulated Atmospheric Conditions: Formation of Nitrosamines, Nitramines, Amides, and Photochemical Oxidant. *Environ. Sci. Technol.* 12 (8), 946–953. <https://doi.org/10.1021/es60144a009>.
- Ramírez, N., Ozel, M.Z., Lewis, A.C., Marcé, R.M., Borrull, F., Hamilton, J.F., 2012. Determination of nicotine and N-nitrosamines in house dust by pressurized liquid extraction and comprehensive gas chromatography-nitrogen chemiluminescence detection. *J. Chromatogr. A* 1219, 180–187. <https://doi.org/10.1016/j.chroma.2011.11.017>.
- Rounbehler, D.P., Reisch, J.W., Coombs, J.R., Fine, D.H., 1980. Nitrosamine air sampling sorbents compared for quantitative collection and artifact formation. *Anal. Chem.* 52 (2), 273–276. <https://doi.org/10.1021/ac50052a014>.
- Schick, S.F., Farraro, K.F., Perrino, C., Sleiman, M., van de Vossenberg, G., Trinh, M.P., Hammond, S.K., Jenkins, B.M., Balmes, J., 2014. Thirdhand cigarette smoke in an experimental chamber: evidence of surface deposition of nicotine, nitrosamines and polycyclic aromatic hydrocarbons and de novo formation of NNK. *Tob. Control* 23. SFTFPM.org, 2018. *San Francisco Tobacco-Free Project Mission. Tobacco related data. Age-standardized total cigarette smoking prevalence, all counties, 1996-2012, USA*. [WWW Document]. URL <http://sanfranciscotobaccofreeproject.org/resources-data/> (accessed 2.8.18).
- Sleiman, M.L., Gundel, L.A., Pankow, J.F., Jacob, P., Singer, B.C., Destailats, H., 2010. Formation of carcinogens indoors by surface-mediated reactions of nicotine with nitrous acid, leading to potential thirdhand smoke hazards. *Proc. Natl. Acad. Sci. U. S. A.* 107, 6576–6581. DOI: 10.1073/pnas.0912820107.
- VandenBoer, T.C., Markovic, M.Z., Sanders, J.E., Ren, X., Pusede, S.E., Browne, E.C., Cohen, R.C., Zhang, L., Thomas, J., Brune, W.H., Murphy, J.G., 2014. Evidence for a nitrous acid (HONO) reservoir at the ground surface in Bakersfield, CA, during CalNex 2010. *J. Geophys. Res. Atmos.* 119, 9093–9106. DOI: DOI: 10.1002/2013JD020971.
- Whitehead, Todd P., Havel, Christopher, Metayer, Catherine, Benowitz, Neal L., Jacob, Peyton, 2015. Tobacco Alkaloids and Tobacco-Specific Nitrosamines in Dust from Homes of Smokeless Tobacco Users, Active Smokers, and Nontobacco Users. *Chem. Res. Toxicol.* 28 (5), 1007–1014. <https://doi.org/10.1021/acs.chemrestox.5b00040>.
- Whitlatch, A., Schick, S., 2019. Thirdhand Smoke at Philip Morris. *Nicotine Tob. Res.* 21, 1680–1688. <https://doi.org/10.1093/ntnr/nty153>.
- World Health Organization, 2019. *WHO report on the global tobacco epidemic, 2019*, Geneva: World Health Organization.