

The PROTEUS-Trials Consortium

Snyder, Claire; Crossnohere, Norah L. ; King, Madeleine; Reeve, Bryce B.; Bottomley, Andrew; Calvert, Melanie; Thorner, Elissa ; Wu, Albert W ; Brundage, Michael; PROTEUS-Trials Consortium

DOI:

[10.1177/17407745221077691](https://doi.org/10.1177/17407745221077691)

License:

Creative Commons: Attribution-NonCommercial (CC BY-NC)

Document Version

Publisher's PDF, also known as Version of record

Citation for published version (Harvard):

Snyder, C, Crossnohere, NL, King, M, Reeve, BB, Bottomley, A, Calvert, M, Thorner, E, Wu, AW, Brundage, M & PROTEUS-Trials Consortium 2022, 'The PROTEUS-Trials Consortium: optimizing the use of patient-reported outcomes in clinical trials', *Clinical Trials*, vol. 19, no. 3, pp. 277-284.
<https://doi.org/10.1177/17407745221077691>

[Link to publication on Research at Birmingham portal](#)

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

The PROTEUS-Trials Consortium: Optimizing the use of patient-reported outcomes in clinical trials

Clinical Trials

1–8

© The Author(s) 2022




Article reuse guidelines:

sagepub.com/journals-permissions

DOI: 10.1177/17407745221077691

journals.sagepub.com/home/ctj

Claire Snyder^{1,2,3}  Norah Crossnohere⁴ Madeleine King⁵ Bryce B Reeve⁶
Andrew Bottomley⁷ Melanie Calvert^{8,9,10,11,12} Elissa Thorner^{1,3}
Albert W Wu^{1,2} and Michael Brundage¹³; for the PROTEUS-Trials
Consortium

Abstract

Background: The assessment of patient-reported outcomes in clinical trials has enormous potential to promote patient-centred care, but for this potential to be realized, the patient-reported outcomes must be captured effectively and communicated clearly. Over the past decade, methodologic tools have been developed to inform the design, analysis, reporting, and interpretation of patient-reported outcome data from clinical trials. We formed the PROTEUS-Trials Consortium (Patient-Reported Outcomes Tools: Engaging Users and Stakeholders) to disseminate and implement these methodologic tools.

Methods: PROTEUS-Trials are engaging with patient, clinician, research, and regulatory stakeholders from 27 organizations in the United States, Canada, Australia, the United Kingdom, and Europe to develop both organization-specific and cross-cutting strategies for implementing and disseminating the methodologic tools. Guided by the Knowledge-to-Action framework, we conducted consortium-wide webinars and meetings, as well as individual calls with participating organizations, to develop a workplan, which we are currently executing.

Results: Six methodologic tools serve as the foundation for PROTEUS-Trials dissemination and implementation efforts: the Standard Protocol Items: Recommendations for Interventional Trials-patient-reported outcome extension for writing protocols with patient-reported outcomes, the International Society for Quality of Life Research Minimum Standards for selecting a patient-reported outcome measure, Setting International Standards in Analysing Patient-Reported Outcomes and Quality of Life Endpoints Data Consortium recommendations for patient-reported outcome data analysis, the Consolidated Standards for Reporting of Trials-patient-reported outcome extension for reporting clinical trials with patient-reported outcomes, recommendations for the graphic display of patient-reported outcome data, and a Clinician's Checklist for reading and using an article about patient-reported outcomes. The PROTEUS-Trials website (www.TheProteusConsortium.org) serves as a central repository for the methodologic tools and associated resources.

¹Johns Hopkins School of Medicine, Baltimore, MD, USA

²Johns Hopkins Bloomberg School of Public Health, Baltimore, MD, USA

³Sidney Kimmel Comprehensive Cancer Center, Johns Hopkins University, Baltimore, MD, USA

⁴College of Medicine, Ohio State University, Columbus, OH, USA

⁵School of Psychology, The University of Sydney, Sydney, NSW, Australia

⁶School of Medicine, Duke University, Durham, NC, USA

⁷European Organisation for the Research and Treatment of Cancer, Brussels, Belgium

⁸Birmingham Health Partners Centre for Regulatory Science and Innovation, University of Birmingham, Birmingham, UK

⁹Centre for Patient Reported Outcomes Research, Institute of Applied Health Research, University of Birmingham, Birmingham, UK

¹⁰National Institute for Health Research (NIHR) Birmingham Biomedical Research Centre, University of Birmingham, Birmingham, UK

¹¹NIHR Applied Research Collaboration West Midlands, University of Birmingham, Birmingham, UK

¹²NIHR Surgical Reconstruction and Microbiology Research Centre University Hospitals Birmingham NHS Foundation Trust and University of Birmingham, Birmingham, UK

¹³Queen's Cancer Research Institute, Queen's University, Kingston, ON, Canada

Corresponding author:

Claire Snyder, Johns Hopkins School of Medicine, 624 N. Broadway, 6th Floor, Baltimore, MD 21205, USA.

Email: csnyder@jhu.edu

To date, we have developed (1) a roadmap to visually display where each of the six methodologic tools applies along the clinical trial trajectory, (2) web tutorials that provide guidance on the methodologic tools at different levels of detail, (3) checklists to provide brief summaries of each tool's recommendations, (4) a handbook to provide a self-guided approach to learning about the tools and recommendations, and (5) publications that address key topics related to patient-reported outcomes in clinical trials. We are also conducting organization-specific activities, including meetings, presentations, workshops, and webinars to publicize the existence of the methodologic tools and the PROTEUS-Trials resources. Work to develop communications strategies to ensure that PROTEUS-Trials reach key audiences with relevant information about patient-reported outcomes in clinical trials and PROTEUS-Trials is ongoing.

Discussion: The PROTEUS-Trials Consortium aims to help researchers generate patient-reported outcome data from clinical trials to (1) enable investigators, regulators, and policy-makers to take the patient perspective into account when conducting research and making decisions; (2) help patients understand treatment options and make treatment decisions; and (3) inform clinicians' discussions with patients regarding treatment options. In these ways, the PROTEUS Consortium promotes patient-centred research and care.

Keywords

Patient-reported outcomes, clinical trials, reporting methods, protocols, measure selection, data visualization

Background

Patients, clinicians, regulators, and policy-makers value data on patient-reported outcomes (PROs), such as symptoms, functioning, and health-related quality of life, from clinical trials to inform decision-making.^{1–8} To be most informative, the PRO methods need to be specified appropriately, the PRO endpoints measured effectively, the PRO data analysed properly, and the PRO results reported clearly to multiple stakeholders. However, there is evidence that these goals are frequently not met at the protocol, analysis, reporting, or application levels.^{6,9–13}

Methodologic tools have been developed to help clinical trialists optimize the capture and communication of PROs.^{14–19} These tools, developed in partnership with patients and other stakeholders, provide guidance on designing the PRO aspects of clinical trials, collecting and analysing the PRO data, and interpreting and reporting the PRO findings. However, these tools require a coordinated, stakeholder-driven implementation and dissemination strategy to ensure their usefulness in practice.

In 2018, we formed the PROTEUS Consortium (Patient-Reported Outcomes Tools: Engaging Users & Stakeholders), first with a focus on advancing the use of PROs in clinical research (PROTEUS-Trials) and, subsequently, expanded to address the use of PROs in clinical practice (PROTEUS-Practice).²⁰ PROTEUS-Trials are partnering with stakeholders to promote the application of existing tools to optimize the capture and communication of PROs in clinical trials. The goal is to give researchers guidance to help them conduct their studies so that patients can make better-informed health decisions; regulators and policy-makers can take the patient perspective into account in their deliberations; and clinicians can have reliable, relevant, and

interpretable PRO data to discuss treatment options with patients. Stated differently, the PROTEUS-Trials Consortium aims to promote patient-centred research and care at multiple levels.

Methods

The work of PROTEUS-Trials is guided by the Knowledge-to-Action framework, an implementation and dissemination, or knowledge translation, model with two parts.²¹ First is knowledge development, which creates knowledge tools and products through inquiry and synthesis. Second is the action cycle, which encourages dissemination of the knowledge tools by identifying knowledge gaps, adapting the knowledge to the local context, assessing facilitators and barriers to knowledge use, and implementing tailored interventions. The cycle continues with monitoring and sustaining knowledge use while evaluating outcomes.

In the case of PROTEUS-Trials, the knowledge tools are the existing methodologic tools for integrating PROs in clinical trials, mentioned above and described in more detail below.^{14–19} Therefore, PROTEUS-Trials focuses primarily on the action cycle. First, the Principal Investigators (C.S. and M.B.), Project Manager (N.C.), and other Steering Committee members (M.K., B.B.R., A.B., M.C., E.T., and A.W.W.) who played key roles in developing the PRO methodologic tools assembled the consortium, engaging relevant patient, clinician, research, and regulatory groups. Second, a consortium kick-off webinar reviewed PROTEUS' objectives and approach. Third, we held calls with representatives from each organization to obtain their input on knowledge translation strategies specific to their contexts and needs, and more generally. Fourth, we held an in-person meeting (June 2019)

Table 1. Organizations with PROTEUS-Trials participants.^a

AcademyHealth
American Cancer Society
American Society of Clinical Oncology
American Society for Radiation Oncology
Australian Clinical Trials Alliance
Canadian Association of Radiation Oncology
Cancer Australia
Consolidated Standards for Reporting of Trials (CONSORT)
Critical Path Institute PRO Consortium
European Medicines Agency-Scientific Advice Working Party/
Dutch Medicines Evaluation Board
European Organisation for the Research and Treatment of
Cancer
Food and Drug Administration
Health Canada
Industry (GlaxoSmithKline)
International Society for Quality of Life Research
ISPOR
Journal editor perspective
Medicines and Healthcare Products Regulatory Agency
National Cancer Institute
National Cancer Research Institute
National Clinical Trials Network PRO representatives
National Coalition for Cancer Survivorship
National Institute for Health and Care Excellence
Oncology Nursing Society
Patient-Centered Outcomes Research Institute
Society for Clinical Trials
Standard Protocol Items: Recommendations for Interventional
Trials (SPIRIT)

^aParticipating in PROTEUS does not imply endorsement of any particular PRO tools or guidance documents.

to develop and prioritize cross-cutting knowledge translation strategies. Fifth and ongoing, we are executing the organization-specific and cross-cutting knowledge translation strategies. The consortium has a cancer focus because some of the methodologic tools were developed in oncology contexts; however, the expectation is that this work is largely applicable across health conditions.

Results

Consortium formation

There are 27 organizations with participants in the PROTEUS-Trials Consortium (Table 1). Membership includes patient and clinician advocacy groups, government and regulatory agencies, clinical trial cooperative groups, organizations focused on research methods, and funding agencies.

Methodologic tools

The consortium's knowledge translation activities focus on these six methodologic tools, which are summarized in Figure 1.

Guidelines for inclusion of PROs in clinical trial protocols: the Standard Protocol Items: Recommendations for Interventional Trials-PRO extension. The Standard Protocol Items: Recommendations for Interventional Trials (SPIRIT)-PRO extension recommends best practices for writing the PRO aspects of randomized controlled trial protocols.¹⁴ It extends the 2013 SPIRIT guidance that identifies the minimum elements generally required in clinical trial protocols.²² The SPIRIT-PRO extension builds on the SPIRIT guidance by addressing the minimum elements related to PROs that should be included in clinical trial protocols. SPIRIT-PRO was developed through a Delphi process and consensus meeting, which was informed by a list of PRO-specific protocol items generated from a systematic review of existing guidance and a survey of international stakeholders. The resulting recommendations provide guidance on PRO issues relating to the trial rationale, objectives, eligibility criteria, concepts used to evaluate the intervention, timepoints for assessment, PRO instrument selection and measurement properties, data collection plan, translation to other languages, proxy completion, strategies to minimize missing data, and whether PRO data will be monitored during the study to inform clinical care.

International Society for Quality of Life Research minimum standards for PRO measures used in patient-centred outcomes and comparative effectiveness research. In 2013, the International Society for Quality of Life Research (ISOQOL) conducted a project to recommend minimum standards for PRO measures in patient-centred outcomes and comparative effectiveness research.¹⁵ To develop these standards, the ISOQOL Task Force reviewed existing guidelines for the selection of PRO measures and surveyed ISOQOL members to obtain their input on potential PRO standards. The final recommendations, which helped to inform the Patient-Centered Outcomes Research Institute's methodology standards related to PROs,²³ address documentation of the conceptual and measurement model for the PRO measure; evidence for reliability and validity; interpretability of scores; translation quality; and patient and investigator burden.

Setting International Standards in Analysing PROs and Quality of Life Endpoints Data Consortium. The European Organisation for the Research and Treatment of Cancer (EORTC) formed the Setting International Standards in Analysing PROs and Quality of Life Endpoints Data (SISAQOL) Consortium to establish international standards in analysing PRO endpoint data.¹⁶ The consortium includes researchers, statisticians, clinician researchers, regulators, patients, and stakeholders with multidisciplinary expertise and international perspectives. The goal is to set standards that

Tool	Clinical Trial Stage	Rationale	Objective	Approach
Guidelines for Inclusion of Patient-Reported Outcomes in Clinical Trial Protocols: The SPIRIT-PRO Extension ¹⁴	Design	To ensure that critical aspects of the PRO substudy are included in the trial protocol for successful conduct	<ul style="list-style-type: none"> To provide international, consensus-based, PRO-specific protocol guidance To address existing weaknesses related to PROs, including poor quality protocols, data, and reporting 	Recommends items to address in clinical trial protocols where PROs are primary or key secondary outcomes
ISOQOL Minimum Standards for PRO Measures Used in Patient-Centered Outcomes and Comparative Effectiveness Research ¹⁵	Design	PROs must be measured in a valid, standardized way using appropriate tools and methods to ensure valid conclusions	<ul style="list-style-type: none"> To develop PRO measure minimum standards for the design and selection of a PRO measure for use in patient-centered outcomes research (PCOR) and comparative effectiveness research (CER) The standards represent the minimum criteria required for a PRO measure to be judged suitable for a PCOR or CER study These standards are intended to promote the appropriate use of PRO measures in PCOR and CER, which in turn can improve the effectiveness and efficiency of healthcare delivery 	Provides guidance for selecting PRO measures for use in patient-centered and comparative effectiveness research
Setting International Standards in Analysing Patient-Reported Outcomes and Quality of Life Endpoints Data (SISAQOL) Consortium ¹⁶	Analysis	To ensure a consistent and methodologically appropriate PRO data analysis	<ul style="list-style-type: none"> To recommend approaches to address issues related to multidimensional outcomes, longitudinal data analysis, and missing data To promote clarity in specifying analytic objectives To standardize terminology 	Recommends statistical approaches for analyzing PRO data
Reporting of Patient-Reported	Reporting	To ensure that the PRO	<ul style="list-style-type: none"> To provide international, consensus-based, PRO-specific reporting guidance 	Identifies the relevant information to include
Recommendations for Graphically Displaying PRO Data ¹⁸	Reporting	To promote consistent presentation of PRO data so that clinicians and patients can understand what the PRO scores mean	<ul style="list-style-type: none"> To address the challenges patients and clinicians report when trying to interpret the meaning and implications of PRO data To create standardized PRO graphic displays to minimize the confusion created by variation in how PRO measures are scored and scaled 	Provides evidence-based recommendations for presenting PRO data clearly to patients and clinicians/researchers
Clinician's Checklist for Reading and Using an Article about Patient-Reported Outcomes ¹⁹	Application	To help clinicians assess the quality of PRO research studies and determine whether findings are useful for clinical practice	<ul style="list-style-type: none"> To address barriers to clinicians' application of PRO results, including a lack of training, variation in PRO measures, and inconsistency in how PRO results are reported To help practicing clinicians apply results of clinical research studies that include PROs in their patient care 	Provides a checklist to evaluate the quality of studies that use PROs

Figure 1. Methodologic tools for implementing patient-reported outcomes (PROs) in clinical trials.

are methodologically rigorous, comprehensive, and practical. Initial SISAQOL recommendations focus on establishing the research objectives a priori, distinguishing between domains with confirmatory analyses (superiority, non-inferiority, and equivalence) versus those that are exploratory/descriptive. The recommendations also address specifying the assumptions, endpoints of interest, recommended statistical models, and handling missing data. The initial SISAQOL work is continuing through the SISAQOL-IMI (Innovative Medicines Initiative) Consortium.²⁴

Reporting of PROs in randomized trials: the Consolidated Standards of Reporting Trials-PRO extension. The CONSORT (Consolidated Standards of Reporting Trials) guidance provides recommendations for publications reporting clinical trial results.²⁵ In 2013, a PRO-specific extension was published, which includes recommendations for identifying the PRO as an outcome in the abstract, providing the background and rationale for PRO assessment, describing the PRO hypothesis, providing evidence of the PRO measure's reliability and validity, detailing the mode of PRO




Figure 2. PROTEUS-Trials roadmap.

completion, describing missing PRO data rates and statistical methods, reporting baseline PRO data and results for each domain and timepoint pre-specified for analysis, and discussing the PRO-specific limitations and generalizability.¹⁷

Recommendations for graphically displaying PRO data. A specific issue related to reporting PRO results is the best way to graphically report the data so that patients and clinicians can easily and accurately interpret the PRO findings.¹⁸ Patients and clinicians value the information provided by PRO results from clinical trials, but due to variations in how PRO measures are scored and scaled, and the different approaches to analysing them (e.g. modelling scores over time and proportions meeting a responder definition (i.e. improved/stable/worsened)), they have also reported challenges interpreting PRO results.⁶ To address these issues, a multi-phase research study explored approaches for displaying clinical trial PRO results graphically to identify formats that are interpreted most accurately and rated clearest.^{26–30} These results informed the development of stakeholder-engaged, evidence-driven recommendations for how to display PRO data to promote understanding and use.¹⁸

Clinician's Checklist for reading and using an article about PROs. As noted above, clinicians value the PRO information from clinical trials, but they also report challenges interpreting the findings so that they can use them in practice.⁶ Another tool developed to address this issue is the Clinician's Checklist for reading and using an article about PROs.¹⁹ The checklist includes five questions: (1) Was the PRO assessment strategy appropriate? (2) Did they measure PRO effectively? (3) Should I believe the results? (4) Were the results placed in clinical context? (5) Do the results apply to my

patients? For each question, the checklist provides a description of what to look for in the article, guiding clinicians who are not expert in PRO research on how to use PRO findings in their practice.

Workplan

Based on our consultations with the PROTEUS-Trials participating organizations individually, and then jointly at the in-person meeting, we developed a prioritized knowledge translation workplan, including the below activities.

Website. www.TheProteusConsortium.org includes background on PROTEUS-Trials, links to the six methodologic tools, and the resources developed to-date, including those listed below.

Roadmap. We created a visual display of the clinical trial continuum and where each of the six methodologic tools apply (Figure 2).

Web tutorials. The PROTEUS-Trials Steering Committee members who developed the tools recorded a series of presentations on PROs, PROTEUS-Trials, and the PROTEUS-Trials tools. High-level presentations are directed at, for example, principal investigators who want to ensure that the researchers responsible for the PRO aspects follow best practices. Intermediate-level presentations provide sufficient detail for someone who is reviewing the PRO aspects of a trial (e.g. a manuscript peer-reviewer), but not someone implementing the recommendations directly. Advanced-level presentations are directed to the researchers responsible for implementing the recommendations themselves.

Checklists. A checklist, with citations, for each of the six tools helps users follow the guidance document recommendations without having to refer to lengthy, academic publications.

Handbook. To complement the web tutorials, a handbook provides a self-guided approach to learning about the tools and recommendations.

Publications. Several papers were prioritized to advance the PROTEUS-Trials objectives. For example, we conducted a project comparing different guidance documents' recommendations for PRO measure selection.³¹ We also developed recommendations for grant applicants and reviewers regarding the key information to include/review in grant proposals.³² A paper that highlights some of the 'greatest hits' (focused on value-added) of PROs in clinical trials is forthcoming.

Meetings/presentations/workshops/webinars. Most of our organization-specific strategies involve meetings, presentations, workshops, and webinars to key groups. We have presented on PROTEUS on over 35 occasions to a variety of audiences, in-person and virtually. These presentations are a critical aspect of engaging with different stakeholders and reaching relevant audiences to disseminate information about the tools.

Communication strategy. We are developing communications strategies to further publicize our work, ensure we reach key audiences, and promote the consortium's objectives.

Discussion

The assessment of PROs in clinical trials has enormous potential to promote patient-centred care, but for this potential to be realized, PRO data must be captured effectively and communicated clearly to diverse audiences. The PROTEUS-Trials Consortium aims to engage with stakeholders to develop organization-specific and cross-cutting strategies for implementing and disseminating methodologic tools for optimizing the use of PROs in clinical trials. It is always advisable to include a researcher with PRO expertise on the clinical trial team. This article summarizes the tools and resources available to these PRO experts, as well as others who engage with PROs at a higher level (e.g. peer-reviewers and principal investigators). Future work will evaluate the impact of PROTEUS-Trials on the capture and communication of PROs from clinical trials.

While the PROTEUS-Trials Consortium has focused on six core tools, users are encouraged to consider other guidance as applicable to their purpose and

jurisdiction. For example, if planning a regulatory submission to the Food and Drug Administration, its PRO guidance would be most applicable.¹ Reassuringly, the PROTEUS paper comparing various guidance documents regarding PRO measure selection found general consistency in their recommendations.³¹

In summary, through the PROTEUS-Trials Consortium, we hope that researchers will be better able to generate PRO data from clinical trials to (1) enable investigators, regulators, and policy-makers to take the patient perspective into account when conducting research and making decisions; (2) help patients understand treatment options and make treatment decisions; and (3) inform clinicians' discussions with patients regarding treatment options. In these ways, the PROTEUS Consortium aims to promote patient-centred research and care.

Declaration of conflicting interests


The author(s) declared the following potential conflicts of interest with respect to the research, authorship, and/or publication of this article: C.S., M.B., and N.C. report an unrestricted grant from Pfizer for a separate PROTEUS project, for which B.B.R., M.C., E.T., and A.W.W. also receive payments as Steering Committee members. C.S., A.W.W., and M.B. have had previous funding from Genentech outside of the submitted work. C.S. has received consulting fees from Janssen, via Health Outcomes Solutions. E.T. has received funding from Pfizer for a separate project. M.C. reports personal fees from Astellas, from Takeda, Glaukos, Merck, Daiichi Sankyo, CIS Oncology, Aparito Ltd., and GSK, and grants from Health Data Research UK, Wellcome Trust, Alan Turing Institute, Research England, and UK Research and Innovation outside the submitted work; M.C. is a National Institute for Health Research (NIHR) senior investigator and received funding from the NIHR Birmingham Biomedical Research Centre, NIHR Surgical Reconstruction and Microbiology Research Centre and NIHR ARC West Midlands at the University of Birmingham and University Hospitals Birmingham National Health Service Foundation Trust, Health Data Research UK, Innovate UK (part of UK Research and Innovation), Macmillan Cancer Support, and UCB Pharma. A.W.W. reports consulting fees from ViiV, Osmotica, Pfizer, GSK, Gilead, and Johnson & Johnson. A.B. is a member of the EORTC; he has research grants from BI, Pfizer, and BMS to support the EORTC. No other authors have conflicts to report. Participating in PROTEUS does not imply endorsement of any particular PRO tools or guidance documents. The views expressed in this publication are those of the author(s) and not necessarily those of the NIHR or the Department of Health and Social Care.

Funding

The author(s) disclosed receipt of the following financial support for the research, authorship, and/or publication of this article: PROTEUS-Trials have received funding through an Engagement Award from the Patient-Centered Outcomes Research Institute (12565-JHU) and through an unrestricted grant from Genentech, a Roche Company. The funders had

no role in the study design, collection, analysis, and interpretation of data, or writing of the report; there were no restrictions regarding publication.

ORCID iD

Claire Snyder  <https://orcid.org/0000-0001-8952-4561>

References

1. U.S. Food and Drug Administration. Guidance for industry: patient reported outcome measures: use in medical product development to support labeling claims. *Federal Register* 2009; 74(35): 65132–65133.
2. Acquadro C, Berzon R, Dubois D, et al. Incorporating the patient's perspective into drug development and communication: an ad hoc task force report of the Patient-Reported Outcomes (PRO) Harmonization Group meeting at the Food and Drug Administration, February 16, 2001. *Value Health* 2003; 6(5): 522–531.
3. Au H-J, Ringash J, Brundage M, et al. Added value of health-related quality of life measurement in cancer clinical trials: the experience of the NCIC CTG. *Expert Rev Pharmacoecon Outcomes Res* 2010; 10(2): 119–128.
4. Till JE, Osoba D, Pater JL, et al. Research on health-related quality of life: dissemination into practical applications. *Qual Life Res* 1994; 3(4): 279–283.
5. Lipscomb J, Gotay CC, Snyder C, et al. *Outcomes assessment in cancer: measures, methods, and applications*. Cambridge: Cambridge University Press, 2005.
6. Brundage M, Bass B, Jolie R, et al. A knowledge translation challenge: clinical use of quality of life data from cancer clinical trials. *Qual Life Res* 2011; 20(7): 979–985.
7. Bezjak A, Ng P, Skeel R, et al. Oncologists' use of quality of life information: results of a survey of Eastern Cooperative Oncology Group physicians. *Qual Life Res* 2001; 10(1): 1–13.
8. Kluetz PG, O'Connor DJ and Soltys K. Incorporating the patient experience into regulatory decision making in the USA, Europe, and Canada. *Lancet Oncol* 2018; 19(5): e267–e274.
9. Kyte D, Retzer A, Ahmed K, et al. Systematic evaluation of patient-reported outcome protocol content and reporting in cancer trials. *J Natl Cancer Inst Monogr* 2019; 111(11): 1170–1178.
10. Efficace F, Fayers P, Pusic A, et al. Quality of patient-reported outcome reporting across cancer randomized controlled trials according to the CONSORT patient-reported outcome extension: a pooled analysis of 557 trials. *Cancer* 2015; 121(18): 3335–3342.
11. Mercieca-Bebber R, Friedlander M, Calvert M, et al. A systematic evaluation of compliance and reporting of patient-reported outcome endpoints in ovarian cancer randomised controlled trials: implications for generalisability and clinical practice. *J Patient Rep Outcomes* 2017; 1(1): 5.
12. Bylicki O, Gan HK, Joly F, et al. Poor patient-reported outcomes reporting according to CONSORT guidelines in randomized clinical trials evaluating systemic cancer therapy. *Ann Oncol* 2014; 26(1): 231–237.
13. St Germain D, Denicoff A, Torres A, et al. Reporting of health-related quality of life endpoints in National Cancer Institute-supported cancer treatment trials. *Cancer* 2020; 126(11): 2687–2693.
14. Calvert M, Kyte D, Mercieca-Bebber R, et al. Guidelines for inclusion of patient-reported outcomes in clinical trial protocols: the SPIRIT-PRO extension. *JAMA* 2018; 319: 483–494.
15. Reeve BB, Wyrwich KW, Wu AW, et al. ISOQOL recommends minimum standards for patient-reported outcome measures used in patient-centered outcomes and comparative effectiveness research. *Qual Life Res* 2013; 22(8): 1889–1905.
16. Coens C, Pe M, Dueck AC, et al. International standards for the analysis of quality of life and patient reported outcomes endpoints in cancer randomised controlled trials: recommendations based on critical reviews of the literature and international multi-expert, multi-stakeholder collaborative process. *Lancet Oncol* 2020; 21: e83–96.
17. Calvert M, Blazeby J, Altman DG, et al. Reporting of patient-reported outcomes in randomized trials: the CONSORT PRO extension. *JAMA* 2013; 309: 814–822.
18. Snyder C, Smith K, Holzner B, et al. Making a picture worth a thousand numbers: recommendations for graphically displaying patient-reported outcomes data. *Qual Life Res* 2019; 28(2): 345–356.
19. Wu AW, Bradford AN, Velanovich V, et al. Clinician's checklist for reading and using an article about patient-reported outcomes. *Mayo Clin Proc* 2014; 89(5): 653–661.
20. The PROTEUS Consortium: patient-reported outcomes tools – engaging users & stakeholders, www.TheProteusConsortium.org (accessed 30 July 2021).
21. Straus SE, Tetroe JM and Graham ID. Knowledge translation is the use of knowledge in health care decision making. *J Clin Epidemiol* 2011; 64: 6–10.
22. Chan AW, Tetzlaff JM, Altman DG, et al. SPIRIT 2013 statement: defining standard protocol items for clinical trials. *Ann Intern Med* 2013; 158: 200–207.
23. Patient-Centered Outcomes Research Institute. Methodology standards associated with patient centeredness, <https://www.pcori.org/research-results/about-our-research/research-methodology/pcori-methodology-standards#Associated%20with%20Patient-Centeredness> (accessed 30 July 2021).
24. SISAQOL-IMI: setting international standards of patient-reported outcomes quality of life endpoints in cancer clinical trials – IMI, <https://www.sisaqol-imi.org/> (accessed 21 December 2021).
25. Schulz KF, Altman DG, Moher D, et al. CONSORT 2010 statement: updated guidelines for reporting parallel group randomised trials. *BMJ* 2010; 340: c332.
26. Brundage MD, Smith KC, Little EA, et al. Communicating patient-reported outcome scores using graphic formats: Results from a mixed methods evaluation. *Qual Life Res* 2015; 24(10): 2457–2472.
27. Smith KC, Brundage MD, Tolbert E, et al. Engaging stakeholders to improve presentation of patient-reported outcomes data in clinical practice. *Support Care Cancer* 2016; 24(10): 4149–4157.

28. Brundage MD, Blackford A, Tolbert E, et al. Presenting comparative study PRO results to clinicians and researchers: beyond the eye of the beholder. *Qual Life Res* 2018; 27(1): 75–90.
29. Tolbert E, Brundage M, Bantug E, et al. Picture this: presenting longitudinal patient-reported outcome research study results to patients. *Med Decis Making* 2018; 38(8): 994–1005.
30. Tolbert E, Brundage M, Bantug E, et al. In proportion: approaches for displaying patient-reported outcome research study results as percentages responding to treatment. *Qual Life Res* 2019; 28(3): 609–620.
31. Crossnohere NL, Brundage M, Calvert MJ, et al. International guidance on the selection of patient-reported outcome measures in clinical trials: a review. *Qual Life Res* 2021; 30(1): 21–40.
32. Snyder C, Gilbert A, Moher D, et al. Recommendations for including or reviewing patient-reported outcome endpoints in grant applications. *BMJ* 2021; 373: n1367.