

Global perspectives on autism acceptance, camouflaging behaviours and mental health in autism spectrum disorder

Keating, Connor Tom; Hickman, Lydia; Geelhand, Philippine; Takahashi, Toru; Leung, Joan; Schuster, Bianca; Rybicki, Alicia; Girolamo, Teresa Marie; Clin, Elise; Papastamou, Fanny; Belenger, Marie; Eigsti, Inge-Marie; Cook, Jennifer Louise; Kosaka, Hirotaka; Osu, Rieko; Okamoto, Yuko; Sowden, Sophie

DOI:

[10.1371/journal.pone.0261774](https://doi.org/10.1371/journal.pone.0261774)

License:

Creative Commons: Attribution (CC BY)

Document Version

Publisher's PDF, also known as Version of record

Citation for published version (Harvard):

Keating, CT, Hickman, L, Geelhand, P, Takahashi, T, Leung, J, Schuster, B, Rybicki, A, Girolamo, TM, Clin, E, Papastamou, F, Belenger, M, Eigsti, I-M, Cook, JL, Kosaka, H, Osu, R, Okamoto, Y & Sowden, S 2021, 'Global perspectives on autism acceptance, camouflaging behaviours and mental health in autism spectrum disorder: A registered report protocol', *PLoS ONE*, vol. 16, no. 12, e0261774. <https://doi.org/10.1371/journal.pone.0261774>

[Link to publication on Research at Birmingham portal](#)

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy


While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

REGISTERED REPORT PROTOCOL

Global perspectives on autism acceptance, camouflaging behaviours and mental health in autism spectrum disorder: A registered report protocol

Connor Tom Keating^{1*}, Lydia Hickman¹, Philippine Geelhand², Toru Takahashi³, Joan Leung⁴, Bianca Schuster¹, Alicia Rybicki¹, Teresa Marie Girolamo⁵, Elise Clin², Fanny Papastamou², Marie Belenger², Inge-Marie Eigsti⁵, Jennifer Louise Cook¹, Hirotaka Kosaka⁶, Rieko Osu³, Yuko Okamoto⁷, Sophie Sowden¹



This is a Registered Report and may have an associated publication; please check the article page on the journal site for any related articles.

1 School of Psychology, University of Birmingham, Birmingham, United Kingdom, **2** ACTE (Autism in Context: Theory and Experiment) at LaDisco (Center for Linguistics Research) and ULB Neuroscience Institute, Université libre de Bruxelles, Brussels, Belgium, **3** Faculty of Human Sciences, Waseda University, Tokorozawa, Japan, **4** School of Psychology, University of Auckland, Auckland, New Zealand, **5** Department of Psychological Sciences, University of Connecticut, Storrs, Mansfield, Connecticut, United States of America, **6** Department of Neuropsychiatry, University of Fukui, Fukui, Japan, **7** Waseda Institute for Advanced Study, Waseda University, Tokyo, Japan

* CXK655@bham.ac.uk

OPEN ACCESS

Citation: Keating CT, Hickman L, Geelhand P, Takahashi T, Leung J, Schuster B, et al. (2021) Global perspectives on autism acceptance, camouflaging behaviours and mental health in autism spectrum disorder: A registered report protocol. PLoS ONE 16(12): e0261774. <https://doi.org/10.1371/journal.pone.0261774>

Editor: Robert Didden, Radboud University, NETHERLANDS

Received: July 9, 2021

Accepted: December 9, 2021

Published: December 30, 2021

Copyright: © 2021 Keating et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: All relevant data is available on the Open Science Framework: <https://doi.org/10.17605/OSF.IO/Q8TEW>.

Funding: This project was supported by the Universitas 21 Researcher Resilience Fund and the Medical Research Council (MRC, United Kingdom) MR/R015813/1.

1. Study information

1.1. Description (optional)

Autism spectrum disorder (hereafter ‘autism’) is a neurodevelopmental condition that is characterised by difficulties in social communication and restricted and repetitive interests [1]. Throughout life, autistic people (‘identity-first’ terminology is used throughout in line with the preferences of the autistic community [2] and guidance for avoiding ableist language [3]) experience a higher risk of psychiatric and mental health disorders [4] and elevated risks of premature mortality by nearly two decades than their non-autistic peers [5]. In a population-derived sample of young autistic people aged 10 to 14, 71% met criteria for at least one mental health disorder, with 41% having additional psychiatric diagnoses [6]. Similarly high rates of mental health problems have been found in other studies of autistic adolescents [7] and autistic adults [8–11]. Two of the most common psychiatric problems for autistic adults are depression and anxiety [12–15], with prevalence rates for depressive and anxiety disorders reaching as high as 47.1% and 54.0%, respectively [16]. It is also possible that prevalence rates of co-occurring psychiatric and mental health difficulties are underestimated, with many autistic individuals being subject to missed, delayed or inaccurate diagnoses [17] and healthcare needs which remain unmet in health systems modelled on the needs of neurotypical individuals [18].

For autistic individuals, mental health difficulties increase the likelihood of poorer long-term outcomes [19] including lower social and adaptive functioning [10], educational and employment difficulties [20, 21], poorer quality of life [22–24], and suicidality [25]. In order to support autistic people in achieving positive long-term outcomes, it is important to further understand the presentation of mental ill-health in autism and to determine *why* autistic individuals are at higher risk of mental health difficulties. This is crucial since the mental health of autistic individuals has been identified as a key priority for research across multiple stakeholder groups [26, 27].

Competing interests: The authors have declared that no competing interests exist.

There are many factors that could contribute to elevated mental health difficulties in autism. For example, since the prevalence of alexithymia (a subclinical condition characterised by difficulties identifying, expressing and differentiating emotions [28]), a transdiagnostic risk factor for numerous psychiatric conditions (e.g., depression, [29], and anxiety disorders, [30]) is higher in the autistic [49.93%] than the non-autistic population (4.89%; [31]), alexithymia may lead to increased risk of mental health difficulties in autism. Alternatively, Botha and Frost [32] advocate the extension of the minority stress model to explain the substantial health disparities between autistic and non-autistic individuals. The minority stress model posits that high stress burdens can negatively impact the mental and physical health of marginalised groups [33]. Such stress burden and resulting poorer health is frequently observed in minority groups including sexual, gender identity and ethnic minorities [34, 35], and minority stressors have recently been described to increase depressive symptoms in those with physical disabilities [36]. Examples of minority stressors include everyday discrimination, internalised stigma, and concealment. In line with this, the current study focuses on the impact of autism acceptance and camouflaging (which are minority stressors) on the mental health of autistic individuals (a marginalised group).

1.1.1. Autism acceptance, camouflaging and mental health. Autism acceptance, as discussed here, can be defined as “an individual feeling accepted or appreciated as an autistic person, with autism positively recognised and accepted by others and the self as an integral part of that individual” [37]. The majority of autistic individuals in the UK feel that society does not accept [43%] or only sometimes accepts [48%] them as an autistic person (in total 91%; [37]). In spite of this, until recently, research had not directly examined autistic people’s experiences of autism acceptance and how this relates to mental health difficulties. The first study to do so confirmed an association between acceptance and mental health, finding that autism acceptance from external sources predicted depression and stress [37] and personal acceptance predicted depression. In addition, individuals who spontaneously reported camouflaging their autistic traits also reported higher symptoms of depression and fewer experiences of acceptance in the past week [37].

Camouflaging can be defined as “the use of strategies by autistic people to minimise the visibility of their autism in social situations” [38]. These strategies can involve compensation, whereby autistic individuals develop behaviours to help them manage social situations (e.g., learning how to use eye contact or developing scripts to help them navigate social interactions; [38, 39]), and masking, whereby autistic individuals aim to hide their autistic traits (e.g., deliberately suppressing stimming behaviours; [38]). Whilst the vast majority of autistic individuals report camouflaging to some degree [38], there are indications that camouflaging may be more common among women [40–48] and those higher in autistic traits [40, 49–51]. There is inconsistent evidence regarding whether camouflaging is associated with broader cognitive ability [40], with some studies finding that a higher level of education is associated with greater camouflaging [49], and others suggesting there is no association between Intelligence Quotient (IQ) and camouflaging [38, 42, 52] or executive functioning and camouflaging [48].

However, a burgeoning literature suggests that camouflaging may be a risk factor for depression and anxiety in autism [25, 37, 53–57], irrespective of gender [58]. Interestingly, Cage and colleagues [37] suggest that camouflaging may mediate the relationship between social stressors (e.g., lack of autism acceptance, bullying) and anxiety and depression. That is, those who experience more social stressors (such as a lack of autism acceptance) are more likely to camouflage their autistic traits, which in turn results in higher levels of anxiety and depression. However, a recent study found that whilst autism-related stigma was associated with lower wellbeing, camouflaging did not mediate this relationship (despite camouflaging being positively associated with autism-related stigma; [59]). Future research should verify

whether camouflaging mediates the relationship between autism acceptance/stigma and anxiety and depression more specifically.

It is important to note that the terms we use to refer to camouflaging vary across the globe [60]. In Japan, the term ‘over-adaptation’ is used to describe excessive efforts made by people (i.e., society more broadly) to meet external demands and expectations, even if this means forcibly suppressing their internal needs [61]. In recent years, this term has been applied to autism to describe the substantial efforts made by autistic people to fit into an environment as a result of differences in interpersonal and social behaviour [62–64]. As such, this concept is highly similar to the construct of camouflaging (see [61] for a discussion around the similarities and differences between the constructs of camouflaging and over-adaptation in autism). Importantly, there is anecdotal evidence that over-adaptation is associated with poorer mental health in autistic individuals in Japan [65]. However, to date no empirical studies have tested this relationship [61], and therefore further study of over-adaptation and camouflaging in Japan is necessary.

1.1.2. Cross-cultural perspectives on autism acceptance, camouflaging and mental health. Culture can be defined as “a set of basic assumptions and values, orientations to life, beliefs, policies, procedures and behavioural conventions that are shared by a group of people” [66]. Unsurprisingly, across different cultures there can be large variation in these beliefs, attitudes and behaviours. In the case of autism, we see cross-cultural variation in attitudes towards autistic people: previous research has found that *non-autistic* college students in the US reported lower levels of autism-related stigma than their Lebanese [67, 68] and Japanese [69] counterparts. Whilst these studies demonstrate that there may be variation in levels of autism stigmatisation among *non-autistic* individuals across cultures, studies have not yet compared *autistic* individuals’ experiences of autism acceptance across different cultures. Given that levels of stigma reported by non-autistic individuals vary cross-culturally, it is plausible that experience of acceptance from the perspective of autistic individuals themselves also vary across cultures. Such cross-cultural differences may exist in both external acceptance (as experiencing stigma may result in autistic individuals feeling less accepted by others) and personal acceptance (as stigma can become internalised resulting in decreased personal acceptance or ‘self-stigma’; [54, 70, 71]).

As Lai and colleagues [72] highlight, it is paramount to determine the levels of camouflaging in the autistic population. Currently, it is unknown how many undiagnosed autistic adults exist [73]. However, it is thought that there are more undiagnosed autistic individuals in some areas of the world than others [74]. Investigating the rates of camouflaging across cultures will help us to work towards determining the true prevalence of autism, and its sex/gender ratio, across the globe. To our knowledge, studies have not yet explored camouflaging in autistic populations outside of the UK (e.g., [58]) and USA (e.g., [75]), though there are anecdotal reports of over-adaptation in Japan. Rather than assuming levels of camouflaging in the UK and USA reflect those globally, researchers should collaborate internationally to compare camouflaging behaviours cross-culturally. Cross-cultural perspectives could have great utility for understanding the mechanisms behind camouflaging, as the social environments in which autistic individuals aim to adapt to vary across cultures. For example, by determining whether the rates of camouflaging are higher within certain cultures, we can work towards identifying socio-cultural factors that may contribute to camouflaging. This is important as there is a consensus among researchers that ‘there is much to be learnt about mechanisms’ [72].

Whilst no research has compared camouflaging across cultures, it seems likely that rates of camouflaging will vary between cultures for a number of reasons. Firstly, in collectivist cultures (which prioritise community interdependence and shared group norms and values; [76]), there may be more pressure relative to individualistic cultures (which prioritise independence

and individualism; [76]) to camouflage one's autistic traits in order to minimise deviation from shared group norms. Thus, those in collectivist cultures may alter their behaviour or hide their autistic traits in order to avoid shame and fit in [62]. In contrast, in regions where there are more discussions about neurodiversity- both as a movement and as an ideology (e.g., the United States, Canada, and the United Kingdom; see [77])- individuals may feel less pressured to camouflage their autistic traits. Finally, since autism-related stigma varies across cultures [67–69] and reduced autism acceptance is associated with increased camouflaging [34, 61], levels of camouflaging may also differ cross-culturally.

Moreover, to the best of our knowledge, studies have not yet compared the level of mental health difficulties in cross-cultural samples of autistic individuals. Like camouflaging, there are reasons to believe there is cross-cultural variation in the level of these difficulties. Firstly, since autism acceptance differs across cultures [67–69], the level of mental health difficulties may also vary because a lack of autism acceptance or stigma is associated with poorer mental health [32, 37]. Secondly, since beliefs about mental health vary across cultures [78], and these beliefs can affect willingness to seek and adhere to treatment [79], there may be cross-cultural differences in levels of mental health difficulties. Finally, given that accessibility of mental health services also differs across cultures [80], individuals from certain regions may be unable to get the help they need, thus leading to poorer mental health. Research is necessary to explore whether the levels of mental health difficulties experienced by autistic individuals vary across cultures.

1.1.3. The current study. In the current study, we aim to replicate the regression analyses from previous research that found a) depression was predicted by external and personal acceptance, and stress was predicted by external acceptance [37], b) camouflaging was predicted by stigma [59], and c) anxiety was predicted by camouflaging [58] (whilst controlling for relevant participant characteristics, e.g., age, gender, etc.) in a more representative sample of autistic adults. In addition, we aim to compare autistic individuals' experiences of autism acceptance, camouflaging and mental health across multiple countries: Australia, Belgium, Canada, Japan, South Africa, the United Kingdom, and the United States. Importantly, exploring these factors across cultures has the potential to a) elucidate the potential routes to greater autism acceptance, reduced need for camouflaging behaviours and improved mental health outcomes, b) identify 'priority regions' for anti-stigma interventions (like those in Ranson and Byrne, [81] and Gillespie-Lynch et al., [82]), and c) highlight areas where greater support for mental health difficulties is needed. In addition, since acceptance is likely to vary across cultures, we are employing an experimental design which naturally maximises variation in autism acceptance, thus giving us the ability to more sensitively assess the impact of autism acceptance on camouflaging and mental health respectively.

In order to fulfil our aims, participants will complete a number of questionnaires including Autism Acceptance Questions [37], the Camouflaging Autistic Traits Questionnaire [51], and the 21-item Depression, Anxiety and Stress Scale (DASS-21; [83]). Participants will also provide demographic information (e.g., age, sex assigned at birth, gender identity, sexual identity, ethnicities, country of birth, country of residence, years lived in country of residence, number of siblings, level of education, income, autism diagnosis, person/team that gave them their diagnosis, age of autism diagnosis, co-occurring diagnoses, how verbal they would consider themselves now, and how verbal they were as a child).

1.2. Hypotheses (required)

1.2.1. Replicating the regression analyses from previous research. First, we will conduct regression analyses on our data as a whole (i.e., participant data will be analysed as one large and diverse group) to replicate the findings from previous studies [37, 58, 59] in a more diverse

sample. In line with Cage and colleagues [37], we predict that external acceptance and personal acceptance will contribute to depression scores, after controlling for age, gender, age of diagnosis, anxiety scores and stress scores. In addition, we predict that external acceptance will contribute to stress scores, after controlling for age, gender, age of diagnosis, anxiety scores and depression scores. As was the case in Perry et al., [59], we hypothesise that autism acceptance will predict camouflaging scores after controlling for age, gender, age of diagnosis, and autistic traits. In line with Hull et al., [58] we predict that camouflaging will contribute to anxiety scores after controlling for age and autistic traits.

1.2.2. Comparing levels of autism acceptance, camouflaging and mental health across cultures. Finally, we will separate participants into groups based on their country and use a between-subjects multivariate analysis of variance (MANOVA) to explore participants' experiences of autism acceptance, camouflaging, and levels of depression, anxiety and stress (dependent variables) across these countries (independent variable). We predict that there will be differences in the level of autism acceptance across countries. In addition, we aim to explore whether there are any differences in the level of camouflaging and mental health difficulties across countries.

2. Design plan

In this section, you will be asked to describe the overall design of your study. Remember that this research plan is designed to register a single study, so if you have multiple experimental designs, please complete a separate preregistration.

2.1. Study type (required)

Observational Study—Data is collected from study subjects that are not randomly assigned to a treatment. This includes surveys, natural experiments and regression discontinuity designs.

2.2. Blinding (required)

No blinding is involved in this study.

2.3. Study design (required)

This study employs a between-subjects design, where we will be comparing participants experiences of personal acceptance, external acceptance, camouflaging, depression, anxiety and stress across cultures.

2.4. Randomization (optional)

No randomization is involved in this study.

3. Sampling plan

In this section we'll ask you to describe how you plan to collect samples, as well as the number of samples you plan to collect and your rationale for this decision. Please keep in mind that the data described in this section should be the actual data used for analysis, so if you are using a subset of a larger dataset, please describe the subset that will actually be used in your study.

3.1. Existing data (required)

Registration prior to creation of data: As of the date of submission of this research plan for pre-registration, the data have not yet been collected, created, or realised.

3.2. Data collection procedures (required)

Participants will be recruited via an existing international autism research database kept by the U21 Autism Research Network. The database was set up as part of a collaboration between seven autism research groups across 4 continents and 7 countries (Australia, Belgium, Hong Kong, Japan, New Zealand, UK, and USA). The aim of this initiative was to diversify samples recruited for autism research and to generate opportunities to answer questions relevant to the global autistic community. In this particular study, we will be recruiting participants from our sites in Belgium, Japan, UK and USA. In addition, we will recruit participants via Prolific, thus allowing us to recruit participants from Australia, Canada, and South Africa (amongst others), and via social media. All participants will provide informed consent. This study has been approved by the Science, Technology, Engineering and Mathematics (STEM) committee at the University of Birmingham (ERN_16-0281AP9D).

3.3. Sample size (required)

The final sample in this study will comprise at least 286 autistic adults (aged over 18) from across the globe.

3.4. Sample size rationale (optional)

3.4.1. Replicating the regression analyses from previous research. In our regression analyses, we will include data for all participants that complete the study in one group (regardless of country of residence) to replicate the findings from previous studies [37, 58, 59] in a large and cross-cultural sample of autistic adults. The chosen sample size for these analyses is based on a series of sample size calculations conducted using G*Power [84], which focus on replicating the regression analyses of interest from Cage et al., [37], Perry et al., [59] and Hull et al., [58]. As a whole, we aim to verify whether experiencing a lack of autism acceptance predicts greater camouflaging and heightened depression and stress, and whether increased camouflaging predicts elevated anxiety (see Fig 1). Further details about the specific analyses can be seen below.




Fig 1. A diagram showing the relationships that we will test in the current study. The plus (+) indicates a positive relationship between the variables and the minus (-) indicates a negative relationship.

<https://doi.org/10.1371/journal.pone.0261774.g001>

Please note that for the following sample size calculations, we used an Bonferroni-adjusted alpha level of 0.0125 to account for multiple testing (4 regression analyses), and a power level of 90% (thereby giving us higher power than is standard in the field [85, 86]). To replicate the finding from Cage et al., that external acceptance and personal acceptance account for 17.5% variance explained in depression scores after controlling for age, gender, age of diagnosis, anxiety and stress ($f^2 = 0.37$), 51 participants are required. In order to replicate the finding that external acceptance and personal acceptance account for 3% variance explained in stress scores after controlling for age, gender, age of diagnosis, anxiety and stress, 246 participants are required. To replicate the finding from Perry et al., that stigma significantly predicts camouflaging ($f^2 = 0.069$) after controlling for age, gender, age of diagnosis, and autistic traits, 211 participants are required. Finally, in order to replicate the finding that anxiety accounts for 4% variance explained in camouflaging scores after controlling for age and autistic traits, 286 participants are required. Therefore, the final sample will comprise at least 286 participants (with full datasets).

3.4.2. Comparing autism acceptance, camouflaging and mental health across cultures.

In the analyses where we will compare participants' experiences of autism acceptance, camouflaging, and levels of depression, anxiety and stress across countries, we will only include data from countries that have at least 30 participants (and therefore we will have sufficient data to draw comparisons between these multiple groups). We selected this cut-off as Sekaran [87] highlights that "where samples are to be broken into sub-samples. . . a minimum sample size of 30 for each category is necessary". Importantly, Sekaran [87] also highlights that "in multivariate research. . . the sample size should be several times (preferably 10 times or more) as large as the number of variables in this study". By recruiting at least 286 participants to our study, our sample is over 47 times larger than the number of variables in our multivariate analysis (6 variables: external acceptance, personal acceptance, camouflaging, depression, anxiety and stress), is as large as necessary based on our power analyses (largest sample size generated = 286), and would facilitate comparisons between participants in up to 9 countries.

4. Variables

In this section you can describe all variables (both manipulated and measured variables) that will later be used in your confirmatory analysis plan. In your analysis plan, you will have the opportunity to describe how each variable will be used. If you have variables which you are measuring for exploratory analyses, you are not required to list them, though you are permitted to do so.

4.1. Manipulated variables (optional)

Group membership—The autistic participants will be divided into groups based on the country in which they live (Australia, Belgium, Canada, Japan, etc.) in our Multivariate Analysis of Variance (MANOVA).

4.2. Measured variables (required)

4.2.1. Demographic information. Participants will be asked a number of demographic questions to establish the nature of the sample. These will include questions about a) age, b) sex assigned at birth, c) gender identity, d) sexual identity, e) ethnicities, f) country of birth, g) country of residence, h) years lived in country of residence, i) level of education, j) how many siblings they have, k) income, l) co-occurring diagnoses m) how verbal they would consider themselves now, and n) how verbal they were as a child (as recommended by Botha, Hanlon and Williams [88]). Participants will also be asked to confirm whether or not they have an

autism diagnosis, which diagnosis they received (e.g., Autism vs. Autism Spectrum Disorder), who gave them their diagnosis (e.g., multidisciplinary team, doctor, clinical psychologist, etc.), and finally the age they received that diagnosis. Those without a formal diagnosis will not be eligible to participate in the study.

4.2.2. Autism acceptance. To measure perceptions of autism acceptance, we will use the six self-report autism acceptance questions outlined in Cage et al., [37]. First, in order to obtain a categorical response for acceptance, participants will be asked whether they feel that society (which will be specified as the general public, made up of people who do not personally know them) generally accepted them, with “yes”, “no”, “sometimes” and “prefer not to say” as response options. Second, participants will be asked to respond to the statement “over the past week, I have felt accepted by society as an autistic person/ person with autism”, on a five-point Likert scale from “strongly agree” to “strongly disagree”. Next, to measure perceptions of autism acceptance from different sources, participants will also be asked: “how accepted by society do you feel as an autistic person?”; “how accepted by your family and friends do you feel as an autistic person?”; and “how much have you personally accepted yourself as an autistic person?”. They will report their responses on a scale from zero, “not at all”, to ten, “completely”. Finally, in order to obtain qualitative responses, participants will have the opportunity to “tell us more about experiences of acceptance or non-acceptance” in an open text box. The English version of this set of autism acceptance questions has acceptable internal consistency (Cronbach’s $\alpha = 0.64$) and good validity [37].

4.2.3. Camouflaging of autistic traits. To measure the extent to which participants camouflage their autistic traits, we will use the Camouflaging Autistic Traits Questionnaire (CAT-Q; [51]). The CAT-Q comprises 25 items (e.g., “In social situations, I feel like I’m pretending to be normal”), rated on a seven-point Likert scale (ranging from one, “Strongly Disagree” to seven, “Strongly Agree”). Total scores on the CAT-Q can range from 25–175, with higher scores reflecting greater camouflaging. This self-report questionnaire assesses three different domains relevant to camouflaging: compensation (strategies used to compensate for socio-communicative difficulties, e.g., “I have spent time learning social skills from television shows and films, and try to use these in my interactions”); masking (strategies used to present a non-autistic or less autistic persona to others, e.g., “I monitor my body language or facial expressions so that I appear interested by the person I am interacting with”), and assimilation (strategies used to fit in to uncomfortable social situations, e.g., “I have to force myself to interact with people when I am in social situations”). The English version of the CAT-Q has strong psychometric properties, including internal consistency (Cronbach’s $\alpha = 0.94$) and test-retest reliability (Pearson’s $r = 0.77$; [51]).

4.2.4. Depression, anxiety and stress. To measure levels of depression, anxiety and stress, we will use the Depression, Anxiety and Stress Scale (DASS-21; [83]). In this 21-item questionnaire, participants judge whether certain statements (e.g., “I found it hard to wind down”) could be applied to their life over the past week on a scale from one to four (1 = “did not apply to me at all; 2 = “applied to me some of the time”; 3 = “applied to me a considerable degree”; 4 = “applied to me very much or most of the time”). The 21 items can be reduced to 7 items each for the depression, anxiety and stress subscales. Total scores for each subscale are calculated and multiplied by two (meaning that scores for each subscale range from 14 to 42), with higher scores representing higher depression, anxiety and stress. The DASS-21 has been previously used with autistic participants (e.g., [37, 89]) and the English [37, 89], French [90], and Japanese [91] versions have good internal consistency and factor validity.

4.2.5. Questionnaires to facilitate our analyses in which we control for other participant characteristics. The following questionnaires are included to facilitate a) the exploration of the relationships found previously between autistic traits and camouflaging, and alexithymic

traits and mental health, and b) our analyses in which we control for these participant characteristics (see Exploratory Analyses).

4.2.6. Autistic traits. The autistic traits of all participants will be assessed via the Autism Quotient (AQ; [92]). This self-report questionnaire assesses five different domains relevant to autistic characteristics (attention switching, attention to detail, communication, social skill and imagination). Participants are instructed to respond to 50 items with one of four responses: 'definitely agree', 'slightly agree', 'slightly disagree' and 'definitely disagree'. These responses are scored using a binary system, where endorsing an autistic trait (either mildly or strongly) is scored as 1, and not endorsing an autistic trait (either mildly or strongly) is scored as 0. As such, scores on the AQ range from 0 to 50, with higher scores representing higher levels of autistic traits. The English [93], French [94], and Japanese [95] versions of the AQ have strong psychometric properties.

4.2.7. Alexithymic traits. The alexithymic traits of participants will be measured via the 20-item Toronto Alexithymia Scale [96]. This self-report questionnaire comprises 20 items rated on a five-point Likert scale (from 1, strongly disagree, to 5, strongly agree). Total scores on the TAS-20 range from 20 to 100, with higher scores representing higher levels of alexithymic traits. The TAS-20 is the most popular tool for assessing alexithymia and the English [96, 97], French [98], and Japanese [99] versions have good psychometric properties.

4.2.8. Translation of relevant questionnaires. Many of the questionnaires used in this study (e.g., the Depression, Anxiety and Stress Scale, the Autism Quotient, the Toronto Alexithymia Scale) have already been translated into the target languages (English, French, Japanese). However, other questionnaires (e.g., the Autism Acceptance Questions and the CAT-Q) are yet to be translated. We will follow the recommendations outlined in Gorecki et al., [100] for conducting translations for cross-cultural research. As such, we will use the forward-backward method of translation that has been used by other researchers in this field (e.g., [69]). The psychometric properties of the translated questionnaires will be assessed and reported in the manuscript (as was the case in Someki et al. [69]). More specifically, following the procedures outlined in Cage et al., [34] and Hull et al., [48], we will assess a) the internal consistency of the translated questionnaires via item analyses, and b) the factor structure of the translated questionnaires via principal component analyses.

5. Analysis plan

You may describe one or more confirmatory analysis in this preregistration. Please remember that all analyses specified below must be reported in the final article, and any additional analyses must be noted as exploratory or hypothesis generating.

A confirmatory analysis plan must state up front which variables are predictors (independent) and which are the outcomes (dependent), otherwise it is an exploratory analysis. You are allowed to describe any exploratory work here, but a clear confirmatory analysis is required.

5.1. Statistical models (required)

First, in our regression analyses, we will include data for all participants that completed the study in one group (collapsing across countries) to explore relationships in a large and cross-cultural sample of autistic adults. For these four regression analyses, we will use a $p < .0125$ threshold to establish whether to reject the null hypothesis (corrected for multiple testing). Then, we will separate participants into groups based on their country of residence and compare participants' experiences of autism acceptance, camouflaging, and levels of depression, anxiety and stress across these groups.

5.1.1. Replicating the regression analyses from previous research. Following the procedures from Cage et al., [37], we will complete two hierarchical regression analyses with depression and stress as the outcome variables respectively. Blockwise entry will be used to analyse the data: in step one, age, age of diagnosis, gender and the other DASS subscale scores will be entered. In the second step, the two types of acceptance (external sources and personal) will be added.

In line with the procedures outlined in Perry et al., [59], we will complete a multiple regression analysis with camouflaging as the dependent variable, and autism acceptance, age, gender, age of diagnosis and autistic traits as the independent variables.

As in Hull et al., [58], we will complete a hierarchical regression analysis with anxiety as the dependent variable. In step one, age and autistic traits will be added into the model. In step two, camouflaging will be entered into the model.

5.1.2. Comparing autism acceptance, camouflaging and mental health across cultures. Finally, in order to compare participants' experiences of autism acceptance, camouflaging, and levels of depression, anxiety and stress across countries, we will use a multivariate analysis of variance (MANOVA). This MANOVA will have the between-subjects factor *country* and the dependent variables *external acceptance*, *personal acceptance*, *depression*, *anxiety*, *stress*, and *camouflaging*. This will facilitate exploration of group differences in the levels of autism acceptance, camouflaging and mental health difficulties.

5.2. Transformations (optional)

In the instance that the data violates any assumptions of the planned statistical analyses, standard transformations will be completed such that the data can be analysed with parametric tests. If assumptions remain unmet, non-parametric analyses will be run. In any analyses that investigate the contribution of multiple predictor variables to a dependent variable, the predictor variables will be z-scored to facilitate comparisons.

5.3. Inference criteria (optional)

For our regression analyses, to account for the four tests we will use a $p < .0125$ significance threshold to determine whether to accept or reject the null hypothesis. For our MANOVA, we will use a $p = .05$ significance threshold.

In our analyses, Bayes Factors (BF_{01}) may be reported, which provide a ratio of the likelihood for the observed data under the null hypothesis compared to the alternative hypothesis [101]. For all Bayesian analyses, we will follow the classification scheme in JASP outlined by Lee and Wagenmakers [102]: BF_{10} values between one and three will be considered as weak evidence, between three and ten as moderate evidence and greater than ten as strong evidence for the alternative hypothesis.

5.4. Data exclusion (optional)

We will exclude any participants that state that they do not have a diagnosis of autism from our analyses. In order to represent the diverse experiences of autistic individuals, we aim to include as much participant data as possible in our analyses. However, we will use Mahalanobis Distance to determine multivariate outliers in our data. Outliers that persist after any necessary transformations and centering (e.g., z-scoring) will be excluded on a case-by-case basis.

5.5. Missing data (optional)

We will use multiple imputation to handle missing data. This statistical technique is a relatively flexible, general purpose approach to dealing with missing data [103].

5.6. Exploratory analysis (optional)

We will explore the relationships between our dependent variables (external acceptance, personal acceptance, depression, anxiety, stress and camouflaging) and various individual differences (e.g., autistic traits, alexithymic traits, gender, level of education, age of diagnosis, income etc.). For continuous variables, we will run simple (Bonferroni-corrected) correlations between our dependent variables and demographic factors (e.g., age, autistic traits). For categorical variables, we will use Bonferroni-corrected independent samples t-tests to compare levels of autism acceptance, camouflaging and mental health difficulties across the demographic factor (e.g., gender). For example, as gender is a categorical variable, we will use Bonferroni-corrected independent samples t-tests to compare levels of acceptance, camouflaging and mental health across genders. Conducting these analyses will allow us to verify whether any individual differences are associated with any of our dependent variables as has been found previously in some instances (e.g., camouflaging and gender; [41]).

Then, we will conduct our MANOVA again whilst controlling for any of the individual differences that are associated with our dependent variables or which differ significantly between groups. This will allow us to determine whether there are cultural differences in autism acceptance, camouflaging and mental health that go beyond these specific individual differences.

Moreover, in order to investigate whether camouflaging mediates any relationships we identify between autism acceptance and mental health outcomes, we will complete mediation analyses. There are several methods used to test the statistical significance of mediated effects [104–106]. In the present study, we will use structural equation modelling (SEM), a powerful statistical technique used to infer causal relationships between variables, for our mediation analyses. We decided to use SEM over standard regression methods (as outlined by Baron and Kenny; [107]) as this technique is regarded as “a more appropriate inference framework for mediation analyses” (see [108]). In addition, we will use bias-corrected bootstrapping to create confidence intervals as there is a consensus that it is the most powerful method for testing mediated effects [104, 109–112]. Whilst we believe that there will be sufficient statistical power to detect an effect, we may supplement this analysis with Bayesian mediation analyses if it appears that we are underpowered.

In addition, we will calculate standardised residuals for the relationships between autism acceptance and mental health, autism acceptance and camouflaging, and camouflaging and mental health, for all participants. Then we will complete an exploratory MANOVAs in which we will compare the standardised residuals across countries. This will allow us to compare the strength of these relationships in each of the countries by comparing the levels of unexplained variance following regression analyses. This is useful because the strengths of these relationships could, in principle, vary across our countries. For example, it could be that in collectivist, relative to individualistic, cultures, camouflaging accounts for a smaller portion of variance in mental health as hiding any traits that deviate from the norm is common (and therefore may not be detrimental to mental health).

Finally, in order to compare the strength of the relationships between our dependent variables across genders, we will complete an additional exploratory MANOVA using the standardised residuals discussed above. This is useful because the strength of the relationships between our dependent variables could vary across genders: it could be the case, for instance, that there is a stronger relationship between autism acceptance and camouflaging in autistic women than autistic men, thus explaining the higher rates of camouflaging displayed by autistic women [40–48]. In addition, it could be that there is a weaker relationship between camouflaging and mental health difficulties in autistic men since they more frequently report positive consequences of camouflaging [38].

6. Other

6.1. Other (optional)

Please note that the authors consulted an autism consultancy group (Birmingham Psychology Autism Research Team Consultancy Committee) and the global U21 Autism Research Network when designing the study in order to confirm that a) the aims of the study were in line with community priorities, b) that the questionnaires were clear and accessible to autistic participants and c) that the aims and methodology were in line with norms for research conduct in each participating country.

Acknowledgments

We thank the Birmingham Psychology Autism Research Team (B-PART) consultancy group for their advice and input on the study. We are really grateful for their fantastic contribution.

Author Contributions

Conceptualization: Connor Tom Keating, Lydia Hickman, Sophie Sowden.

Data curation: Connor Tom Keating, Lydia Hickman, Philippine Geelhand, Toru Takahashi, Joan Leung, Bianca Schuster, Alicia Rybicki, Teresa Marie Girolamo, Elise Clin, Fanny Papastamou, Marie Belenger, Inge-Marie Eigsti, Jennifer Louise Cook, Hirotaka Kosaka, Rieko Osu, Yuko Okamoto, Sophie Sowden.

Formal analysis: Connor Tom Keating.

Funding acquisition: Connor Tom Keating, Lydia Hickman, Bianca Schuster, Sophie Sowden.

Investigation: Connor Tom Keating.

Methodology: Connor Tom Keating, Lydia Hickman, Philippine Geelhand, Toru Takahashi, Elise Clin, Fanny Papastamou, Marie Belenger, Sophie Sowden.

Project administration: Connor Tom Keating, Alicia Rybicki, Sophie Sowden.

Resources: Connor Tom Keating.

Supervision: Sophie Sowden.

Visualization: Connor Tom Keating.

Writing – original draft: Connor Tom Keating.

Writing – review & editing: Connor Tom Keating, Lydia Hickman, Philippine Geelhand, Toru Takahashi, Joan Leung, Bianca Schuster, Alicia Rybicki, Teresa Marie Girolamo, Elise Clin, Fanny Papastamou, Marie Belenger, Inge-Marie Eigsti, Jennifer Louise Cook, Hirotaka Kosaka, Rieko Osu, Yuko Okamoto, Sophie Sowden.

References

1. American Psychiatric Association. Diagnostic and Statistical Manual of Mental Disorders, 5th Edition (DSM-5). 5th ed. BMC Medicine. 2013:133–137
2. Kenny L, Hattersley C, Molins B, Buckley C, Povey C, Pellicano E. Which terms should be used to describe autism? Perspectives from the UK autism community. *Autism*. 2016; 20(4):442–62. <https://doi.org/10.1177/1362361315588200> PMID: 26134030
3. Bottema-Beutel K, Kapp SK, Lester JN, Sasson NJ, Hand BN. Avoiding Ableist Language: Suggestions for Autism Researchers. *Autism in Adulthood*. 2021; 3(1):18–29.

4. Gillberg IC, Helles A, Billstedt E, Gillberg C. Boys with Asperger Syndrome Grow Up: Psychiatric and Neurodevelopmental Disorders 20 Years After Initial Diagnosis. *Journal of Autism and Developmental Disorders*. 2016; 46(1):74–82. Available from: <https://link.springer.com/article/10.1007/s10803-015-2544-0> PMID: 26210519
5. Hirvikoski T, Mittendorfer-Rutz E, Boman M, Larsson H, Lichtenstein P, Bölte S. Premature mortality in autism spectrum disorder. *British Journal of Psychiatry*. 2016; 208(3):232–8. Available from: <https://www.cambridge.org/core>. <https://doi.org/10.1192/bjp.bp.114.160192> PMID: 26541693
6. Simonoff E, Pickles A, Charman T, Chandler S, Loucas T, Baird G. Psychiatric disorders in children with autism spectrum disorders: Prevalence, comorbidity, and associated factors in a population-derived sample. *Journal of the American Academy of Child and Adolescent Psychiatry*. 2008; 47(8):921–9. <https://doi.org/10.1097/CHI.0b013e318179964f> PMID: 18645422
7. Eaves LC, Ho HH. Young Adult Outcome of Autism Spectrum Disorders. *Journal of Autism and Developmental Disorders*. 2008; 38(4): 739–47 <https://doi.org/10.1007/s10803-007-0441-x> PMID: 17764027
8. Griffiths S, Allison C, Kenny R, Holt R, Smith P, Baron-Cohen S. The Vulnerability Experiences Quotient (VEQ): A Study of Vulnerability, Mental Health and Life Satisfaction in Autistic Adults. *Autism Research*. 2019; 12(10):1516–28. <https://doi.org/10.1002/aur.2162> PMID: 31274233
9. Lever AG, Geurts HM. Psychiatric Co-occurring Symptoms and Disorders in Young, Middle-Aged, and Older Adults with Autism Spectrum Disorder. *Journal of Autism and Developmental Disorders*. 2016; 46(6):1916–30. <https://doi.org/10.1007/s10803-016-2722-8> PMID: 26861713
10. Moss P, Howlin P, Savage S, Bolton P, Rutter M. Self and informant reports of mental health difficulties among adults with autism findings from a long-term follow-up study. *Autism*. 2015; 19(7):832–41. <https://doi.org/10.1177/1362361315585916> PMID: 26014841
11. Sedgewick F, Leppanen J, Tchanturia K. Gender differences in mental health prevalence in autism. *Advances in Autism*. 2020.
12. Croen LA, Zerbo O, Qian Y, Massolo ML, Rich S, Sidney S, et al. The health status of adults on the autism spectrum. *Autism*. 2015; 19(7):814–23. <https://doi.org/10.1177/1362361315577517> PMID: 25911091
13. Gotham K, Brunwasser SM, Lord C. Depressive and anxiety symptom trajectories from school age through young adulthood in samples with autism spectrum disorder and developmental delay. *Journal of the American Academy of Child and Adolescent Psychiatry*. 2015; 54(5):369–376.e3. Available from: <https://doi.org/10.1016/j.jaac.2015.02.005> PMID: 25901773
14. Levy A, Perry A. Outcomes in adolescents and adults with autism: A review of the literature. *Research in Autism Spectrum Disorders*. 2011; 5(4):1271–82. Available from: <http://dx.doi.org/10.1016/j.rasd.2011.01.023>
15. Lugnegård T, Hallerbäck MU, Gillberg C. Psychiatric comorbidity in young adults with a clinical diagnosis of Asperger syndrome. *Research in Developmental Disabilities*. 2011; 32(5):1910–7. <https://doi.org/10.1016/j.ridd.2011.03.025> PMID: 21515028
16. Hossain MM, Khan N, Sultana A, Ma P, McKyer ELJ, Ahmed HU, et al. Prevalence of comorbid psychiatric disorders among people with autism spectrum disorder: An umbrella review of systematic reviews and meta-analyses. Vol. 287, *Psychiatry Research*. Elsevier Ireland Ltd; 2020. p. 112922. <https://doi.org/10.1016/j.psychres.2020.112922> PMID: 32203749
17. Bauman ML. Medical comorbidities in autism: Challenges to diagnosis and treatment. *Neurotherapeutics*. 2010; 7(3):320–7. Available from: <https://link.springer.com/article/10.1016/j.nurt.2010.06.001> PMID: 20643385
18. Nicolaidis C, Raymaker D, McDonald K, Dern S, Boisclair WC, Ashkenazy E, et al. Comparison of healthcare experiences in autistic and non-autistic adults: A cross-sectional online survey facilitated by an academic-community partnership. *Journal of General Internal Medicine*. 2013; 28(6):761–9. <https://doi.org/10.1007/s11606-012-2262-7> PMID: 23179969
19. Lai MC, Kasseh C, Besney R, Bonato S, Hull L, Mandy W, et al. Prevalence of co-occurring mental health diagnoses in the autism population: a systematic review and meta-analysis. *The Lancet Psychiatry*. 2019; 6(10):819–29. Available from: [https://doi.org/10.1016/S2215-0366\(19\)30289-5](https://doi.org/10.1016/S2215-0366(19)30289-5) PMID: 31447415
20. Keen D, Webster A, Ridley G. How well are children with autism spectrum disorder doing academically at school? An overview of the literature. *Autism*. 2016; 20(3):276–94. <https://doi.org/10.1177/1362361315580962> PMID: 25948598
21. Taylor JL, Henninger NA, Mailick MR. Longitudinal patterns of employment and postsecondary education for adults with autism and average-range IQ. *Autism*. 2015; 19(7):785–93. <https://doi.org/10.1177/1362361315585643> PMID: 26019306

22. Adams D, Clark M, Keen D. Using self-report to explore the relationship between anxiety and quality of life in children on the autism spectrum. *Autism Research*. 2019; 12(10):1505–15. <https://doi.org/10.1002/aur.2155> PMID: 31207183
23. Knüppel A, Telléus GK, Jakobsen H, Lauritsen MB. Characteristics of Young Adults with Autism Spectrum Disorder Performing Different Daytime Activities. *Journal of Autism and Developmental Disorders*. 2019; 49(2):542–55. Available from: <https://doi.org/10.1007/s10803-018-3730-7> PMID: 30151782
24. Robertson SM. Neurodiversity, Quality of Life, and Autistic Adults: Shifting Research and Professional Focuses onto Real-Life Challenges. *Disability Studies Quarterly*. 2009; 30(1). Available from: <https://dsq-sds.org/article/view/1069/1234>
25. Cassidy S, Bradley L, Shaw R, Baron-Cohen S. Risk markers for suicidality in autistic adults. *Molecular Autism*. 2018; 9(1):1–14. <https://doi.org/10.1186/s13229-018-0226-4> PMID: 30083306
26. Cusack J, Sterry R. Your questions: shaping future autism research. London: Autistica. 2016.
27. Roche L, Adams D, Clark M. Research priorities of the autism community: A systematic review of key stakeholder perspectives. *Autism*. 2021; 25(2):336–48. <https://doi.org/10.1177/1362361320967790> PMID: 33143455
28. Nemiah JC. Alexithymia: a view of the psychosomatic process. *Modern Trends in Psychosomatic Medicine*. 1976; 3:430–9. Available from: <https://ci.nii.ac.jp/naid/10029440327>
29. Celikel FC, Kose S, Erkorkmaz U, Sayar K, Cumurcu BE, Cloninger CR. Alexithymia and temperament and character model of personality in patients with major depressive disorder. *Comprehensive Psychiatry*. 2010; 51(1):64–70. <https://doi.org/10.1016/j.comppsy.2009.02.004> PMID: 19932828
30. Frewen PA, Lanius RA, Dozois DJA, Neufeld RWJ, Pain C, Hopper JW, et al. Clinical and Neural Correlates of Alexithymia in Posttraumatic Stress Disorder. *Journal of Abnormal Psychology*. 2008; 117(1):171–81. Available from: /record/2008-01363-014 <https://doi.org/10.1037/0021-843X.117.1.171> PMID: 18266495
31. Kinnaird E, Stewart C, Tchanturia K. Investigating alexithymia in autism: A systematic review and meta-analysis. *European Psychiatry*. 2019; 55:80. Available from: /pmc/articles/PMC6331035/ <https://doi.org/10.1016/j.eurpsy.2018.09.004> PMID: 30399531
32. Botha M, Frost DM. Extending the Minority Stress Model to Understand Mental Health Problems Experienced by the Autistic Population. *Society and Mental Health*. 2020; 10(1):20–34. Available from: <http://journals.sagepub.com/doi/10.1177/2156869318804297>
33. Meyer IH. Prejudice, social stress, and mental health in lesbian, gay, and bisexual populations: Conceptual issues and research evidence. *Psychology of Sexual Orientation and Gender Diversity*. 2013; 1(S):3–26.
34. Frost DM, Lehavot K, Meyer IH. Minority stress and physical health among sexual minority individuals. *Journal of Behavioral Medicine*. 2015; 38(1):1–8. Available from: www.columbia.edu/~im15/. <https://doi.org/10.1007/s10865-013-9523-8> PMID: 23864353
35. Williams DR, Williams-Morris R. Racism and mental health: The African American experience. *Ethnicity & Health*. 2000; 5(3–4):243–68. Available from: <https://www.tandfonline.com/doi/abs/10.1080/713667453> PMID: 28712877
36. Brown RL. Functional limitation and depressive symptomatology: Considering perceived stigma and discrimination within a stress and coping framework. *Stigma and Health*. 2017; 2(2):98–109. Available from: /pmc/articles/PMC5421991/ <https://doi.org/10.1037/sah0000042> PMID: 28497112
37. Cage E, Di Monaco J, Newell V. Experiences of Autism Acceptance and Mental Health in Autistic Adults. *Journal of Autism and Developmental Disorders*. 2018; 48(2):473–84. Available from: <https://doi.org/10.1007/s10803-017-3342-7> PMID: 29071566
38. Hull L, Petrides K V., Allison C, Smith P, Baron-Cohen S, Lai MC, et al. “Putting on My Best Normal”: Social Camouflaging in Adults with Autism Spectrum Conditions. *Journal of Autism and Developmental Disorders*. 2017; 47(8):2519–34. <https://doi.org/10.1007/s10803-017-3166-5> PMID: 28527095
39. Cook J, Crane L, Bourne L, Hull L, Mandy W. Camouflaging in an everyday social context: An interpersonal recall study. *Autism*. 2021; 136236132199264. Available from: <http://journals.sagepub.com/doi/10.1177/1362361321992641>
40. Cook J, Hull L, Crane L, Mandy W. Camouflaging in autism: A systematic review. *Clinical Psychology Review*. 2021; 89:102080. <https://doi.org/10.1016/j.cpr.2021.102080> PMID: 34563942
41. Cassidy SA, Gould K, Townsend E, Pelton M, Robertson AE, Rodgers J. Is Camouflaging Autistic Traits Associated with Suicidal Thoughts and Behaviours? Expanding the Interpersonal Psychological Theory of Suicide in an Undergraduate Student Sample. *Journal of Autism and Developmental Disorders*. 2020; 50(10):3638–48. Available from: <https://doi.org/10.1007/s10803-019-04323-3> PMID: 31820344

42. Lai MC, Lombardo M V., Ruigrok ANV, Chakrabarti B, Auyeung B, Szatmari P, et al. Quantifying and exploring camouflaging in men and women with autism. *Autism*. 2017; 21(6):690–702. <https://doi.org/10.1177/1362361316671012> PMID: 27899710
43. Hull L, Lai MC, Baron-Cohen S, Allison C, Smith P, Petrides K V., et al. Gender differences in self-reported camouflaging in autistic and non-autistic adults. *Autism*. 2020; 24(2):352–63. Available from: <http://journals.sagepub.com/doi/10.1177/1362361319864804> PMID: 31319684
44. Parish-Morris J, Liberman MY, Cieri C, Herrington JD, Yerys BE, Bateman L, et al. Linguistic camouflage in girls with autism spectrum disorder. *Molecular Autism*. 2017; 8(1):48. Available from: <https://molecularautism.biomedcentral.com/articles/10.1186/s13229-017-0164-6> PMID: 29021889
45. Ratto AB, Kenworthy L, Yerys BE, Bascom J, Wieckowski AT, White SW, et al. What About the Girls? Sex-Based Differences in Autistic Traits and Adaptive Skills. *Journal of Autism and Developmental Disorders*. 2018; 48(5):1698–711. Available from: <https://doi.org/10.1007/s10803-017-3413-9> PMID: 29204929
46. Wood-Downie H, Wong B, Kovshoff H, Mandy W, Hull L, Hadwin JA. Sex/Gender Differences in Camouflaging in Children and Adolescents with Autism. *Journal of Autism and Developmental Disorders*. 2021; 51(4):1353–64. Available from: <https://doi.org/10.1007/s10803-020-04615-z> PMID: 32691191
47. Rynkiewicz A, Schuller B, Marchi E, Piana S, Camurri A, Lassalle A, et al. An investigation of the “female camouflage effect” in autism using a computerized ADOS-2 and a test of sex/gender differences. *Molecular Autism*. 2016; 7(1):10. Available from: <http://www.molecularautism.com/content/7/1/10>
48. Schuck RK, Flores RE, Fung LK. Brief Report: Sex/Gender Differences in Symptomology and Camouflaging in Adults with Autism Spectrum Disorder. *Journal of Autism and Developmental Disorders*. 2019; 49(6):2597–604. Available from: <https://doi.org/10.1007/s10803-019-03998-y> PMID: 30945091
49. Livingston LA, Shah P, Milner V, Happé F. Quantifying compensatory strategies in adults with and without diagnosed autism. *Molecular Autism*. 2020; 11(1):15. Available from: <https://molecularautism.biomedcentral.com/articles/10.1186/s13229-019-0308-y> PMID: 32051011
50. Brown CM, Attwood T, Garnett M, Stokes MA. Am I Autistic? Utility of the Girls Questionnaire for Autism Spectrum Condition as an Autism Assessment in Adult Women. *Autism in Adulthood*. 2020; 2(3):216–26. Available from: <https://www.liebertpub.com/doi/10.1089/aut.2019.0054>
51. Hull L, Mandy W, Lai MC, Baron-Cohen S, Allison C, Smith P, et al. Development and Validation of the Camouflaging Autistic Traits Questionnaire (CAT-Q). *Journal of Autism and Developmental Disorders*. 2019; 49(3):819–33. Available from: <https://doi.org/10.1007/s10803-018-3792-6> PMID: 30361940
52. Corbett BA, Schwartzman JM, Libsack EJ, Muscatello RA, Lerner MD, Simmons GL, et al. Camouflaging in Autism: Examining Sex-Based and Compensatory Models in Social Cognition and Communication. *Autism Research*. 2021; 14(1):127–42. <https://doi.org/10.1002/aur.2440> PMID: 33220170
53. Bargiela S, Steward R, Mandy W. The Experiences of Late-diagnosed Women with Autism Spectrum Conditions: An Investigation of the Female Autism Phenotype. *Journal of Autism and Developmental Disorders*. 2016; 46(10):3281–94. <https://doi.org/10.1007/s10803-016-2872-8> PMID: 27457364
54. Cage E, Troxell-Whitman Z. Understanding the Reasons, Contexts and Costs of Camouflaging for Autistic Adults. *Journal of Autism and Developmental Disorders*. 2019; 49(5):1899–911. Available from: <https://doi.org/10.1007/s10803-018-03878-x> PMID: 30627892
55. Tierney S, Burns J, Kilbey E. Looking behind the mask: Social coping strategies of girls on the autistic spectrum. *Research in Autism Spectrum Disorders*. 2016; 23:73–83. Available from: <http://dx.doi.org/10.1016/j.rasd.2015.11.013>
56. Tubío-Funqueiriño M, Cruz S, Sampaio A, Carracedo A, Fernández-Prieto M. Social Camouflaging in Females with Autism Spectrum Disorder: A Systematic Review. *Journal of Autism and Developmental Disorders*. 2021; 51(7):2190–9. Available from: <https://doi.org/10.1007/s10803-020-04695-x> PMID: 32926304
57. Allely CS. Understanding and recognising the female phenotype of autism spectrum disorder and the “camouflage” hypothesis: a systematic PRISMA review. *Advances in Autism*. 2019; 5(1):14–37.
58. Hull L, Levy L, Lai MC, Petrides K V., Baron-Cohen S, Allison C, et al. Is social camouflaging associated with anxiety and depression in autistic adults? *Molecular Autism*. 2021; 12(1):13. Available from: <https://molecularautism.biomedcentral.com/articles/10.1186/s13229-021-00421-1> PMID: 33593423
59. Perry E, Mandy W, Hull L, Cage E. Understanding Camouflaging as a Response to Autism-Related Stigma: A Social Identity Theory Approach. *Journal of Autism and Developmental Disorders*. 2021; 1–11. Available from: <https://doi.org/10.1007/s10803-020-04506-3> PMID: 32350790
60. 水澤慶緒里, 中澤清. 過剰適応の日豪比較:メルボルン大学他との研究交流から. *臨床教育心理学研究*. 2012; 38:11–8.

61. 若菜千田, 智岡田. 自閉スペクトラム症における過剰適応とカモフラージュの臨床的意義. 北海道大学大学院教育学研究院附属子ども発達臨床研究センター. 2021; 15:57–66.
62. 杉山登志郎, 高橋脩. 就労に挫折した自閉症青年の臨床的検討. 発達障害研究. 1994; 16(3):198–207.
63. 米田衆介. アスペルガーの人はなぜ生きづらいのか? 大人の発達障害を考える. 2011;
64. 本田秀夫. 自閉スペクトラムの人たちにみられる過剰適応的対人関係. 精神科治療学. 2018; 33:456–8.
65. 横田圭司, 千田若菜. 知的障害と発達障害をめぐるトランジション. 児童青年精神医学とその近接領域. 2018; 59(5):566–76.
66. Matsumoto DR. Culture and psychology. Pacific Grove: Brooks/Cole Pub. Co.; 1996.
67. Gillespie-Lynch K, Daou N, Sanchez-Ruiz MJ, Kapp SK, Obeid R, Brooks PJ, et al. Factors underlying cross-cultural differences in stigma toward autism among college students in Lebanon and the United States. *Autism*. 2019; 23(8):1993–2006. Available from: <http://journals.sagepub.com/doi/10.1177/1362361318823550> PMID: 30939906
68. Obeid R, Daou N, DeNigris D, Shane-Simpson C, Brooks PJ, Gillespie-Lynch K. A Cross-Cultural Comparison of Knowledge and Stigma Associated with Autism Spectrum Disorder Among College Students in Lebanon and the United States. *Journal of Autism and Developmental Disorders*. 2015; 45(11):3520–36. Available from: <https://pubmed.ncbi.nlm.nih.gov/26084712/> PMID: 26084712
69. Someki F, Torii M, Brooks PJ, Koeda T, Gillespie-Lynch K. Stigma associated with autism among college students in Japan and the United States: An online training study. *Research in Developmental Disabilities*. 2018; 76:88–98. <https://doi.org/10.1016/j.ridd.2018.02.016> PMID: 29602160
70. Bachmann CJ, Höfer J, Kamp-Becker I, Küpper C, Poustka L, Roepke S, et al. Internalised stigma in adults with autism: A German multi-center survey. *Psychiatry Research*. 2019; 276:94–9. <https://doi.org/10.1016/j.psychres.2019.04.023> PMID: 31030006
71. Corrigan PW, Watson AC. Understanding the impact of stigma on people with mental illness. *World Psychiatry*. 2002; 1(1):16–20. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/16946807> PMID: 16946807
72. Lai M, Hull L, Mandy W, Chakrabarti B, Nordahl CW, Lombardo M V., et al. Commentary: ‘Camouflaging’ in autistic people—reflection on Fombonne (2020). *Journal of Child Psychology and Psychiatry*. 2020;jcpp.13344. Available from: <https://onlinelibrary.wiley.com/doi/10.1111/jcpp.13344> PMID: 33289092
73. Fombonne E. Camouflage and autism. *Journal of Child Psychology and Psychiatry and Allied Disciplines*. 2020; 61(7):735–8. <https://doi.org/10.1111/jcpp.13296> PMID: 32658354
74. Elsabbagh M, Divan G, Koh YJ, Kim YS, Kauchali S, Marcín C, et al. Global Prevalence of Autism and Other Pervasive Developmental Disorders. *Autism Research*. 2012; 5(3):160–79. Available from: <http://www.autismepidemiology.net> <https://doi.org/10.1002/aur.239> PMID: 22495912
75. Jorgenson C, Lewis T, Rose C, Kanne S. Social Camouflaging in Autistic and Neurotypical Adolescents: A Pilot Study of Differences by Sex and Diagnosis. *Journal of Autism and Developmental Disorders*. 2020; 50(12):4344–55. Available from: <https://pubmed.ncbi.nlm.nih.gov/32270386/> <https://doi.org/10.1007/s10803-020-04491-7> PMID: 32270386
76. Triandis H. Collectivism v. Individualism: A Reconceptualisation of a Basic Concept in Cross-cultural Social Psychology. In: *Cross-Cultural Studies of Personality, Attitudes and Cognition*. Palgrave Macmillan UK; 1988. p. 60–95. Available from: https://link.springer.com/chapter/10.1007/978-1-349-08120-2_3
77. Cascio MA. Neurodiversity as a Conceptual Lens and Topic of Cross-Cultural Study. *Cult Mind, Brain*. 2020;477–93. Available from: <https://www.cambridge.org/core/books/culture-mind-and-brain/neurodiversity-as-a-conceptual-lens-and-topic-of-crosscultural-study/8F5F3D1E247B38F5F695F3CB357BBD95>
78. Leong FTL, Kalibatseva Z. Cross-cultural barriers to mental health services in the United States. *Cerebrum*. 2011; 2011:5. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/23447774> PMID: 23447774
79. Nieuwsma JA, Pepper CM, Maack DJ, Birgenheir DG. Indigenous perspectives on depression in rural regions of India and the United States. *Transcultural Psychiatry*. 2011; 48(5):539–68. Available from: <https://pubmed.ncbi.nlm.nih.gov/22021105/> <https://doi.org/10.1177/1363461511419274> PMID: 22021105
80. Volpe U, Mihai A, Jordanova V, Sartorius N. The pathways to mental healthcare worldwide. *Current Opinion in Psychiatry*. 2015; 28(4):299–306. Available from: <http://journals.lww.com/00001504-201507000-00007> <https://doi.org/10.1097/YCO.0000000000000164> PMID: 26001921
81. Ranson NJ, Byrne MK. Promoting Peer Acceptance of Females with Higher-functioning Autism in a Mainstream Education Setting: A Replication and Extension of the Effects of an Autism Anti-Stigma

- Program. *Journal of Autism and Developmental Disorders*. 2014; 44(11):2778–96. Available from: <https://link.springer.com/article/10.1007/s10803-014-2139-1> PMID: 24816870
82. Gillespie-Lynch K, Brooks PJ, Someki F, Obeid R, Shane-Simpson C, Kapp SK, et al. Changing College Students' Conceptions of Autism: An Online Training to Increase Knowledge and Decrease Stigma. *Journal of Autism and Developmental Disorders*. 2015; 45(8):2553–66. Available from: <https://pubmed.ncbi.nlm.nih.gov/25796194/> <https://doi.org/10.1007/s10803-015-2422-9> PMID: 25796194
83. Lovibond SH, Lovibond PF. *Manual for the depression anxiety stress scales*. Sydney Psychology Foundation. 1995;
84. Faul F, Erdfelder E, Lang AG, Buchner A. G*Power 3: A flexible statistical power analysis program for the social, behavioral, and biomedical sciences. *Behavior Research Methods*. 2007; 39(2):175–91. Available from: <https://link.springer.com/article/10.3758/BF03193146> PMID: 17695343
85. Perugini M, Gallucci M, Costantini G. A Practical Primer To Power Analysis for Simple Experimental Designs. *International Review of Social Psychology*. 2018; 31(1):1–23. Available from: <http://www.rips-irsp.com/articles/10.5334/irsp.181/>
86. Milfont TL, Klein RA. Replication and Reproducibility in Cross-Cultural Psychology. *Journal of Cross Cultural Psychology*. 2018; 49(5):735–50.
87. Sekaran U. *Research Methods for Business: A Skill-Building Approach*. Wiley & Sons, New York; 2003.
88. Botha M, Hanlon J, Williams GL. Does Language Matter? Identity-First Versus Person-First Language Use in Autism Research: A Response to Vivanti. *Journal of Autism and Developmental Disorders*. 2021; 1–9. Available from: <https://doi.org/10.1007/s10803-020-04506-3> PMID: 32350790
89. Maddox BB, White SW. Comorbid Social Anxiety Disorder in Adults with Autism Spectrum Disorder. *Journal of Autism and Developmental Disorders*. 2015; 45(12):3949–60. Available from: <https://doi.org/10.1007/s10803-015-2531-5> PMID: 26243138
90. Ramasawmy L. Validation of the French Depression Anxiety Stress Scales (DASS-21) and predictors of depression in an adolescent Mauritian population. 2015. (Doctoral Dissertation, Aix-Marseille).
91. Hayashi K, Takahashi H. A study of the Japanese version of the depression anxiety stress scale. *Journal of the Australian and New Zealand Student Services Association*. 2001; 18:1–7.
92. Baron-Cohen S, Wheelwright S, Skinner R, Martin J, Clubley E. The Autism-Spectrum Quotient (AQ): Evidence from Asperger Syndrome/High-Functioning Autism, Males and Females, Scientists and Mathematicians. *Journal of Autism and Developmental Disorders*. 2001; 31(1):5–17. Available from: <https://doi.org/10.1023/a:1005653411471> PMID: 11439754
93. Stevenson JL, Hart KR. Psychometric properties of the autism-spectrum quotient for assessing low and high levels of autistic traits in college students. *Journal of Autism and Developmental Disorders*. 2017; 47(6):1838–53. Available from: <https://doi.org/10.1007/s10803-017-3109-1> PMID: 28349365
94. Sonié S, Kassai B, Pirat E, Bain P, Robinson J, Gomot M, et al. The French version of the autism-spectrum quotient in adolescents: A cross-cultural validation study. *Journal of Autism and Developmental Disorders*. 2013; 43(5):1178–83. Available from: <https://doi.org/10.1007/s10803-012-1663-0> PMID: 23015111
95. Kurita H, Koyama T, Osada H. Autism-Spectrum Quotient-Japanese version and its short forms for screening normally intelligent persons with pervasive developmental disorders. *Psychiatry and Clinical Neurosciences*. 2005; 59(4):490–6. Available from: <https://doi.org/10.1111/j.1440-1819.2005.01403.x> PMID: 16048456
96. Bagby RM, Parker JDA, Taylor GJ. The twenty-item Toronto Alexithymia scale-I. Item selection and cross-validation of the factor structure. *Journal of Psychosomatic Research*. 1994; 38(1):23–32. [https://doi.org/10.1016/0022-3999\(94\)90005-1](https://doi.org/10.1016/0022-3999(94)90005-1) PMID: 8126686
97. Taylor GJ, Bagby RM, Parker JDA. The 20-Item Toronto Alexithymia Scale: IV. Reliability and factorial validity in different languages and cultures. *Journal of Psychosomatic Research*. 2003; 55(3):277–83. [https://doi.org/10.1016/s0022-3999\(02\)00601-3](https://doi.org/10.1016/s0022-3999(02)00601-3) PMID: 12932803
98. Loas G, Parker JDA, Otmani O, Verrier A, Fremaux D. Confirmatory factor analysis of the french translation of the 20-item Toronto Alexithymia scale. *Perceptual and Motor Skills*. 1997; 85(2):1018. Available from: <https://doi.org/10.2466/pms.1997.85.3.1018> PMID: 9399312
99. Komaki G, Maeda M, Arimura T, Nakata A, Shinoda H, Ogata I, et al. The reliability and factorial validity of the Japanese version of the 20-item Toronto Alexithymia Scale. *Journal of Psychosomatic Research*. 2003; 55(2):143.
100. Gorecki C, Brown JM, Briggs M, Coleman S, Dealey C, McGinnis E, et al. Language translation & cross-cultural adaptation guideline. Recommendations for language translation and cross-cultural adaption of the PU-QOL questionnaire. 2014.

101. Dienes Z. Using Bayes to get the most out of non-significant results. *Frontiers in Psychology*. 2014; 5:781. Available from: <https://doi.org/10.3389/fpsyg.2014.00781> PMID: 25120503
102. Lee MD, Wagenmakers EJ. *Bayesian Cognitive Modeling: A Practical Course*. Cambridge University Press; 2013: 1–264. Available from: /record/2014-14454-000
103. Sterne JAC, White IR, Carlin JB, Spratt M, Royston P, Kenward MG, et al. Multiple imputation for missing data in epidemiological and clinical research: potential and pitfalls. *British Medical Journal*. 2009; 338(7713):157–60. Available from: <https://doi.org/10.1136/bmj.b2393> PMID: 19564179
104. Hayes AF, Scharkow M. The Relative Trustworthiness of Inferential Tests of the Indirect Effect in Statistical Mediation Analysis: Does Method Really Matter? *Psychological Science*. 2013; 24(10):1918–27. Available from: <https://doi.org/10.1177/0956797613480187> PMID: 23955356
105. MacKinnon DP, Lockwood CM, Hoffman JM, West SG, Sheets V. A comparison of methods to test mediation and other intervening variable effects. *Psychological Methods*. 2002; 7(1):83–104. Available from: /record/2002-00925-005 <https://doi.org/10.1037/1082-989x.7.1.83> PMID: 11928892
106. Shrout PE, Bolger N. Mediation in experimental and nonexperimental studies: New procedures and recommendations. *Psychological Methods*. 2002; 7(4):422–45. Available from: /record/2002-11349-006 PMID: 12530702
107. Baron RM, Kenny DA. The moderator-mediator variable distinction in social psychological research: Conceptual, strategic, and statistical considerations. *Journal of Personality and Social Psychology*. 1986; 51(6):1173–82. Available from: <https://doi.org/10.1037//0022-3514.51.6.1173> PMID: 3806354
108. Gunzler D, Chen T, Wu P, Zhang H. Introduction to mediation analysis with structural equation modeling. *Shanghai Archives of Psychiatry*. 2013; 25(6):390–4. Available from: /pmc/articles/PMC4054581/ <https://doi.org/10.3969/j.issn.1002-0829.2013.06.009> PMID: 24991183
109. Cheung MWL. Comparison of Approaches to Constructing Confidence Intervals for Mediating Effects Using Structural Equation Models. *Structural Equation Modeling: A Multidisciplinary Journal*. 2007; 14(2):227–46. Available from: <http://www.tandfonline.com/doi/abs/10.1080/10705510709336745>
110. Fritz MS, MacKinnon DP. Required sample size to detect the mediated effect. *Psychological Science*. 2007; 18(3):233–9. Available from: <https://doi.org/10.1111/j.1467-9280.2007.01882.x> PMID: 17444920
111. MacKinnon DP, Lockwood CM, Williams J. Confidence limits for the indirect effect: Distribution of the product and resampling methods. *Multivariate Behavioral Research*. 2004; 39(1):99–128. Available from: https://doi.org/10.1207/s15327906mbr3901_4 PMID: 20157642
112. Valente MJ, Gonzalez O, Miočević M, MacKinnon DP. A Note on Testing Mediated Effects in Structural Equation Models: Reconciling Past and Current Research on the Performance of the Test of Joint Significance. *Educational and Psychological Measurement*. 2016; 76(6):889–911. Available from: <https://doi.org/10.1177/0013164415618992> PMID: 27833175