UNIVERSITY^{OF} BIRMINGHAM University of Birmingham Research at Birmingham

Cooking outdoors or with cleaner fuels does not increase malarial risk in children under 5 years

Woolley, Katherine E; Bartington, Suzanne E; Pope, Francis D; Greenfield, Sheila M; Tusting, Lucy S; Price, Malcolm J; Thomas, G Neil

DOI: 10.1186/s12936-022-04152-3

License: Creative Commons: Attribution (CC BY)

Document Version Publisher's PDF, also known as Version of record

Citation for published version (Harvard):

Woolley, KÉ, Bartington, SE, Pope, FD, Greenfield, SM, Tusting, LS, Price, MJ & Thomas, GN 2022, 'Cooking outdoors or with cleaner fuels does not increase malarial risk in children under 5 years: a cross-sectional study of 17 sub-Saharan African countries', *Malaria Journal*, vol. 21, no. 1, 133. https://doi.org/10.1186/s12936-022-04152-3

Link to publication on Research at Birmingham portal

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

•Users may freely distribute the URL that is used to identify this publication.

•Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.

•User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?) •Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

RESEARCH



Cooking outdoors or with cleaner fuels does not increase malarial risk in children under 5 years: a cross-sectional study of 17 sub-Saharan African countries



Katherine E. Woolley¹, Suzanne E. Bartington^{1*}, Francis D. Pope², Sheila M. Greenfield¹, Lucy S. Tusting^{3,4}, Malcolm J. Price^{1,5}, and G. Neil Thomas¹

Abstract

Background: Smoke from solid biomass cooking is often stated to reduce household mosquito levels and, therefore, malarial transmission. However, household air pollution (HAP) from solid biomass cooking is estimated to be responsible for 1.67 times more deaths in children aged under 5 years compared to malaria globally. This cross-sectional study investigates the association between malaria and (i) cleaner fuel usage; (ii) wood compared to charcoal fuel; and, (iii) household cooking location, among children aged under 5 years in sub-Saharan Africa (SSA).

Methods: Population-based data was obtained from Demographic and Health Surveys (DHS) for 85,263 children within 17 malaria-endemic sub-Saharan countries who were who were tested for malaria with a malarial rapid diagnostic test (RDT) or microscopy. To assess the independent association between malarial diagnosis (positive, negative), fuel type and cooking location (outdoor, indoor, attached to house), multivariable logistic regression was used, controlling for individual, household and contextual confounding factors.

Results: Household use of solid biomass fuels and kerosene cooking fuels was associated with a 57% increase in the odds ratio of malarial infection after adjusting for confounding factors (RDT adjusted odds ratio (AOR):1.57 [1.30–1.91]; Microscopy AOR: 1.58 [1.23–2.04]) compared to cooking with cleaner fuels. A similar effect was observed when comparing wood to charcoal among solid biomass fuel users (RDT AOR: 1.77 [1.54–2.04]; Microscopy AOR: 1.21 [1.08–1.37]). Cooking in a separate building was associated with a 26% reduction in the odds of malarial infection (RDT AOR: 0.74 [0.66–0.83]; Microscopy AOR: 0.75 [0.67–0.84]) compared to indoor cooking; however no association was observed with outdoor cooking. Similar effects were observed within a sub-analysis of malarial mesoendemic areas only.

Conclusion: Cleaner fuels and outdoor cooking practices associated with reduced smoke exposure were not observed to have an adverse effect upon malarial infection among children under 5 years in SSA. Further mixed-methods research will be required to further strengthen the evidence base concerning this risk paradigm and to support appropriate public health messaging in this context.

¹ Institute of Applied Health Research, University of Birmingham, Edgbaston, Birmingham, UK

Full list of author information is available at the end of the article



© The Author(s) 2022. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit http://creativecommons.org/ficenses/by/4.0/. The Creative Commons Public Domain Dedication waiver (http://creativecommons.org/publicdomain/zero/1.0/) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

^{*}Correspondence: s.bartington@bham.ac.uk

Keywords: Malaria, Household air pollution, Children under 5 years, Low and middle-income country, Sub-Saharan Africa, Biomass

Background

Smoke arising from solid biomass cooking (wood, dung, charcoal, crop residue) is widely perceived to act as a mosquito repellent among communities [1–3], therefore protecting against mosquito-borne disease. However, solid biomass cooking produces health harming levels of household air pollution (HAP), estimated to be responsible for around 450,000 deaths in children aged under 5 years worldwide [4], compared to only 274,000 estimated deaths from malaria in 2019 [1]. This discordance in perceived compared to actual health risks associated with malarial transmission could impact upon uptake of structural interventions (e.g., cleaner fuel transition [LPG, electricity, biogas]) and behavioural changes intended to reduce harmful HAP exposure, notably among those living in endemic malarial regions.

Sub-Saharan Africa (SSA) has the highest malarial prevalence globally with 94% of cases and deaths, caused by predominantly by *Plasmodium falciparum* [5]. Identified risk factors for malarial infection include poor household construction [6–8] (e.g., open eaves), animals sleeping in the house [9] and presence of standing water near the house [10, 11]. The use of mosquito nets, household insecticidal spraying, and larval source management [12] have become common practice advocated in malarial prevention, driven in part by the World Health Organization's (WHO) coordinated response [5]. Another, common preventive measure is use of mosquito repellent smoke from the burning of certain types of plant materials, such as *churai* in West Africa [2, 13].

There is little evidence supporting reduced mosquito infiltration [14, 15] or malarial transmission associated with solid biomass fuel cooking [2, 16]; although there is some evidence that solid biomass cooking reduces the risk of arboviruses in Guatemala [17]. Therefore, to better understand this disease risk paradigm, this study investigates the association of malarial acquisition among children aged under 5 years with regard to: (i) cleaner or solid biomass fuels and kerosene cooking; (ii) charcoal or wood fuel usage; and (iii) indoor and outdoor cooking, within households in 17 SSA countries using the population-based Demographic and Health Survey (DHS) data.

Methods

Data sources

This cross-sectional study uses publicly available survey data for 17 malarial-endemic SSA countries with

available malarial data (Fig. 1), obtained from the DHS program supported by the United States Agency for International Development (USAID) within the last 10 years (2010-2020). The DHS undertakes full surveys every 5 years, and intermediate Malaria Indicators Surveys (MIS) [18]; only some of the full DHS survey modules undertake malarial testing. For those DHS surveys including malaria modules, malarial testing is undertaken by trained fieldworkers on a sub-sample of eligible children aged 6-59 months using a malarial rapid diagnostic test (RDT) [18]. A two-stage stratified sampling technique was employed to obtain a representative population-based sample, with residential households randomly selected. Eligible households included those with an ever-married (married, widowed or divorced) woman aged between 15 and 49 years and resident the night before the survey. Ethical approval for data collection was gained from the relevant government authority [18], and authorization for data access was provided by the DHS.

Malarial endemicity was generated for each cluster by assessment of malarial prevalence obtained from the open source Malaria Atlas Project [19] within eligible countries, and defined as holoendemic (>75%), hyperendemic (51–75%), mesoendemic (11–50%), hypoendemic (<10%) [20]. Those data points that fell within hypoendemic areas were excluded from the analysis due to lower rate of malarial infection and testing. Malarial prevalence data were geocoded to the cluster geographic coordinates using the spatial analyst tool in ArcMAP 10.7 [21]; a method that has been previously used for this purpose [22].

As the wealth index provided by DHS contains cooking fuel as an indicator variable, a new modified wealth index was calculated in SPSS [23] using principal component analysis [24] to prevent circularity [8]. The index indicator variables included source of drinking water, house construction material, provision of a toilet facility and household assets, which varied by country (Additional file 1).

Predictor and outcome variables

Proxies for household air pollution (HAP) exposure levels

Three analyses were undertaken (Table 1), undertaking comparisons by the main type of cooking fuel used and cooking location respectively: cleaner vs solid biomass fuels and kerosene fuels; charcoal vs wood fuels; outdoor vs indoor cooking (indoors, in a separate building).



Measure of malarial diagnosis

A malarial infection was determined by a positive RDT (n=17 countries) and in some countries a subsequent blood smear test via microscopy taken at the point of interview (n=11 countries), both of which were modelled as a binary (negative, positive) outcome variable, in separate analysis within this study. The RDT was undertaken using the SD BIOLINE Malaria Ag test, in all countries, which has estimated sensitivity of 99.7% and specificity of 99.5% [25]. Whereas, only certain countries collected blood samples which were collected with the parasites detected in the blood at time of survey using microscopy [18], with estimated sensitivity of 95.7% and specificity of 97.9% [26].

Explanatory variables

Covariates were included for the relevant contextual, household and individual factors identified as influencing both HAP exposure and malarial risk. Covariates were included in regression models as categorical variables other than household altitude, which was modelled as a continuous variable. Regional level variables were: malarial endemicity (mesoendemic, hyperendemic and holoendemic), season (dry, wet), rural or urban residence (rural, urban), cluster altitude (metres). Household level variables were: number of household members $(\leq 6, > 6)$, household smoking (no, yes), modified wealth index (lowest, low, middle, high, highest), biomass cooking fuel type (where applicable; kerosene, coal/lignite, charcoal, wood, straw/shrubs/grass, agricultural crop, animal dung), household insecticide spraying within the last 12 months (no, yes) and dwelling construction (traditional, modern). Child variables were: age (<1, 1, 2, 3, 4 years), birth order (first born, not first born), child's sex (male, female), slept under mosquito net last night (no, yes-treated (ITN) net, yes-untreated net). The season variable is created using regional and country level information from the CIA fact book [27] and the World Bank climate change knowledge portal [28]. The household construction variable is a composite variable derived from the wall, roof and floor material. Firstly, each of the three materials were categorized into natural, rudimentary and finished construction material using the criteria outlined by Tusting et al. [8], followed by the creation of the household construction variable where modern household construction was define as wall, roof and floor being made of finished materials.

Table 1	Analyses, sub-analyse:	s and exploratory analyses undertaken wi	th detail on categorisation of the exposu	re of interest	
Analysis	Exposure of interest	Categories	Adjusted for	Sub-analysis	Exploratory analysis controlling for [‡]
Analysis 1	Biomass usage	Cleaner (electricity, LPG, natural gas, biogas) Solid biomass fuels and kerosene (kerosene, coal/lignite, charcoal, wood, straw/ shrubs/grass, agricultural crop, animal dung)	Child's age, child's gender, birth order, Child slept under slept under mosquito net last night, modified wealth index, number of household members, place of residence, malarial endemicity, season, cluster altitude and cooking location	• Urban areas only • Rural areas only • Mesoendemic areas only	 Household insecticidal spraying Household smoking and cooking location
Analysis 2	Biomass fuel type*	• Charcoal • Wood	Child's age, child's gender, birth order, Child slept under slept under mosquito net last night, modified wealth index, number of household members, place of residence, malarial endemicity, season, cluster altitude and cooking location	• Urban areas only • Rural areas only • Mesoendemic areas only	 Household insecticidal spraying Household smoking and cooking location
Analysis 3	Cooking location ⁺	• Outdoors • In a separate building • Indoors	Child's age, child's gender, birth order, Child slept under slept under mosquito net last night, modified wealth index, number of household members, place of residence, malarial endemicity, season, cluster altitude and biomass cooking fuel type	 Urban areas only Rural areas only Mesoendemic areas only Wood cooking only 	 Household insecticidal spraying Household smoking and cooking location
* Charcoal a † Only Solid	and wood are the most corr biomass fuels and kerosen	imonly used type of biomass fuel and are next to e. ie (kerosene, coal/lignite, charcoal, wood, straw/shr	sch other on fuel ladder, with charcoal being relati ubs/grass, agricultural crop, animal dung) were inc	vely less polluting luded in the analysis and includ	ed as a covariate

of into f + h .; . 4 -Jati i+h rtalv 4 _ + _ -4 _ _ < Tahle 1 ⁺ Countries excluded due to the variable being incomplete, high level of missing or low cell counts. For household insecticidal spraying excluded countries were: Burkina Faso 2017–2018, Cameron 2018, DRC 2013–2014, Malawi 2017, Mali 2018, Tanzania 2017 and Togo 2017. For household smoking and cooking location excluded countries were: Burkina Faso 2017–2018, Cameron 2018, DRC 2013–2014, 2018 and Sierra Leone 2018, Tanzania 2017 and Togo 2017. For household smoking and cooking location excluded countries were: Burkina Faso 2017–2018, Cameron 2018, Malawi 2017, Mozambique 2018 and Sierra Leone 2016.

Data analysis

Data preparation and analysis was undertaken in R studio [29]. Each variable was described within the combined dataset using number of cases (n), and percentage (%) and median and Interquartile range (IQR) for continuous variables. The level of missing data ranged from 0 to 48% of clinically relevant variables at a country level, which was imputed using the MICE package [30] with 50 iterations [31, 32]; to prevent bias from list-wise deletion [33]. To test the association between cooking practices and malarial infection, multivariable logistic regression using the survey package [34], was used to account for the complex sampling strategy; reporting adjusted odds ratios (AOR) and 95% confidence intervals (95% CI). The MIS survey did not contain information on cooking location and household smoking, therefore a sub-analysis was undertaken using countries where these variables were available for analysis. Sub-analyses were also undertaken for rural, urban, wood cooking fuel houses and mesoendemic areas. In addition, the analysis was repeated to include additional covariates among a sub-set of countries where additional variables of interest were available. This enabled investigation of the influence of (i) household cooking location; (ii) household smoking; and (iii) household insecticidal spraying, as some of the variables are missing from certain countries.

Results

This study identified 85,263 children aged under 5 years children living in 17 participating countries (DHS=9, MIS=7) from 2011 to 2019, with a total of 74,461 RDT and 48,491 microscopy test results. Within the pooled full dataset, median child age was 3 years (IQR: 2–4). The proportion of girls ranged from 48.0% in Guinea (2012) to 51.0% in Cote d'Ivoire (2011–2012), with overall 49.4% in the pooled dataset (Table 2).

Malarial infection was positively identified by RDT among 34.6% of children in the combined dataset at the time of testing, with the highest point prevalence in Guinea 2012 (51.8%) and lowest in Tanzania 2017 (7.07%) (Table 3). However, where microscopy was undertaken malarial infection was identified in 28.2% of children, with the highest prevalence in Guinea 2012 (48.7%) and lowest in Uganda 2018–2019 (11.3%). Of the areas surveyed, most were in mesoendemic areas (Fig. 2), with holoendemicity in Cote d'Ivoire 2011–2012, DRC 2013–2014, Guinea 2012 and Liberia 2016. Of those children with a positive malarial RDT result, 1.3% resided in cleaner cooking households. Whereas, 35.2% in outdoor cooking households and 35.7% in a household where

cooking was typically undertaken in a separate building (Table 3).

Analysis 1—Solid biomass fuel usage and risk of malarial infection

In pooled analyses, cooking with solid biomass fuels and kerosene fuels was observed to be independently associated with a 57% increase in the adjusted odds ratio for malarial infection, compared to cleaner cooking (electricity, LPG) (Fig. 3) (RDT AOR: 1.57 [1.30-1.914]; Microscopy AOR: 1.58 [1.23-2.04]) (Table 3). A 61% increase in adjusted odds ratio was also observed when investigating the effect of cooking location and household smoking with solid biomass fuels and kerosene compared to cleaner cooking fuels (RDT AOR: 1.61 [1.28-2.02]; Microscopy AOR: 1.61 [1.20-2.15]. The increased malarial infection adjusted odds ratio associated with solid biomass fuels and kerosene cooking remained in the stratified sub-analysis among rural locations (RDT AOR: 1.41 [1.02-1.95]; Microscopy AOR: 2.10 [1.34-3.32]), urban locations (RDT AOR: 1.58 [1.24-2.03] only) and mesoendemic regions (RDT AOR: 1.58 [1.28-1.95]; Microscopy AOR: 1.59 [1.21-2.08]) (Table 4).

Analysis 2—Biomass fuel type and risk of malarial infection Among biomass fuel households only, use of wood compared to charcoal fuel was associated with an increased adjusted odds ratio of malarial infection (RDT AOR: 1.77 [1.54–2.04]; Microscopy AOR: 1.21 [1.08–1.37]) (Fig. 4), with a similar effect being observed in the exploratory analysis controlling for cooking location and household smoking (RDT AOR: 1.26 [1.10–1.46] only) and in mesoendemic areas (RDT AOR: 1.77 [1.49–2.09]; Microscopy AOR: 1.26; [1.10–1.44]) (Table 4). In the stratified sub-analysis it was observed that urban areas had a greater adjusted odds ratio of malarial infection associated with wood compared to charcoal cooking

(RDT AOR: 2.25 [1.79-2.78]), in comparison to rural

Analysis 3—Household cooking location and risk of malarial infection

areas (RDT AOR: 1.43 [1.21-1.70]).

No significant association was observed between household cooking location and malaria adjusted odds ratio (RDT AOR: 0.94 [0.83–1.05]; Microscopy AOR: 0.97 [95% CI 0.83–1.05]) (Fig. 5). In comparison, cooking in a separate building was associated with a reduced adjusted odds ratio of malarial infection by 74% compared to indoor cooking (Fig. 5) (RDT AOR: 0.74 [0.66–0.83]; Microscopy AOR: 0.75 [0.67–0.84]). The same reduced malarial infection adjusted odds ratio associated with cooking in a separate building was observed in stratified

Country	Survey	z	Positive	Positive	Child's age (ye	ars) n (%)				Females (%)
			KDI (%)*	microscopy (%)*	~	-	2	3	4	
Central Africa										
Cameroon 2018	DHS	4417	23.9	I	567 (12.8%)	873 (19.8%)	1056 (23.9%)	1002 (22.7%)	919 (20.8%)	48.7
DRC 2013-2014	DHS	6359	35.9	28.3	868 (13.6%)	1263 (19.9%)	1515 (23.8%)	1390 (21.9%)	1324 (20.8%)	50.1
East Africa										
Burundi 2016–2017	DHS	4309	47.4	33.4	604 (14.0%)	901 (20.9%)	935 (21.7%)	921 (21.4%)	948 (22.0%)	49.5
Malawi 2017	MIS	1929	41.3	I	229 (11.9%)	374 (19.4%)	438 (22.7%)	406 (21.1%)	480 (24.9%)	49.3
Mozambique 2018	MIS	384	45.4	I	507 (13.4%)	769 (20.3%)	944 (25.0%)	810 (21.4%)	753 (19.9%)	49.3
Tanzania 2017	MIS	5882	7.1	I	782 (13.3%)	1197 (20.3%)	1383 (23.5%)	1308 (22.2%)	1212 (20.6%)	49.7
Uganda 2018–2019	MIS	5282	21.0	11.3	631 (11.9%)	1011 (19.1%)	1281 (24.3%)	1228 (23.2%)	1131 (21.4%)	49.5
West Africa										
Benin 2017–2018	DHS	11,981	36.4	39.3	1747 (14.6%)	2390 (19.9%)	2705 (22.6%)	2699 (22.5%)	2440 (20.4%)	49.2
Burkina Faso 2017–2018	MIS	4839	20.8	17.1	645 (13.3%)	877 (18.1%)	1175 (24.3%)	1149 (23.7%)	992 (20.5%)	49.2
Cote d'Ivoire 2011–2012	DHS	3679	50	17.6	550 (14.9%)	749 (20.4%)	932 (25.3%)	808 (22.0%)	640 (17.4%)	51.0
Ghana 2019	MIS	2143	25.9	I	269 (12.6%)	407 (19.0%)	565 (26.4%)	457 (21.3%)	445 (20.8%)	49.1
Guinea 2021	DHS	3022	51.8	48.4	394 (13.0%)	580 (19.2%)	660 (21.8%)	729 (24.1%)	659 (21.8%)	48.0
Liberia 2016	DHS	3074	45.0	I	388 (12.6%)	581 (18.9%)	711 (23.1%)	712 (23.2%)	682 (22.2%)	49.1
Mali 2018	DHS	5159	26.4	Ι	664 (12.9%)	1117 (21.7%)	1224 (23.7%)	1126 (21.8%)	1028 (19.9%)	49.5
Nigeria 2018	DHS	9791	34.8	21.9	1335 (13.6%)	2017 (20.6%)	2273 (23.2%)	2153 (22.0%)	2013 (20.6%)	49.2
Sierra Leone 2016	MIS	6763	52.7	40.1	946 (14.0%)	1226 (18.1%)	1594 (23.6%)	1587 (23.5%)	1411 (20.9%)	50.0
Togo 2017	MIS	2850	44.3	28.8	401 (14.1%)	566 (19.8%)	666 (23.4%)	630 (22.1%)	588 (20.6%)	50.3
N: Number of child observations,	DHS: Demogra	aphic and Hea	ith Survey, MIS: N	Aalaria Indicators Sur	vey, n: number of ch	ild observation with	each category			

* Percentage for positive results based on those children who received a conclusive result from malaria test

Table 3 Descriptive statistics for the combined dataset (N = 85,263)

	Malaria RDT re	sult (N = 74,461)		Malaria Micros	copy results (N =	48,491)
	Negative N = 48,699 (65.4%)	Positive N = 25,761 (34.6%)	p value	Negative N = 34,802 (71.8%)	Positive N = 13,689 (28.2%)	p value
Proxies for HAP exposure levels						
Cooking fuel			< 0.001			< 0.001
Electricity	247 (0.5%)	47 (0.2%)		196 (0.6%)	22 (0.2%)	
LPG	2404 (4.9%)	295 (1.1%)		1287 (3.7%)	98 (0.7%)	
Natural gas	305 (0.6%)	9 (0.0%)		201 (0.6%)	7 (0.1%)	
Biogas	38 (0.1%)	8 (0.0%)		16 (0.0%)	7 (0.0%)	
Kerosene	1256 (2.6%)	220 (0.9%)		927 (2.7%)	107 (0.8%)	
Coal, lignite	155 (0.3%)	42 (0.2%)		103 (0.3%)	24 (0.2%)	
Charcoal	10,043 (20.7%)	2297 (8.9%)		6368 (18.3%)	1500 (11.0%)	
Wood	33,799 (69.5%)	22,397 (87.0%)		25,288 (72.8%)	11,602 (84.8%)	
Other biomass	370 (0.8%)	417 (1.6%)		358 (1.0%)	307 (2.2%)	
No food cooked in house	15 (0.0%)	8 (0.0%)		7 (0.0%)	2 (0.0%)	
Missing	68	22		50	13	
Cooking location			< 0.001			< 0.001
In the house	7108 (29.0%)	4129 (29.1%)		6326 (31.2%)	2830 (32.2%)	
In a separate building	9170 (37.5%)	5068 (35.7%)		6468 (31.9%)	2627 (29.9%)	
Outdoors	8196 (33.5%)	4994 (35.2%)		7482 (36.9%)	3321 (37.8%)	
Missing	24,226	11,571		14,526	4911	
Contextual and contextual variables						
Place of residence			< 0.001			< 0.001
Urban	17,582 (36.1%)	4683 (18.2%)		11,635 (33.4%)	2669 (19.5%)	
Season			< 0.001			< 0.001
Dry	25,169 (51.7%)	11,776 (45.7%)		20,750 (59.6%)	6583 (48.1%)	
Malarial endemicity			< 0.001			< 0.001
Mesoendemic	42,772 (87.8%)	19,018 (73.8%)		29,351 (84.3%)	9457 (69.1%)	
Hyperendemic	5729 (11.8%)	6286 (24.4%)		5116 (14.7%)	3971 (29.0%)	
Holoendemic	198 (0.4%)	457 (1.8%)		335 (1.0%)	261 (1.9%)	
Cluster altitude			< 0.001			< 0.001
Median IQR	294 (85, 596)	321 (156, 590)		322 (149, 764)	324 (149, 588)	
Household level variables						
Modified Wealth Index			< 0.001			< 0.001
Lowest	8669 (17.8%)	7714 (29.9%)		6633 (19.1%)	3976 (29.0%)	
Low	9618 (19.7%)	7306 (28.4%)		6925 (19.9%)	3722 (27.2%)	
Middle	9919 (20.4%)	5698 (22.1%)		6949 (20.0%)	2908 (21.2%)	
High	10,886 (22.4%)	3802 (14.8%)		7724 (22.2%)	2225 (16.3%)	
Highest	9608 (19.7%)	1241 (4.8%)		6569 (18.9%)	859 (6.3%)	
Household smoking			< 0.001			< 0.001
No	20,049 (81.6%)	10,852 (76.1%)		16,195 (79.5%)	6631 (75.1%)	
Missing	24,119	11,497		14,430	4860	
Number of household members			< 0.001			< 0.001
≤ 6	26,538 (54.6%)	13,007 (50.6%)		18,579 (53.5%)	6631 (48.5%)	
Missing	68	44		51	31	
Household insecticide spraying within last 12 months			< 0.001			< 0.001
No	18,189 (91.1%)	13,044 (94.9%)		17,582 (93.3%)	8527 (95.6%)	
Yes	1779 (8.9%)	703 (5.1%)		1260 (6.7%)	394 (4.4%)	
Missing	28,731	12,014		15,960	4768	
House construction			< 0.001			< 0.001

Table 3 (continued)

Malaria RDT res	sult (N = 74,461)		Malaria Micros	copy results (N $=$	48,491)
Negative N = 48,699 (65.4%)	Positive N = 25,761 (34.6%)	p value	Negative N = 34,802 (71.8%)	Positive N = 13,689 (28.2%)	p value
28,361 (58.2%)	19,352 (75.1%)		20,902 (60.1%)	10,056 (73.5%)	
20,338 (41.8%)	6410 (24.9%)		13,900 (39.9%)	3634 (26.5%)	
		< 0.001			< 0.001
7643 (15.7%)	2282 (8.9%)		5319 (15.3%)	1272 (9.3%)	
10,335 (21.2%)	4404 (17.1%)		7359 (21.1%)	2186 (16.0%)	
11,266 (23.1%)	6328 (24.6%)		8127 (23.4%)	3239 (23.7%)	
10,254 (21.1%)	6497 (25.2%)		7425 (21.3%)	3568 (26.1%)	
9201 (18.9%)	6251 (24.3%)		6572 (18.9%)	3424 (25.0%)	
		< 0.001			
14,376 (33.7%)	5338 (24.5%)		9392 (30.8%)	2553 (22.1%)	
6102	3993		4306	2164	
		0.068			
24,535 (50.4%)	13,112 (50.9%)		17,489 (50.3%)	6971 (50.9%)	
		< 0.001			< 0.001
20,615 (42.3%)	12,078 (46.9%)		15,858 (45.6%)	6942 (50.7%)	
26,991 (55.4%)	13,204 (51.3%)		18,320 (52.6%)	6525 (47.7%)	
1093 (2.2%)	480 (1.9%)		624 (1.8%)	222 (1.6%)	
	Malaria RDT res Negative N = 48,699 (65.4%) 28,361 (58.2%) 20,338 (41.8%) 7643 (15.7%) 10,335 (21.2%) 11,266 (23.1%) 10,254 (21.1%) 9201 (18.9%) 14,376 (33.7%) 6102 24,535 (50.4%) 20,615 (42.3%) 20,615 (42.3%) 1093 (2.2%)	Malaria RDT result (N = 74,461) Negative N = 48,699 (65.4%) Positive N = 25,761 (34.6%) 28,361 (58.2%) 19,352 (75.1%) 20,338 (41.8%) 6410 (24.9%) 7643 (15.7%) 2282 (8.9%) 10,335 (21.2%) 4404 (17.1%) 11,266 (23.1%) 6328 (24.6%) 10,254 (21.1%) 6497 (25.2%) 9201 (18.9%) 6251 (24.3%) 14,376 (33.7%) 5338 (24.5%) 6102 3993 24,535 (50.4%) 13,112 (50.9%) 20,615 (42.3%) 12,078 (46.9%) 20,991 (55.4%) 13,204 (51.3%) 1093 (2.2%) 480 (1.9%)	Malaria RDT result (N = 74,461) Negative N = 48,699 (65.4%) Positive N = 25,761 (34.6%) p value 28,361 (58.2%) 19,352 (75.1%) 4403 (10 (24.9%) 20,338 (41.8%) 6410 (24.9%) 20,338 (41.8%) 6410 (24.9%) 7643 (15.7%) 2282 (8.9%) 10,335 (21.2%) 4404 (17.1%) 11,266 (23.1%) 6328 (24.6%) 9201 (18.9%) 6251 (24.3%) 9201 (18.9%) 5338 (24.5%) 6102 3993 14,376 (33.7%) 5338 (24.5%) 6102 3993 24,535 (50.4%) 13,112 (50.9%) 20,615 (42.3%) 12,078 (46.9%) 20,991 (55.4%) 13,204 (51.3%) 1093 (2.2%) 480 (1.9%)	Malaria RDT result (N = 74,461) Malaria Microsol Negative N = 48,699 (65.4%) Positive N = 25,761 (34.6%) p value Negative N = 34,802 (71.8%) 28,361 (58.2%) 19,352 (75.1%) 20,902 (60.1%) 20,338 (41.8%) 6410 (24.9%) 13,900 (39.9%) 28,361 (55.7%) 2282 (8.9%) 5319 (15.3%) 10,335 (21.2%) 4404 (17.1%) 7359 (21.1%) 11,266 (23.1%) 6328 (24.6%) 8127 (23.4%) 10,254 (21.1%) 6497 (25.2%) 7425 (21.3%) 9201 (18.9%) 6251 (24.3%) 6572 (18.9%) 14,376 (33.7%) 5338 (24.5%) 9392 (30.8%) 6102 3993 4306 24,535 (50.4%) 13,112 (50.9%) 17,489 (50.3%) 20,615 (42.3%) 12,078 (46.9%) 18,320 (52.6%) 20,991 (55.4%) 13,204 (51.3%) 18,320 (52.6%) 1093 (2.2%) 480 (1.9%) 624 (1.8%)	Malaria RDT result (N = 74,461) Malaria Microscy results (N = Negative N = 48,699 (65.4%) Positive N = 25,761 (34.6%) p value Negative N = 34,802 (71.8%) Positive N = 13,689 (28.2%) 28,361 (58.2%) 19,352 (75.1%) 20,902 (60.1%) 10,056 (73.5%) 20,338 (41.8%) 6410 (24.9%) 13,900 (39.9%) 3634 (26.5%) 20,338 (41.8%) 6410 (24.9%) 5319 (15.3%) 1272 (9.3%) 7643 (15.7%) 2282 (8.9%) 5319 (15.3%) 1272 (9.3%) 10,335 (21.2%) 4404 (17.1%) 7359 (21.1%) 2186 (16.0%) 11,266 (23.1%) 6328 (24.6%) 8127 (23.4%) 3239 (23.7%) 10,254 (21.1%) 6497 (25.2%) 7425 (21.3%) 3424 (25.0%) 9201 (18.9%) 6251 (24.3%) 6572 (18.9%) 3424 (25.0%) 6102 3993 <0.068

N: Number of observations; %: column percentage for categorical variables; IQR: interquartile range; ITN: insecticide-treated nets; RDT: rapid diagnostic test





sub-analyses for wood cooking (RDT AOR: 0.75 [0.67–0.85]; Microscopy AOR: 0.77 [0.67–0.87]), rural (RDT AOR: 0.70 [0.62–0.80]; Microscopy AOR: 0.73 [0.64–0.84]) and mesoendemic areas (RDT AOR: 0.73 [0.65–0.82]; Microscopy AOR: 0.74 [0.65–0.83) only (Table 4).

Discussion

This large exploratory study of over 85,000 children aged under 5 years living in 17 malaria-endemic SSA found no evidence to suggest that use of cleaner fuels (e.g., LPG, electricity, biogas), charcoal vs wood, or outdoor cooking location are associated with an increased risk of malarial infection. Indeed, the findings suggest that solid biomass fuel usage may be associated with a higher incidence of malarial infection among children in SSA. There are a number of factors that may account for the increase in infections, such as the longer cooking times and thus of carbon dioxide production [35], a major mosquito attractant [36], found with solid biomass fuel cooking [37]. Additionally, the use of solid biomass fuels, particularly wood, crop residue and dung, require women, to typically collect fallen or harvest branches from woods and forests where mosquitoes commonly reside, often taking children under 5 years on their backs, thereby increasing risk of mosquito bites.

It is highly likely that risk of within household acquisition of malaria is also influenced by socioeconomic factors such as household construction characteristics (eaves space, wall type) and living conditions [8, 38–41] which are not fully captured in the DHS composite wealth index. It is also recognized that use of cleaner domestic energy sources, cooking in a separate building and selection of biomass cooking fuel type may reflect socio-economic determinants, also related to malarial microepidemiology at the household level [42, 43]. The child's age is also a key factor in malarial infection risk, with an observed increased risk with increasing age, potentially reflecting behavioural, nutritional or exposure differences. In terms of modifiable factors for malarial infection prevention and control, there is strong evidence supporting the sustained use of ITN bed nets, larval source management and household insecticide spraying [12]; of which only ITN bed nets could be controlled for in the main analyses. The importance of household insecticide spraying can be seen in the subsidiary analysis undertaken among countries for which this information was available, identifying that there was no association with type of biomass fuel and malarial infection risk (RDT: AOR 1.23 [0.94-1.61]; Microscopy AOR: 1.07 [0.77-1.47]; Table 4); however, this sub-analysis is likely to be underpowered and should be interpreted with caution.

The analyses presented also did not explore broader contextual factors associated with household or village level clustering of malarial transmission, including position of households in relation to mosquito sites and local attitudes to malarial treatment which are recognized to influence local variations in malarial prevalence [44]. The DHS dataset did not contain information on cooking practices such as timing or duration, both of which

			במרוו הססמוו וא שומר			והוווכם ממומזר	י, באטוטומנטו א מוזי	מוומ מחכיו	cicki				
Analysis	Outcome	Analysis 1 Biomass vs cle	eaner cooking			Analysis 2 Wood vs charc	oal cooking:			Analysis 3 Cooking location			
		Cooking fuel	AOR [95% CI]	<i>p</i> value	z	Cooking fuel	AOR [95% CI]	<i>p</i> value	z	Type of cooking location	AOR [95% CI]	<i>p</i> value	z
Combined	RDT	Cleaner	Ref.			Charcoal	Ref.			Indoor	Ref.		
dataset*		Biomass	1.57 [1.30–1.91]	< 0.001	43,759	Wood	1.77 [1.54–2.04]	< 0.001	73,072	In a separate building	0.74 [0.66–0.83]	< 0.001	23,754
										Outdoor	0.94 [0.83-1.05]	0.26	
	Microscopy	Cleaner	Ref.			Charcoal	Ref.			Indoor	Ref.		
		Biomass	1.58 [1.23–2.04]	< 0.001	30,007	Wood	1.21 [1.08–1.37]	0.001	46,206	ln a separate building	0.75 [0.67–0.84]	< 0.001	21,383
Sub-analysis										Outdoor	0.97 [0.86–1.09]	0.58	
Rural areas	RDT	Cleaner	Ref.			Charcoal	Ref.			Indoor	Ref.		
		Biomass	1.41 [1.02–1.95]	0.04	31,100	Wood	1.43 [1.21–1.70]	< 0.001	54,473	ln a separate building	0.70 [0.62–0.80]	< 0.001	16,988
										Outdoor	0.91 [0.79–1.04]	0.17	
	Microscopy	Cleaner	Ref.			Charcoal	Ref.			Indoor	Ref.		
		Biomass	2.10 [1.34–3.32]	0.001	20,290	Wood	1.09 [0.91–1.30]	0.36	34,693	ln a separate building	0.73 [0.64–0.84]	< 0.001	15,193
										Outdoor	0.92 [0.80–1.07]	0.28	
Urban areas	RDT	Cleaner	Ref.			Charcoal	Ref.			Indoor	Ref.		
		Biomass	1.58 [1.24–2.03]	< 0.001	12,659	Wood	2.23 [1.79–2.78]	< 0.001	18,599	In a separate building	0.96 [0.78–1.19]	0.72	6766
										Outdoor	0.99 [0.81–1.21]	0.90	
	Microscopy	Cleaner	Ref.			Charcoal	Ref.			Indoor	Ref.		
		Biomass	1.30 [0.96–1.76]	0.09	9717	Wood	1.40 [1.20–1.64]	< 0.001	11,513	In a separate building	0.86 [0.69–1.07]	0.17	6190
										Outdoor	1.08 [0.90–1.31]	0.40	
Mesoendemic	RDT	Cleaner	Ref.			Charcoal	Ref.			Indoor	Ref.		
areas		Biomass	1.58 [1.28–1.95]	< 0.001	35,167	Wood	1.77 [1.49–2.09]	< 0.001	57,814	In a separate building	0.73 [0.65–0.82]	< 0.001	20,349
										Outdoor	0.92 [0.81–1.05]	0.22	
	Microscopy	Cleaner	Ref.			Charcoal	Ref.			Indoor	Ref.		
		Biomass	1.59 [1.21–2.08]	0.001	23,519	Wood	1.26 [1.10–1.44]	0.001	35,898	In a separate building	0.73 [0.65–0.83]	< 0.001	18,209
										Outdoor	0.94 [0.83–1.08]	0.37	

_ data 2.2 4 ç ÷ . ÷ J J . _ J ÷. 044 5

Ambuic	Current C	Ambraic 1			ColordonA				c sinde			
Analysis	Outcome	Analysis I Biomass vs c	leaner cooking:		Analysis z Wood vs cha	ırcoal cooking		ξ Ω	ialysis 3 oking location			
		Cooking fue	el AOR [95% CI]	<i>p</i> value N	Cooking fue	el AOR [95% CI]	<i>p</i> value N	<u>2 0</u>	pe of cooking cation	AOR [95% CI]	<i>p</i> value	z
Wood only	RDT							Inc	door	Ref.		
								u nq	a separate iilding	0.75 [0.67–0.85]	< 0.001	19,406
								0	ıtdoor	0.90 [0.79–1.02]	0.10	
	Microscopy							Inc	door	Ref.		
								nd U	a separate iilding	0.77 [0.67–0.87]	< 0.001	17,244
								O	ıtdoor	0.94 [0.82–1.08]	0.36	
Exploratory analysis		ī			ī					, t		
Controlling for	RDT	Cleaner	Ref.		Charcoal	Ref.		Inc	door	Ref.		
household mos- quito spraying [†]		Biomass	1.23 [0.94–1.61]	0.14 26,7	78 Wood	1.94 [1.62–2.33]	< 0.001 3€	nl 898,i bu	a separate iilding	0.85 [0.73–0.99]	0.03	9951
								O	utdoor	0.95 [0.81–1.10]	0.47	
	Microscopy	. Cleaner	Ref.		Charcoal	Ref.		Inc	door	Ref.		
		Biomass	1.07 [0.77–1.47]	0.69 18,1	32 Wood	1.30 [1.13–1.49]	< 0.001 27	7,115 In bu	a separate iilding	0.76 [0.65–0.88]	< 0.001	9676
								O	ıtdoor	0.92 [0.79–1.07]	0.29	
AOR Adjusted odds ra Table S3.1	atio, <i>95% Cl</i> 95%	confidence inter	rval, N Number of obs	ervations, <i>RDT</i> Rap	oid diagnostic test,	Ref Reference group. R	esults in bold a	are statisti	cally significant. Un.	adjusted results are in	Additiona	I file 3:
* Controlled for: Child	′'s age, child's g∈	ender, birth order	r, Child slept under sle	pt under mosquit	o net last night, mo	odified wealth index, nu	umber of hous	ehold mei	mbers, place of resic	dence, malarial endem	iicity, seaso	on and
⁺ Burkina Faso 2017–2 level of missing or low	2018, Cameron v cell counts	2018, DRC 2013-	-2014, Malawi 2017, M	ali 2018, Nigeria 2	018, Tanzania 2017	' and Togo 2017 were e	xcluded due to	o the hous	ehold mosquito spi	aying variable being i	ncomplete	, high

Woolley et al. Malaria Journal (2022) 21:133





influence the amount of smoke produced and therefore HAP exposure, and may also generate higher localized levels of indoor CO_2 [35] thereby attracting mosquitoes into the home [36]. In addition, season could only be accounted for at country or broader regional level, which does not take into account microclimates, in addition, the DHS is normally undertaken in the dry season and the MIS in the wet season when the malarial transmission risk is increased [18]. HAP interventions should be developed to include activities which communicate that cooking practices which produce less smoke do not increase risk of malaria transmission to residents. It is also important to reinforce health protection advice regarding evidence-based measures for mosquito control. Further qualitative and quantitative research is merited, for a detailed investigation of the relationships between cooking location, fuel choice and risk of malarial acquisition, considering a wider range of transmission risk factors at a local level.

The rural-urban differences in cooking activity patterns, which can be most clearly noted within the differences observed in distribution between fuel types, is likely to reflect relative economic development, improved access to cleaner fuel sources in urban areas and reduced potential for cohabitation with livestock [45]. However, the rural-urban divide was not as distinct within the cleaner fuel or cooking location sub-analysis, indicating that other contextual and compositional factors exist which may influence malarial infection risk (e.g., nutrition). Although season, malarial endemicity and altitude were captured as confounding factors within our analyses, information was not available for other contextual factors of relevance to malarial infection risk, such as temperature [46].

Additionally, although the cooking practices are reported at the time of interview, this survey question does not take into consideration longer-term trends which may vary on a seasonal basis. Further prospective research is required to better understand environmental influences upon malarial microepidemiology including objective pollutant exposure assessment, capture of household design characteristics, lifestyle and time-activity factors to assess relationships with mosquito breeding conditions, malarial parasitaemia and outcomes among adults and children.

Conclusion

This large-scale observational study suggests that use of cleaner fuels and outdoor cooking practices typically associated with lower levels of household smoke, were not associated with an increased malarial acquisition risk among children living in SSA. Further mixed-methods research is required to better understand the relationships between cooking practices, cooking fuel emissions, mosquito activity and risk of malarial acquisition at household and community levels in this world region.

Abbreviations

AOR: Adjusted odds ratio; DHS: Demographic and Health Survey; DRC: Democratic Republic of Congo; HAP: Household air pollution; ITN: Insecticidetreated net; LPG: Liquefied petroleum gas; MICE: Multiple imputation by chained equations; MIS: Malaria indicator Survey; RDT: Rapid diagnostic test; SSA: Sub-Saharan Africa; USAID: United States Agency for International Development; 95% CI: 95% Confidence interval.

Supplementary Information

The online version contains supplementary material available at https://doi. org/10.1186/s12936-022-04152-3.

Additional file 1: Table S1.1. Predictors included with the PCA analysis for the modified wealth index by country.

Additional file 2: Table S2.1. Unadjusted and adjusted odds ratio of malarial infection with solid biomass fuels and kerosene cooking compared to cleaner cooking—Analysis 1. **Table S2.2.** Unadjusted and adjusted odds ratio of malarial infection with wood cooking compared to charcoal cooking—Analysis 2. **Table S2.3.** Unadjusted and adjusted odds ratio of malarial infection with cooking location (outdoor, in a separate building) compared to indoors—Analysis 3.

Additional file 3: Table S3.1. Unadjusted odds ratio of malarial infection for each cooking practices for the combined dataset, exploratory and sub-analysis.

Acknowledgements

We are grateful for the access to and use of the DHS data.

Author contributions

KEW: Conceptualization; methodology; data curation, formal analysis, visualization and roles/writing—original draft. SEB and GNT: conceptualization; supervision and writing—review and editing. MJP: methodology and writing—review and editing. FDP and SG: supervision and writing—review and editing. LST: writing—review and editing. All authors read and approved the final manuscript.

Funding

KEW is funded by a University of Birmingham Global Challenges Scholarship. MJP is supported by the NIHR Birmingham Biomedical Research Centre. The views expressed are those of the authors and not necessarily those of the NHS, the NIHR or the Department of Health and Social Care.

Availability of data and materials

The data that support the findings of this study freely and publicly available from https://dhsprogram.com/data/ and https://malariaatlas.org/.

Declarations

Ethics approval and consent to participate

Not applicable as no data was collected as part of this study; authorization was given by DHS to access the online data archive. Details on ethical approval and consent are available from https://dhsprogram.com.

Consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests.

Author details

¹ Institute of Applied Health Research, University of Birmingham, Edgbaston, Birmingham, UK. ²School of Geography, Earth and Environmental Sciences, University of Birmingham, Edgbaston, Birmingham, UK. ³Department of Disease Control, London School of Hygiene and Tropical Medicine, London, UK. ⁴ Centre on Climate Change and Planetary Health, London School of Hygiene & Tropical Medicine, London, UK. ⁵NIHR Birmingham Biomedical Research Centre, University Hospitals Birmingham NHS Foundation Trust and University of Birmingham, UK.

Received: 21 October 2021 Accepted: 6 April 2022 Published online: 27 April 2022

References

- Tamire M, Addissie A, Skovbjerg S, Andersson R, Lärstad M. Socio-cultural reasons and community perceptions regarding indoor cooking using biomass fuel and traditional stoves in rural Ethiopia: a qualitative study. Int J Environ Res Public Health. 2018;15:2035.
- Biran A, Smith L, Lines J, Ensink J, Cameron M. Smoke and malaria: are interventions to reduce exposure to indoor air pollution likely to increase exposure to mosquitoes? Trans R Soc Trop Med Hyg. 2007;101:1065–71.
- Devakumar D, Qureshi Z, Mannell J, Baruwal M, Sharma N, Rehfuess E, et al. Women's ideas about the health effects of household air pollution, developed through focus group discussions and artwork in Southern Nepal. Int J Environ Res Public Health. 2018;15:248.
- 4. Global Burden of Disease. GBD Compare, IHME Viz Hub. 2019. https:// vizhub.healthdata.org/gbd-compare/: Accessed Dec 2020.
- WHO. Malaria. Geneva: World Health Organization; 2020. https://www. who.int/news-room/fact-sheets/detail/malaria: Accessed Feb 2021.
- Gunawardena DM, Wickremasinghe AR, Muthuwatta L, Weerasingha S, Rajakaruna J, Senanayaka T, et al. Malaria risk factors in an endemic region of Sri Lanka, and the impact and cost implications of risk factor-based interventions. Am J Trop Med Hyg. 1998;58:533–42.
- Konradsen F, Amerasinghe P, Van Der Hoek W, Amerasinghe F, Perera D, Piyaratne M. Strong association between house characteristics and malaria vectors in Sri Lanka. Am J Trop Med Hyg. 2003;68:177–81.
- Tusting LS, Bottomley C, Gibson H, Kleinschmidt I, Tatem AJ, Lindsay SW, et al. Housing improvements and malaria risk in Sub-Saharan Africa: a multi-country analysis of survey data. PLoS Med. 2017;14:e1002234.
- 9. Ghebreyesus TA, Haile M, Witten KH, Getachew A, Yohannes M, Lindsay SW, et al. Household risk factors for malaria among children in the Ethiopian highlands. Trans R Soc Trop Med Hyg. 2000;94:17–21.
- Hajison PL, Feresu SA, Mwakikunga BW. Malaria in children under-five: a comparison of risk factors in lakeshore and highland areas, Zomba district, Malawi. PLoS ONE. 2018;13:e0207207.
- Abossie A, Yohanes T, Nedu A, Tafesse W, Damitie M. Prevalence of malaria and associated risk factors among febrile children under five years: a cross-sectional study in arba minch zuria district, south Ethiopia. Infect Drug Resist. 2020;13:363–72.
- 12. Tizifa TA, Kabaghe AN, McCann RS, van den Berg H, Van Vugt M, Phiri KS. Prevention efforts for malaria. Curr Trop Med Rep. 2018;5:41–50.
- Vernède R, van Meer M, Alpers M. Smoke as a form of personal protection against mosquitos, a field study in Papua New Guinea. Southeast Asian J Trop Med Public Health. 1994;25:771–5.
- 14. Kaindoa EW, Mkandawile G, Ligamba G, Kelly-Hope LA, Okumu FO. Correlations between household occupancy and malaria vector biting risk in rural Tanzanian villages: implications for high-resolution spatial targeting of control interventions. Malar J. 2016;15:199.
- 15 Hennessee I, Kirby M, Misago X, Umupfasoni J, Clasen T, Kitron U, et al. Assessing the effects of cooking fuels on *Anopheles* mosquito behavior: an experimental study in rural Rwanda. Am J Trop Med Hyg. 2022;106:1196–208 (online ahead of print).
- Snow RW, Bradley AK, Hayes R, Byass P, Greenwood BM. Does woodsmoke protect against malaria? Ann Trop Med Parasitol. 1987;81:449–51.
- Madewell ZJ, Madewell ZJ, Madewell ZJ, López MR, Espinosa-Bode A, Brouwer KC, et al. Inverse association between dengue, chikungunya, and Zika virus infection and indicators of household air pollution in

Santa Rosa, Guatemala: a case-control study, 2011–2018. PLoS ONE. 2020;15:e0234399.

- Croft T, Marshall AMJ, Courtney AK, et al. Guide to DHS statistics. Rockville: ICF. 2018. http://www.measuredhs.com/pubs/pdf/DHSG1/Guide_to_ DHS_Statistics_29Oct2012_DHSG1.pdf. http://citeseerx.ist.psu.edu/viewd oc/download?doi=10.1.1.431.8235&rep=rep1&type=pdf. Accessed Jan 2020.
- Malaria Atlas Project. Welcome to the Malaria Atlas Project—MAP. 2021. https://malariaatlas.org/. Accessed Feb 2021.
- WHO. Report on the malaria conference in Equatorial Africa, held under the joint auspices of the World Health Organization and the commission for technical co-operation in Africa South of the Sahara. Kampala, Uganda, 27 November–9 December 1950. Geneva: World Health Organization; 1951. https://apps.who.int/iris/handle/10665/40153. Accessed Feb 2021.
- 21. ESRI. ArcMap 10.7. Redlands: Environmental Systems Research Institute; 2019.
- Florey L, Taylor C. Using household survey data to explore the effects of improved housing conditions on malaria infection in children in Sub-Saharan Africa. ICF International; 2016. https://dhsprogram.com/pubs/ pdf/AS61/AS61.pdf: Accessed Feb 2021.
- IBM Corp. IBM SPSS statistics for Windows. Version 27.0. Armonk: IBM Corp; 2020.
- 24 Rutstein SO. Steps to constructing the new DHS Wealth Index. Rockville: ICF International; 2015.
- MedicalExpo. Standard diagnostics. https://www.medicalexpo.com/ prod/standard-diagnostics-70168.html. Accessed Mar 2021.
- Rodulfo H, De Donato M, Mora R, González L, Contreras CE. Comparison of the diagnosis of malaria by microscopy, immunochromatography and PCR in endemic areas of Venezuela. Braz J Med Biol Res. 2007;40:535–43.
- Central Intelligence Agency (CIA). The world factbook. 2019. https:// www.cia.gov/library/publications/the-world-factbook/. Accessed Jan 2020.
- 28. World Bank Group. World Bank climate change knowledge portal. https:// climateknowledgeportal.worldbank.org/. Accessed July 2021.
- 29. R Core Team. R: a language and environment for statistical computing. R version 3.6.0. Vienna: R Foundation for Statistical Computing; 2020.
- van Buuren S, Groothuis-Oudshoorn K. mice: Multivariate imputation by chained equations in R. J Stat Softw. 2011;45:1–67.
- 31. Bodner TE. What improves with increased missing data imputations? Struct Equ Model A Multidiscip J. 2008;15(651–75):2.
- 32. White IR, Royston P, Wood AM. Multiple imputation using chained equations: issues and guidance for practice. Stat Med. 2011;30:377–99.
- Jakobsen JC, Gluud C, Wetterslev J, Winkel P. When and how should multiple imputation be used for handling missing data in randomised clinical trials—a practical guide with flowcharts. BMC Med Res Methodol. 2017;17:162.
- 34. Lumley T. Survey: analysis of complex survey samples. R package version 4.0. 2020.
- 35. Sharma D, Jain S. Impact of intervention of biomass cookstove technologies and kitchen characteristics on indoor air quality and human exposure in rural settings of India. Environ Int. 2019;123:240–55.
- 36. Cardé RT. Multi-cue integration: how female mosquitoes locate a human host. Curr Biol. 2015;25:R793–5.
- Chakraborty D, Mondal NK, Datta JK. Indoor pollution from solid biomass fuel and rural health damage: a micro-environmental study in rural area of Burdwan, West Bengal. Int J Sustain Built Environ. 2014;3:262–71.
- Snyman K, Mwangwa F, Bigira V, Kapisi J, Clark TD, Osterbauer B, et al. Poor housing construction associated with increased malaria incidence in a cohort of young Ugandan children. Am J Trop Med Hyg. 2015;92:1207–13.
- Jatta E, Jawara M, Bradley J, Jeffries D, Kandeh B, Knudsen JB, et al. How house design affects malaria mosquito density, temperature, and relative humidity: an experimental study in rural Gambia. Lancet Planet Health. 2018;2:e498-508.
- 40 Wanzirah H, Tusting LS, Arinaitwe E, Katureebe A, Maxwell K, Rek J, et al. Mind the gap: house structure and the risk of malaria in Uganda. PLoS ONE. 2015;10:e0117396.
- Tusting LS, Ippolito MM, Willey BA, Kleinschmidt I, Dorsey G, Gosling RD, et al. The evidence for improving housing to reduce malaria: a systematic review and meta-analysis. Malar J. 2015;14:209.

- 42. Custodio E, Descalzo MÁ, Villamor E, Molina L, Snchez I, Lwanga M, et al. Nutritional and socio-economic factors associated with *Plasmodium falciparum* infection in children from Equatorial Guinea: results from a nationally representative survey. Malar J. 2009;8:225.
- Guerra M, de Sousa B, Ndong-Mabale N, Berzosa P, Arez AP. Malaria determining risk factors at the household level in two rural villages of mainland Equatorial Guinea. Malar J. 2018;17:203.
- 44. Greenwood BM. The microepidemiology of malaria and its importance to malaria control. Trans R Soc Trop Med Hyg. 1989;83(Suppl):25–9.
- Colbeck I, Nasir ZA, Ali Z. Characteristics of indoor/outdoor particulate pollution in urban and rural residential environment of Pakistan. Indoor Air. 2010;20:40–51.
- Dabaro D, Birhanu Z, Negash A, Hawaria D, Yewhalaw D. Effects of rainfall, temperature and topography on malaria incidence in elimination targeted district of Ethiopia. Malar J. 2021;20:104.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions

