

# The difference between effect modification and covariate confounding

Marlin, Nadine; Allotey, John

DOI:

[10.1111/1471-0528.16703](https://doi.org/10.1111/1471-0528.16703)

License:

Creative Commons: Attribution (CC BY)

*Document Version*

Publisher's PDF, also known as Version of record

*Citation for published version (Harvard):*

Marlin, N & Allotey, J 2021, 'The difference between effect modification and covariate confounding', *BJOG: An International Journal of Obstetrics & Gynaecology*, vol. 128, no. 10, pp. 1574-1574. <https://doi.org/10.1111/1471-0528.16703>

[Link to publication on Research at Birmingham portal](#)

## General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

## Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact [UBIRA@lists.bham.ac.uk](mailto:UBIRA@lists.bham.ac.uk) providing details and we will remove access to the work immediately and investigate.



## The difference between effect modification and covariate confounding

**NADINE MARLIN**, PRAGMATIC CLINICAL TRIALS UNIT, BARTS AND THE LONDON SCHOOL OF MEDICINE AND DENTISTRY, QUEEN MARY UNIVERSITY

LONDON, LONDON, UK **JOHN ALLOTEY**, WOMEN'S HEALTH RESEARCH UNIT, BARTS AND THE LONDON SCHOOL OF MEDICINE AND DENTISTRY, QUEEN MARY UNIVERSITY OF LONDON, LONDON, UK

BJOG  
PERSPECTIVES

BJOG

RESEARCH METHODS GUIDES  
Series edited by Karla Hemming  
and Natalie AM Cooper

### Learning points

- When effect modification is present the effect of a risk factor (such as having gestational diabetes) on an outcome (such as cardiovascular disease) will be different in different subgroups (for example by those who are overweight).
- Effect modification can also be investigated in randomised trials to identify if the effect of a treatment differs between subgroups.

Fadl et al.<sup>1</sup> describe a case-control study evaluating whether gestational diabetes mellitus (GDM) is associated with increased risk of future cardiovascular disease (CVD) and whether this risk differs for different subgroups of women. One subgroup they looked at was weight status (normal or overweight). It was concluded that GDM is indeed a useful marker of CVD risk, and the increase in risk of CVD indicated by GDM is higher in women who are overweight. It was, therefore, concluded that weight status is an effect modifier.

Effect modification differs from confounding. Confounding occurs when a variable to some extent accounts for the observed effect of a risk factor. In our example, this would mean that the association between GDM and CVD could be explained by the fact that women with GDM are more likely to be overweight whereas women without GDM are more likely to be of normal weight. Under this scenario, the observed increase in risk of women with GDM would actually be representing the association between weight status and CVD. This can

be modelled by including weight status (overweight or normal weight) as an independent variable in a regression model. Such a model would assume that having GDM increases the CVD risk to the same degree for normal and overweight women.

If, however, having GDM increases the risk of CVD more in overweight women than in normal weight women then weight status is an effect modifier. For example, the value of GDM as a predictor of risk might be stronger in overweight compared with normal weight women. Effect modification can be investigated informally by performing linear regression on separate subgroups. In our example, we would perform two separate linear regressions of the characteristic GDM on CVD, one for 'overweight' women and a second for 'normal weight' women (our two subgroups). In a more formal approach, effect modification can be tested for in multivariable linear regression by including the 'interaction between weight status (overweight or normal weight) and GDM' as an independent variable (an interaction test). Then the *P* value for an interaction test indicates whether the two effect sizes are significantly different, i.e. whether weight status is an effect modifier.

Effect estimates are usually presented within subgroups (or stratified by subgroups), for example the odds ratio for GDM as a risk factor of CVD within normal weight women and separately within overweight women.

Analysis models accounting for effect modifiers allow the researcher to identify population groups with higher risk or, in a

clinical trial setting, participants for whom an intervention might be more effective. For example, we might wish to evaluate whether labetalol is less effective for treating gestational hypertension in different ethnic subgroups. Subgroups of interest should be specified a priori, clinically plausible and limited, thereby avoiding spurious results due to chance.

### Useful resources

- <https://www.youtube.com/watch?v=stCCd7CUTVY>
- TJ Vander Weele. Confounding and effect modification: distribution and measure. *Epidemiol Method* 2012;1(1):55–82

### Disclosure of interest

Completed disclosure of interests form available to view online as supporting information.

### Supporting Information

Additional supporting information may be found online in the Supporting Information section at the end of the article. ■

### Reference

- 1 Fadl H, Magnuson A, Östlund I, Montgomery S, Hanson U, Schwarcz E. Gestational diabetes mellitus and later cardiovascular disease: a Swedish population based case-control study. *BJOG* 2014;121:1530–6.