

Are multiple mitochondrial related signalling pathways involved in carotid body oxygen sensing?

Holmes, Andrew P; Swiderska, Agnieszka; Nathanael, Demitris; Aldossary, Hayyaf S; Ray, Clare J; Coney, Andrew M; Kumar, Prem

DOI:

[10.3389/fphys.2022.908617](https://doi.org/10.3389/fphys.2022.908617)

License:

Creative Commons: Attribution (CC BY)

Document Version

Publisher's PDF, also known as Version of record

Citation for published version (Harvard):

Holmes, AP, Swiderska, A, Nathanael, D, Aldossary, HS, Ray, CJ, Coney, AM & Kumar, P 2022, 'Are multiple mitochondrial related signalling pathways involved in carotid body oxygen sensing?', *Frontiers in Physiology*, vol. 13, 908617. <https://doi.org/10.3389/fphys.2022.908617>

[Link to publication on Research at Birmingham portal](#)

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.



Are Multiple Mitochondrial Related Signalling Pathways Involved in Carotid Body Oxygen Sensing?

Andrew P. Holmes^{1*}, Agnieszka Swiderska², Demitris Nathanael¹, Hayyaf S. Aldossary^{1,3}, Clare J. Ray¹, Andrew M. Coney¹ and Prem Kumar^{1*}

¹School of Biomedical Sciences, Institute of Clinical Sciences, College of Medical and Dental Sciences, University of Birmingham, Birmingham, United Kingdom, ²Unit of Cardiac Physiology, Division of Cardiovascular Sciences, Faculty of Biology, Medicine and Health, University of Manchester, Manchester, United Kingdom, ³College of Medicine, Basic Medical Sciences, King Saud bin Abdulaziz University for Health Sciences, Riyadh, Saudi Arabia

OPEN ACCESS

Edited by:

James Duffin,
University of Toronto, Canada

Reviewed by:

Lin Gao,
University of Seville, Spain
Patrice G. Guyenet,
University of Virginia, United States

*Correspondence:

Andrew P. Holmes
a.p.holmes@bham.ac.uk
Prem Kumar
p.kumar@bham.ac.uk

Specialty section:

This article was submitted to
Respiratory Physiology and
Pathophysiology,
a section of the journal
Frontiers in Physiology

Received: 30 March 2022

Accepted: 09 May 2022

Published: 31 May 2022

Citation:


Holmes AP, Swiderska A, Nathanael D, Aldossary HS, Ray CJ, Coney AM and Kumar P (2022) Are Multiple Mitochondrial Related Signalling Pathways Involved in Carotid Body Oxygen Sensing? *Front. Physiol.* 13:908617. doi: 10.3389/fphys.2022.908617

It is generally acknowledged that the carotid body (CB) type I cell mitochondria are unique, being inhibited by relatively small falls in P_{aO_2} well above those known to inhibit electron transport in other cell types. This feature is suggested to allow for the CB to function as an acute O_2 sensor, being stimulated and activating systemic protective reflexes before the metabolism of other cells becomes compromised. What is less clear is precisely how a fall in mitochondrial activity links to type I cell depolarisation, a process that is required for initiation of the chemotransduction cascade and post-synaptic action potential generation. Multiple mitochondrial/metabolic signalling mechanisms have been proposed including local generation of mitochondrial reactive oxygen species (mitoROS), a change in mitochondrial/cellular redox status, a fall in MgATP and an increase in lactate. Although each mechanism is based on compelling experimental evidence, they are all not without question. The current review aims to explore the importance of each of these signalling pathways in mediating the overall CB response to hypoxia. We suggest that there is unlikely to be a single mechanism, but instead multiple mitochondrial related signalling pathways are recruited at different P_{aO_2} s during hypoxia. Furthermore, it still remains to be determined if mitochondrial signalling acts independently or in partnership with extra-mitochondrial O_2 -sensors.

Keywords: carotid body, hypoxia, mitochondria, succinate, arterial chemoreceptor, O_2 sensor, metabolism, mitochondrial inhibitors

INTRODUCTION—A ROLE FOR MITOCHONDRIA IN CAROTID BODY O_2 SENSING

The carotid body (CB) type I or glomus cell has an extraordinary ability to be stimulated by relatively small falls in P_{aO_2} , well above those that start to impact on the metabolism of other cell types (Kumar and Prabhakar, 2012). This allows the CB to act as a protective peripheral chemoreceptor that detects and responds to arterial hypoxia, leading to downstream reflex activation (Kumar, 2009). Of the proposed O_2 sensors within the CB, including the mitochondria, haem-oxygenase 2, NADPH-oxidase and O_2 -dependent K^+ ion channels, only the mitochondria have been identified as having a unique phenotype in comparison with other O_2 -insensitive cell types (Holmes et al., 2018a) and it has become increasingly apparent that the type I cell mitochondria have an important role in CB O_2 sensing.



O_2 binding within mitochondria takes place within the CuB/haem a_3 (cytochrome a_3) binuclear centre of cytochrome c oxidase (complex IV). Multiple groups have shown that the CB mitochondrial electron transport chain is inhibited as the O_2 drops below an unusually and uniquely high PO_2 threshold (Duchen and Biscoe, 1992; Buckler and Turner, 2013). The K_m of cytochrome c oxidase for O_2 in the CB is remarkably high (Buckler and Turner, 2013) and cytochrome a_3 reduction status is excessive over the physiological range of PO_2 s known to stimulate the type I cell (Mills and Jobsis, 1970; Mills and Jobsis, 1972). Emerging evidence also suggests that the CB type I cell has a unique mitochondrial gene expression signature, characterised by relatively high mRNA levels of *Ndufa4l2*, *Cox4i2*, and *Cox8b* (Zhou et al., 2016; Gao et al., 2017), with

the latter two coding for important cytochrome c oxidase proteins. These genes are regulated in part by HIF2 α , a critical transcription factor that underpins CB development, adult function, and plasticity in response to chronic hypoxia (Hodson et al., 2016; Macias et al., 2018; Moreno-Dominguez et al., 2020). Conditional deletion of *Cox4i2* in tyrosine hydroxylase (TH)-positive CB type I cells diminishes both the type I cell and ventilatory response to hypoxia in the mouse (Moreno-Dominguez et al., 2020). Indeed, competitive inhibition of O_2 binding into cytochrome c oxidase using NO donors mimics hypoxia by causing intense chemoafferent stimulation when delivered at high concentrations, and augmented chemosensitivity at more moderate levels (Iturriaga et al., 2000; Mosqueira and Iturriaga, 2002; Holmes et al., 2016).

Furthermore, saturating concentrations of exogenous NaCN, a cytochrome c oxidase inhibitor, act to abolish any further K⁺ channel inhibition induced by hypoxia (Wyatt and Buckler, 2004). Collectively, these findings point to impaired binding of O₂ into cytochrome c oxidase and a subsequent run-down in mitochondrial activity as key events of hypoxia sensing in the CB.

An important consideration is how a fall in mitochondrial activity might then lead to activation of the CB chemotransduction cascade? This has been an important research focus over recent years and a better understanding of these specific mechanisms could offer new strategies to manipulate CB function in health and disease (Chang, 2017). A number of mitochondrial/metabolic signalling pathways have been proposed including an elevation in mitochondrial reactive oxygen species (mitoROS) generation (Fernandez-Aguera et al., 2015; Arias-Mayenco et al., 2018; Swiderska et al., 2021), an alteration in nicotinamide adenine dinucleotide (NAD⁺) and flavoprotein (Fp) redox status (Bernardini et al., 2020), a fall in MgATP (Varas et al., 2007) and an increase in lactate production (Chang et al., 2015) (Figure 1). Although there is compelling evidence for each of these processes, they are all not without question. The aim of this review is to explore the relative importance of each pathway and to consider the possibility that at least two or more of these may be needed to co-operate to produce full CB chemoexcitation during hypoxia.


ARE MITOCHONDRIAL ROS AND CHANGES IN NAD⁺/FP REDOX STATUS REQUIRED FOR CB ACTIVATION IN HYPOXIA?

The mitochondria are the main generators of ROS in most mammalian cells, with complex I and III considered to be the primary sources (Murphy, 2009). Using redox-sensitive green fluorescent protein (roGFP) targeted to selective mitochondrial compartments, Fernández-Agüera and colleagues demonstrated that ROS were elevated during hypoxia in the mitochondrial intermembrane space (IMS) (Fernandez-Aguera et al., 2015). Importantly, the authors showed that the increase in mitochondrial IMS ROS occurred over a similar physiological PO₂ range known to cause type I cell excitation (Arias-Mayenco et al., 2018). This rise was not apparent in mouse type I cells that had deletion of the *Ndufs2* gene, suggesting that production of ROS in hypoxia was solely at complex I. In contrast, it was found that mitochondrial matrix ROS was decreased upon hypoxic exposure in both wild-type (WT) and *Ndufs2* deficient type I cells. Consistent with the idea that complex I derived ROS is an important effector signalling molecule, mice with conditional or inducible deletion of *Ndufs2* (*Ndufs2*-KO) displayed a significantly attenuated rise in respiratory frequency when challenged by acute exposure to 10% environmental O₂ (Fernandez-Aguera et al., 2015; Arias-Mayenco et al., 2018). Type I cells isolated from these mice also exhibited strong depression of neurosecretory and Ca²⁺ responses to acute hypoxia. These cells did, however, respond to hypercapnia,

confirming functionality and selective depletion in O₂ sensing capabilities. The *Ndufs2* gene appears to be of particular importance as genetic deletion of *Mt-nd6*, (the mitochondrial encoded subunit six of the NADH dehydrogenase complex), did not impair CB or ventilatory responses to hypoxia (Gomez-Nino et al., 2018). Intracellular delivery of exogenous H₂O₂ has been shown to increase input resistance, indicative of an inhibitory action on leak K⁺ channels (Fernandez-Aguera et al., 2015). The specific identity of such channels is still to be elucidated and the effect of ROS on voltage-dependent K⁺ channels remains to be further studied. A key future experiment will be to examine whether mitochondrial derived ROS can cause complete or partial block of TASK and/or BK_{Ca}, two key K⁺ channels thought to be involved in rodent and human CB chemostimulation (Wyatt and Peers, 1995; Peers and Wyatt, 2007; Mkrtchian et al., 2012; Buckler, 2015). It will also be interesting to examine the distance between mitochondria and effector ion channels, which could, potentially, form discreet and tightly controlled redox-sensitive signalling microdomains.

If mitochondrial complex I derived ROS are essential, then it would be expected that pharmacological targeting of complex I should abolish or severely reduce CB responses to hypoxia. In CB slices isolated from rats, application of rotenone, a mitochondrial complex I inhibitor, has been shown to inhibit secretory responses to hypoxia in a concentration dependent manner (Ortega-Saenz et al., 2003). Similar results have been observed in terms of rotenone being able to abolish any further depression of background K⁺ currents caused by hypoxia (Wyatt and Buckler, 2004). However, in these latter experiments, a similar attenuation of hypoxic sensing was also observed by NaCN and FCCP (a mitochondrial uncoupler), which act independently of complex I. Furthermore, robust rises in Ca²⁺ induced by hypoxia have been observed in the presence of rotenone, with the concurrent application of ascorbate and TMPD, a cocktail suggested to deliver electrons directly to cytochrome c and subsequently maintain electron flow through to cytochrome c oxidase (Wyatt and Buckler, 2004). A similar restoration of the responses to NO donors (thought to excite the CB by preventing O₂ binding into cytochrome c oxidase) can be achieved using ascorbate and TMPD following blockade of upstream mitochondrial complexes (Mosqueira and Iturriaga, 2002). Interestingly, it has also been reported by many groups that the CB can still be excited under conditions of complete anoxia, when the generation of ROS would theoretically be zero (Buckler and Turner, 2013; Peng et al., 2019). Whether or not complete removal of O₂ from an *ex vivo* experimental set up can be achieved is debatable. Nevertheless, these experiments do point towards the existence of a substantial signalling pathway activated upon inhibition of cytochrome c oxidase that is independent of complex I and ROS generation.

To probe the importance of mitochondrial ROS further, we have recently measured rat CB chemoafferent activity and ventilation in the presence of the mitochondrial antioxidants, MitoTEMPO and SKQ1 (Swiderska et al., 2021). Both agents caused a moderate decrease (20%–50%) of the chemoafferent response to hypoxia, although a significant component remained intact (Figure 2). MitoTEMPO and SKQ1 produced a similar



attenuation of the rise in respiratory frequency upon exposure to 10% environmental O_2 . SKQ1, a mitochondrial antioxidant targeted to the IMS, also led to an approximately 20% decrease in the hypoxic ventilatory response, whilst the response to hypercapnia was completely preserved. We cannot rule out the possibility that some mitoROS were not completely removed by these agents and there was also the potential for non-selective actions, e.g. on the post-synaptic afferent fibres and brain stem neurones. However, these data are more consistent with the idea that mitoROS are only partially responsible for eliciting CB stimulation during hypoxia.

The exact mechanism of mitochondrial ROS production is another important consideration. The idea that complex I ROS can be generated in particularly large amounts following excessive succinate metabolism at complex II and reverse electron transport (RET) gained significant attention due to its association with mediating cardiac myocyte cell death following ischemia-reperfusion (Chouchani et al., 2014). The strong link between succinate dehydrogenase (SDH; complex

II) mutations and paraganglioma (tumour of the CB) does suggest that succinate metabolism has a heightened importance in the CB (Favier et al., 2015; Kluckova and Tennant, 2018). Interestingly, the CB type I cell does appear to contain an unusually high concentration of succinate in comparison with brain and adrenal medulla (Fernandez-Aguera et al., 2015). Conditional homozygous deletion of the *Sdh* gene leads to a powerful attenuation of the Ca^{2+} response to hypoxia in isolated type I cells (Gao et al., 2017). However, these cells also exhibited similar depressions in response to high K^+ , possibly indicative of a more general impairment of CB function. In contrast, less severe heterozygous deletion of *Sdh* was found to augment basal neurosecretion and have no impact on the hypoxic sensitivity (Piruat et al., 2004).

In our own experiments, exogenous application of the cell permeable form of succinate (diethyl succinate, DESucc) did lead to chemoafferent stimulation in the isolated rat CB, but this measured only approximately 10–30% of that induced by a subsequent severe hypoxic stimulus (Swiderska et al., 2021).

Addition of a high concentration of dimethyl malonate (DMM; a competitive inhibitor of SDH) also produced a significant rise in basal chemoafferent frequency and caused partial (20%–50%) reduction of the activity in hypoxia. This was consistent with the level of hypoxic inhibition we observed in the presence of MitoTEMPO and SKQ1. It could be argued that DMM failed to completely block SDH activity in these experiments, although this concentration was sufficient to completely prevent the rise in chemoafferent activity induced by exogenous DESucc. Other groups have also seen partial attenuation of hypoxic neurosecretion in the presence of DMM, with a significant proportion of the response still being preserved (Arias-Mayenco et al., 2018). DMM also caused significant, but not complete, attenuation of IMS ROS generation evoked by hypoxia. Taken together, these data do suggest that succinate metabolism is important for ROS generation and CB stimulation during hypoxia but is not the sole mechanism. Furthermore, it is possible that there is an alternative mode of IMS mitochondrial ROS production that is independent of succinate metabolism. An intriguing mechanism is the potential generation of ROS from mitochondrial complex III. The functional importance of hypoxia-induced mitochondrial complex III ROS production has been proposed for pulmonary artery smooth muscle cells (PASMCS) (Waypa et al., 2013). Deletion of the Rieske iron-sulphur protein of Complex III from isolated PASMCS abolished mitochondrial ROS generation and increases in $[Ca^{2+}]_i$ during hypoxia. A similar depletion of hypoxic sensitivity was observed in PASMCS isolated from mice expressing a tunicate alternative oxidase, which allowed for electron flux to continue during concomitant inhibition of mitochondrial complexes III and IV (Sommers et al., 2020). It will be interesting to investigate whether or not complex III ROS have a parallel function in mediating CB type I cell activation in hypoxia. A summary of the estimated importance of succinate metabolism and mitochondrial ROS production in mediating chemoafferent and ventilatory responses to hypoxia is shown in **Figure 2**.

Whilst mitochondrial IMS ROS are suggested to increase in hypoxia, there is also a significant rise in NAD(P)H (Duchen and Biscoe, 1992; Bernardini et al., 2020). The rise in NAD(P)H occurs over a PO_2 range approximately equal to that which is known to stimulate the type I cell (Buckler and Turner, 2013) and is dependent on mitochondrial complex I activity (Arias-Mayenco et al., 2018). An elevation in local mitochondrial NADH would likely trigger an increase in cytosolic NADH through action of the malate/aspartate shuttle. A number of reports have indicated that pyridine nucleotides are capable of modifying voltage-gated K^+ channel activation/inactivation kinetics in primary cells and expression systems (Tipparaju et al., 2005; Kilfoil et al., 2019). In the CB type I cell it has been observed that addition of NADH to the patch pipette prevents hypoxic inhibition of K_v currents (Fernandez-Aguera et al., 2015). The identity of specific channels that are regulated by NADH, its functional importance and the mechanism of NADH channel modulation are areas that need to be addressed in the future. However, this data does indicate that NADH may have an important signalling role in the CB during hypoxia.

NAD(P)H, $FADH_2$ and $FMNH_2$ are key sources of electrons for both mitochondrial and cytosolic redox reactions. Thus, their relative concentration in comparison with their oxidised forms (NAD⁺, FAD and FMN) can be used to indicate the overall mitochondrial and/or cellular redox status (Bernardini et al., 2020). In the intact CB preparation, Bernardini and colleagues have shown that flavoprotein (Fp) fluorescence decreases rapidly during hypoxia and is accompanied by a concurrent rise in NAD(P)H (Bernardini et al., 2020). As Fp fluorescence represents the oxidised forms of flavoproteins, this data is consistent with an increase in $FADH_2$ and $FMNH_2$ groups and an overall switch to a more reduced cellular redox status. The authors suggest that such a redox shift leads to stimulation of the redox-sensitive protein PIN1 (located at the cell membrane) which in turn promotes NADPH oxidase activity. Accordingly, pharmacological inhibition of PIN1 was found to partially inhibit membrane depolarisation caused by hypoxia. However, mice lacking the *Gp91 phox* gene exhibit full CB and ventilatory sensitivity to hypoxia (Roy et al., 2000). Thus, there may be alternative signalling pathways subject to changes in the cellular redox status that are still to be revealed. Investigating any direct action of NAD(P)H on ion channel function in the CB could be a logical next step.

IS A FALL IN MgATP Coupled to CB Excitation in Hypoxia?

A fall in mitochondrial activity is likely to modify the adenine nucleotide concentrations within the type I cell and this itself has been suggested to link either directly or indirectly to cellular excitation. In most mammalian cells the majority of ATP is complexed with Mg^{2+} , forming MgATP (Gout et al., 2014). Many early studies aimed to measure whole CB adenine nucleotide concentration, including ATP, before and after hypoxia and following addition of mitochondrial poisons. The results were somewhat variable in relation to the effect of hypoxia, with reports of a slight fall in ATP (approximately 20%) by some groups (Obeso et al., 1985), but no significant change observed by others (Acker and Starlinger, 1984; Verna et al., 1990). These data perhaps indicated that a fall in ATP and/or an increase in AMP were unlikely to be important signalling molecules in the type I cell. That said, studies using ATP depleting agents such as 2-deoxy-D-glucose and cyanide have reported a clear negative correlation between ATP content and neurosecretion and/or chemoafferent activity (Obeso et al., 1985; Obeso et al., 1986). All of these experiments measured total ATP content of the CB, which is likely to include contributions from type II cells, neurons, blood vessels and interstitial fluid, in addition to type I cells. ATP is also an important neurotransmitter released by type I cells during hypoxia (Zhang et al., 2000), and this itself can undergo further degradation into ADP, AMP and adenosine by ectonucleotidases including NTPDase1,2,3 and CD73 (Conde and Monteiro, 2004; Salman et al., 2017; Holmes et al., 2018b). Therefore, the total ATP content measured would be dependent on a combination of metabolically derived ATP, the level of stored and released ATP as a neurotransmitter and the

rate of extracellular ATP degradation. As such, it is difficult to say, with confidence, that total CB ATP content is an accurate reflection of type I cell MgATP concentration.

To date, no study has been able to make direct measurements of type I cell ATP or MgATP. However, using Mag-indo-1, a Mg^{2+} sensitive fluorescent probe, Varas and colleagues were able to demonstrate that free Mg^{2+} within isolated rat type I cells significantly increased upon exposure to hypoxia (Varas et al., 2007). They concluded that this was a direct consequence of a fall in ATP production and a decrease in MgATP. Similar rises in Mg^{2+} were observed upon exposure to mitochondrial poisons including CN^- and oligomycin (a H^+ -ATP synthase inhibitor). It does therefore appear that MgATP falls in the rat type I cell upon exposure to hypoxia. However, it is not apparent if this takes place over the same physiological PO_2 range known to stimulate the type I cell, and further studies are needed to validate results in other species including mouse and human.

Numerous investigations have reported that oligomycin is capable of activating many aspects of the CB chemotransduction cascade including inhibition of K^+ channels, rises in $[Ca^{2+}]_i$ and chemoafferent excitation (Mulligan et al., 1981; Wyatt and Buckler, 2004). This strongly supports the idea that a fall in mitochondrial ATP production is sufficient to cause rapid CB stimulation. Chemoafferent excitation evoked by hypoxia is also severely suppressed following sustained oligomycin application (Mulligan et al., 1981; Shirahata et al., 1987). This is perhaps not completely surprising as some mitochondrial ATP production is likely required to support the increase in type I cell and chemoafferent activity during hypoxia (Bardsley et al., 2021).

It is important to consider precisely how a fall in MgATP may couple to type I cell depolarisation. TASK channels are highly expressed in the CB type I cell and are thought to play a major role in evoking type I cell depolarisation in hypoxia (Buckler et al., 2000; Kim et al., 2009; Mkrtchian et al., 2012; Zhou et al., 2016). MgATP can directly activate TASK-like channels in the type I cell with an EC_{50} of approximately 2.3 mM and saturating concentration of just above 10 mM, both of which are within an estimated physiological range (Williams and Buckler, 2004; Varas et al., 2007). In contrast, substances including free ATP, ADP, AMP and NADH do not appear to directly modulate TASK-like channel activity. Cell attached patches of CB type I cells demonstrate a high degree of single K^+ channel activity at a membrane potential of $-70mV$, largely attributable to these TASK-like channels (Varas et al., 2007; Zhou et al., 2016). When these patches are excised into the inside-out configuration, channel activity rapidly decreases to a level of about 10% of control. Interestingly, this run-down can be partially alleviated by 5 mM MgATP, with approximately 50% of channel activity being preserved (Varas et al., 2007). These data suggest that in normoxic physiological conditions, intracellular MgATP acts to maintain the high level of TASK-like channel activity. In hypoxia, the depletion in MgATP would lead to a fall in activity, sufficient to cause cellular depolarisation. However, these data also imply that there are other important intracellular factors, in addition to MgATP, that contribute to TASK channel opening in normoxia, which may be reduced under hypoxic conditions. Such substances are still to be identified and warrant further

consideration. It is also not clear precisely how MgATP regulates TASK channels in the type I cell, since they are not known to contain a specific ATP binding pocket and/or be intrinsically sensitive to nucleotides. Thus, there could be a type I cell specific adapter protein, coupled to TASK channels which confers sensitivity to MgATP. Aiming to reveal the identity of this type of protein could be an interesting area for future research.

Though important, TASK channels are unlikely to be the only ion channels involved in CB O_2 sensing. In TASK1/3 deficient murine CB type I cells, Ca^{2+} elevations and augmented catecholamine release in response to hypoxia are still strongly preserved (Ortega-Saenz et al., 2010; Turner and Buckler, 2013). In rat type I cells, an array of different TASK channel inhibitors all produced significant chemostimulation, but in each case this was well below that elicited by severe hypoxia (O'donohoe et al., 2018). Other potential K^+ channel candidates that could be regulated by adenine nucleotides include TREK-1 and K_{ATP} . Similar to TASK, TREK-1 is known to be activated by physiological intracellular concentrations of MgATP (Enyeart et al., 2002). It has been observed that TREK-1 is expressed in the rat type I cell (Yamamoto and Taniguchi, 2006) but preliminary findings suggest an absence in the human CB (Mkrtchian et al., 2012). Thus, its relevance in CB O_2 sensing still needs to be defined. An ATP- and glibenclamide-sensitive K^+ channel has been identified in the type I cell, but it was shown to be inhibited by ATP, therefore making it an unlikely candidate to be responsible for coupling a fall in mitochondrial ATP production with cellular depolarisation in hypoxia (Kim et al., 2011).

An alternative mechanism to cause type I cell depolarisation could be through activation of inward Na^+ , Ca^{2+} or Mg^{2+} currents, carried through TRP channels (Kumar et al., 2006; Kim et al., 2015). Multiple classes of TRP channels have been reported in the type I cell and/or the chemoafferent fibres including TRPV, TRPC and TRPM (Buniel et al., 2003; Roy et al., 2012; Jendzjowsky et al., 2018; Shin et al., 2019). TRPM7 is suggested to be the most highly expressed TRP channel within the mouse CB type I cell, indicative of functional importance (Zhou et al., 2016). TRPM7 has been linked to chemoexcitation upon exposure to leptin, either given exogenously or when elevated during obesity (Shin et al., 2019; Shin et al., 2021). However, its significance in mediating CB type I cell stimulation in hypoxia has not been studied to date. TRPM7 channel activity can be modified by numerous intracellular factors, including the Mg^{2+} -complexed nucleotides, MgATP and MgGTP (Nadler et al., 2001; Demeuse et al., 2006). In expression systems, it has been reported that TRPM7 activity significantly increases when MgATP drops below a threshold of approximately 1–2 mM (Nadler et al., 2001). Therefore, it is possible that TRPM7 could be activated in the CB type I cell during hypoxia, in response to a fall in MgATP, and this clearly warrants further investigation.

A number of observations counter the idea that MgATP depletion is necessary to activate the CB during hypoxia. Type I cell Ca^{2+} responses to rotenone alone and rotenone plus hypoxia are non-additive (Ortega-Saenz et al., 2003; Arias-Mayenco et al., 2018). Both of these stimuli would decrease MgATP and, potentially, to a greater extent when applied together. Thus, it would appear that any greater fall in MgATP does not lead to any

further type I cell stimulation. However, there is also the possibility that saturating concentrations of rotenone may be sufficient to prevent any further fall in MgATP caused by hypoxia. Direct measurements of MgATP are needed for clarification. The finding that hypoxic responses are robustly maintained in murine type I cells deficient in ATP sensitive TASK1/3 channels also points towards the presence of a non-ATP sensitive mechanism (Ortega-Saenz et al., 2010). Finally, it has been observed that inhibition of voltage-dependent K⁺ current by hypoxia is still apparent in type I cells dialyzed with 3–5 mM MgATP (Lopez-Lopez et al., 1989; Perez-Garcia et al., 2004). Again, this does indicate that there is an O₂ sensitive signalling mechanism in the type I cell that is independent of MgATP. Evaluation of [Ca²⁺]_i and neurosecretory responses to hypoxia measured under conditions of preserved/clamped intracellular MgATP may provide further insight into this matter.

In addition to directly impacting on ion channels, variations in adenine nucleotide concentrations could signal through an intermediate protein/pathway which in turn couples to the cell membrane. A likely candidate would be the AMP-activated protein kinase (AMPK), whose phosphorylation (and thus activation) status is enhanced by an increase in the AMP/ATP ratio. Robust expression of AMPK subunits have been reported in both rodent type I cells (Wyatt et al., 2007; Zhou et al., 2016) and human whole CBs (Mkrтчian et al., 2012). Initial experiments using AICAR, an AMP-mimetic that activates AMPK, were very promising. In rat CBs, AICAR was observed to inhibit background Ba²⁺ sensitive K⁺ currents, cause a rise in [Ca²⁺]_i and generate an increase in chemoafferent nerve activity within minutes of drug application (Evans et al., 2005; Wyatt et al., 2007). Compound C, an inhibitor of AMPK, decreased both the type I cell and sensory nerve response to hypoxia. However, these results have been countered by experiments where both AICAR and A769662 (a more direct AMPK activator) were without effect on type I cell TASK channel currents and [Ca²⁺]_i (Kim et al., 2014). Furthermore, it has now been suggested that a CB with conditional deletion of both α1 and α2 subunits of AMPK still retains full chemoafferent sensitivity to hypoxia (Mahmoud et al., 2016). This finding was reported from a single experiment and a more complete dataset is still necessary to draw firm conclusions. There is also the possibility that an elevation in alternative O₂ sensing pathways may have countered the absence of AMPK signalling. However, on current evidence, it does appear that AMPK is not required to activate the CB in hypoxia. Potential roles for AMPK in mediating CB plasticity in cardiorespiratory pathology remain to be defined.

IS AN ELEVATION IN LACTATE AND STIMULATION OF OLFACTORY RECEPTOR 78 (OLFR78) NECESSARY TO ACHIEVE FULL CB CHEMOSTIMULATION IN HYPOXIA?

When oxidative phosphorylation becomes restricted during low O₂ conditions, ATP can be partially preserved through an increase in glycolytic flux. Glycolysis is highly sensitive to the ATP/AMP ratio and is maximally stimulated by oligomycin (an

ATP-synthase inhibitor) (Tanner et al., 2018; Yagi et al., 2021). To maintain high levels of glycolysis, generated pyruvate is rapidly converted to lactate by lactate dehydrogenase (LDH) (Rogatzki et al., 2015). Exaggerated glucose uptake and lactate removal are also necessary to sustain augmented glycolytic rates (Tanner et al., 2018).

In the CB, it has been hypothesized that an elevation in lactate provides the link between mitochondrial run-down and cellular stimulation during hypoxia (Chang et al., 2015). In support of this idea, it has been reported that rates of glucose uptake in the CB are enhanced during hypoxia (Obeso et al., 1993). Indeed, multiple groups have observed that lactate can also directly stimulate the CB in a concentration dependent manner, with a threshold of approximately 5–10 mM and maximum response achieved at around 30 mM (Chang et al., 2015; Peng et al., 2020; Torres-Torrelo et al., 2021). Arterial blood lactate concentrations can rise to approximately 6–7 mM during exposure to hypoxia (10% inspired) in mice and it is debatable whether this would be sufficient to excite the CB (Chang et al., 2015; Torres-Torrelo et al., 2021). It is, however, possible that local extracellular concentrations of lactate within the CB may be much higher, especially if type I cells have a much higher rate of glycolysis and lactate excretion. Experiments measuring tissue concentrations of lactate in the CB during exposure to hypoxia will be essential to clarify this. Some reports have demonstrated that the peak excitation produced by lactate is similar to or greater than that induced by hypoxia (Chang et al., 2015; Torres-Torrelo et al., 2021). However, others suggest that responses to lactate, even at 30 mM, are relatively modest, measuring only around 30% of that induced by hypoxia (Peng et al., 2020). The difference is likely a consequence of the severity of the hypoxic stimulus intensity used between the different studies, although reduced accessibility to type I cells due to diffusion limitations in the whole CB preparations may also be a contributing factor. It has also been observed that some isolated rat type I cells are unresponsive to 20 mM lactate, as evidenced by no change in membrane potential or outward K⁺ currents (Spiller et al., 2021). The apparent lack of a response could be due to loss of intracellular signalling as experiments were performed in the whole cell patch clamp configuration. To counter this, in the same study, administration of lactate to the *in-situ* CB preparation via the perfusate failed to evoke any chemoafferent excitation at 5 and 20 mM: a situation where lactate would have sufficient access to the tissue and intracellular signalling would be fully preserved. Alternatively, it could be that lactate only excites a proportion of type I cells, but not all. The idea that different type I cells can respond to some but not all stimuli need further exploration. In our own experiments, we have detected that inhibition of glycolysis using iodoacetate leads to rapid and sustained chemoafferent excitation (Holmes et al., 2014). This indicates that there exists a signalling pathway in the CB that can cause chemoafferent excitation independent of an increase in lactate production. Similar measurements have been reported using other inhibitors of glycolysis including 2-deoxy-D-glucose (Obeso et al., 1986). We have also seen that lactate (5 mM) can decrease the elevated chemoafferent activity caused by prolonged glucose deprivation, probably by helping to preserve

ATP (Holmes et al., 2014). Thus, under different experimental conditions it has been shown that lactate can stimulate, inhibit, or have no effect on CB function. Its specific role in mediating CB hypoxic sensitivity needs further clarification.

It was originally proposed by Chang and colleagues that OLF78 was necessary for lactate to activate the CB in hypoxia (Chang et al., 2015). They revealed that *Olf78* deficient mice failed to display any significant hypoxic ventilatory response. CBs isolated from these mice were also unresponsive to hypoxia and high concentrations of exogenous lactate (30 mM). Another group has reported contradictory results, whereby single CB type I cells isolated from *Olf78*^{-/-} mice produced neurosecretory responses to hypoxia that were indistinguishable from WT littermates (Torres-Torrel et al., 2018). These mice also displayed a full hypoxic ventilatory response. The discrepancies between the two groups are hard to reconcile but could be dependent on genetic drift, redundancy mechanisms, animal husbandry and/or background strain.

A third group have recently suggested that *Olf78*^{-/-} mice have partially attenuated CB responses to mild but not severe hypoxia (Peng et al., 2020). In *ex vivo* experiments, when the hypoxic stimulus was mild, a significant chemoafferent response was still observed, measuring approximately 40% of WT controls. In whole animal breathing experiments, ventilation in response to 12% O₂ was slightly reduced in *Olf78*^{-/-} mice when normalized to O₂ consumption. There was also a considerable fall in P_aCO₂ consistent with hyperventilation. Interestingly, this group also showed that CBs from *Olf78* KO mice were still sensitive to lactate. Thus, they concluded that even though OLF78 played a role in CB O₂ sensing, its activation in mild hypoxia must be dependent on a substance other than lactate. OLF78 can respond to many substances including the short chain fatty acids acetate, propionate and butyrate, but so far stimulation of the CB by these substances has only been observed at supraphysiological concentrations (Peng et al., 2020). Identification of a substance that can stimulate OLF78 in the CB at physiological concentrations could be of considerable importance, especially given that *Olf78* is one of the most highly expressed genes in the CB (Chang et al., 2015).

For lactate that is produced intracellularly to activate OLF78 it is assumed that it must first leave the cell to gain access to the extracellular facing binding site. It has been shown that monocarboxylate transporters 2 and 4 (MCT-2 and MCT-4) are expressed in type I cells, but not MCT-1, providing potential route for lactate both into and out of the cells (Torres-Torrel et al., 2021). In isolated type I cell preparations, the small volume recording chamber is constantly renewed by the external perfusate, thereby preventing any extracellular lactate accumulation. Yet, many studies have demonstrated that isolated type I cells retain intrinsic O₂ sensitivity and are capable of eliciting rises in mitoROS, increases in [Ca²⁺]_i and neurosecretion (Fernandez-Aguera et al., 2015; Arias-Mayenco et al., 2018). This counters the idea of autocrine/paracrine lactate signalling being essential to stimulate the type I cells in hypoxia.

Collectively, these data do not rule out lactate being involved in CB O₂ sensing but do suggest that it most likely does not act through OLF78. If lactate does reach high mM concentrations in CB tissue during hypoxia, then it could act by alternative




FIGURE 3 | Potential for numerous mitochondrial related signalling mechanisms to co-operate to produce the full carotid body response to hypoxia. Different mitochondrial related signalling mechanisms could be responsible for the rise in chemoafferent discharge at different levels of hypoxia, with an alteration in cellular redox status and mitochondrial reactive oxygen species acting during mild-moderate hypoxia, a decrease in MgATP being important during moderate-severe hypoxia and an increase in lactate contributing during severe hypoxia. There may well be significant overlap between these signalling pathways and potential for modification/adaptation.

mechanisms such as causing intracellular acidification (Torres-Torrel et al., 2021). What is apparent is that a very large rise in lactate concentration would be required to stimulate the CB. This would likely only occur when falls in MgATP are very severe, possibly limiting its role to the terminal hypoxia section of the chemoafferent PO₂-response curve.

CONCLUSION-PUTTING TOGETHER THE FULL RESPONSE TO HYPOXIA

Given the non-linear shape of the chemoafferent and ventilatory response curves to O₂ (Kumar and Prabhakar, 2012), the existence might be construed of a single sensor mechanism with a similarly complex and varying sensitivity across the entire physiological range of PO₂. Alternatively, a number of sensor mechanisms, each with its own particular, potentially overlapping sensitivity, might act together synergistically to provide the entire response. The conflicting data to date, tends to suggest the latter as the most probable means by which the CB type I cell enables and maintains its exquisite O₂ sensitivity between the extremes of hyperoxia and anoxia. As suggested by Prabhakar, this may involve multiple O₂ sensor mechanisms, both mitochondrial and extra-mitochondrial, acting upon multiple neurotransmitter systems within a so-called “chemosome,” to generate both the high and broad-range sensitivity as well as the rapid response that characterizes this

receptor system (Prabhakar, 2006). In addition, inhibitory mechanisms within the chemosome could act in a protective manner during periods of prolonged stimulation, such as the chronic hypoxia associated with numerous cardiorespiratory pathologies. We suggest here, that such a chemosome might exist within the distinctive CB type I mitochondria themselves, with perhaps mitoROS and cellular redox signalling contributing more towards the higher PO₂ portion of the response curve, with a fall in MgATP and/or lactate contributing more towards the severe hypoxia/anoxic portions of the curve (Figure 3).

Mitochondrial signalling may act either independently or in partnership with extra-mitochondrial O₂-sensors, to generate the full, physiological response to acute hypoxia and perhaps even to the remodelling plasticity observed in the CB and its reflexes with chronic hypoxia. One of the most influential hypotheses of CB hypoxic chemotransduction involves O₂ dependent generation of H₂S (Olson et al., 2008; Peng et al., 2010). Compelling evidence has been presented demonstrating that mice deficient in the H₂S generating enzyme cystathionine γ -lyase (CSE) have an almost completely absent hypoxic ventilatory response (Peng et al., 2010). CBs isolated from these mice have a diminished chemoafferent response to hypoxia whilst retaining normal hypercapnic sensitivity. The rise in H₂S in hypoxia is thought to be a consequence of reduced CO synthesis subsequent to a decrease in the haem-oxygenase 2 activity (Yuan et al., 2015). Although H₂S is able to rapidly and reversibly inhibit BK_{Ca} channels (Telezhkin et al., 2010) it is also a classical inhibitor of cytochrome c oxidase. The calculated concentrations of H₂S required to excite isolated rat type I cells also evoke an increase in NADH, consistent with concurrent mitochondrial inhibition (Buckler, 2011). This could provide a unifying link between these two hypotheses. The endogenous role of H₂S in modifying the O₂ sensitivity of the unique CB type I cell mitochondria requires further exploration as does the intriguing possibility of a novel functional role of mitochondrial generated reactive sulphur species (Olson, 2021).

In this review we have discussed some of the key mitochondrial related signalling mechanisms in the CB known

to date. It is likely that novel mitochondrial and extra-mitochondrial signalling mechanisms will emerge in the future as we move towards a more complete understanding. Indeed, very recent data suggests that local mitochondrial temperature transients contribute to type I cell membrane depolarisation (Rakoczy et al., 2022). Furthermore, a fail-safe mechanism has been recently proposed based on model testing, whereby low O₂ leads to disruption of transporter activity involved in maintenance of intracellular pH (Duffin, 2020). The ensuing intracellular acidification is predicted to be sufficient to cause type I activation. The model also indicates that such a mechanism could account for significant CO₂-O₂ stimulus interaction. Clearly, experimental data is now required to validate these model data. Given the emerging prominence of the CB hyperactivity in initiating hypertension and cardiac arrhythmia (Iturriaga et al., 2021) future studies are needed to investigate how mitochondrial O₂ sensitivity and signalling is altered in pathology.

AUTHOR CONTRIBUTIONS

All authors listed have made a substantial, direct, and intellectual contribution to the work and approved it for publication.

FUNDING

This work was funded by the British Heart Foundation (FS/PhD/20/29093) to AC, DN, PK, and AH and a Scholarship from King Saud bin Abdulaziz University for Health Sciences (Riyadh, Saudi Arabia) provided to HA, through the Saudi Arabian Cultural Bureau in London.

ACKNOWLEDGMENTS

Figures were created using BioRender.com.

REFERENCES

- Acker, H., and Starlinger, H. (1984). Adenosine Triphosphate Content in the Cat Carotid Body under Different Arterial O₂ and CO₂ Conditions. *Neurosci. Lett.* 50, 175–179. doi:10.1016/0304-3940(84)90482-8
- Arias-Mayenco, I., González-Rodríguez, P., Torres-Torrel, H., Gao, L., Fernández-Agüera, M. C., Bonilla-Henao, V., et al. (2018). Acute O₂ Sensing: Role of Coenzyme QH₂/Q Ratio and Mitochondrial ROS Compartmentalization. *Cell Metab.* 28, 145–158. doi:10.1016/j.cmet.2018.05.009
- Bardsley, E. N., Pen, D. K., McBryde, F. D., Ford, A. P., and Paton, J. F. R. (2021). The Inevitability of ATP as a Transmitter in the Carotid Body. *Aut. Neuroscience-Basic Clin.* 234, 102815. doi:10.1016/j.autneu.2021.102815
- Bernardini, A., Wolf, A., Brockmeier, U., Riffkin, H., Metzen, E., Acker-Palmer, A., et al. (2020). Carotid Body Type I Cells Engage Flavoprotein and Pin1 for Oxygen Sensing. *Am. J. Physiology-Cell Physiology* 318, C719–C731. doi:10.1152/ajpcell.00320.2019
- Buckler, K. J. (2011). Effects of Exogenous Hydrogen Sulphide on Calcium Signalling, Background (TASK) K Channel Activity and Mitochondrial Function in Chemoreceptor Cells. *Pflugers Arch.* 463, 743–754. doi:10.1007/s00424-012-1089-8
- Buckler, K. J. (2015). TASK Channels in Arterial Chemoreceptors and Their Role in Oxygen and Acid Sensing. *Pflugers Arch. - Eur. J. Physiol.* 467, 1013–1025. doi:10.1007/s00424-015-1689-1
- Buckler, K. J., and Turner, P. J. (2013). Oxygen Sensitivity of Mitochondrial Function in Rat Arterial Chemoreceptor Cells. *J. Physiology* 591, 3549–3563. doi:10.1113/jphysiol.2013.257741
- Buckler, K. J., Williams, B. A., and Honore, E. (2000). An Oxygen-, Acid- and Anaesthetic-Sensitive TASK-like Background Potassium Channel in Rat Arterial Chemoreceptor Cells. *J. Physiology* 525, 135–142. doi:10.1111/j.1469-7793.2000.00135.x
- Buniel, M. C. F., Schilling, W. P., and Kunze, D. L. (2003). Distribution of Transient Receptor Potential Channels in the Rat Carotid Chemosensory Pathway. *J. Comp. Neurol.* 464, 404–413. doi:10.1002/cne.10798
- Chang, A. J. (2017). Acute Oxygen Sensing by the Carotid Body: from Mitochondria to Plasma Membrane. *J. Appl. Physiology* 123, 1335–1343. doi:10.1152/jappphysiol.00398.2017
- Chang, A. J., Ortega, F. E., Riegler, J., Madison, D. V., and Krasnow, M. A. (2015). Oxygen Regulation of Breathing through an Olfactory Receptor Activated by Lactate. *Nature* 527, 240–244. doi:10.1038/nature15721

- Chouchani, E. T., Pell, V. R., Gaude, E., Aksentijević, D., Sundier, S. Y., Robb, E. L., et al. (2014). Ischaemic Accumulation of Succinate Controls Reperfusion Injury through Mitochondrial ROS. *Nature* 515, 431–435. doi:10.1038/nature13909
- Conde, S. V., and Monteiro, E. C. (2004). Hypoxia Induces Adenosine Release from the Rat Carotid Body. *J. Neurochem.* 89, 1148–1156. doi:10.1111/j.1471-4159.2004.02380.x
- Demeuse, P., Penner, R., and Fleig, A. (2006). TRPM7 Channel Is Regulated by Magnesium Nucleotides via its Kinase Domain. *J. General Physiology* 127, 421–434. doi:10.1085/jgp.200509410
- Duchen, M. R., and Biscoe, T. J. (1992). Mitochondrial Function in Type I Cells Isolated from Rabbit Arterial Chemoreceptors. *J. Physiol.* 450, 13–31. doi:10.1113/jphysiol.1992.sp019114
- Duffin, J. (2020). Fail-safe Aspects of Oxygen Supply. *J. Physiol.* 598, 4859–4867. doi:10.1113/jp280301
- Enyeart, J. J., Xu, L., Danthi, S., and Enyeart, J. A. (2002). An ACTH- and ATP-Regulated Background K⁺ Channel in Adrenocortical Cells Is TREK-1. *J. Biol. Chem.* 277, 49186–49199. doi:10.1074/jbc.m207233200
- Evans, A. M., Mustard, K. J. W., Wyatt, C. N., Peers, C., Dipp, M., Kumar, P., et al. (2005). Does AMP-Activated Protein Kinase Couple Inhibition of Mitochondrial Oxidative Phosphorylation by Hypoxia to Calcium Signaling in O₂-Sensing Cells? *J. Biol. Chem.* 280, 41504–41511. doi:10.1074/jbc.m510040200
- Favier, J., Amar, L., and Gimenez-Roqueplo, A.-P. (2015). Paraganglioma and Pheochromocytoma: from Genetics to Personalized Medicine. *Nat. Rev. Endocrinol.* 11, 101–111. doi:10.1038/nrendo.2014.188
- Fernández-Agüera, M. C., Gao, L., González-Rodríguez, P., Pintado, C. O., Arias-Mayenco, I., García-Flores, P., et al. (2015). Oxygen Sensing by Arterial Chemoreceptors Depends on Mitochondrial Complex I Signaling. *Cell Metab.* 22, 825–837. doi:10.1016/j.cmet.2015.09.004
- Gao, L., Bonilla-Henao, V., García-Flores, P., Arias-Mayenco, I., Ortega-Sáenz, P., and López-Barneo, J. (2017). Gene Expression Analyses Reveal Metabolic Specifications in Acute O₂-Sensing Chemoreceptor Cells. *J. Physiol.* 595, 6091–6120. doi:10.1113/jp274684
- Gomez-Niño, A., Docio, I., Prieto-Lloret, J., Simarro, M., De La Fuente, M. A., and Rocher, A. (2018). Mitochondrial Complex I Dysfunction and Peripheral Chemoreflex Sensitivity in a FASTK-Deficient Mouse Model. *Adv. Exp. Med. Biol.* 1071, 51–59. doi:10.1007/978-3-319-91137-3_6
- Gout, E., Rébeillé, F., Douce, R., and Bligny, R. (2014). Interplay of Mg²⁺, ADP, and ATP in the Cytosol and Mitochondria: Unravelling the Role of Mg²⁺ in Cell Respiration. *Proc. Natl. Acad. Sci. U. S. A.* 111, E4560–E4567. doi:10.1073/pnas.1406251111
- Hodson, E. J., Nicholls, L. G., Turner, P. J., Llyr, R., Fielding, J. W., Douglas, G., et al. (2016). Regulation of Ventilatory Sensitivity and Carotid Body Proliferation in Hypoxia by the PHD2/HIF-2 Pathway. *J. Physiol.* 594, 1179–1195. doi:10.1113/jp271050
- Holmes, A. P., Ray, C. J., Coney, A. M., and Kumar, P. (2018a). Is Carotid Body Physiological O₂ Sensitivity Determined by a Unique Mitochondrial Phenotype? *Front. Physiol.* 9, 562. doi:10.3389/fphys.2018.00562
- Holmes, A. P., Ray, C. J., Pearson, S. A., Coney, A. M., and Kumar, P. (2018b). Ecto-5'-nucleotidase (CD73) Regulates Peripheral Chemoreceptor Activity and Cardiorespiratory Responses to Hypoxia. *J. Physiol.* 596, 3137–3148. doi:10.1113/jp274498
- Holmes, A. P., Turner, P. J., Buckler, K. J., and Kumar, P. (2016). Moderate Inhibition of Mitochondrial Function Augments Carotid Body Hypoxic Sensitivity. *Pflugers Arch. - Eur. J. Physiol.* 468, 143–155. doi:10.1007/s00424-015-1745-x
- Holmes, A. P., Turner, P. J., Carter, P., Leadbeater, W., Ray, C. J., Hauton, D., et al. (2014). Glycogen Metabolism Protects against Metabolic Insult to Preserve Carotid Body Function during Glucose Deprivation. *J. Physiol.* 592, 4493–4506. doi:10.1113/jphysiol.2014.276105
- Iturriaga, R., Alcayaga, J., Chapleau, M. W., and Somers, V. K. (2021). Carotid Body Chemoreceptors: Physiology, Pathology, and Implications for Health and Disease. *Physiol. Rev.* 101, 1177–1235. doi:10.1152/physrev.00039.2019
- Iturriaga, R., Villanueva, S., and Mosquera, M. (2000). Dual Effects of Nitric Oxide on Cat Carotid Body Chemoreception. *J. Appl. Physiology* 89, 1005–1012. doi:10.1152/jappl.2000.89.3.1005
- Jendzjowsky, N. G., Roy, A., Barioni, N. O., Kelly, M. M., Green, F. H. Y., Wyatt, C. N., et al. (2018). Preventing Acute Asthmatic Symptoms by Targeting a Neuronal Mechanism Involving Carotid Body Lysophosphatidic Acid Receptors. *Nat. Commun.* 9, 4030. doi:10.1038/s41467-018-06189-y
- Kilfoil, P. J., Chapalamadugu, K. C., Hu, X., Zhang, D., Raucci, F. J., Tur, J., et al. (2019). Metabolic Regulation of Kv Channels and Cardiac Repolarization by Kvβ2 Subunits. *J. Mol. Cell. Cardiol.* 137, 93–106. doi:10.1016/j.yjmcc.2019.09.013
- Kim, D., Cavanaugh, E. J., Kim, I., and Carroll, J. L. (2009). Heteromeric TASK-1/TASK-3 Is the Major Oxygen-Sensitive Background K⁺channel in Rat Carotid Body Glomus Cells. *J. Physiol.* 587, 2963–2975. doi:10.1113/jphysiol.2009.171181
- Kim, D., Kang, D., Martin, E. A., Kim, I., and Carroll, J. L. (2014). Effects of Modulators of AMP-Activated Protein Kinase on TASK-1/3 and Intracellular Ca²⁺ Concentration in Rat Carotid Body Glomus Cells. *Respir. Physiology Neurobiol.* 195, 19–26. doi:10.1016/j.resp.2014.01.020
- Kim, D., Kim, I., Papreck, J. R., Donnelly, D. F., and Carroll, J. L. (2011). Characterization of an ATP-Sensitive K⁽⁺⁾ Channel in Rat Carotid Body Glomus Cells. *Respir. Physiology Neurobiol.* 177, 247–255. doi:10.1016/j.resp.2011.04.015
- Kim, I., Fite, L., Donnelly, D. F., Kim, J. H., and Carroll, J. L. (2015). “Possible Role of TRP Channels in Rat Glomus Cells,” in *Arterial Chemoreceptors in Physiology and Pathophysiology*. Editors C. Peers, P. Kumar, C. N. Wyatt, E. Gauda, C. A. Nurse, and N. Prabhakar (Berlin: Springer-Verlag Berlin), 227–232. doi:10.1007/978-3-319-18440-1_25
- Kluckova, K., and Tennant, D. A. (2018). Metabolic Implications of Hypoxia and Pseudohypoxia in Pheochromocytoma and Paraganglioma. *Cell Tissue Res.* 372, 367–378. doi:10.1007/s00441-018-2801-6
- Kumar, P., Pearson, S., and Gu, Y. C. (2006). A Role for TRP Channels in Carotid Body Chemotransduction? *FASEB J.* 20, A1229. doi:10.1096/fasebj.20.5.a1229
- Kumar, P., and Prabhakar, N. R. (2012). Peripheral Chemoreceptors: Function and Plasticity of the Carotid Body. *Compr. Physiol.* 2, 141–219. doi:10.1002/cphy.c100069
- Kumar, P. (2009). Systemic Effects Resulting from Carotid Body Stimulation-Invited Article. *Adv. Exp. Med. Biol.* 648, 223–233. doi:10.1007/978-90-481-2259-2_26
- López-Barneo, J., and Ortega-Sáenz, P. (2022). Mitochondrial Acute Oxygen Sensing and Signaling. *Crit. Rev. Biochem. Mol. Biol.* 57, 205–225. doi:10.1080/10409238.2021.2004575
- López-López, J., González, C., Ureña, J., and López-Barneo, J. (1989). Low pO₂ Selectively Inhibits K Channel Activity in Chemoreceptor Cells of the Mammalian Carotid Body. *J. Gen. Physiol.* 93, 1001–1015. doi:10.1085/jgp.93.5.1001
- Macias, D., Cowburn, A. S., Torres-Torrel, H., Ortega-Sáenz, P., López-Barneo, J., and Johnson, R. (2018). Correction: HIF-2α Is Essential for Carotid Body Development and Function. *Elife* 7, e34681. doi:10.7554/eLife.38781
- Mahmoud, A. D., Lewis, S., Juričić, L., Udoh, U.-A., Hartmann, S., Jansen, M. A., et al. (2016). AMP-activated Protein Kinase Deficiency Blocks the Hypoxic Ventilatory Response and Thus Precipitates Hypoventilation and Apnea. *Am. J. Respir. Crit. Care Med.* 193, 1032–1043. doi:10.1164/rccm.201508-1667oc
- Mills, E., and Jöbsis, F. F. (1972). Mitochondrial Respiratory Chain of Carotid Body and Chemoreceptor Response to Changes in Oxygen Tension. *J. Neurophysiology* 35, 405–428. doi:10.1152/jn.1972.35.4.405
- Mills, E., and Jöbsis, F. F. (1970). Simultaneous Measurement of Cytochrome A3 Reduction and Chemoreceptor Afferent Activity in the Carotid Body. *Nature* 225, 1147–1149. doi:10.1038/2251147a0
- Mkrtchian, S., Kählin, J., Ebberyd, A., Gonzalez, C., Sanchez, D., Balbir, A., et al. (2012). The Human Carotid Body Transcriptome with Focus on Oxygen Sensing and Inflammation - a Comparative Analysis. *J. Physiology-London* 590, 3807–3819. doi:10.1113/jphysiol.2012.231084
- Moreno-Domínguez, A., Ortega-Sáenz, P., Gao, L., Colinas, O., García-Flores, P., Bonilla-Henao, V., et al. (2020). Acute O₂ Sensing through HIF2α-dependent Expression of Atypical Cytochrome Oxidase Subunits in Arterial Chemoreceptors. *Sci. Signal* 13, eaay9452. doi:10.1126/scisignal.aay9452
- Mosqueira, M., and Iturriaga, R. (2002). Carotid Body Chemoreceptor Excitation Induced by Nitric Oxide: Involvement of Oxidative Metabolism. *Respir. Physiol. Neurobiol.* 131, 175–187. doi:10.1016/s1569-9048(02)00020-4
- Mulligan, E., Lahiri, S., and Storey, B. T. (1981). Carotid Body O₂ Chemoreception and Mitochondrial Oxidative Phosphorylation. *J. Appl. Physiology* 51, 438–446. doi:10.1152/jappl.1981.51.2.438

- Murphy, M. P. (2009). How Mitochondria Produce Reactive Oxygen Species. *Biochem. J.* 417, 1–13. doi:10.1042/bj20081386
- Nadler, M. J. S., Hermosura, M. C., Inabe, K., Perraud, A.-L., Zhu, Q., Stokes, A. J., et al. (2001). LTRPC7 Is a Mg-ATP-Regulated Divalent Cation Channel Required for Cell Viability. *Nature* 411, 590–595. doi:10.1038/35079092
- O'donohoe, P. B., Huskens, N., Turner, P. J., Pandit, J. J., and Buckler, K. J. (2018). A1899, PK-THPP, ML365, and Doxapram Inhibit Endogenous TASK Channels and Excite Calcium Signaling in Carotid Body Type-1 Cells. *Physiol. Rep.* 6, e13876. doi:10.14814/phy2.13876
- Obeso, A., Almaraz, L., and Gonzalez, C. (1985). Correlation between Adenosine Triphosphate Levels, Dopamine Release and Electrical Activity in the Carotid Body: Support for the Metabolic Hypothesis of Chemoreception. *Brain Res.* 348, 64–68. doi:10.1016/0006-8993(85)90360-9
- Obeso, A., Almaraz, L., and Gonzalez, C. (1986). Effects of 2-Deoxy-D-Glucose on *In Vitro* Cat Carotid Body. *Brain Res.* 371, 25–36. doi:10.1016/0006-8993(86)90806-1
- Obeso, A., Gonzalez, C., Rigual, R., Dinger, B., and Fidone, S. (1993). Effect of Low O₂ on Glucose Uptake in Rabbit Carotid Body. *J. Appl. Physiology* 74, 2387–2393. doi:10.1152/jappl.1993.74.5.2387
- Olson, K. R. (2021). A Case for Hydrogen Sulfide Metabolism as an Oxygen Sensing Mechanism. *Antioxidants* 10, 1650. doi:10.3390/antiox10111650
- Olson, K. R., Healy, M. J., Qin, Z., Skovgaard, N., Vulesevic, B., Duff, D. W., et al. (2008). Hydrogen Sulfide as an Oxygen Sensor in Trout Gill Chemoreceptors. *Am. J. Physiology-Regulatory, Integr. Comp. Physiology* 295, R669–R680. doi:10.1152/ajpregu.00807.2007
- Ortega-Sáenz, P., Levitsky, K. L., Marcos-Almaraz, M. T., Bonilla-Henao, V., Pascual, A., and López-Barneo, J. (2010). Carotid Body Chemosensory Responses in Mice Deficient of TASK Channels. *J. Gen. Physiol.* 135, 379–392. doi:10.1085/jgp.200910302
- Ortega-Saenz, P., Pardal, R., Garcia-Fernandez, M., and Lopez-Barneo, J. (2003). Rotenone Selectively Occludes Sensitivity to Hypoxia in Rat Carotid Body Glomus Cells. *J. Physiology* 548, 789–800. doi:10.1113/jphysiol.2003.039693
- Peers, C., and Wyatt, C. N. (2007). The Role of maxiK Channels in Carotid Body Chemotransduction. *Respir. Physiology Neurobiol.* 157, 75–82. doi:10.1016/j.resp.2006.10.010
- Peng, Y.-J., Gridina, A., Wang, B., Nanduri, J., Fox, A. P., and Prabhakar, N. R. (2020). Olfactory Receptor 78 Participates in Carotid Body Response to a Wide Range of Low O₂ Levels but Not Severe Hypoxia. *J. Neurophysiology* 123, 1886–1895. doi:10.1152/jn.00075.2020
- Peng, Y.-J., Makarenko, V. V., Gridina, A., Chupikova, I., Zhang, X., Kumar, G. K., et al. (2019). H2S Mediates Carotid Body Response to Hypoxia but Not Anoxia. *Respir. Physiology Neurobiol.* 259, 75–85. doi:10.1016/j.resp.2018.08.001
- Peng, Y.-J., Nanduri, J., Raghuraman, G., Souvannakitti, D., Gadalla, M. M., Kumar, G. K., et al. (2010). H₂S Mediates O₂ Sensing in the Carotid Body. *Proc. Natl. Acad. Sci. U.S.A.* 107, 10719–10724. doi:10.1073/pnas.1005866107
- Pérez-García, M. T., Colinas, O., Miguel-Velado, E., Moreno-Domínguez, A., and López-López, J. R. (2004). Characterization of the Kv Channels of Mouse Carotid Body Chemoreceptor Cells and Their Role in Oxygen Sensing. *J. Physiol.* 557, 457–471. doi:10.1113/jphysiol.2004.062281
- Piruat, J. I., Pintado, C. O., Ortega-Sa'enz, P., Roche, M., and López-Barneo, J. (2004). The Mitochondrial SDHD Gene Is Required for Early Embryogenesis, and its Partial Deficiency Results in Persistent Carotid Body Glomus Cell Activation with Full Responsiveness to Hypoxia. *Mol. Cell Biol.* 24, 10933–10940. doi:10.1128/mcb.24.24.10933-10940.2004
- Prabhakar, N. R. (2006). O₂sensing at the Mammalian Carotid Body: Why Multiple O₂sensors and Multiple Transmitters? *Exp. Physiol.* 91, 17–23. doi:10.1113/expphysiol.2005.031922
- Rakoczy, R. J., Schiebrel, C. M., and Wyatt, C. N. (2022). Acute Oxygen-Sensing via Mitochondria-Generated Temperature Transients in Rat Carotid Body Type I Cells. *Front. Physiol.* 13, 874039. doi:10.3389/fphys.2022.874039
- Rogatzi, M. J., Ferguson, B. S., Goodwin, M. L., and Gladden, L. B. (2015). Lactate Is Always the End Product of Glycolysis. *Front. Neurosci.* 9, 22. doi:10.3389/fnins.2015.00022
- Roy, A., Mandadi, S., Fiamma, M.-N., Rodikova, E., Ferguson, E. V., Whelan, P. J., et al. (2012). Anandamide Modulates Carotid Sinus Nerve Afferent Activity via TRPV1 Receptors Increasing Responses to Heat. *J. Appl. Physiology* 112, 212–224. doi:10.1152/japplphysiol.01303.2010
- Roy, A., Rozanov, C., Mokashi, A., Daudu, P., Al-Mehdi, A. B., Shams, H., et al. (2000). Mice Lacking in Gp91 Phox Subunit of NAD(P)H Oxidase Showed Glomus Cell [Ca(2+)](i) and Respiratory Responses to Hypoxia. *Brain Res.* 872, 188–193. doi:10.1016/s0006-8993(00)02458-6
- Salman, S., Vollmer, C., McClelland, G. B., and Nurse, C. A. (2017). Characterization of Ectonucleotidase Expression in the Rat Carotid Body: Regulation by Chronic Hypoxia. *Am. J. Physiology-Cell Physiology* 313, C274–C284. doi:10.1152/ajpcell.00328.2016
- Shin, M.-K., Erasó, C. C., Mu, Y.-P., Gu, C., Yeung, B. H. Y., Kim, L. J., et al. (2019). Leptin Induces Hypertension Acting on Transient Receptor Potential Melastatin 7 Channel in the Carotid Body. *Circ. Res.* 125, 989–1002. doi:10.1161/circresaha.119.315338
- Shin, M.-K., Mitrut, R., Gu, C., Kim, L. J., Yeung, B. H. Y., Lee, R., et al. (2021). Pharmacological and Genetic Knockdown of Trpm7 in the Carotid Body Treats Obesity-Induced Hypertension. *Hypertension* 78, 104–114. doi:10.1161/hypertensionaha.120.16527
- Shirahata, M., Andronikou, S., and Lahiri, S. (1987). Differential Effects of Oligomycin on Carotid Chemoreceptor Responses to O₂ and CO₂ in the Cat. *J. Appl. Physiology* 63, 2084–2092. doi:10.1152/jappl.1987.63.5.2084
- Sommer, N., Alebrahimdehkordi, N., Pak, O., Knoepp, F., Strielkov, I., Scheibe, S., et al. (2020). Bypassing Mitochondrial Complex III Using Alternative Oxidase Inhibits Acute Pulmonary Oxygen Sensing. *Sci. Adv.* 6, eaba0694. doi:10.1126/sciadv.aba0694
- Spiller, P. F., Da Silva, M. P., and Moraes, D. J. A. (2021). Lactate Does Not Activate the Carotid Body of Wistar Rat. *Respir. Physiology Neurobiol.* 285, 103593. doi:10.1016/j.resp.2020.103593
- Swiderska, A., Coney, A. M., Alzahrani, A. A., Aldossary, H. S., Batis, N., Ray, C. J., et al. (2021). Mitochondrial Succinate Metabolism and Reactive Oxygen Species Are Important but Not Essential for Eliciting Carotid Body and Ventilatory Responses to Hypoxia in the Rat. *Antioxidants* 10, 840. doi:10.3390/antiox10060840
- Tanner, L. B., Goglia, A. G., Wei, M. H., Sehgal, T., Parsons, L. R., Park, J. O., et al. (2018). Four Key Steps Control Glycolytic Flux in Mammalian Cells. *Cell Syst.* 7, 49–62. doi:10.1016/j.cels.2018.06.003
- Telezkhin, V., Brazier, S. P., Cayzac, S. H., Wilkinson, W. J., Riccardi, D., and Kemp, P. J. (2010). Mechanism of Inhibition by Hydrogen Sulfide of Native and Recombinant BKCa Channels. *Respir. Physiology Neurobiol.* 172, 169–178. doi:10.1016/j.resp.2010.05.016
- Tipparaju, S. M., Saxena, N., Liu, S.-Q., Kumar, R., and Bhatnagar, A. (2005). Differential Regulation of Voltage-Gated K⁺channels by Oxidized and Reduced Pyridine Nucleotide Coenzymes. *Am. J. Physiology-Cell Physiology* 288, C366–C376. doi:10.1152/ajpcell.00354.2004
- Torres-Torreló, H., Ortega-Sáenz, P., Gao, L., and López-Barneo, J. (2021). Lactate Sensing Mechanisms in Arterial Chemoreceptor Cells. *Nat. Commun.* 12, 4166. doi:10.1038/s41467-021-24444-7
- Torres-Torreló, H., Ortega-Sáenz, P., Macías, D., Omura, M., Zhou, T., Matsunami, H., et al. (2018). The Role of Olf78 in the Breathing Circuit of Mice. *Nature* 561, E33–E40. doi:10.1038/s41586-018-0545-9
- Turner, P. J., and Buckler, K. J. (2013). Oxygen and Mitochondrial Inhibitors Modulate Both Monomeric and Heteromeric TASK-1 and TASK-3 Channels in Mouse Carotid Body Type-1 Cells. *J. Physiol.* 591, 5977–5998. doi:10.1113/jphysiol.2013.262022
- Varas, R., Wyatt, C. N., and Buckler, K. J. (2007). Modulation of TASK-like Background Potassium Channels in Rat Arterial Chemoreceptor Cells by Intracellular ATP and Other Nucleotides. *J. Physiol.* 583, 521–536. doi:10.1113/jphysiol.2007.135657
- Verna, A., Talib, N., Roumy, M., and Pradet, A. (1990). Effects of Metabolic Inhibitors and Hypoxia on the ATP, ADP and AMP Content of the Rabbit Carotid Body *In Vitro*: the Metabolic Hypothesis in Question. *Neurosci. Lett.* 116, 156–161. doi:10.1016/0304-3940(90)90402-u
- Waypa, G. B., Marks, J. D., Guzy, R. D., Mungai, P. T., Schriewer, J. M., Dokic, D., et al. (2013). Superoxide Generated at Mitochondrial Complex III Triggers Acute Responses to Hypoxia in the Pulmonary Circulation. *Am. J. Respir. Crit. Care Med.* 187, 424–432. doi:10.1164/rccm.201207-1294oc
- Williams, B. A., and Buckler, K. J. (2004). Biophysical Properties and Metabolic Regulation of a TASK-like Potassium Channel in Rat Carotid Body Type I Cells. *Am. J. Physiology-Lung Cell. Mol. Physiology* 286, L221–L230. doi:10.1152/ajplung.00010.2003

- Wyatt, C. N., and Peers, C. (1995). Ca(2+)-activated K⁺ Channels in Isolated Type I Cells of the Neonatal Rat Carotid Body. *J. Physiol.* 483 (Pt 3), 559–565. doi:10.1113/jphysiol.1995.sp020606
- Wyatt, C. N., and Buckler, K. J. (2004). The Effect of Mitochondrial Inhibitors on Membrane Currents in Isolated Neonatal Rat Carotid Body Type I Cells. *J. Physiol.* 556, 175–191. doi:10.1113/jphysiol.2003.058131
- Wyatt, C. N., Mustard, K. J., Pearson, S. A., Dallas, M. L., Atkinson, L., Kumar, P., et al. (2007). AMP-activated Protein Kinase Mediates Carotid Body Excitation by Hypoxia. *J. Biol. Chem.* 282, 8092–8098. doi:10.1074/jbc.m608742200
- Yagi, H., Kasai, T., Rioual, E., Ikeya, T., and Kigawa, T. (2021). Molecular Mechanism of Glycolytic Flux Control Intrinsic to Human Phosphoglycerate Kinase. *Proc. Natl. Acad. Sci.* 118, e2112986118. doi:10.1073/pnas.2112986118
- Yamamoto, Y., and Taniguchi, K. (2006). Expression of Tandem P Domain K⁺ Channel, TREK-1, in the Rat Carotid Body. *J. Histochem Cytochem.* 54, 467–472. doi:10.1369/jhc.5a6755.2005
- Yuan, G., Vasavda, C., Peng, Y. J., Makarenko, V. V., Raghuraman, G., Nanduri, J., et al. (2015). Protein Kinase G-Regulated Production of H₂S Governs Oxygen Sensing. *Sci. Signal* 8, ra37. doi:10.1126/scisignal.2005846
- Zhang, M., Zhong, H., Vollmer, C., and Nurse, C. A. (2000). Co-release of ATP and ACh Mediates Hypoxic Signalling at Rat Carotid Body Chemoreceptors. *J. Physiology* 525, 143–158. doi:10.1111/j.1469-7793.2000.t01-1-00143.x
- Zhou, T., Chien, M.-S., Kaleem, S., and Matsunami, H. (2016). Single Cell Transcriptome Analysis of Mouse Carotid Body Glomus Cells. *J. Physiol.* 594, 4225–4251. doi:10.1113/jp271936

Conflict of Interest: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Publisher's Note: All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

Copyright © 2022 Holmes, Swiderska, Nathanael, Aldossary, Ray, Coney and Kumar. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.