# **UNIVERSITY** OF BIRMINGHAM University of Birmingham Research at Birmingham

# Social facilitation of energy intake in adult women is sustained over three days in a crossover laboratory experiment and is not compensated for under freeliving conditions

Ruddock, Helen K.; Brunstrom, Jeffrey M.; Vartanian, Lenny R.; Higgs, Suzanne

DOI 10.1016/j.appet.2022.106141

License: Creative Commons: Attribution-NonCommercial-NoDerivs (CC BY-NC-ND)

Document Version Peer reviewed version

*Citation for published version (Harvard):* Ruddock, HK, Brunstrom, JM, Vartanian, LR & Higgs, S 2022, 'Social facilitation of energy intake in adult women is sustained over three days in a crossover laboratory experiment and is not compensated for under free-living conditions', Appetite, vol. 176, 106141. https://doi.org/10.1016/j.appet.2022.106141

Link to publication on Research at Birmingham portal

#### **General rights**

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

•Users may freely distribute the URL that is used to identify this publication.

•Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.

•User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?) •Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

#### Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

# Social facilitation of energy intake in adult women is sustained over three days in a crossover laboratory experiment and is not compensated for under free-living conditions

Helen K. Ruddock<sup>1</sup>, Jeffrey M. Brunstrom<sup>2, 3</sup>, Lenny R. Vartanian<sup>4</sup>, Suzanne Higgs<sup>1</sup>

School of Psychology, University of Birmingham, Birmingham, United Kingdom (HR, SH)
 Nutrition and Behaviour Unit, School of Psychological Science, University of Bristol,
 Bristol, United Kingdom (JMB)
 NIHR Bristol Biomedical Research Centre, University Hospitals Bristol and Weston NHS
 Foundation Trust and University of Bristol (JMB)
 School of Psychology, UNSW Sydney, Sydney, Australia (LV)

Corresponding author: Professor Suzanne Higgs, School of Psychology, University of Birmingham, Birmingham, United Kingdom E-Mail: s.higgs.1@bham.ac.uk Tel: +441214144907

**Pre-registration**: The study methods and analysis plan were registered prior to analysis on the Open Science Framework website: Study 1 (<u>https://doi.org/10.17605/OSF.IO/HMABE</u>), Study 2 <u>https://doi.org/10.17605/OSF.IO/FA3PN</u>

#### Sources of support for the work

The research was funded by an Economic and Social Research Council grant awarded to SH, JB, and LV. Project reference: ES/P01027X/1.

This study was supported by the National Institute for Health and Care Research Bristol Biomedical Research Centre. The views expressed are those of the author(s) and not necessarily those of the NIHR or the Department of Health and Social Care Running title; Sustained social facilitation of intake

#### Abstract

2 People eat more when they eat a meal with familiar others than they do when eating alone. 3 However, it is unknown whether eating socially impacts intake over the longer-term. The aim of Study 1 was to examine whether socially facilitated intake is sustained across all meals and 4 5 across three consecutive days. The aim of Study 2 was to examine whether increased intake 6 during a social meal taken in the laboratory is compensated for under free-living conditions. 7 In Study 1, adult women (n = 26) ate all their meals across three days either with a friend or 8 alone in a counterbalanced cross-over design. In Study 2 adult women (n = 63) consumed a 9 meal in the laboratory either alone or with two friends and then recorded everything they ate 10 and drank for the next three days using electronic food diary software. In Study 1 intake 11 across 3 days was significantly greater in the Social (M = 7310 kcal, SD = 1114) than in the 12 Alone condition (M = 6770 kcal, SD = 974) (F(1,423) = 16.10, p < .001, d = 0.51). In Study 13 2 participants consumed significantly more in the laboratory when eating with their friends 14 (M = 1209 kcal, SD = 340) than when eating alone (M = 962 kcal, SD = 301) (F(1,63) = 15 13.28, p = .001, d = 0.77). Analysis of food diary data plus laboratory intake showed that 16 intake remained significantly greater in the Social (M = 6396 kcal, SD = 1470) than in the Alone condition after 4 days (M = 5776 kcal, SD = 1182) (F(1,59) = 5.59, p = .021, d =17 18 0.05). These results show that social facilitation of eating is sustained over three days and 19 suggest that people fail to compensate for the social facilitation of eating. 20

21 Key words: Social influences; Food intake; Compensation; Food diary; Laboratory study.

- 22
- 23

#### Introduction

25 Food intake is strongly influenced by environmental cues. For example, people eat more 26 when presented with a greater variety of foods (McCrory et al., 1999) and when served a larger 27 portion size (Rolls, Morris, & Roe, 2002). Another important, and yet often overlooked, 28 external influence on food intake is social context. The mere act of eating socially exerts a 29 particularly powerful influence on food intake (de Castro et al., 1990). Known as the 'social 30 facilitation of eating,' research using food diaries, covert observation, and experimental 31 manipulations have all shown that people eat more when eating with others, especially when 32 eating with friends and family, relative to when dining alone (Ruddock, Brunstrom, Vartanian, 33 & Higgs, 2019).

34 Although we see evidence that energy intake is influenced by environmental cues, almost all studies have focused on food intake at a single occasion. But what happens if the cue is 35 36 presented over a longer period? Is the increase energy intake sustained over several days? Or 37 does the effect wane over time? Even if the effect is sustained with repeated presentations of 38 the cue, what happens in between eating occasions? Do people compensate for the effect of the 39 cue by reducing their food intake in other meals? If they do, perhaps the net result is no overall 40 increase in intake? Despite their importance, these questions remain largely unexplored. A rare 41 exception can be found in the work of Rolls et al. (2006, 2007), who found that participants ate 42 more when they were provided with larger portions, and that this pattern was sustained across 43 all meals consumed over two (Rolls, Roe, & Meengs, 2006) and even 11 consecutive days 44 (Rolls, Roe, & Meengs, 2007). These findings are important because they suggest that 45 environmental cues may contribute to longer-term increases in energy intake which, over time, 46 could contribute to weight gain and possibly to the development of obesity.

The social facilitation of intake is even larger than the effect of portion size (Ruddock et al., 2019) but it is unknown whether the effect is sustained over time and whether or not people compensate for socially facilitated intake. To find out whether the social facilitation of eating affects energy intake over the longer-term, experimental research is required in which social context is systematically manipulated in the laboratory. Examining the social facilitation of eating under controlled conditions is important because it eliminates extraneous explanations for such effects (e.g. differences in setting, portion sizes, etc.), and provides insight into the *causal* relationship between social context and energy intake.

55 The aim of the current research was to examine whether the social facilitation of eating 56 is sustained over several days and whether people compensate for socially facilitated intake. In 57 Study 1, we tested the hypothesis that participants would eat more when eating with a friend, 58 relative to when eating alone, and that this effect would be sustained across breakfast, lunch, 59 and dinner, and across three consecutive days. In Study 2, we examined whether increased 60 intake during a social meal with friends was compensated for under free-living conditions. 61 Study 2 tested the hypothesis that participants who consume a social meal with friends in the 62 lab would eat more than participants who eat alone, and that this difference in intake would not 63 be compensated for by eating less at subsequent meals consumed within real-world settings.

64

65

#### Study 1

Method

#### 66 **Participants**

67 Pairs of friends were recruited via social media and poster advertisements which were placed around the University of Birmingham campus. Only female participants were 68 69 recruited to reduce error variance related to sex/gender differences in amounts consumed, and because women eating with men is associated with reduced intake due to impression 70 71 management concerns (Brindal, Wilson, Mohr, & Wittert, 2015; Vartanian, 2015). Both men and women have been observed to show social facilitation of intake (Ruddock et al., 2019). 72 The study was advertised as examining the effect of 'time of day and group working on 73 74 problem solving ability'. Participants were eligible to take part if they met the following

criteria which were listed on the study advertisement: 1) were aged over 18 years, 2) were 75 76 occasional social eaters (1-3 meals per week), 3) had a self-reported BMI between 18-25  $kg/m^2$ , 4) liked and were willing to eat the test foods, and 5) were willing to refrain from 77 consuming calorie-containing food and drinks outside of those provided during the study. 78 79 Participants were excluded if they were on any medication known to affect appetite, had been 80 diagnosed with an eating disorder, were regular smokers, were following a weight-loss diet, 81 were an athlete in training, were pregnant or breastfeeding, or had any food allergies or 82 intolerances. Using G\*Power (Faul, Erdfelder, Lang, & Buchner, 2007), we calculated that a 83 sample size of 26 participants was required to provide 80% power to detect medium-sized 84 main effects of eating condition (social versus alone) and interactions between condition and day/meal type (f=.30) using a repeated-measures design (alpha = .05). The findings from a 85 86 meta-analysis of the effects of social facilitation on eating (Ruddock et al., 2019) suggest a 87 large effect size but, given that it is unknown whether the effects are maintained over time, 88 we took a more conservative approach and predicted a medium effect size. Participants 89 received cash in exchange for taking part. The study methods and analysis plans were 90 registered after data collection but before analysis on the Open Science Framework website (https://doi.org/10.17605/OSF.IO/HMABE). The study protocol was approved by the 91 92 University of Birmingham's Research Ethics Committee and was conducted in line with 93 ethical standards stated in the Declaration of Helsinki 1975.

94

#### 95 Design

A within-subjects counter-balanced crossover design was used in which participants attended two phases of 3 consecutive days (weekdays only). In one phase, participants attended alone for 3 days and in the other phase they attended with a friend (also a participant). The two 3-day phases were separated by a washout period of 14 days. The order 100 of Social versus Alone phase was randomly determined by the researcher (HR) using the

101 random integer generator available at: https://www.random.org/integers/.

#### 102 Measures

103 Three Factor Eating Questionnaire-18.

The Three-Factor Eating Questionnaire Revised 18-item version (TFEQ-18) was 104 105 included to assess dietary behaviour (Karlsson, Persson, Sjöström, & Sullivan, 2000). The 106 instrument is a shortened and revised version of the original 51-item TFEQ (Stunkard & 107 Messick, 1985), and it comprises the following three subscales: 1) dietary restraint (i.e. 108 attempts to restrict food intake in order to control body weight; six items), 2) uncontrolled 109 eating (i.e. tendency to experience a loss of control over eating; nine items), and 3) emotional 110 eating (i.e. eating in response to negative moods; three items). In each case, a higher score 111 reflects a tendency to exhibit the associated construct.

112

#### 113 Food menus

The three daily menus are presented in **Table 1**. The same meals were provided for the Social and Alone conditions but the order of menus was randomised within phase. One litre of water was provided at each meal. At breakfast, participants were offered a choice of either tea or coffee, along with the option to add up to 50 ml of semi-skimmed milk and 15 g of sugar. The amount of each food provided was fixed and so participants could not ask for more of the individual foods. However, sufficient food was provided overall, such that participants could not consume all of it (and none did).

121

#### 122 **Procedure**

Testing took place between February 2019 and August 2019. All eligible participants
were tested. On each day, participants came to the eating behaviour laboratory at the
University of Birmingham for breakfast (between 8-10am), lunch (between 12-2pm), and

dinner (between 5-7pm), and were instructed to refrain from eating or drinking any caloriecontaining drinks, other than those provided during the test days. Meal timings were
scheduled to allow 4 hours between breakfast and lunch, and 5 hours between lunch and
dinner, and participants were free to leave the lab between meals. On each of the three days a
different menu was served, and the order of these menus was counterbalanced across
participants.

132 Before each meal, participants completed a short questionnaire in which they were 133 asked whether they had felt ill since their last meal, whether they had taken any medication 134 which may have affected their appetite, and whether they had consumed any other 135 foods/caloric beverages since their last meal. Participants who answered positively to the 136 latter question were asked to record a) what and how much they ate, b) the time that they ate, c) where they ate, and c) how many people were present when they ate. Before breakfast, 137 138 participants were also asked to record the amount of time (in minutes) that they had spent 139 engaging in light, moderate, and vigorous activities in the past 24 hours. Before each meal, 140 participants completed hunger and fullness ratings (see supplementary materials for a 141 description of these measures).

142 Participants were then seated in a dining room (a room in the laboratory furnished with 143 a table/tablecloth, table lamp and dining chairs) either alone (Alone condition) or with their 144 friend (Social condition) and were provided with the meal which was laid out on the table. In 145 the Social condition, both participants were presented with the same foods, though each 146 participant had their own meal (i.e. they did not share a meal). Participants were invited to eat 147 as they normally would, i.e. communication was not prohibited, and were told they could eat 148 as much as they wished and to notify the experimenter once they had finished eating. The 149 researcher covertly recorded the duration of the meal (see supplementary materials), and food intake was determined by covertly weighing foods before and after each meal. Following the 150 151 meal, participants completed measures of hunger and fullness, food liking, and mood (see

152 supplementary materials for a description of these measures). To reinforce the believability of 153 the cover story, participants were then given five minutes to complete a word- or number-154 based problem-solving activity. They also completed a short questionnaire about how 155 difficult they found the activity, whether they thought the time of day had affected their 156 performance, and the strategy that they had used to complete the task with their friend (if 157 applicable).

158 At the end of the study (i.e. after dinner on day 3, phase 2), demand characteristics were 159 assessed by asking participants to write down what they thought were the aims of the study. 160 Measures of friendship closeness were also taken by asking participants how long they had 161 known their friend (in months), how well they think they know their friend (using a scale ranging from 1 to 10 with anchor points 'Not very well' and 'Very well' respectively), and 162 163 how close they feel to their friend (1-10 scale with anchor points 'Not very close' and 'Very 164 close', respectively). Participants also indicated their age and ethnicity, and then completed 165 the TFEQ. Finally, the experimenter assessed the participant's height and weight, which was 166 used to calculate BMI, and participants were fully debriefed as to the true aims of the study. 167

#### 168 Data analysis

169 For the main variable of interest (i.e. calorie intake), outlying values were identified 170 using Hoaglin and Iglewicz's (1987) outlier labelling rule. Six participants reported feeling ill 171 prior to at least 1 meal occasion, and one participant reported taking medication which may 172 affect appetite prior to two meal occasions (both social meals). However, the amount eaten 173 by these participants was within the normal range (i.e. none were identified as outliers) and 174 so their data was retained within subsequent analyses. Because observations were non-175 independent (i.e. participants signed up to the study in pairs), data were analysed using a multilevel model (MLM). Condition (i.e. Alone vs Social), condition order (i.e. Social first vs 176 177 Alone first), day (i.e. day 1, day 2, day 3), and meal (breakfast, lunch, and dinner) were

entered as fixed effects predictors of calorie intake. In the multi-level analyses, we planned to
include covariates in cases where a variable correlated significantly with the dependent
variable. Variables tested for correlations were age, BMI, and hunger. Statistical analyses
were conducted using SPSS version 27.0 (IBM Corp, 2020). For exploratory analyses of
effects of social context on food liking, appetite and mood change, see Supplementary
analyses.

#### 184

#### Results

185 **Participants** 

A total of 26 participants (13 friend pairs) took part in the study. Participant 186 187 characteristics are provided in Table 2. No participants guessed the true aims of the study. 188 The majority (n=21) confirmed that they had not eaten or consumed any calorie-containing 189 drinks, other than those provided to them, across the three days. Five participants reported 190 that they had consumed additional food on at least one occasion during the six test days. Of 191 these, two had consumed extra food during the Alone phase, two consumed additional food 192 during the Social phase, and one participant consumed additional food during both Social and 193 Alone conditions. Removal of these participants did not affect the overall findings and so 194 their data were included in the final analysis.

Initial inspection of the calorie intake data revealed one outlying value (295 kcal consumed at lunch, day 1, Social condition). However, removing this datapoint had no material impact on outcomes of the statistical analyses, and so the results are reported with this datapoint included. Datapoints from one participant-pair (dinner, day 3, Social condition) were removed due to a failure to follow instructions.

200 In support of our hypothesis, participants ate significantly more calories in the Social

201 condition (M=7310 kcal, SD=1114) relative to the Alone condition (M= 6770 kcal, SD=974),

202 F(1,423)=16.10, p<.001, d=0.51 (Figure 1). Figure 1 presents mean calories consumed as a

203 function of condition, day, and meal. There was also a main effect of day on food intake,

204	F(2,423)=7.05, p<.001, such that participants ate less on day 1 than on day 2 (p=.008) and
205	day 3 (p<.001). Calories consumed did not differ between days 2 and 3 (p=.32), and there
206	was neither a significant day $\times$ condition interaction, F(2,423)=0.08, p=.92, nor a significant
207	meal type × condition interaction, $F(2,423)=2.33$ , p=.098.
208	
209	Interim discussion
210	The results from Study 1 support our hypothesis that participants would eat more when
211	eating with a friend than when eating alone, and that this effect would be sustained across
212	breakfast, lunch, and dinner, as well as across three consecutive study days. This is important
213	because it suggests that the social facilitation of eating persists across multiple meals,
214	producing a sustained increase in energy intake over time.
215	In Study 2, we build on these findings by examining whether participants compensate
216	for the social facilitation of eating by reducing their energy intake at subsequent meals in a
217	real-world setting. In Study 1, participants ate all their meals either socially or alone, and so it
218	is unclear whether increased intake at a social meal might be offset by a reduction in intake at
219	the next eating opportunity. Therefore, in Study 2 we examined self-reported free-living
220	intake over four consecutive days immediately after participants had eaten a buffet lunch in
221	the lab either alone (Alone condition) or with two friends (Social condition). In line with
222	evidence of inadequate energy compensation following changes in energy intake (Levitsky,
223	2005; Levitsky et al., 2019), we hypothesised that participants in the Social condition would
224	eat more than those in the Alone condition, and this would not be compensated for by eating
225	less at subsequent meals.

#### Study 2

#### 12

#### 228

#### Method

#### 229 Participants

230 Participants were recruited via social media and poster advertisements which were 231 placed around the University of Birmingham campus. Participants signed up to the study in 232 groups of three friends. As in Study 1, only female participants were recruited. The study was 233 advertised as examining the effect of 'mood on eating behaviour.' The inclusion/exclusion 234 criteria were the same as for Study 1 except that there was no requirement for participants to 235 refrain from eating anything outside of the lab. Using G\*Power, and based on the results of 236 Study 1, we calculated that a sample size of 60 participants would be required to provide 80% 237 power to detect medium-sized main effects (f=.37) between the Social and Alone conditions 238 for total caloric intake (alpha = .05) in a between-subjects design. We predicted a slightly 239 larger effect size for Study 2 because the participants are eating with 2 friends rather than 1 as 240 they did for Study 1 and social facilitation is known to be enhanced when there are more people present (De Castro and Brewer 1992). Additional participants were recruited to 241 242 account for attrition. In total, 69 took part. The study method and analysis plan were 243 preregistered after data collection but prior to analysis on the Open Science Framework 244 website (https://doi.org/10.17605/OSF.IO/FA3PN). The study protocol was approved by the 245 University of Birmingham's Research Ethics Committee and was conducted in line with ethical standards stated in the Declaration of Helsinki 1975. All eligible participants were 246 247 tested.

248

#### 249 Design

A between subjects (Social versus alone condition) design was used in which participants took part in either the alone or social eating condition and then completed a food diary for the remainder of that day and the next three days. As participants signed up for the study as a group of three friends, each trio was randomly allocated to either the Alone or
Social condition by the researcher (HR) using the random integer generator available at:
https://www.random.org/integers/. Participants in the Alone condition ate the buffet lunch in
the laboratory alone, while those in the Social condition ate the same buffet with their two
friends.

258

259 Materials

260 Buffet lunch

Participants were provided with a buffet lunch comprising 1952 kcal. **Table 3** provides a full list of foods provided to each participant. The amount of each food provided was fixed and so participants could not ask for more of the individual foods. However, sufficient food was provided overall, such that participants could not consume all of it (and none did).

266

267 Food diary

268 Participants used Myfood24 software (2016) to record everything that they ate and 269 drank for the 4 days following their initial lab session. MyFood24 is a 24-hour dietary recall tool that provides a valid and user-friendly measure of food intake (Carter et al., 2015; Wark 270 271 et al., 2018). After each eating episode (breakfast, lunch, dinner, and snacks), participants 272 recorded the foods and drinks consumed, and their respective portion size. To minimise 273 under-reporting, Myfood24 also includes prompts for commonly forgotten foods, and 274 participants are asked to review their diary before submitting it. The output is generated by drawing on a nutritional information database of 40,274 food items and it provides a 275 276 summary of daily calories consumed by each participant. After submitting a food diary, 277 participants were automatically directed to a follow-up questionnaire (using Qualtrics software) in which they were asked to record how many people they ate with during each 278

meal or snack. If participants indicated that they had eaten a meal or snack with one or more people, they were asked to record how well they knew each person. Specifically, for each individual at the meal, they indicated whether the person was a friend, a family member, a romantic partner, an acquaintance, or a stranger. To obscure the true purpose of the study, and consistent with the cover story, participants were then asked to choose words that described their mood during each meal or snack (i.e. happy, angry, annoyed, sad/depressed, excited, content, anxious).

#### 286 **Procedure**

287 Testing took place between October 2019 and February 2020, and in the same room as in Study 1. Participants were invited to attend the Eating behaviour laboratory at the 288 289 University of Birmingham between 12-2pm (to coincide with normal lunch hours), and were 290 instructed to refrain from eating or drinking any calorie-containing drinks for at least three 291 hours before the start of their session. Participants arrived at the lab with their two friends. 292 Before the meal, participants completed VAS measures of hunger and fullness, and then completed a measure of food cravings (see supplementary materials for descriptions of 293 294 these measures). They were then offered the buffet lunch and were instructed to eat as much 295 as they wished. Food was laid out on a table and each participant was given their own buffet 296 (i.e. friends did not share). Participants were invited to eat as they normally would, i.e. 297 communication was not prohibited, and they were told they could eat as much as they wished 298 and to notify the experimenter once they had finished eating. Meal duration was recorded 299 covertly, and foods were weighed covertly before and after eating to determine food intake. 300 Following the meal, participants were placed in separate rooms and completed VAS 301 measures of appetite, food liking, mood, and overall meal enjoyment (see supplementary 302 materials).

After completing the questionnaires, participants were shown how to record their food intake using the Myfood24 software. They were instructed to record everything that they ate and drank for the remainder of that day (day 1), and for three subsequent days (days 2-4).

306 Alcohol and non-alcoholic drinks were recorded but due to the high social intake of 307 alcohol in this population we analysed the data with and without calories from alcohol. 308 Between five and 14 days following the first lab session, participants returned to the lab to 309 complete the following assessments: 1) demand characteristics were assessed by asking 310 participants to write down what they thought the aims of the study were; 2) dietary restraint, 311 uncontrolled eating, and emotional eating, were assessed using the TFEQ-18 (Karlsson et al., 312 2000) (described in Study 1); 3) friendship familiarity was assessed by asking participants to write down how long they have known each of the friends with whom they had participated 313 314 (open ended question), and how well they felt they know these friends (1-10 scale anchored 315 by 'Not very well' and 'Very well', respectively); and 4) other demographics, including age 316 and ethnicity. Height and weight were then measured by the researcher to calculate BMI, and 317 participants were fully debriefed as to the true aims of the study.

318

#### 319 **Data analysis**

320 For the main variables of interest (calorie intakes in the lab and across days), outlying 321 values were identified using Hoaglin and Iglewicz's (1987) outlier labelling rule. Because 322 observations are non-independent (i.e. participants signed up to the study in groups of three), 323 data were analysed using MLMs. In three separate analyses, Condition (Alone vs Social) was 324 entered as a fixed-effects predictor of food consumed (kcal) during the lab session, of total 325 intake (kcal) at the end of day 1 (lab intake + food diary intake for day 1) and of total intake 326 at the end of day 4 (lab intake + food diary intake for days 1 to 4). Potential covariates were entered into a bivariate correlation matrix with the dependent variables. Variables that were 327

328	significantly correlated with a dependent variable were included as covariates in the multi-
329	level analyses. Statistical analyses were conducted using SPSS version 27.0 (IBM, 2020).
330	
331	Results
332	Initial data checks
333	Initial inspection of the data revealed that calorie intake on day 1 (food diary data) was
334	not normally distributed (skewness=3.02, SE=0.29; kurtosis=15.06, SE=0.58). Using Hoaglin
335	and Iglewicz's (1987) outlier labelling rule, two participants were identified as outliers,
336	having consumed over 1837 kcal on day 1 after the lab session (i.e. not including calories
337	consumed within the lab or before lunch). Removing these participants corrected the
338	distribution, and they were therefore excluded from subsequent analyses. No participants
339	correctly guessed the aim of the study. Initial inspection of the data revealed that BMI
340	correlated positively with the amount consumed during the lab session, r=.283, p=.021, total
341	day 1 intake (i.e. lab intake + food diary intake for day 1), r=.322, p=.009, and total intake
342	across all four days, r=.383, p=.002. BMI was therefore included as a covariate in the main
343	analyses.

#### 345 **Participants**

Participant characteristics are shown in **Table 4**. A MANOVA revealed no betweencondition differences in the participants' age, BMI, TFEQ-subscale scores, or friendship familiarity, F(6,59)=0.63, p=.630. MLM analyses also revealed no differences between conditions on appetite ratings prior to the meal, F(1,65)=0.24, p=.623 (Social: M=78.1 SD=13.1; Alone: 76.6, SD=12.0).

#### 352 Effect of condition on food intake in the lab

353 The MLM revealed a significant effect of condition on food intake, F(1,63)=13.28,

354 p=.001, d=0.77, such that participants in the Social condition consumed significantly more

calories than did those in the Alone condition (Social: M=1209 kcal, SD=340; Alone: M=962

- 356 kcal, SD=301).
- 357

#### 358 Effect of condition on day 1 intake

Food diary data for day 1 were obtained from 65 participants (Alone n=33; Social

n=32). There was a significant effect of condition on day 1 total intake, F(1,61)=5.79,

361 p=.019, d=0.50. Participants consumed significantly more in the Social condition (M=1990

362 kcal, SD=468) than in the Alone condition (M=1756 kcal, SD=460). This result did not

363 change when adding calories from alcohol (Social: M=2080 kcal, SD=525; Alone: M=1845

364 kcal, SD=482; F(1,61)=4.37, p=.041).

Further analyses revealed that there were no between-condition differences in calories consumed *after* the lab session on day 1, F(1, 61)=0.03, p=.875, or the number of meals eaten socially F(1,65)=0.24, p=.877. These findings are important because they suggest that the difference in total calorie intake at the end of day 1 was due to differences in intake that occurred during the lab meal.

370

#### 371 Effect of condition on total four-day calorie intake

Total four-day intake (i.e. lab calories + *all* food diary data) was obtained from 63 participants (Alone n=31; Social n=32). Participants in the Social condition consumed

374 significantly more calories over the four days than did those in the Alone condition (Social:

375 M=6396 kcal, SD=1470; Alone: M=5776 kcal, SD=1182), F(1,59)=5.59, p=.021, d=0.46.

376 This result did not change when adding calories from alcohol (Social: M=6712 kcal,

377 SD=1600;Alone: M=5980 kcal, SD=1228; F(1,59)=6.33, p=.015).

378Removing the food consumed during the lab session showed that the amount consumed379during the four days *following* the lab session did not differ between conditions,380F(1,59)=2.32, p=.133, and there were no between-condition differences in the number of381meals that were consumed socially after the lab session, F(1,64)=0.30, p=.589. These382findings suggest that participants in the Social condition did not compensate for additional383food consumed during the lab session by eating less over subsequent meals (see Figure 2).

- 384
- 385

#### Discussion

386 Across two controlled studies, we provide the first evidence that the social facilitation 387 of eating is sustained across several days (Study 1), and that people fail to compensate for 388 additional calories consumed during social meals under free-living conditions (Study 2). In 389 Study 1, across three consecutive days, participants consumed an additional 539 kcal when 390 they ate all their meals with a friend (relative to eating alone). These findings were extended 391 in Study 2, in which we examined whether participants would compensate for the social 392 facilitated increase in energy intake in the lab by reducing their energy intake at subsequent 393 real-world meals. Those who ate a social meal in the lab consumed a larger lunch (additional 394 247 calories) than did those who ate alone, and there was no evidence for compensation 395 across the following four days. Together, the findings from Studies 1 and 2 suggest that the 396 social facilitation of eating is sustained over time and that people fail to compensate for the 397 social facilitation of eating. These findings are important because they suggest that eating 398 socially may lead to greater energy intake over the longer-term. On average social meals 399 were around 150 calories larger than non-social meals. For a woman with an average height 400 and weight, relative to eating alone, consuming one social meal per day could result in weight 401 gain of around 4 kg over a year (Hall et al., 2011).

402 By experimentally manipulating social context, we can rule out other explanations for 403 the social facilitation of eating (e.g. differences in the type of food available, context, etc.) 404 and show that eating socially has a *causal* effect on energy intake, which persisted over 405 several days. Our findings are also consistent with research demonstrating sustained effects 406 of other environmental cues (i.e. portion size) on intake (Rolls et al., 2006, 2007). Together, 407 these findings provide further support for the idea that at least over a period of days, 408 stimulation of intake by external factors does not induce active regulatory appetite 409 mechanisms to counteract increased consumption (Levitsky, 2005; Levitsky et al., 2019), and 410 that day-to-day intake is not under tight biological regulation (Casanova, Finlayson, Blundell, 411 & Hopkins, 2019).

412 Evidence for the persistent effects of social context on intake over several days is 413 consistent with the idea that the social facilitation of eating may be a hard-wired 414 psychological phenomenon. Indeed, evidence for the social facilitation of eating has been 415 observed across a range of non-human animals (Forkman, 1991; Harlow & Yudin, 1933; 416 Rajecki, Kidd, Wilder, & Jaeger, 1975; Tolman, 1964), suggesting that it may serve an 417 important evolutionary purpose. As we have discussed in detail elsewhere (Ruddock et al., 418 2019), one possibility is that the social facilitation of eating evolved as a strategy to ensure 419 that we obtain maximum personal resources while sharing limited food resources with other 420 group members.

421 There are implications of the present results for healthy eating and nutritional 422 interventions. Social eating might be used to increase the food intake of undernourished 423 populations e.g. elderly people with reduced appetite. People who wish to avoid overeating, 424 might wish to develop strategies that allow them to experience the benefits of social eating 425 (Dunbar, 2017) while at the same time mitigating the effects of social context on excess 426 calorie intake. One strategy may be to actively compensate for socially facilitated food intake 427 by eating smaller meals before or after a social meal. Another strategy may be to advise people to plan their meal in advance of a social occasion. Indeed, in a recent study (Ruddock, 428 429 Long, Brunstrom, Vartanian, & Higgs, 2021) we found that participants who served

themselves *before* eating with a friend consumed significantly fewer calories than those who
served themselves *during* the meal. Thus, pre-ordering food or serving oneself before the
start of a meal, may help people to avoid unintentionally overeating during social meals.

A strength of the present studies is that we examined food intake within laboratory-433 434 and real-world settings, and so we were able to establish the *causal* effect of social eating on 435 longer-term calorie intake, while maximising the applicability of our findings to real-world 436 contexts. A further methodological strength was that food intake was monitored for several 437 days after a social meal (Study 2). The results also suggest that laboratory-based 438 demonstrations of the social facilitation of eating are unlikely to be explained by the novelty 439 of eating with a friend in a context in which free food is available, because such an effect might be expected to wear off over time. 440

441 A limitation of the present studies is that we did not measure energy expenditure and 442 so we cannot rule out the possibility that participants compensated for additional calorie 443 intake by expending more energy. In Study 1, participants recorded the amount of time that they had spent engaging in light, moderate, and vigorous exercise during the 24 hours prior to 444 445 each test day. Analysis of these data revealed no significant main effect of condition (Alone vs Social) on exercise duration, suggesting that participants did not compensate for socially 446 447 facilitated food intake by engaging in more physical activity (see supplementary materials). However, future research could incorporate other more precise measures of energy 448 expenditure (such as actigraphy). In addition, recruitment was restricted to women with a 449 450 BMI within the normal weight range. It is therefore important for future research to establish the generalisability of our findings to other populations. To date there has been no systematic 451 study of the moderating effects of weight status and/or sex/gender on social facilitation of 452 453 eating. However data from self-report and observational studies indicate that people with overweight may show a weaker effect, perhaps because concerns about portraying a 454

455 particular impression to others overrides social facilitation effects in these contexts (Salvy, de

456 la Haye, Bowker, & Hermans, 2012).

To conclude, our findings provide compelling evidence that the social facilitation of eating leads to an uncompensated increase in intake that is sustained over several days. Future research should establish the extent to which social eating contributes to weight gain and to develop strategies to help people manage social eating situations to allow them to reach their health goals.

462

#### 463 Acknowledgments

- 464 myfood24 was developed through Medical Research Council funding, grant G110235.
- 465 myfood24 is now being supported by spinout company Dietary Assessment Ltd. Requests to

466 use myfood24 should be made to <u>enquiries@myfood24.org</u>.

467

468 We thank following students for assistance with the data collection: Elaina Dudley, Keira

469 Perks, Milene Fernandes, Sarah Miller, Harriet Abbot, Arielle Kaufman (Study 1) and

470 Xuebing Huang, Jinyu Liu, Ilayda Yondem, Macarena Perales, Georgina Hobden, Emma

471 Williams, Cerys Cherry (Study 2).

472

- 473 Conflict of Interest and Funding Disclosure
- 474 Helen K. Ruddock "no conflicts of interest."
- 475 Jeff M. Brunstrom "no conflicts of interest."

476 Lenny R. Vartanian "no conflicts of interest."

477 Suzanne Higgs "no conflicts of interest."

478

#### 479 Author Contributions

- 480 SH, JB, LV & HR designed the research. Testing and data collection were performed by HR.
- 481 HR performed the data analyses. SH, JB, LV & HR wrote the paper. All authors approved the

- 482 final version of the manuscript for submission. SH had primary responsibility for final
- 483 content.
- 484
- 485 **Data Sharing.**
- 486 Data described in the manuscript will be made publicly and freely available without
- 487 restriction at <u>https://reshare.ukdataservice.ac.uk/</u>.

489

#### References

- 491 Brindal, E., Wilson, C., Mohr, P., & Wittert, G. (2015). Eating in groups: Do multiple social
- 492 influences affect intake in a fast-food restaurant? *Journal of Health Psychology*, 20(5),
- 493 483–489. https://doi.org/10.1177/1359105315576607
- 494 Carter, M. C., Albar, S. A., Morris, M. A., Mulla, U. Z., Hancock, N., Evans, C. E., ...
- Consortium, myfood24. (2015). Development of a UK Online 24-h Dietary Assessment
  Tool: myfood24. *Nutrients*, 7(6), 4016–4032. https://doi.org/10.3390/nu7064016
- 497 Casanova, N., Finlayson, G., Blundell, J. E., & Hopkins, M. (2019). Biopsychology of human
- 498 appetite understanding the excitatory and inhibitory mechanisms of homeostatic
- 499 control. *Current Opinion in Physiology*, *12*, 33–38.
- 500 https://doi.org/https://doi.org/10.1016/j.cophys.2019.06.007
- 501 De Castro, J. M., Brewer, E. M., Elmore, D. K., & Orozco, S. (1990). Social facilitation of
- the spontaneous meal size of humans occurs regardless of time, place, alcohol or snacks.
  Appetite, 15(2), 89-101.
- 504 De Castro, J. M., & Brewer, E. M. (1992). The amount eaten in meals by humans is a power
- 505 function of the number of people present. Physiology & behavior, 51(1), 121-125.
- 506 Dunbar, R. I. M. (2017). Breaking Bread: the Functions of Social Eating. *Adaptive Human*
- 507 *Behavior and Physiology*, *3*(3), 198–211. https://doi.org/10.1007/s40750-017-0061-4
- 508 Faul, F., Erdfelder, E., Lang, A. G., & Buchner, A. (2007). G\*Power 3: A flexible statistical
- 509 power analysis program for the social, behavioral, and biomedical sciences. *Behavior*
- 510 *Research Methods*, *39*, 175–191.
- 511 Forkman, B. (1991). Social facilitation is shown by gerbils when presented with novel but not
- 512 with familiar food. *Animal Behaviour*, 42(5), 860–861.
- 513 https://doi.org/https://doi.org/10.1016/S0003-3472(05)80132-0
- 514 Hall, K. D., Sacks, G., Chandramohan, D., Chow, C. C., Wang, Y. C., Gortmaker, S. L., &
- 515 Swinburn, B. A. (2011). Quantification of the effect of energy imbalance on

- 516 bodyweight. *The Lancet*, *378*(9793), 826–837.
- 517 Harlow, H. F., & Yudin, H. C. (1933). Social behavior of primates. I. Social facilitation of
- 518 feeding in the monkey and its relation to attitudes of ascendance and submission.
- 519 *Journal of Comparative Psychology*, *16*(2), 171–185. https://doi.org/10.1037/h0071690
- 520 Hoaglin, D. C., & Iglewicz, B. (1987). Fine-Tuning Some Resistant Rules for Outlier
- 521 Labeling. *Journal of the American Statistical Association*, 82(400), 1147–1149.
- 522 https://doi.org/10.2307/2289392
- 523 IBM SPSS Statistics for Windows. (2020). Armonk, NY: IBM Corp.
- 524 Karlsson, J., Persson, L. O., Sjöström, L., & Sullivan, M. (2000). Psychometric properties
- and factor structure of the Three-Factor Eating Questionnaire (TFEQ) in obese men and
- 526 women. Results from the Swedish Obese Subjects (SOS) study. *International Journal of*

527 *Obesity*, 24(12), 1715–1725. https://doi.org/10.1038/sj.ijo.0801442

528 Levitsky, D. A. (2005). The non-regulation of food intake in humans: Hope for reversing the

529 epidemic of obesity. *Physiology and Behavior*, *86*(5), 623–632.

- 530 https://doi.org/10.1016/j.physbeh.2005.08.053
- 531 Levitsky, D. A., Sewall, A., Zhong, Y., Barre, L., Shoen, S., Agaronnik, N., ... Pacanowski,
- 532 C. (2019). Quantifying the imprecision of energy intake of humans to compensate for
- 533 imposed energetic errors: A challenge to the physiological control of human food intake.
- 534 *Appetite*, *133*, 337–343. https://doi.org/10.1016/j.appet.2018.11.017
- 535 McCrory, M. A., Fuss, P. J., McCallum, J. E., Yao, M., Vinken, A. G., Hays, N. P., &
- 536 Roberts, S. B. (1999). Dietary variety within food groups: association with energy intake
- and body fatness in men and women. *The American Journal of Clinical Nutrition*, 69(3),
- 538 440–447. https://doi.org/10.1093/ajcn/69.3.440
- 539 MyFood24. (2016). Leeds, UK: University of Leeds.
- 540 Rajecki, D. W., Kidd, R. F., Wilder, D. A., & Jaeger, J. (1975). Social factors in the
- 541 facilitation of feeding in chickens: Effects of imitation, arousal, or disinhibition? *Journal*

- 542 *of Personality and Social Psychology*, *32*(3), 510–518.
- 543 https://doi.org/10.1037/h0077065
- 544 Rolls, B. J., Morris, E. L., & Roe, L. S. (2002). Portion size of food affects energy intake in
- 545 normal-weight and overweight men and women. *The American Journal of Clinical*
- 546 *Nutrition*, 76(6), 1207–1213. https://doi.org/10.1093/ajcn/76.6.1207
- 547 Rolls, B. J., Roe, L. S., & Meengs, J. S. (2006). Larger portion sizes lead to a sustained
- 548 increase in energy intake over 2 days. *Journal of the American Dietetic Association*,
- 549 106(4), 543–549. https://doi.org/10.1016/j.jada.2006.01.014
- 550 Rolls, B. J., Roe, L. S., & Meengs, J. S. (2007). The effect of large portion sizes on energy
- 551 intake is sustained for 11 days. *Obesity*, *15*(6), 1535–1543.
- 552 https://doi.org/10.1038/oby.2007.182
- Ruddock, H.K, Brunstrom, J. M., Vartanian, L. R., & Higgs, S. (2019). A systematic review
  and meta-analysis of the social facilitation of eating. *The American Journal of Clinical*

555 *Nutrition*, *110*(4), 842–861. https://doi.org/10.1093/ajcn/nqz155

- 556 Ruddock, H.K., Long, E. V., Brunstrom, J. M., Vartanian, L. R., & Higgs, S. (2021). People
- serve themselves larger portions before a social meal. *Scientific Reports*, *11*, 11072.
- 558 Salvy, S. J., de la Haye, K., Bowker, J. C., & Hermans, R. C. J. (2012). Influence of peers
- and friends on children's and adolescents' eating and activity behaviors. *Physiology and Behavior*, *106*, 369–378.
- 561 Stunkard, A. J., & Messick, S. (1985). The Three-Factor Eating Questionnaire to Measure
- 562 Dietary Restraint, Disinhibition and Hunger. *Journal of Psychosomatic Research*, 29(1),
  563 71–83.
- 564 Tolman, C. W. (1964). Social facilitation of feeding behaviour in the domestic chick. *Animal*
- 565 Behaviour, 12(2), 245–251. https://doi.org/https://doi.org/10.1016/0003-3472(64)90008-
- 566 9
- 567 Vartanian, L. R. (2015). Impression management and food intake. Current directions in

- 569 Wark, P. A., Hardie, L. J., Frost, G. S., Alwan, N. A., Carter, M., Elliott, P., ... Cade, J. E.
- 570 (2018). Validity of an online 24-h recall tool (myfood24) for dietary assessment in
- 571 population studies: Comparison with biomarkers and standard interviews. *BMC*
- 572 *Medicine*, 16(1), 1–14. https://doi.org/10.1186/s12916-018-1113-8
- 573
- 574

### **Figure legends**

<ul> <li>represent the standard error of the mean.</li> <li><i>Figure 2</i>. Mean calories consumed as a function of condition and day. Error bars represent the standard error of the mean.</li> </ul>	575	Figure 1. Mean calories consumed as a function of meal, day, and condition. Error bars
<ul> <li>577</li> <li>578 <i>Figure 2.</i> Mean calories consumed as a function of condition and day. Error bars repres</li> <li>579 the standard error of the mean.</li> </ul>	576	represent the standard error of the mean.
<ul> <li><i>Figure 2.</i> Mean calories consumed as a function of condition and day. Error bars repres</li> <li>the standard error of the mean.</li> </ul>	577	
579 the standard error of the mean.	578	Figure 2. Mean calories consumed as a function of condition and day. Error bars represent
	579	the standard error of the mean.

	Menu 1 (total kcal= 3589)	Menu 2 (total kcal = 3699)	Menu 3 (total kcal = 3843)
Breakfast	<ul> <li>2 x wholemeal toast with 40 g hazelnut chocolate spread (433 kcal)</li> <li>150 g strawberry yogurt (123 kcal)</li> <li>207 g canned fruit with juice (101 kcal)</li> <li>150 g orange juice (70 kcal)</li> </ul>	<ul> <li>Bagel with 60 g soft cheese spread (425 kcal)</li> <li>150 g strawberry yogurt (123 kcal)</li> <li>207 g canned fruit with juice (101 kcal)</li> <li>150 g orange juice (70 kcal)</li> </ul>	<ul> <li>80 g granola (353 kcal)</li> <li>200 g semi-skimmed milk (100 kcal)</li> <li>150 g strawberry yogurt (123 kcal)</li> <li>207 g canned fruit with juice (101 kcal)</li> <li>150 g orange juice (70 kcal)</li> </ul>
	kcal = 727	kcal = 719	kcal = 747
Lunch	<ul> <li>200 g cheese &amp; onion quiche (521 kcal)</li> <li>150 g new potatoes (114 kcal)</li> <li>35 g green salad (8 kcal)</li> <li>75 g brownie bites (291 kcal)</li> <li>50 g salted crisps (272 kcal)<sup>1</sup></li> </ul>	<ul> <li>2 x bean burgers (458 kcal)</li> <li>White bread roll with 10 g margarine (247 kcal)</li> <li>60 g millionaire bites (300 kcal)</li> <li>70 g cheese tortilla chips (349 kcal)</li> </ul>	<ul> <li>Cheese sandwich comprising 3 pieces of wholemeal bread, 20 g margarine, 60 g cheddar cheese (742 kcal)</li> <li>70 g flapjack bites (313 kcal)<sup>2</sup></li> <li>50 g salt &amp; pepper crisps (311 kcal)<sup>1</sup></li> </ul>
	kcal = 1206	kcal = 1354	kcal = 1366
Dinner	<ul> <li>100 g (uncooked weight) pasta mixed with 250 g tomato pasta sauce, 30 g cheddar cheese (580 kcal)</li> <li>200 g tiramisu (500 kcal)</li> <li>110 g milk chocolate buttons (576 kcal)</li> </ul>	<ul> <li>300 g cheese &amp; tomato pizza (767 kcal)</li> <li>35 g salad (8 kcal)</li> <li>200 g chocolate dessert (270 kcal)</li> <li>110 g milk chocolate pieces (581 kcal)</li> </ul>	<ul> <li>450 g vegetarian lasagne (408 kcal)</li> <li>200 g (frozen weight) chips (358 kcal)</li> <li>150 g strawberry cheesecake (416 kcal)</li> <li>110 g milk chocolate (548 kcal)</li> </ul>
	kcal = 1656	kcal = 1626	kcal = 1730

ıps;

	Mean (SD)
Age (years)	20.8(2.8)
BMI $(kg/m^2)$	23.0(2.9)
TFEQ-restraint	14.1(4.0)
TFEQ-uncontrolled	22.9(4.7)
TFEQ-Emotional	7.2(2.4)
Friendship duration	21.1(29.4)
(months)	
Friendship 'How well'*	8.1(1.7)
Friendship 'How close'†	8.0(1.8)
* On a scale of $1-10$ (with anchor poin	ts 'Not very well' and 'Very well' respectiv

586 Table 2. Participant characteristics in Study 1

587 \* On a scale of 1-10 (with anchor points 'Not very well' and 'Very well', respectively), how well do you think you know your friend?

588 †On a scale of 1-10 (with anchor points 'Not very close' and 'Very close', respectively), how close do you feel you are with your friend?

589

## 

# 593 Table 3. Foods provided during the buffet lunch (per participant) in Study 2.

	Portion size (g)	kcal
Tesco cheese & onion quiche	200	524
Tesco salted crisps <sup>1</sup>	25	136
Cadburys dairy milk chocolate buttons	60	321
Tesco stuffed crust cheese pizza	215	550
Brownies	50	192
Flapjacks <sup>2</sup>	50	224

594 UK to US translation: <sup>1</sup>Chips; <sup>2</sup>Oat bars

Alone condition (n=34)	Social condition (n=33)	Univariate test statistic
Mean (SD)	Mean (SD)	
19.4(1.1)	19.4(1.1)	F(1,64)=0.01, p=.911
22.4(2.9)	21.7(2.5)	F(1,64)=1.04, p=.311
12.7(3.9)	13.5(2.8)	F(1,64)=0.88, p=.351
22.4(4.1)	22.2(3.2)	F(1,64)=0.07, p=.788
7.9(1.7)	7.9(2.2)	F(1,64)=0.00, p=.999
7.2(1.6)	7.5(2.1)	F(1,64)=0.25, p=.620
	Alone condition (n=34) Mean (SD) 19.4(1.1) 22.4(2.9) 12.7(3.9) 22.4(4.1) 7.9(1.7) 7.2(1.6)	Alone condition (n=34)       Social condition (n=33)         Mean (SD)       Mean (SD)         19.4(1.1)       19.4(1.1)         22.4(2.9)       21.7(2.5)         12.7(3.9)       13.5(2.8)         22.4(4.1)       22.2(3.2)         7.9(1.7)       7.9(2.2)         7.2(1.6)       7.5(2.1)

Table 4. Participant characteristics in Study 2.

601 \* On a scale of 1-10 (with anchor points 'Not very well' and 'Very well', respectively), how well do you think you know your friend?