

# Hardy uncertainty principle and unique continuation properties of covariant Schrödinger flows

Antonio Barcelo, Juan; Fanelli, Luca; Gutierrez, Susana; Ruiz, Alberto; Cruz Vilela, Mari

DOI:

[10.1016/j.jfa.2013.02.017](https://doi.org/10.1016/j.jfa.2013.02.017)

License:

None: All rights reserved

*Document Version*

Peer reviewed version

*Citation for published version (Harvard):*

Antonio Barcelo, J, Fanelli, L, Gutierrez, S, Ruiz, A & Cruz Vilela, M 2013, 'Hardy uncertainty principle and unique continuation properties of covariant Schrödinger flows', *Journal of Functional Analysis*, vol. 264, no. 10, pp. 2386-2415. <https://doi.org/10.1016/j.jfa.2013.02.017>

[Link to publication on Research at Birmingham portal](#)

## **Publisher Rights Statement:**

NOTICE: this is the author's version of a work that was accepted for publication in the above journal. Changes resulting from the publishing process, such as peer review, editing, corrections, structural formatting, and other quality control mechanisms may not be reflected in this document. Changes may have been made to this work since it was submitted for publication. A definitive version was subsequently published in *Journal of Functional Analysis*, [VOL 264, ISSUE 10, 2013. DOI: 10.1016/j.jfa.2013.02.017]

## **General rights**

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

## **Take down policy**

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact [UBIRA@lists.bham.ac.uk](mailto:UBIRA@lists.bham.ac.uk) providing details and we will remove access to the work immediately and investigate.

# HARDY UNCERTAINTY PRINCIPLE AND UNIQUE CONTINUATION PROPERTIES OF COVARIANT SCHRÖDINGER FLOWS

J.A. BARCELÓ, L. FANELLI, S. GUTIÉRREZ, A. RUIZ, AND M.C. VILELA

ABSTRACT. We prove a logarithmic convexity result for exponentially weighted  $L^2$ -norms of solutions to electromagnetic Schrödinger equation, without needing to assume smallness of the magnetic potential. As a consequence, we can prove a unique continuation result in the style of the Hardy uncertainty principle, which generalize the analogous theorems which have been recently proved by Escauriaza, Kenig, Ponce and Vega.

## 1. INTRODUCTION

This paper is concerned with sharp decay profiles, at two distinct times, of  $L^2$ -solutions to an electromagnetic Schrödinger equation of the type

$$(1.1) \quad \partial_t u = i(\Delta_A + V)u,$$

where  $u = u(x, t) : \mathbb{R}^{n+1} \rightarrow \mathbb{C}$ ,  $A = A(x) : \mathbb{R}^n \rightarrow \mathbb{R}^n$ ,  $V = V(x, t) : \mathbb{R}^n \rightarrow \mathbb{C}$ , and we use the notations

$$\nabla_A := \nabla - iA, \quad \Delta_A := (\nabla - iA)^2.$$

Our main goal is to start with a project devoted to understand sufficient conditions on solutions to (1.1), the coefficients  $A, V$ , and the behavior of the solutions at two different times which ensure the rigidity  $u \equiv 0$ . This follows a program which has been developed for the magnetic free case  $A \equiv 0$  by Escauriaza, Kenig, Ponce and Vega in the last few years in the sequel of papers [4, 5, 6, 7, 8], and more recently with Cowling in [2]. Their main motivations is the connection between unique continuation properties of Schrödinger evolutions and the so called *Hardy uncertainty principle* (see e.g. [17]), which can be briefly stated as follows:

*If  $f(x) = O(e^{-|x|^2/\beta^2})$  and its Fourier transform  $\hat{f}(\xi) = O(e^{-4|\xi|^2/\alpha^2})$ , then*

$$\alpha\beta < 4 \Rightarrow f \equiv 0$$

$$\alpha\beta = 4 \Rightarrow f \text{ is a constant multiple of } e^{-\frac{|x|^2}{\beta^2}}.$$

---

*Date:* March 19, 2013.

*2000 Mathematics Subject Classification.* 35J10, 35L05.

*Key words and phrases.* Schrödinger equation, electromagnetic potentials, Carleman estimates, uncertainty principle.

The first and fourth authors were supported by the Spanish projects MTM2008-02568 and MTM2011-28198, the third by the Spanish projects MTM2007-62186 and MTM2011- 24054, and finally, the fifth by Spanish projects MTM2007-62186 and MTM2011-28198.

Due to the strict connection between the Fourier transform  $\mathcal{F}$  and the solution to the free Schrödinger equation with initial datum  $f$  in  $L^2$ , namely

$$u(x, t) := e^{it\Delta} f(x) = (2\pi it)^{-\frac{n}{2}} e^{i\frac{|x|^2}{4t}} \mathcal{F} \left( e^{i\frac{|\cdot|^2}{4t}} f \right) \left( \frac{x}{2t} \right),$$

the Hardy uncertainty principle has a PDE's-counterpart, which can be stated as follows:

If  $u(x, 0) = O\left(e^{-|x|^2/\beta^2}\right)$  and  $u(x, T) := e^{iT\Delta} u(x, 0) = O\left(e^{-|x|^2/\alpha^2}\right)$ , then

$$\alpha\beta < 4T \Rightarrow u \equiv 0$$

$$\alpha\beta = 4T \Rightarrow u(x, 0) \text{ is a constant multiple of } e^{-\left(\frac{1}{\beta^2} + \frac{i}{4T}\right)|x|^2}.$$

The corresponding  $L^2$ -versions of the previous results were proved in [16] and affirm the following:

$$\begin{aligned} e^{|x|^2/\beta^2} f \in L^2, \quad e^{4|\xi|^2/\alpha^2} \hat{f} \in L^2, \quad \alpha\beta \leq 4 \Rightarrow f \equiv 0 \\ e^{|x|^2/\beta^2} u(x, 0) \in L^2, \quad e^{|x|^2/\alpha^2} e^{iT\Delta} u(x, 0) \in L^2, \quad \alpha\beta \leq 4T \Rightarrow u \equiv 0. \end{aligned}$$

Obviously, without loss of generality, we might restrict our attention to the case  $T = 1$ . An interesting survey about this topic can be found in [1].

One of the major contributions of the authors of [2, 4, 5, 6, 7, 8] was to deeply understand the relation between these kind of properties and the logarithmic convexity property of exponentially weighted  $L^2$ -norms of solutions to Schrödinger equations (see also [3], [9] for analogous results concerning unique continuation from the infinity). This permits to perform purely real analytical proofs, and then allows rough coefficients in the differential equations, which are difficult to be handled by Fourier transform or general complex analysis tools. For example, in [6] and [7] the authors considered any bounded potential of the form  $V = V_1(x) + V_2(x, t)$ , possibly being  $V_2$  complex-valued, without assuming any Sobolev regularity and any smallness condition on the two components; in this situation, they were able to establish the analog to the above statements, in the cases  $\alpha\beta < 2$  first ([6]), and the sharp  $\alpha\beta < 4$  later ([7]), for  $T = 1$ . The strategy can be roughly summarized as follows:

- Assume  $e^{|x|^2/\beta^2} u(x, 0)$ ,  $e^{|x|^2/\alpha^2} e^{i\Delta} u(x, 1) \in L^2$  and prove a logarithmic convexity estimate for the quantity  $H(t) := \|e^{g(x,t)} u(t)\|_{L^2}$  of the type  $H(t) \lesssim H(0)^t H(1)^{1-t}$ , where  $g$  is a suitable function, bounded with respect to  $t$  and quadratically growing with respect to  $x$ . This shows that a gaussian decay at times 0 and 1 is preserved (and in fact improved) for intermediate times.
- Start a self-improvement argument, by suitably moving the center of the gaussian as  $e^{a(t)|x+b(t)|^2}$ , based on analytical estimates (Carleman estimates; this leads up to the non-sharp result  $\alpha\beta < 2$ , see [6]) or on the logarithmic convexity itself (this leads to the sharp result  $\alpha\beta < 4$ , see [7]), which finally gives the rigidity  $u \equiv 0$ .

Amazingly, proving these results in a rigorous way represents not just a considerable technical difficulty, but also a conceptual obstacle which, if avoided, could bring to misleading results. To overcome this problem, the above mentioned authors introduce a small artificial dissipation term in the equation which turns out to be fundamental, and finally let it tend to zero.

It is quite natural to claim that, with some efforts, some small first-order terms can be introduced in the argument by Escauriaza, Kenig, Ponce and Vega without losing the results. The aim of this paper is to understand in which way the uncertainty can be described in the presence of first-order perturbations in covariant form, as in (1.1) when  $A \neq 0$ . Precisely, our goal here is to obtain similar results without assuming any smallness conditions on  $A$  and possibly respecting the mathematical properties of the quantities which are behind these kinds of models.

We continue to introduce the terminology and notation required to state the main results of this paper.

Let  $A = (A^1(x), \dots, A^n(x)) : \mathbb{R}^n \rightarrow \mathbb{R}^n$ , with  $n \geq 2$ , be a vector field, which we will usually refer to as a magnetic potential, and denote the magnetic field by  $B(x) = DA(x) - DA^t(x)$ , the anti-symmetric gradient of  $A$ , namely

$$B \in \mathcal{M}_{n \times n}(\mathbb{R}), \quad B_{jk}(x) := A_j^k(x) - A_k^j(x).$$

From now on, given a scalar function  $f$ , we always use the notation  $f_k(x) = \partial_{x_k} f(x)$ , while an upper index will denote the component of a vector. In dimension  $n = 2$ ,  $B$  is identified with the scalar function  $B \cong \text{curl } A = A_2^1 - A_1^2$ ; the same identification holds in dimension  $n = 3$ , where now  $\text{curl } A$  is a vector field and

$$v^t B = v \times \text{curl } A \quad \forall v \in \mathbb{R}^3,$$

the cross denoting the vectorial product in  $\mathbb{R}^3$ . Since equation (1.1) is gauge invariant (see Section 2.1 below), it is always important to keep in mind that the physically meaningful quantity is the magnetic field  $B$ , while the potential  $A$  is a mathematical construction. This fact has to be considered when we state a theorem, since a meaningful result should not depend on a particular choice of the gauge.

As it will be clear in the sequel, another relevant object is the vector-field  $\Psi(x) := x^t B(x)$ ; in 3D, it can be interpreted, modulo its intensity, as a tangential projection of  $\text{curl } A$ , since

$$x^t B(x) = x \times \text{curl } A(x) = |x| B_\tau(x), \quad n = 3$$

following the notation  $B_\tau = \frac{x^t}{|x|} B$  introduced in [10]. As we see in (1.5) below,  $x^t B$  is essentially the only component of  $B$  on which one needs to make suitable assumptions, in order to obtain a Hardy uncertainty principle.

We can now state the main result of this paper.

**Theorem 1.1.** *Let  $n \geq 3$ , and let  $u \in \mathcal{C}([0, 1]; L^2(\mathbb{R}^n))$  be a solution to*

$$(1.2) \quad \partial_t u = i(\Delta_A + V_1(x) + V_2(x, t))u$$

*in  $\mathbb{R}^n \times [0, 1]$ , with  $A = A(x) : \mathbb{R}^n \rightarrow \mathbb{R}^n$ ,  $V_1 = V_1(x) : \mathbb{R}^n \rightarrow \mathbb{R}$ ,  $V_2 = V_2(x, t) : \mathbb{R}^{n+1} \rightarrow \mathbb{C}$ . Assume that*

$$(1.3) \quad \int_0^1 A(sx) ds \in \mathbb{R}^n$$

*is well defined at almost every  $x \in \mathbb{R}^n$ . Moreover, denote by  $B = B(x) = DA - DA^t$ ,  $B_{jk} = A_j^k - A_k^j$  and assume that there exists a unit vector  $v \in \mathcal{S}^{n-1}$  such that*

$$(1.4) \quad v^t B(x) \equiv 0.$$

In addition, assume that

$$(1.5) \quad \|x^t B\|_{L^\infty}^2 := \frac{M_A}{4} < \infty$$

$$(1.6) \quad \|V_1\|_{L^\infty} := M_1 < \infty$$

$$(1.7) \quad \sup_{t \in [0,1]} \left\| e^{\frac{|\cdot|^2}{(\alpha t + \beta(1-t))^2}} V_2(\cdot, t) \right\|_{L^\infty} e^{\sup_{t \in [0,1]} \|\Im V_2(\cdot, t)\|_{L^\infty}} := M_2 < \infty$$

$$(1.8) \quad \left\| e^{\frac{|\cdot|^2}{\beta^2}} u(\cdot, 0) \right\|_{L^2} + \left\| e^{\frac{|\cdot|^2}{\alpha^2}} u(\cdot, 1) \right\|_{L^2} < \infty,$$

for some  $\alpha, \beta > 0$ . Then,  $\alpha\beta < 2$  implies  $u \equiv 0$ .

**Remark 1.2.** Among various consequences, conditions (1.3), (1.5) and (1.6) imply the self-adjointness in  $L^2$  of the hamiltonian  $H_A := -\Delta_A + V_1$ , with form-domain  $H^1(\mathbb{R}^n)$ , after a suitable reduction to the so called Crönstrom (or transversal) gauge (see Section 2.1 and Proposition 2.6 below). Hence the Schrödinger flow  $e^{itH_A}$  is well defined for any  $t \in \mathbb{R}$  by the Spectral Theorem, and unitary in  $L^2$ , so that given  $u(x, 0) \in L^2$  there exists a unique solution  $u \in \mathcal{C}([0, 1]; L^2(\mathbb{R}^n))$  of the integral equation

$$u(\cdot, t) = e^{itH_A} u(\cdot, 0) + \int_0^t e^{i(t-s)H_A} V_2(\cdot, s) u(\cdot, s) ds,$$

provided (1.7).

In addition, also the heat flow  $e^{tH_A}$  is well defined for positive times, and this will be used in the sequel.

**Remark 1.3.** Notice that no smallness conditions on  $A, V_1, V_2$  are required in the statement of Theorem 1.1. On the other hand, condition (1.4) naturally comes into play once we prove a Carleman estimate (Lemma 4.1 below), which is one of the tools to prove Theorem 1.1. We remark that we cannot prove the result in dimension  $n = 2$ , which remains as an open question, since there are no  $2 \times 2$ -antisymmetric matrices with non-trivial kernel.

The clearest examples of fields  $B$  satisfying our assumptions can be constructed as follows. Denote by

$$M_{2k-1} = \begin{pmatrix} J & & & \\ & \ddots & & \\ & & J & \\ & & 0 & 0 \end{pmatrix}, \quad M_{2k} = \begin{pmatrix} J & & & \\ & \ddots & & \\ & & J & \\ & & 0 & 0 \\ & & 0 & 0 \end{pmatrix},$$

with  $J := \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$  and  $k \geq 2$ . Now define for  $n \geq 3$

$$B(x) = \frac{x^t}{|x|^2} M_n$$

and notice that the assumptions of Theorem 1.1 are satisfied. In particular, in dimension  $n = 3$  we can identify the above  $B$  with

$$B(x) = |x|^{-2}(-x_2, x_1, 0) = \text{curl} \left( \frac{x_1 x_3}{|x|^2}, \frac{x_2 x_3}{|x|^2}, \frac{-x_1^2 - x_2^2}{|x|^2} \right).$$

This is, as far as we understand, also a quite interesting hint about the fact that Theorem 1.1 is presumably not true, in total generality, for any magnetic field  $B$ , and it will be matter of future work.

**Remark 1.4.** The constraint  $\alpha\beta < 2$  in Theorem 1.1 is far from the sharp  $\alpha\beta < 4$  obtained in [7] in the magnetic-free case  $A \equiv 0$ . Actually, we use here the argument introduced in [6], involving the use of a Carleman estimate, which cannot lead to a better result. In addition, already at this level, we see the necessity of the condition (1.4), which is a quite interesting fact. Presumably, when looking for the sharp result, some additional phenomena, involving the presence a non trivial magnetic field, should come into play. This will hopefully be matter of future work.

Theorem 1.1 has several consequences regarding uniqueness of solutions to (1.1), both in the linear and nonlinear cases (i.e.  $V_2(x, t) = |u(x, t)|^p$ ), which we will not investigate in this paper (see [6, 7] for details).

The main tool to prove theorem 1.1 is the following logarithmic convexity result.

**Theorem 1.5** (logarithmic convexity). *Let  $n \geq 2$ , and consider a solution  $u \in \mathcal{C}([0, 1]; L^2(\mathbb{R}^n))$  to*

$$(1.9) \quad \partial_t u = i(\Delta_A + V_1(x) + V_2(x, t))u$$

in  $\mathbb{R}^n \times [0, 1]$ , with  $A = A(x) : \mathbb{R}^n \rightarrow \mathbb{R}^n$ ,  $V_1 = V_1(x) : \mathbb{R}^n \rightarrow \mathbb{R}$ ,  $V_2 = V_2(x, t) : \mathbb{R}^{n+1} \rightarrow \mathbb{C}$ . Denote by  $B = B(x) = DA - DA^t$ ,  $B_{jk} = A_j^k - A_k^j$  and assume that

$$(1.10) \quad \int_0^1 A(sx) ds \in \mathbb{R}^n$$

is well defined at almost every  $x \in \mathbb{R}^n$ . Moreover, assume that

$$(1.11) \quad \|x^t B\|_{L^\infty}^2 := \frac{M_A}{4} < \infty$$

$$(1.12) \quad \|V_1\|_{L^\infty} := M_1 < \infty$$

$$(1.13) \quad \sup_{t \in [0, 1]} \left\| e^{\frac{|\cdot|^2}{(\alpha t + \beta(1-t))^2}} V_2(\cdot, t) \right\|_{L^\infty} e^{\sup_{t \in [0, 1]} \|\Im V_2(\cdot, t)\|_{L^\infty}} := M_2 < \infty$$

$$(1.14) \quad \left\| e^{\frac{|\cdot|^2}{\beta^2}} u(\cdot, 0) \right\|_{L^2} + \left\| e^{\frac{|\cdot|^2}{\alpha^2}} u(\cdot, 1) \right\|_{L^2} < \infty,$$

for some  $\alpha, \beta > 0$ . Then, the function

$$\Theta(t) := \log \left\| e^{\frac{|\cdot|^2}{(\alpha t + \beta(1-t))^2}} u(\cdot, t) \right\|_{L^2}^{\alpha t + \beta(1-t)}$$

is convex in  $[0, 1]$ . In addition, there exists a constant  $N = N(\alpha, \beta) > 0$  such that

$$(1.15) \quad \left\| e^{\frac{|\cdot|^2}{(\alpha t + \beta(1-t))^2}} u(\cdot, t) \right\|_{L^2} \leq e^{N[M_A + M_1 + M_2 + M_1^2 + M_2^2]} \left\| e^{\frac{|\cdot|^2}{\beta^2}} u(\cdot, 0) \right\|_{L^2}^{\frac{\beta(1-t)}{\alpha t + \beta(1-t)}} \left\| e^{\frac{|\cdot|^2}{\alpha^2}} u(\cdot, 1) \right\|_{L^2}^{\frac{\alpha t}{\alpha t + \beta(1-t)}}$$

$$(1.16) \quad \left\| \sqrt{t(1-t)} e^{\frac{|\cdot|^2}{(\alpha t + \beta(1-t))^2}} \nabla_A u(x, t) \right\|_{L^2(\mathbb{R}^n \times [0, 1])} \leq e^{N[M_A + M_1 + M_2 + M_1^2 + M_2^2]} \left( \left\| e^{\frac{|\cdot|^2}{\beta^2}} u(\cdot, 0) \right\|_{L^2} + \left\| e^{\frac{|\cdot|^2}{\alpha^2}} u(\cdot, 1) \right\|_{L^2} \right).$$

**Remark 1.6.** Notice that in this case condition (1.4) is not needed; as a consequence, we can also handle the 2D case, which is included in the statement. We finally remark that both Theorems 1.1 and 1.5 hold in dimension  $n = 1$ , since in this case any reasonable magnetic potential can be gauged away by the Fundamental Theorem of Calculus.

The strategy of the proof of Theorem 1.5 is as follows:

- by gauge transformation, we reduce to the case in which  $x \cdot A \equiv 0$  (see section 2.1) below;
- we add a small dissipation term which regularizes the solution and gives a useful preservation property for the exponentially weighted  $L^2$ -norms of the solution (Lemma 2.10);
- by conformal (or Appell) transformation (see Lemma 2.7), we reduce to the case  $\alpha = \beta$ ;
- we prove Theorem 1.5 in the case  $\alpha = \beta$  (Lemmata 2.14, 2.16);
- we translate the result in terms of the original solution, by inverting the conformal transformation, obtaining the final result.

Once Theorem 1.5 is proved, then Theorem 1.1 follows as an application of a Carleman inequality (Lemma 4.1).

**Acknowledgements.** The authors wish to thank Luis Escauriaza and Luis Vega for some useful discussions about the topic of the paper, and Vladimir Georgiev for addressing them to the topic of Section 2.1 below.

## 2. PRELIMINARIES

We devote this section to collect some preliminary results which will be needed in the proofs of our main results.

In order to prove the main theorems in a rigorous way, we need to add a dissipation term to equation (1.1), which permits to assure that a gaussian decay at time 0 is preserved during the time evolution. For this reason, we study in this section some abstract properties regarding the solutions to

$$(2.1) \quad \partial_t u = (a + ib) (\Delta_A u + V(x, t)u + F(x, t)),$$

with  $a, b \in \mathbb{R}$ ,  $A = A(x, t) : \mathbb{R}^{n+1} \rightarrow \mathbb{R}^n$ ,  $V(x, t), F(x, t) : \mathbb{R}^{n+1} \rightarrow \mathbb{C}$ .

**2.1. The Cronström gauge.** Our first tool is the gauge invariance of equation (2.1). We need to review some algebraic properties of magnetic Schrödinger operators, pointing our attention on the so called Cronström (or transversal) gauge.

Equation (2.1) is *gauge invariant* in the following sense: if  $u$  solves (2.1), and we denote by  $\tilde{A} = A + \nabla\varphi$ , with  $\varphi = \varphi(x) : \mathbb{R}^n \rightarrow \mathbb{R}$ , then the function  $\tilde{u} = e^{i\varphi}u$  is a solution to

$$(2.2) \quad \partial_t \tilde{u} = (a + ib) (\Delta_{\tilde{A}} \tilde{u} + V(x, t)\tilde{u} + e^{i\varphi}F(x, t)).$$

Indeed, it is quite simple to verify that  $\Delta_{\tilde{A}}(e^{i\varphi}u) = e^{i\varphi}\Delta_A u$ .

**Definition 2.1.** A connection  $\nabla - iA(x)$  is said to be in the *Cronström gauge* (or *transversal gauge*) if  $A \cdot x = 0$ , for any  $x \in \mathbb{R}^n$ .

The following Lemma shows the transformation which permits to reduce a suitable potential to the Cronström gauge.

**Lemma 2.2.** *Let  $A = A(x) = (A^1(x), \dots, A^n(x)) : \mathbb{R}^n \rightarrow \mathbb{R}^n$ , for  $n \geq 2$  and denote by  $B = DA - DA^t \in \mathcal{M}_{n \times n}(\mathbb{R})$ ,  $B_{jk} = A_j^k - A_k^j$ , and  $\Psi(x) := x^t B(x) \in \mathbb{R}^n$ . Assume that the two vector quantities*

$$(2.3) \quad \int_0^1 A(sx) ds \in \mathbb{R}^n, \quad \int_0^1 \Psi(sx) ds \in \mathbb{R}^n$$

*are finite, for almost every  $x \in \mathbb{R}^n$ ; moreover, define the (scalar) function*

$$(2.4) \quad \varphi(x) := x \cdot \int_0^1 A(sx) ds \in \mathbb{R}.$$

*Then, the following two identities hold:*

$$(2.5) \quad \tilde{A}(x) := A(x) - \nabla \varphi(x) = - \int_0^1 \Psi(sx) ds$$

$$(2.6) \quad x^t D \tilde{A}(x) = -\Psi(x) + \int_0^1 \Psi(sx) ds.$$

*Proof.* A simple proof of identity (2.5) can be found e.g. in [12]. For the sake of completeness, we write it below. A direct computation shows that

$$\begin{aligned} \varphi_j(x) &= \frac{\partial}{\partial x_j} \varphi(x) = \int_0^1 A^j(sx) ds + \int_0^1 \sum_{k=1}^n s x_k A_j^k(sx) ds \\ &= \int_0^1 A^j(sx) ds + \int_0^1 \sum_{k=1}^n s x_k A_k^j(sx) ds + \int_0^1 \sum_{k=1}^n s x_k B_{jk}(sx) ds \\ &= \int_0^1 A^j(sx) ds + \int_0^1 s \frac{d}{ds} [A^j(sx)] ds + \int_0^1 \Psi^j(sx) ds. \end{aligned}$$

Integrating by parts now yields (2.5).

We now pass to the proof of (2.6). By (2.5), we can now compute

$$\begin{aligned} [D \tilde{A}(x)]_{kj} &= [D(A - \nabla \varphi)]_{kj}(x) \\ &= - \frac{\partial}{\partial x_k} \int_0^1 \sum_{h=1}^n s x_h (A_h^j(sx) - A_j^h(sx)) ds \\ &= - \int_0^1 s (A_k^j(sx) - A_j^k(sx)) ds - \sum_{h=1}^n \int_0^1 s^2 x_h (A_h^j - A_j^h)_k(sx) ds. \end{aligned}$$

Integrating by parts we obtain

$$\begin{aligned} [x^t D \tilde{A}(x)]^j &= \sum_{k=1}^n x_k [D(A - \nabla \varphi)]_{kj}(x) \\ &= - \sum_{k=1}^n \int_0^1 s x_k (A_k^j(sx) - A_j^k(sx)) ds - \sum_{h=1}^n \sum_{k=1}^n \int_0^1 s^2 x_h x_k (A_h^j - A_j^h)_k(sx) ds \\ &= - \sum_{k=1}^n \int_0^1 s x_k (A_k^j(sx) - A_j^k(sx)) ds - \sum_{h=1}^n \int_0^1 s^2 x_h \frac{d}{ds} [A_h^j(sx) - A_j^h(sx)] ds \\ &= \sum_{k=1}^n \int_0^1 s x_k (A_k^j(sx) - A_j^k(sx)) ds - \sum_{k=1}^n x_k (A_k^j(x) - A_j^k(x)), \end{aligned}$$

which proves (2.6).  $\square$



**Corollary 2.3.** *Under the same assumptions of Lemma 2.2, we have:*

$$(2.7) \quad x \cdot \tilde{A}(x) \equiv 0, \quad x \cdot x^t D\tilde{A}(x) \equiv 0.$$

*Proof.* The proof is a quite immediate consequence of (2.5), (2.6), and the fact that  $B$  is an anti-symmetric matrix.  $\square$

**Remark 2.4.** Notice that conditions (1.10) and (1.11) in Theorem 1.5 obviously imply (2.3), hence Lemma 2.2 and Corollary 2.3 are applicable under the assumptions of our main Theorems.

**Example 2.5** (Aharonov-Bohm). The following is possibly the most relevant example of a 2D-magnetic potential for which Lemma 2.2 and Corollary 2.3 do not apply. Define the 2D-Aharonov-Bohm potential as

$$A(x) = |x|^{-2}(-x_2, x_1).$$

In dimension  $n = 2$ , the antisymmetric gradient  $B = DA - DA^t$  is identified with the scalar quantity  $B = \text{curl } A = A_2^1 - A_1^2$ . Writing

$$A(x) = \nabla^\perp \log(|x|),$$

where  $\nabla^\perp$  is the orthogonal gradient, chosen with the correct orientation, gives  $B = \text{curl } A = \Delta \log(|x|) = 2\pi\delta$ . This shows that  $\Psi(x) = x^t B(x) \equiv 0$ ; if formula (2.5) were true in this case, it would give that  $A \equiv 0$ , which is a contradiction. In fact, (2.3) does not hold in this case, since  $A$  is too singular.

In similar ways, it is possible to construct such examples of potentials  $A$ , in every dimension, satisfying  $x^t B = 0$  with  $A \neq 0$ , which are not in contradiction with identity (2.5) since they do not satisfy (2.3).

**2.2. Self-adjointness.** We now state a standard result about the self-adjointness of  $H_A = -\Delta_A - V_1$ .

**Proposition 2.6.** *Let  $A = A(x) = (A^1(x), \dots, A^n(x)) : \mathbb{R}^n \rightarrow \mathbb{R}^n$ ,  $V_1 = V_1(x) : \mathbb{R}^n \rightarrow \mathbb{R}$  and denote by  $B = DA - DA^t \in \mathcal{M}_{n \times n}(\mathbb{R})$ ,  $B_{jk} = A_j^k - A_k^j$ , and  $\Psi(x) := x^t B(x) \in \mathbb{R}^n$ , for  $n \geq 2$ . Assume that*

$$(2.8) \quad \int_0^1 A(sx) ds \in \mathbb{R}^n,$$

*is finite, for almost every  $x \in \mathbb{R}^n$ ; moreover, assume that*

$$(2.9) \quad V_1(x) \in L^\infty \quad x^t B(x) \in L^\infty,$$

*and define  $\tilde{A}$  by (2.5). Finally, consider the quadratic form*

$$\tilde{q}(\varphi, \psi) := \int \nabla_{\tilde{A}} \varphi \cdot \overline{\nabla_{\tilde{A}} \psi} dx + \int V_1 \varphi \overline{\psi} dx.$$

*Then  $\tilde{q}$  is the form associated to a unique self-adjoint operator  $H_{\tilde{A}} = -\Delta_{\tilde{A}} - V_1(x)$ , with form domain  $H^1(\mathbb{R}^n)$ .*

*Proof.* The proof is completely standard. Indeed, notice that both  $q$  and  $\tilde{q}$  are well defined on  $H^1$ , since  $V_1 \in L^\infty$  and  $\tilde{A} \in L^\infty$ , thanks to (2.9) and Lemma 2.2. Moreover, the norm

$$\|\psi\|^2 := \tilde{q}(\psi, \psi) + C\|\psi\|_{L^2}^2$$

is equivalent to the  $H^1$ -norm, for some  $C > 0$  sufficiently large, by the same reasons as above; this show that the form  $\tilde{q}$  is closed. Finally, the form is semibounded, i.e.

$$\tilde{q}(\psi, \psi) \geq -C \|\psi\|_{L^2}^2,$$

by the same arguments. In conclusion, the thesis follows from Theorem VIII.15 in [14].  $\square$

**2.3. The Appell transformation.** Following the strategy in [6], we now introduce a conformal transformation, usually referred to as the *Appell transformation*, as another tool for the proofs of our main results. As we see in the sequel, it permits to reduce matters in Theorem 1.5 to the situation in which  $u(0)$  and  $u(1)$  have the same gaussian decay, namely  $\alpha = \beta$ .

**Lemma 2.7.** *Let  $A = A(y, s) = (A^1(y, s), \dots, A^n(y, s)) : \mathbb{R}^{n+1} \rightarrow \mathbb{R}^n$ ,  $V = V(y, s), F = F(y, s) : \mathbb{R}^n \rightarrow \mathbb{C}$ ,  $u = u(y, s) : \mathbb{R}^n \times [0, 1] \rightarrow \mathbb{C}$  be a solution to*

$$(2.10) \quad \partial_s u = (a + ib) (\Delta_A u + V(y, s)u + F(y, s)),$$

with  $a + ib \neq 0$ , and define, for any  $\alpha, \beta > 0$ , the function

$$(2.11) \quad \tilde{u}(x, t) := \left( \frac{\sqrt{\alpha\beta}}{\alpha(1-t) + \beta t} \right)^{\frac{n}{2}} u \left( \frac{x\sqrt{\alpha\beta}}{\alpha(1-t) + \beta t}, \frac{t\beta}{\alpha(1-t) + \beta t} \right) e^{-\frac{(\alpha-\beta)|x|^2}{4(a+ib)(\alpha(1-t)+\beta t)}}.$$

Then  $\tilde{u}$  is a solution to

$$(2.12) \quad \partial_t \tilde{u} = (a + ib) \left( \Delta_{\tilde{A}} \tilde{u} + i \frac{(\alpha - \beta) \tilde{A} \cdot x}{(a + ib)(\alpha(1-t) + \beta t)} \tilde{u} + \tilde{V}(x, t) \tilde{u} + \tilde{F}(x, t) \right),$$

where

$$(2.13) \quad \tilde{A}(x, t) = \frac{\sqrt{\alpha\beta}}{\alpha(1-t) + \beta t} A \left( \frac{x\sqrt{\alpha\beta}}{\alpha(1-t) + \beta t}, \frac{t\beta}{\alpha(1-t) + \beta t} \right)$$

$$(2.14) \quad \tilde{V}(x, t) = \frac{\alpha\beta}{(\alpha(1-t) + \beta t)^2} V \left( \frac{x\sqrt{\alpha\beta}}{\alpha(1-t) + \beta t}, \frac{t\beta}{\alpha(1-t) + \beta t} \right)$$

$$(2.15) \quad \tilde{F}(x, t) = \left( \frac{\sqrt{\alpha\beta}}{\alpha(1-t) + \beta t} \right)^{\frac{n}{2}+2} F \left( \frac{x\sqrt{\alpha\beta}}{\alpha(1-t) + \beta t}, \frac{t\beta}{\alpha(1-t) + \beta t} \right) e^{-\frac{(\alpha-\beta)|x|^2}{4(a+ib)(\alpha(1-t)+\beta t)}}.$$

*Proof.* The proof is basically an explicit computation. Let us denote by

$$g(t) := \frac{\sqrt{\alpha\beta}}{\alpha(1-t) + \beta t}, \quad c := \sqrt{\frac{\beta}{\alpha}}, \quad y = xg(t), \quad s = ctg(t)$$

$$h(x, t) := \frac{(\alpha - \beta)|x|^2}{4(a + ib)(\alpha(1-t) + \beta t)} = \frac{(\alpha - \beta)c}{4(a + ib)\beta} g(t)|x|^2.$$

With these notations, we easily get

$$\tilde{u}(x, t) = g^{\frac{n}{2}} e^h u(y, s)$$

$$\partial_t \tilde{u} = g^{\frac{n}{2}} e^h \left[ g^2 \partial_s u + 2(a + ib)g \nabla_x h \cdot \nabla_y u + \frac{(\alpha - \beta)c}{\beta} g \left( \frac{n}{2} + h \right) u \right].$$

On the other hand, we have

$$\nabla_x \tilde{u} = g^{\frac{n}{2}} e^h (g \nabla_y u + u \nabla_x h),$$

and therefore

$$g^{\frac{n}{2}} e^h g \nabla_y u = \nabla_x \tilde{u} - \tilde{u} \nabla_x h.$$

Moreover,

$$\begin{aligned} \Delta_x \tilde{u} &= g^{\frac{n}{2}} e^h [g^2 \Delta_y u + 2g \nabla_x h \cdot \nabla_y u + (|\nabla_x h|^2 + \Delta_x h) u] \\ g^{\frac{n}{2}} e^h g^2 \Delta_y u &= \Delta_x \tilde{u} - 2 \nabla_x h \cdot \nabla_x \tilde{u} + |\nabla_x h|^2 \tilde{u} - \tilde{u} \Delta_x h. \end{aligned}$$

Now expand the operator  $\Delta_A = (\nabla - iA) \cdot (\nabla - iA)$ , in order to rewrite equation (2.10) as

(2.16)

$$\partial_s u = (a + ib) (\Delta_y u - i(\operatorname{div}_y A)u - 2iA \cdot \nabla_y u - |A|^2 u + V(y, s)u + F(y, s));$$

finally, since

$$\tilde{A} = gA, \quad \operatorname{div}_x \tilde{A} = g^2 \operatorname{div}_y A, \quad \tilde{V} = g^2 V, \quad \tilde{F} = g^{\frac{n}{2}+2} e^h F,$$

the thesis (2.12) follows from (2.16) and the above identities.  $\square$

**Corollary 2.8.** *With the same notations of Lemma 2.7, denoting by*

$$(2.17) \quad y =: \frac{\sqrt{\alpha\beta}x}{\alpha(1-t) + \beta t}, \quad s =: \frac{\beta t}{\alpha(1-t) + \beta t},$$

we have, for any  $\gamma \in \mathbb{R}$ ,

(2.18)

$$\left\| e^{\gamma|\cdot|^2} \tilde{u}(\cdot, t) \right\|_{L^2} = \left\| e^{\left[ \frac{\gamma\alpha\beta}{(\alpha s + \beta(1-s))^2} + \frac{(\alpha-\beta)a}{4(a^2+b^2)(\alpha s + \beta(1-s))} \right] |\cdot|^2} u(\cdot, s) \right\|_{L^2}$$

(2.19)

$$\left\| e^{\gamma|\cdot|^2} \tilde{F}(\cdot, t) \right\|_{L^2} = \frac{\alpha\beta}{(\alpha(1-t) + \beta t)^2} \left\| e^{\left[ \frac{\gamma\alpha\beta}{(\alpha s + \beta(1-s))^2} + \frac{(\alpha-\beta)a}{4(a^2+b^2)(\alpha s + \beta(1-s))} \right] |\cdot|^2} F(\cdot, s) \right\|_{L^2}$$

(2.20)

$$\begin{aligned} \left\| \sqrt{t(1-t)} e^{\gamma|x|^2} \nabla_{\tilde{A}} \tilde{u} \right\|_{L^2(\mathbb{R}^n \times [0,1])} &= \left\| \sqrt{s(1-s)} e^{\left[ \frac{\gamma\alpha\beta}{(\alpha s + \beta(1-s))^2} + \frac{(\alpha-\beta)a}{4(a^2+b^2)(\alpha s + \beta(1-s))} \right] |y|^2} \right. \\ &\quad \times \left. \left( \frac{\alpha s + \beta(1-s)}{\sqrt{\alpha\beta}} \nabla_A u + \frac{(\alpha-\beta)y}{2(a+ib)\sqrt{\alpha\beta}} u \right) \right\|_{L^2(\mathbb{R}^n \times [0,1])} \end{aligned}$$

(2.21)

$$\begin{aligned} &\left\| \sqrt{t(1-t)} e^{\gamma|x|^2} |x| \tilde{u} \right\|_{L^2(\mathbb{R}^n \times [0,1])} \\ &= \left\| \sqrt{s(1-s)} e^{\left[ \frac{\gamma\alpha\beta}{(\alpha s + \beta(1-s))^2} + \frac{(\alpha-\beta)a}{4(a^2+b^2)(\alpha s + \beta(1-s))} \right] |y|^2} \frac{|y| \sqrt{\alpha\beta}}{\alpha s + \beta(1-s)} u \right\|_{L^2(\mathbb{R}^n \times [0,1])}. \end{aligned}$$

We omit here the details of the proof of the previous Corollary, which are straightforward after Lemma 2.7.

**2.4. Logarithmic convexity.** We now pass to study, from an abstract point of view, the evolution of weighted solutions to (2.1) with gaussian weights.

**Lemma 2.9.** *Let  $u = u(x, t) : \mathbb{R}^{n+1} \rightarrow \mathbb{C}$  be a solution to (2.1) where  $a, b \in \mathbb{R}$ ,  $A = A(x, t) : \mathbb{R}^{n+1} \rightarrow \mathbb{R}^n$ ,  $V, F : \mathbb{R}^{n+1} \rightarrow \mathbb{C}$ , and denote by  $v := e^\varphi u$ , with  $\varphi = \varphi(x, t) : \mathbb{R}^{n+1} \rightarrow \mathbb{R}$ . Then  $v$  solves*

$$(2.22) \quad \partial_t v = (\mathcal{S} + \mathcal{A})v + (a + ib)(V(x, t)v + e^\varphi F(x, t)),$$

where

$$(2.23) \quad \mathcal{S} = a(\Delta_A + |\nabla_x \varphi|^2) - ib(\Delta_x \varphi + 2\nabla_x \varphi \cdot \nabla_A) + \varphi_t$$

$$(2.24) \quad \mathcal{A} = ib(\Delta_A + |\nabla_x \varphi|^2) - a(\Delta_x \varphi + 2\nabla_x \varphi \cdot \nabla_A).$$

In addition, the following identities hold:

$$(2.25) \quad \mathcal{S}_t = 2a(\Im A_t \cdot \nabla_A + \nabla_x \varphi \cdot \nabla_x \varphi_t) + 2b(\Im \nabla_x \varphi_t \cdot \nabla_A - \nabla_x \varphi \cdot A_t) + \varphi_{tt}$$

$$(2.26)$$

$$\begin{aligned} \int_{\mathbb{R}^n} [\mathcal{S}, \mathcal{A}] f \bar{f} dx &= (a^2 + b^2) \left( 4 \int_{\mathbb{R}^n} \nabla_A f \cdot D_x^2 \varphi \overline{\nabla_A f} dx - \int_{\mathbb{R}^n} |f|^2 \Delta_x^2 \varphi dx \right. \\ &\quad \left. + 4 \int_{\mathbb{R}^n} |f|^2 \nabla_x \varphi \cdot D_x^2 \varphi \nabla_x \varphi dx - 4 \Im \int_{\mathbb{R}^n} f (\nabla_x \varphi)^t B \cdot \overline{\nabla_A f} dx \right) \\ &\quad + 2b \Im \int_{\mathbb{R}^n} \bar{f} \nabla_x \varphi_t \cdot \nabla_A f dx + 2a \int_{\mathbb{R}^n} |f|^2 \nabla_x \varphi \cdot \nabla_x \varphi_t dx, \end{aligned}$$

where  $\mathcal{S}_t := (\partial_t \mathcal{S})$ ,  $(D_x^2 \varphi)_{jk} = \frac{\partial^2 \varphi}{\partial x_j \partial x_k}$ ,  $\Delta_x^2 \varphi := \Delta_x(\Delta_x \varphi)$ ,  $B = DA - DA^t$ ,  $B_{jk} = A_j^k - A_k^j$  and  $[\mathcal{S}, \mathcal{A}] = \mathcal{S}\mathcal{A} - \mathcal{A}\mathcal{S}$  denotes the commutator between  $\mathcal{S}$  and  $\mathcal{A}$ .

Notice that  $\mathcal{S}$  is a symmetric operator and  $\mathcal{A}$  is skew-symmetric, with respect to the inner product in  $L^2$ . The proof of Lemma 2.9 is based on explicit computations and will be omitted. We mention the paper [10] for the computation of  $[\Delta_A, \Delta_x \varphi + 2\nabla_x \varphi \cdot \nabla_A]$ , which is the only term in  $[\mathcal{S}, \mathcal{A}]$  one has to compute with a bit of care.

We now prove a dissipation result for equation (2.1), which depends on the fact that  $a > 0$ , and which permits to justify the proofs of the results in the sequel.

**Lemma 2.10.** *Let  $-\Delta_A$  be self-adjoint in  $L^2$  and let  $u \in L^\infty([0, 1]; L^2(\mathbb{R}^n)) \cap L^2([0, 1]; H^1(\mathbb{R}^n))$  be a solution to*

$$(2.27) \quad \partial_t u = (a + ib)(\Delta_A u + V(x, t)u + F(x, t)),$$

in  $\mathbb{R}^n \times [0, 1]$ , with  $a > 0$ ,  $b \in \mathbb{R}$ ,  $A = A(x) : \mathbb{R}^n \rightarrow \mathbb{R}^n$ , and  $V, F : \mathbb{R}^{n+1} \rightarrow \mathbb{C}$ . Then, for any  $\gamma \geq 0$ ,  $T \in [0, 1]$ , we have

$$(2.28) \quad e^{-M_T} \left\| e^{\frac{\gamma a}{a+4\gamma(a^2+b^2)T}|\cdot|^2} u(\cdot, T) \right\|_{L^2} \leq \|e^{\gamma|\cdot|^2} u(\cdot, 0)\|_{L^2} + \sqrt{a^2 + b^2} \left\| e^{\frac{\gamma a}{a+4\gamma(a^2+b^2)t}|x|^2} F(x, t) \right\|_{L^1([0, T]; L^2(\mathbb{R}^n))},$$

with  $M_T := \|a(\Re V)^+ - b\Im V\|_{L^1([0, T]; L^\infty(\mathbb{R}^n))}$ ,  $(\Re V)^+$  being the positive part of  $\Re V$ .

*Proof.* The proof is based on a standard energy method. First notice that, since  $-\Delta_A$  is self-adjoint, solutions  $u \in L^\infty([0, 1]; L^2(\mathbb{R}^n)) \cap L^2([0, 1]; H^1(\mathbb{R}^n))$  to (2.27) do exist by means of the Duhamel principle.

Let  $v = e^{\varphi(x,t)}u$ , satisfying (2.22) by Lemma 2.9. Formally, multiplying (2.22) by  $\bar{v}$ , integrating in  $dx$  and taking the real parts, we obtain by (2.23), (2.24) that

$$(2.29) \quad \begin{aligned} \frac{1}{2} \frac{d}{dt} \|v\|_{L^2}^2 &= \Re \int \mathcal{S}v \bar{v} \, dx + \Re \left\{ (a+ib) \int (|v|^2 V + e^\varphi F \bar{v}) \, dx \right\} \\ &= -a \int |\nabla_A v|^2 \, dx + a \int |\nabla_x \varphi|^2 |v|^2 \, dx + \int \varphi_t |v|^2 \, dx \\ &\quad + 2b \Im \int \bar{v} \nabla_x \varphi \cdot \nabla_A v \, dx + \Re(a+ib) \int (|v|^2 V + e^\varphi F \bar{v}) \, dx. \end{aligned}$$

We can easily estimate

$$(2.30) \quad \Re(a+ib) \int |v|^2 V \, dx \leq \|a(\Re V)^+ - b(\Im V)\|_{L^\infty} \|v\|_{L^2}^2$$

$$(2.31) \quad \Re(a+ib) \int e^\varphi F \bar{v} \, dx \leq \sqrt{a^2 + b^2} \|e^\varphi F\|_{L^2} \|v\|_{L^2}.$$

Analogously, by Cauchy-Schwartz we have

$$(2.32) \quad 2b \Im \int \bar{v} \nabla_x \varphi \cdot \nabla_A v \, dx \leq a \int |\nabla_A v|^2 \, dx + \frac{b^2}{a} \int |\nabla_x \varphi|^2 |v|^2 \, dx;$$

as a consequence, by (2.29) and (2.32) we obtain

$$(2.33) \quad \Re \int \mathcal{S}v \bar{v} \, dx \leq \int \left\{ \left( a + \frac{b^2}{a} \right) |\nabla_x \varphi|^2 + \varphi_t \right\} |v|^2 \, dx,$$

and the choice

$$(2.34) \quad \varphi(x, t) = \frac{\gamma a}{a + 4\gamma(a^2 + b^2)t} |x|^2 \quad \Rightarrow \quad \varphi_t(x, t) = - \left( a + \frac{b^2}{a} \right) |\nabla_x \varphi|^2$$

gives in turn that

$$(2.35) \quad \Re \int \mathcal{S}v \bar{v} \, dx \leq 0.$$

By (2.29), (2.30), (2.31), (2.35), with the choice (2.34), we finally obtain

$$\begin{aligned} &\frac{d}{dt} \|v(\cdot, t)\|_{L^2}^2 \\ &\leq 2 \|a(\Re V)^+ - b(\Im V)\|_{L^\infty} \|v(\cdot, t)\|_{L^2}^2 + 2\sqrt{a^2 + b^2} \|e^\varphi F\|_{L^2} \|v(\cdot, t)\|_{L^2}, \end{aligned}$$

which implies (2.28).

In order to make the previous argument rigorous, since the exponentially weighted  $L^2$ -norms involved in the integration by parts are not finite in principle, it is sufficient to work with truncated and mollified weights of the following form:

$$\varphi_{R,\epsilon}(x, t) = \begin{cases} \varphi(x, t), & \text{if } |x| < R \\ \varphi(R, t), & \text{if } |x| \geq R, \end{cases} \quad \varphi_{R,\epsilon} := (\theta_\epsilon * \varphi_R)(x),$$

$\theta_\epsilon(x)$  being a radial mollifier. Then the result can be obtained by performing the same computation as above and then letting  $\epsilon$  go to 0 and  $R$  to  $\infty$ ; we omit straightforward details.  $\square$

**Remark 2.11.** Notice that the dissipation estimate (2.28) has been proved for stationary magnetic potentials  $A = A(x)$ . In the time-dependent case  $A = A(x, t)$ , the same result would require some additional assumptions on the time derivative  $A_t$ , since we need the self-adjointness property, which at this level seem quite artificial.

The next result, proved by Escauriaza, Kenig, Ponce and Vega in [5, 6], is the abstract core of Theorem 1.5. It is concerned with the connection between the positivity of  $\mathcal{S}_t + [\mathcal{S}, \mathcal{A}]$  and the logarithmic convexity of weighted  $L^2$ -norms with gaussian weights.

**Lemma 2.12** (logarithmic convexity). *Let  $\mathcal{S}$  be a symmetric operator,  $\mathcal{A}$  a skew-symmetric one, both with coefficients depending on  $x$  and  $t$ ,  $f = f(x, t) : \mathbb{R}^{n+1} \rightarrow \mathbb{C}$  be a sufficiently regular function,  $G$  a positive function, and denote by*

$$(2.36) \quad H(t) = \int_{\mathbb{R}^n} |f|^2 dx.$$

Assume that

$$(2.37) \quad |\partial_t f - (\mathcal{S} + \mathcal{A})f| \leq M_1 |f| + G \text{ in } \mathbb{R}^n \times [0, 1], \quad \mathcal{S}_t + [\mathcal{S}, \mathcal{A}] \geq -M_0,$$

for some  $M_0, M_1 \geq 0$  and

$$(2.38) \quad M_2 := \sup_{t \in [0, 1]} \frac{\|G(t)\|_{L^2}}{\|f(t)\|_{L^2}} < \infty.$$

Then the function  $\psi(t) := \log H(t)$  is convex in  $[0, 1]$ . In particular, if

$$(2.39) \quad H(0) < \infty \quad \Rightarrow \quad H(t) < \infty \text{ for any } t \in [0, 1],$$

then there exist a universal constant  $N \geq 0$  such that

$$(2.40) \quad H(t) \leq e^{N(M_0 + M_1 + M_2 + M_1^2 + M_2^2)} H(0)^{1-t} H(1)^t,$$

for any  $t \in [0, 1]$ .

**Remark 2.13.** The proof of Lemma 2.12 is based on the computation of the time derivatives  $\dot{H}(t), \ddot{H}(t)$ . An explicit (formal) computation gives

$$(2.41) \quad \begin{aligned} \frac{d^2}{dt^2} H(t) &= 2\partial_t \Re \int \bar{v}(\partial_t - \mathcal{S} - \mathcal{A})v dx + 2 \int \bar{v}(\mathcal{S}_t + [\mathcal{S}, \mathcal{A}])v dx \\ &\quad + \|\partial_t v - \mathcal{A}v + \mathcal{S}v\|_{L^2}^2 - \|\partial_t v - \mathcal{A}v - \mathcal{S}v\|_{L^2}^2. \end{aligned}$$

This, together with the computation of the first derivative  $\dot{H}(t)$ , shows that, under conditions (2.37), (2.38), the second derivative  $\frac{d^2}{dt^2} \log(H(t))$  is positive. Assumption (2.39) is then the essential information one needs in order to conclude the convexity inequality (2.40). The validity of condition (2.39) depends on an energy estimate of the type (2.28) and needs to be checked each time when Lemma 2.12 is applied to explicit operators  $\mathcal{S}, \mathcal{A}$ , as we see in the following results.

The proof of Lemma 2.12 can be found in [5, 6].

We can finally prove the main results of this section.

**Lemma 2.14.** *Let  $u \in L^\infty([0, 1]; L^2(\mathbb{R}^n)) \cap L^2([0, 1]; H^1(\mathbb{R}^n))$  be a solution to*

$$(2.42) \quad \partial_t u = (a + ib)(\Delta_A u + V(x, t)u + F(x, t)),$$

in  $\mathbb{R}^n \times [0, 1]$ , with  $a > 0$ ,  $b \in \mathbb{R}$ ,  $A = A(x, t) : \mathbb{R}^{n+1} \rightarrow \mathbb{R}^n$ , and  $V, F : \mathbb{R}^{n+1} \rightarrow \mathbb{C}$ . Assume that

$$(2.43) \quad x \cdot A(x) \equiv 0 \equiv x \cdot A_t(x).$$

Moreover, let  $\gamma > 0$  and assume that

$$(2.44) \quad \sup_{t \in [0,1]} \|V(t, \cdot)\|_{L^\infty} := M_1 < \infty, \quad \sup_{t \in [0,1]} \frac{\|e^{\gamma|\cdot|^2} F(\cdot, t)\|_{L^2}}{\|u(\cdot, t)\|_{L^2}} := M_2 < \infty;$$

in addition, denote by  $B = B(x, t) = D_x A - D_x A^t$  and assume

$$(2.45) \quad \frac{1}{\gamma} \sup_{t \in [0,1]} \|A_t(\cdot, t)\|_{L^\infty}^2 + 4\gamma(a^2 + b^2) \sup_{t \in [0,1]} \|x^t B(\cdot, t)\|_{L^\infty}^2 := M_A < \infty.$$

Finally, assume

$$(2.46) \quad \left\| e^{\gamma|\cdot|^2} u(\cdot, 0) \right\|_{L^2} + \left\| e^{\gamma|\cdot|^2} u(\cdot, 1) \right\|_{L^2} < \infty;$$

finally, define  $H(t) = \left\| e^{\gamma|\cdot|^2} u(\cdot, t) \right\|_{L^2}$  and assume that (2.39) holds. Then,  $H(t)$  is finite and logarithmically convex in  $[0, 1]$ ; in particular, there exists a constant  $N = N(\gamma, a, b)$  such that

$$(2.47) \quad H(t) \leq e^{N[M_A + \sqrt{a^2 + b^2}(M_1 + M_2) + (a^2 + b^2)(M_1^2 + M_2^2)]} \left\| e^{\gamma|\cdot|^2} u(\cdot, 0) \right\|_{L^2}^{1-t} \left\| e^{\gamma|\cdot|^2} u(\cdot, 1) \right\|_{L^2}^t,$$

for any  $t \in [0, 1]$ .

**Remark 2.15.** Before the proof, we need another remark about condition (2.39) in the statement. The result ensuring, in concrete situations, the validity of (2.39), is Lemma 2.10. Notice that in the statement of Lemma 2.14 we work with magnetic potentials  $A = A(x, t)$  which possibly depend on time, while the time dependence is not permitted in Lemma 2.10. In fact, as we see in the next section, in the proof of Theorem 1.5, after applying the Appell transformation, a natural time dependence of the magnetic potential appears. On the other hand, condition (2.39) will hold in the the next section as a heritage of the same property before the Appell transformation, and no additional assumptions on  $\partial_t A$  will be needed. This explains why we prefer to assume (2.39) in the previous statement without giving explicit conditions under which it is satisfied.

*Proof of Lemma 2.14.* We need to check that Lemma 2.12 is applicable.

Denote again by  $v = e^{\varphi(x,t)} u$ , with  $\varphi(x, t) = \varphi(x) := \gamma|x|^2$ . By Lemma 2.9,  $v$  satisfies

$$\partial_t v = \mathcal{S}v + \mathcal{A}v + (a + ib)(V(x, t)v + e^\varphi F),$$

where  $\mathcal{S}$  and  $\mathcal{A}$  are given by (2.23), (2.24), respectively. We can estimate

$$(2.48) \quad |\partial_t v - (\mathcal{S} + \mathcal{A})v| \leq \sqrt{a^2 + b^2} (M_1|v| + e^\varphi|F|),$$

which proves the first of the two conditions in (2.37), with  $G := \sqrt{a^2 + b^2} e^\varphi |F|$ . Hence we just need to check the second condition in (2.37). By formulas (2.25) and (2.26) with the choice  $\varphi(x) = \gamma|x|^2$  we obtain

$$(2.49) \quad \int \bar{v} (\mathcal{S}_t + [\mathcal{S}, \mathcal{A}]) v dx = 2a\Im \int \bar{v} A_t \cdot \nabla_A v dx - 4b\gamma \int |v|^2 x \cdot A_t dx \\ + (a^2 + b^2) \left\{ 8\gamma \int |\nabla_A v|^2 dx + 8\gamma\Im \int \bar{v} x^t B \cdot \nabla_A v dx + 32\gamma^3 \int |v|^2 |x|^2 dx \right\}.$$

The second term at the right-hand side of (2.49) vanishes, due to (2.43). By Cauchy-Schwartz, we can estimate

$$(2.50) \quad \left| 2a \Im \int \bar{v} A_t \cdot \nabla_A v \, dx \right| \leq \frac{1}{\gamma} \int |A_t|^2 |v|^2 \, dx + \gamma a^2 \int |\nabla_A v|^2 \, dx$$

$$(2.51) \quad \left| 8\gamma(a^2 + b^2) \Im \int \bar{v} x^t B \cdot \nabla_A v \, dx \right| \leq 4\gamma(a^2 + b^2) \int |x^t B|^2 |v|^2 \, dx \\ + 4\gamma(a^2 + b^2) \int |\nabla_A v|^2 \, dx;$$

by (2.49), (2.50), (2.51) it turns out that

$$(2.52) \quad \int \bar{v} (\mathcal{S}_t + [\mathcal{S}, \mathcal{A}]) v \, dx \geq 3\gamma(a^2 + b^2) \int |\nabla_A v|^2 \, dx + 32\gamma^3(a^2 + b^2) \int |v|^2 |x|^2 \\ - \left( \frac{1}{\gamma} \sup_{t \in [0,1]} \|A_t\|_{L^\infty}^2 + 4\gamma(a^2 + b^2) \sup_{t \in [0,1]} \|x^t B\|_{L^\infty}^2 \right) \int |v|^2 \, dx.$$

Neglecting the positive terms in the last inequality, we have proved that

$$(2.53) \quad \mathcal{S}_t + [\mathcal{S}, \mathcal{A}] \geq -\frac{1}{\gamma} \sup_{t \in [0,1]} \|A_t\|_{L^\infty}^2 - 4\gamma(a^2 + b^2) \sup_{t \in [0,1]} \|x^t B\|_{L^\infty}^2 = -M_A.$$

In addition, we have

$$(2.54) \quad \sup_{t \in [0,1]} \frac{\sqrt{a^2 + b^2} \left\| e^{\gamma|\cdot|^2} F(\cdot, t) \right\|_{L^2}}{\|v(\cdot, t)\|_{L^2}} \leq \sqrt{a^2 + b^2} M_2.$$

The thesis now follows by Lemma 2.12.

In order to obtain a completely rigorous proof of Lemma 2.14 we need a last remark. The positive dissipation  $a > 0$  provides the sufficient interior regularity for Lemma 2.10 to hold. In the next section, when we apply Lemma 2.14 to a concrete situation, in order to justify all the above computations we need to work with the following multipliers. Given  $a > 0$  and  $\rho \in (0, 1)$ , define

$$\varphi_a(x) = \begin{cases} \gamma|x|^2, & \text{if } |x| < 1 \\ \gamma \frac{2|x|^{2-a} - a}{2-a} & \text{if } |x| \geq 1 \end{cases}$$

and replace  $\varphi = \gamma|x|^2$  by  $\varphi_{a,\rho} = \theta_\rho \star \varphi_a$ , being  $\theta_\rho$  a smooth delta-sequence. One can easily check that all the above computations are then justified as a limit when  $a, \rho \rightarrow 0$ . See [6] for further details.  $\square$

In an analogous way, we prove the following result:

**Lemma 2.16.** *Under the same assumptions as in Lemma 2.14, there exists a constant  $N = N\left(\frac{1}{\gamma}, \frac{1}{a^2 + b^2}\right) > 0$  such that*

$$(2.55) \quad \left\| \sqrt{t(1-t)} e^{\gamma|x|^2} \nabla_A u(x, t) \right\|_{L^2(\mathbb{R}^n \times [0,1])} + \gamma \left\| \sqrt{t(1-t)} e^{\gamma|x|^2} |x| u(x, t) \right\|_{L^2(\mathbb{R}^n \times [0,1])} \\ \leq N \left[ (M_1 + \sqrt{M_A} + 1) \sup_{t \in [0,1]} \left\| e^{\gamma|\cdot|^2} u(\cdot, t) \right\|_{L^2} + \sup_{t \in [0,1]} \left\| e^{\gamma|\cdot|^2} F(\cdot, t) \right\|_{L^2} \right].$$



*Proof.* Denote again by  $v = e^{\gamma|x|^2}u$ ; we can hence write

$$\nabla_A u = -2\gamma x e^{-\gamma|x|^2}v + e^{-\gamma|x|^2}\nabla_A v.$$

Consequently, we can estimate

$$(2.56) \quad \begin{aligned} & \left\| \sqrt{t(1-t)}e^{\gamma|x|^2}\nabla_A u(x, t) \right\|_{L^2(\mathbb{R}^n \times [0,1])} + \gamma \left\| \sqrt{t(1-t)}e^{\gamma|x|^2}|x|u(x, t) \right\|_{L^2(\mathbb{R}^n \times [0,1])} \\ & \leq 3\gamma \left\| \sqrt{t(1-t)}|x|v(x, t) \right\|_{L^2(\mathbb{R}^n \times [0,1])} + \left\| \sqrt{t(1-t)}\nabla_A v(x, t) \right\|_{L^2(\mathbb{R}^n \times [0,1])}. \end{aligned}$$

By (2.41), we easily estimate

$$(2.57) \quad \begin{aligned} \frac{d^2}{dt^2}H(t) & \geq 2\partial_t \Re \int \bar{v}(\partial_t - \mathcal{S} - \mathcal{A})v \, dx + 2 \int \bar{v}(\mathcal{S}_t + [\mathcal{S}, \mathcal{A}])v \, dx \\ & \quad - \|\partial_t v - \mathcal{A}v - \mathcal{S}v\|_{L^2_x}^2. \end{aligned}$$

On the other hand, integrating twice by parts we get

$$(2.58) \quad \int_0^1 t(1-t) \frac{d^2}{dt^2}H(t) \, dt = H(1) + H(0) - 2 \int_0^1 H(t) \, dt \leq 2 \sup_{t \in [0,1]} \|v(\cdot, t)\|_{L^2}^2,$$

since  $H(t) \geq 0$ . Integrating by parts and applying Cauchy-Schwartz and estimate (2.48), we obtain

$$(2.59) \quad \begin{aligned} & 2 \int_0^1 \int t(1-t) \partial_t \Re \bar{v}(\partial_t - \mathcal{S} - \mathcal{A})v \, dx \, dt \\ & = -2 \int_0^1 \int (1-2t) \Re \bar{v}(\partial_t - \mathcal{S} - \mathcal{A})v \, dx \, dt \\ & \geq - \left( \sup_{t \in [0,1]} \|\partial_t v - \mathcal{S}v - \mathcal{A}v\|_{L^2}^2 + \sup_{t \in [0,1]} \|v(\cdot, t)\|_{L^2}^2 \right) \\ & \geq -\frac{1}{2} \left\{ [(a^2 + b^2)M_1^2 + 1] \sup_{t \in [0,1]} \|v(\cdot, t)\|_{L^2}^2 + (a^2 + b^2) \sup_{t \in [0,1]} \left\| e^{\gamma|\cdot|^2} F(\cdot, t) \right\|_{L^2}^2 \right\}. \end{aligned}$$

On the other hand, by (2.52) we get

$$(2.60) \quad \begin{aligned} & 2 \int_0^1 \int t(1-t) \bar{v}(\mathcal{S}_t + [\mathcal{S}, \mathcal{A}])v \, dx \, dt \geq -\frac{M_A}{3} \sup_{t \in [0,1]} \|v(\cdot, t)\|_{L^2}^2 \\ & \quad + 2\gamma(a^2 + b^2) \left\{ \left\| \sqrt{t(1-t)}\nabla_A v \right\|_{L^2(\mathbb{R}^n \times [0,1])}^2 + \gamma^2 \left\| \sqrt{t(1-t)}|x|v \right\|_{L^2(\mathbb{R}^n \times [0,1])}^2 \right\} \end{aligned}$$

while by (2.48) we conclude that

$$\begin{aligned}
(2.61) \quad & - \int_0^1 t(1-t) \|\partial_t v - \mathcal{S}v - \mathcal{A}v\|_{L^2}^2 dt \\
& \geq - \sup_{t \in [0,1]} \|\partial_t v - \mathcal{S}v - \mathcal{A}v\|_{L^2}^2 \int_0^1 t(1-t) dt \\
& \geq -\frac{1}{6} (a^2 + b^2) \left\{ M_1^2 \sup_{t \in [0,1]} \|v(\cdot, t)\|_{L^2}^2 + \sup_{t \in [0,1]} \left\| e^{\gamma|\cdot|^2} F(\cdot, t) \right\|_{L^2}^2 \right\}.
\end{aligned}$$

Collecting (2.57), (2.58), (2.59), (2.60), (2.61) we have

$$\begin{aligned}
& \left\| \sqrt{t(1-t)} \nabla_A v \right\|_{L^2(\mathbb{R}^n \times [0,1])}^2 + \gamma^2 \left\| \sqrt{t(1-t)} |x| v \right\|_{L^2(\mathbb{R}^n \times [0,1])}^2 \\
& \leq \left[ \frac{M_1^2}{3\gamma} + \frac{15 + 2M_A}{12\gamma(a^2 + b^2)} \right] \sup_{t \in [0,1]} \|v(\cdot, t)\|_{L^2}^2 + \frac{1}{3\gamma} \sup_{t \in [0,1]} \left\| e^{\gamma|\cdot|^2} F(\cdot, t) \right\|_{L^2}^2,
\end{aligned}$$

which, together with (2.56), proves the claim (2.55).

Also in this case, the proof can be made rigorous by a quite standard argument in the spirit of the one in Lemma 2.14.  $\square$

All the tools we need to prove Theorem 1.5 are now ready.

### 3. PROOF OF THEOREM 1.5

For the proof of Theorem 1.5, we now put together the informations we got in the previous Section. It is sufficient to prove the result in the case  $\alpha < \beta$ ; for the proof in the case  $\alpha > \beta$  replace  $u(x, t)$  by  $\bar{u}(x, 1-t)$ , while in the case  $\alpha = \beta$  the proof essentially reduces to Lemma 2.14 and 2.16 (see Remark 3.3 below). Therefore, from now on we assume

$$\alpha < \beta.$$

We divide the proof of Theorem 1.5 into four steps.

**3.1. Step I: the gauge reduction.** Thanks to assumption (2.43) and Lemma 2.2, it is now sufficient to prove Theorem 1.5 for the function  $\tilde{u} = e^{i\varphi} u$ , where  $\varphi$  is the gauge change defined in (2.4). The new potential is  $\tilde{A}$ , defined in (2.5). By abuse of notations, we will skip the tildes; hence, from now on, the additional (and not restrictive) assumption

$$(3.1) \quad x \cdot A \equiv 0$$

holds, together with the identities (2.5), (2.6), which in our new notations read as

$$(3.2) \quad A(x) = - \int_0^1 \Psi(sx) ds; \quad x^t DA(x) = -\Psi(x) + \int_0^1 \Psi(sx) ds,$$

with  $\Psi(x) = x^t B(x) = x^t (DA(x) - DA^t(x))$ , which also gives

$$(3.3) \quad x \cdot x^t DA \equiv 0.$$

In particular, (1.11) and (3.2) also imply that

$$(3.4) \quad \|A\|_{L^\infty}^2 + \|x^t DA\|_{L^\infty}^2 + \|x^t B\|_{L^\infty}^2 \leq M_A.$$

**3.2. Step II: the heat regularization.** We now regularize equation (1.9) adding a small dissipation term. Denote by

$$H_A := -\Delta_A - V_1$$

and rewrite equation (1.9) as

$$(3.5) \quad \partial_t u = -i(H_A u - F(x, t)), \quad F(x, t) := V_2(x, t)u.$$

Since  $H_A$  is self-adjoint by Proposition 2.6, we can define, by Spectral Theorem, the mixed flow  $e^{(\epsilon+i)tH_A}$ , for any  $\epsilon > 0$ . This gives, by parabolic regularity, the function

$$(3.6) \quad u_\epsilon(\cdot, t) := e^{(\epsilon+i)tH_A} u(\cdot, 0) = e^{\epsilon t H_A} u(t) \in L^\infty([0, 1]; L^2(\mathbb{R}^n)) \cap L^2([0, 1]; H^1(\mathbb{R}^n)),$$

solving (uniquely) the equation

$$(3.7) \quad \begin{cases} \partial_t u_\epsilon = (\epsilon + i)(\Delta_A u_\epsilon + V_1(x)u_\epsilon + F_\epsilon(x, t)), \\ u_\epsilon(0) = u(0), \end{cases}$$

with  $F_\epsilon(\cdot, s) := \frac{i}{\epsilon+i} e^{\epsilon t H_A} (V_2 u(\cdot, s))$  (see e.g. [13, 15]). The positive dissipation now permits to apply Lemma 2.10, which is useful in the sequel to make rigorous the applications of Lemma 2.14. We can now prove the following simple result.

**Lemma 3.1.** *Denote by*

$$(3.8) \quad \alpha_\epsilon^2 = \alpha^2 + 4\epsilon \quad \beta_\epsilon^2 = \beta^2 + 4\epsilon.$$

*The function  $u_\epsilon$  defined in (3.6) satisfies the following inequalities:*

$$(3.9) \quad \left\| e^{\frac{|\cdot|^2}{\beta_\epsilon^2}} u_\epsilon(\cdot, 0) \right\|_{L^2} \leq \left\| e^{\frac{|\cdot|^2}{\beta^2}} u(\cdot, 0) \right\|_{L^2}$$

$$(3.10) \quad \left\| e^{\frac{|\cdot|^2}{\alpha_\epsilon^2}} u_\epsilon(\cdot, 1) \right\|_{L^2} \leq e^{\epsilon \|V_1\|_{L^\infty}} \left\| e^{\frac{|\cdot|^2}{\alpha^2}} u(\cdot, 1) \right\|_{L^2}$$

$$(3.11) \quad \|u_\epsilon(\cdot, t)\|_{L^2} \leq e^{\epsilon \|V_1\|_{L^\infty}} \|u(\cdot, t)\|_{L^2}$$

$$(3.12) \quad \|F_\epsilon(\cdot, t)\|_{L^2} \leq e^{\epsilon \|V_1\|_{L^\infty}} \|V_2\|_{L^\infty} \|u(\cdot, t)\|_{L^2}$$

$$(3.13) \quad \left\| e^{\frac{|\cdot|^2}{(\alpha_\epsilon t + \beta_\epsilon(1-t))^2}} F_\epsilon(\cdot, t) \right\|_{L^2} \leq e^{\epsilon \|V_1\|_{L^\infty}} \left\| e^{\frac{|\cdot|^2}{(\alpha t + \beta(1-t))^2}} V_2(\cdot, t) \right\|_{L^\infty} \|u(\cdot, t)\|_{L^2},$$

for any  $t \in [0, 1]$ .

*Proof.* Inequality (3.9) is immediate.

In order to prove (3.10), let us introduce the function  $w(\cdot, t) := e^{-\epsilon t H_A} u(\cdot, 1)$ , solving the equation

$$\partial_t w = -\epsilon H_A w = \epsilon(\Delta_A w + V_1 w).$$

Then (3.10) follows applying inequality (2.28) to  $w$ , with  $\gamma := \frac{1}{\alpha^2}$  and  $T = 1$ .

To prove (3.11) write  $u_\epsilon(\cdot, t) := e^{\epsilon t H_A} u(t)$  and apply again (2.28), with  $\gamma = 0$  and  $T = t$ .

For the proof of (3.12), introduce the function  $w(\cdot, t) := e^{\epsilon t H_A} (V_2 u(\cdot, t))$  and apply again (2.28), with  $\gamma = 0$ ,  $T = t$ . Finally, by the application of inequality (2.28) to the same function, with  $\gamma = \frac{1}{(\alpha t + \beta(1-t))^2}$  and  $T = t$ , the proof of (3.13) easily follows.  $\square$

**3.3. Step III: the Appell transformation.** We now apply the Appell transformation to the function  $u_\epsilon$ . Let  $\alpha_\epsilon, \beta_\epsilon$  be the same as in (3.8) and define

$$(3.14) \quad \begin{aligned} \tilde{u}_\epsilon(x, t) &:= \left( \frac{\sqrt{\alpha_\epsilon \beta_\epsilon}}{\alpha_\epsilon(1-t) + \beta_\epsilon t} \right)^{\frac{n}{2}} \times \\ &\times u_\epsilon \left( \frac{\sqrt{\alpha_\epsilon \beta_\epsilon}}{\alpha_\epsilon(1-t) + \beta_\epsilon t} x, \frac{\beta_\epsilon}{\alpha_\epsilon(1-t) + \beta_\epsilon t} t \right) e^{\frac{\alpha_\epsilon - \beta_\epsilon}{4(\epsilon+i)(\alpha_\epsilon(1-t) + \beta_\epsilon t)} |x|^2}. \end{aligned}$$

Since  $x \cdot A \equiv 0$  due to step I, by Lemma 2.7 we have that  $\tilde{u}_\epsilon$  solves

$$(3.15) \quad \partial_t \tilde{u}_\epsilon = (\epsilon + i) \left( \Delta_{\tilde{A}_\epsilon} \tilde{u}_\epsilon + \tilde{V}_\epsilon(x, t) \tilde{u}_\epsilon + \tilde{F}_\epsilon(x, t) \right),$$

where

$$(3.16) \quad \tilde{A}_\epsilon(x, t) = \frac{\sqrt{\alpha_\epsilon \beta_\epsilon}}{\alpha_\epsilon(1-t) + \beta_\epsilon t} A \left( \frac{\sqrt{\alpha_\epsilon \beta_\epsilon}}{\alpha_\epsilon(1-t) + \beta_\epsilon t} x \right)$$

$$(3.17) \quad \tilde{V}_\epsilon(x, t) = \frac{\alpha_\epsilon \beta_\epsilon}{(\alpha_\epsilon(1-t) + \beta_\epsilon t)^2} V_1 \left( \frac{\sqrt{\alpha_\epsilon \beta_\epsilon}}{\alpha_\epsilon(1-t) + \beta_\epsilon t} x \right)$$

$$(3.18) \quad \begin{aligned} \tilde{F}_\epsilon(x, t) &= \left( \frac{\sqrt{\alpha_\epsilon \beta_\epsilon}}{\alpha_\epsilon(1-t) + \beta_\epsilon t} \right)^{\frac{n}{2}+2} \times \\ &\times F_\epsilon \left( \frac{\sqrt{\alpha_\epsilon \beta_\epsilon}}{\alpha_\epsilon(1-t) + \beta_\epsilon t} x, \frac{\beta_\epsilon}{\alpha_\epsilon(1-t) + \beta_\epsilon t} t \right) e^{\frac{(\alpha_\epsilon - \beta_\epsilon) |x|^2}{4(\epsilon+i)(\alpha_\epsilon(1-t) + \beta_\epsilon t)}}. \end{aligned}$$

In addition, by Corollary 2.8, for any  $\gamma \in \mathbb{R}$  we have

$$(3.19) \quad \left\| e^{\gamma|\cdot|^2} \tilde{u}_\epsilon(\cdot, t) \right\|_{L^2} = \left\| e^{\left[ \frac{\gamma \alpha_\epsilon \beta_\epsilon}{(\alpha_\epsilon s + \beta_\epsilon(1-s))^2} + \frac{(\alpha_\epsilon - \beta_\epsilon) \epsilon}{4(\epsilon^2 + 1)(\alpha_\epsilon s + \beta_\epsilon(1-s))} \right] |\cdot|^2} u_\epsilon(\cdot, s) \right\|_{L^2}$$

$$(3.20) \quad \begin{aligned} \left\| e^{\gamma|\cdot|^2} \tilde{F}_\epsilon(\cdot, t) \right\|_{L^2} &= \frac{\alpha_\epsilon \beta_\epsilon}{(\alpha_\epsilon(1-t) + \beta_\epsilon t)^2} \times \\ &\times \left\| e^{\left[ \frac{\gamma \alpha_\epsilon \beta_\epsilon}{(\alpha_\epsilon s + \beta_\epsilon(1-s))^2} + \frac{(\alpha_\epsilon - \beta_\epsilon) \epsilon}{4(\epsilon^2 + 1)(\alpha_\epsilon s + \beta_\epsilon(1-s))} \right] |\cdot|^2} F_\epsilon(\cdot, s) \right\|_{L^2}, \end{aligned}$$

for  $s = \frac{\beta_\epsilon t}{\alpha_\epsilon(1-t) + \beta_\epsilon t}$ .

The goal is to apply Lemma 2.14 to the function  $\tilde{u}_\epsilon$ . In order to do this, we now need two more results regarding the evolution of the  $L_x^2$ -norms of  $u$  and  $\tilde{u}_\epsilon$ .

**Lemma 3.2.** *Denote by*

$$(3.21) \quad N_1 := e^{\sup_{t \in [0,1]} \|\Im V_2(\cdot, t)\|_{L^\infty}}.$$

*The following inequalities hold*

$$(3.22)$$

$$\frac{1}{N_1} \|u(\cdot, 0)\|_{L^2} \leq \|u(\cdot, t)\|_{L^2} \leq N_1 \|u(\cdot, 0)\|_{L^2}$$

$$(3.23)$$

$$\frac{d}{dt} \|\tilde{u}_\epsilon(\cdot, t)\|_{L^2} \leq \epsilon \frac{\beta}{\alpha} e^{\|V_1\|_{L^\infty}} N_1 \|u(\cdot, 0)\|_{L^2} \left( \|V_1\|_{L^\infty} + \sup_{t \in [0,1]} \|V_2(\cdot, t)\|_{L^\infty} \right),$$

for any  $t \in [0, 1]$ , where  $u$  is a solution to (1.9) and  $\tilde{u}_\epsilon$  is the function defined in (3.14).

*Proof.* Formally, multiplying (1.9) by  $\bar{u}$ , integrating in  $dx$  and taking the real part of the resulting identity, (3.22) immediately follows. This argument is rigorous for solutions  $u \in C([0, 1]; H^1)$ ; a standard approximation argument permits to conclude the same for  $L^2$ -solutions.

With the same argument, which is now rigorous since  $\tilde{u}_\epsilon$  is in  $H^1$ , by equation (3.15) we easily obtain

$$(3.24) \quad \frac{d}{dt} \|\tilde{u}_\epsilon(\cdot, t)\|_{L^2} \leq \epsilon \left( \left\| \tilde{V}_\epsilon(\cdot, t) \right\|_{L^\infty} \|\tilde{u}_\epsilon(\cdot, t)\|_{L^2} + \left\| \tilde{F}_\epsilon(\cdot, t) \right\|_{L^2} \right),$$

and by (3.17) we easily estimate

$$(3.25) \quad M_{1,\epsilon} := \left\| \tilde{V}_\epsilon(\cdot, t) \right\|_{L^\infty} \leq \sup_{t \in [0,1]} \frac{\alpha_\epsilon \beta_\epsilon}{(\alpha_\epsilon t + \beta_\epsilon(1-t))^2} \|V_1\|_{L^\infty} \leq \frac{\beta}{\alpha} M_1 < \infty,$$

$M_1$  being the constant defined in (1.12). Taking  $\gamma = 0$  in (3.19), since  $\alpha < \beta$  we get

$$\|\tilde{u}_\epsilon(\cdot, t)\|_{L^2} \leq \|u_\epsilon(\cdot, s)\|_{L^2},$$

and by the last inequality, together with (3.11) and (3.22) we conclude that

$$(3.26) \quad \|\tilde{u}_\epsilon(\cdot, t)\|_{L^2} \leq e^{\epsilon \|V_1\|_{L^\infty}} \|u(\cdot, s)\|_{L^2} \leq e^{\epsilon \|V_1\|_{L^\infty}} N_1 \|u(\cdot, 0)\|_{L^2}.$$

Arguing in a similar way, by (3.20) with  $\gamma = 0$ , (3.12) and (3.22) we get

$$(3.27) \quad \begin{aligned} \left\| \tilde{F}_\epsilon(\cdot, t) \right\|_{L^2} &\leq \frac{\beta}{\alpha} \|F_\epsilon(\cdot, s)\|_{L^2} \leq \frac{\beta}{\alpha} e^{\epsilon \|V_1\|_{L^\infty}} \|V_2(\cdot, s)\|_{L^\infty} \|u(\cdot, s)\|_{L^2} \\ &\leq \frac{\beta}{\alpha} e^{\epsilon \|V_1\|_{L^\infty}} \sup_{[t \in 0,1]} \|V_2(\cdot, s)\|_{L^\infty} N_1 \|u(\cdot, 0)\|_{L^2}. \end{aligned}$$

Inequality (3.23) now follows from (3.24), (3.25), (3.26) and (3.27).  $\square$

We are finally ready to check the applicability of Lemma 2.14 to  $\tilde{u}_\epsilon$ .

First, taking  $\gamma = \frac{1}{\alpha_\epsilon \beta_\epsilon} =: \gamma_\epsilon$  in (3.19), since  $\alpha < \beta$  we get

$$(3.28) \quad \left\| e^{\gamma_\epsilon |\cdot|^2} \tilde{u}_\epsilon(\cdot, 0) \right\|_{L^2} \leq \left\| e^{\frac{|\cdot|^2}{\beta_\epsilon^2}} u_\epsilon(\cdot, 0) \right\|_{L^2} < \infty,$$

by (3.9) and (1.14) (here we also used the fact that  $s = 0$  when  $t = 0$ ).

Analogously, by (3.10), (1.14) and the fact that  $s = 1$  when  $t = 1$ , we obtain

$$(3.29) \quad \left\| e^{\gamma_\epsilon |\cdot|^2} \tilde{u}_\epsilon(\cdot, 1) \right\|_{L^2} \leq \left\| e^{\frac{|\cdot|^2}{\alpha_\epsilon^2}} u_\epsilon(\cdot, 1) \right\|_{L^2} < \infty.$$

Taking now  $\gamma = \gamma_\epsilon$  in (3.20), by (3.13) and (3.22) we easily estimate

$$(3.30) \quad \left\| e^{\gamma_\epsilon |\cdot|^2} \tilde{F}_\epsilon(\cdot, t) \right\|_{L^2} \leq \frac{\beta}{\alpha} e^{\epsilon \|V_1\|_{L^\infty}} \left\| e^{\frac{|\cdot|^2}{(\alpha_\epsilon s + \beta_\epsilon(1-s))^2}} V_2(\cdot, s) \right\|_{L^\infty} N_1 \|u(\cdot, 0)\|_{L^2}.$$

On the other hand, taking  $\gamma = 0$  in (3.19) gives

$$(3.31) \quad \lim_{\epsilon \rightarrow 0} \|\tilde{u}_\epsilon(\cdot, t)\|_{L^2} = \lim_{\epsilon \rightarrow 0} \left\| e^{\frac{(\alpha_\epsilon - \beta_\epsilon)\epsilon}{4(\epsilon^2 + 1)(\alpha_\epsilon s + \beta_\epsilon(1-s))} |\cdot|^2} u_\epsilon(\cdot, s) \right\|_{L^2} = \|u(\cdot, s)\|_{L^2}.$$

Now, by Lemma 3.2

$$(3.32) \quad \frac{d}{dt} \|\tilde{u}_\epsilon(\cdot, t)\|_{L^2} \leq C,$$

for some  $C = C\left(\epsilon, \alpha, \beta, \|V_1\|_{L^\infty}, \|u(\cdot, 0)\|_{L^2}, \sup_{t \in [0, 1]} \|V_2(\cdot, t)\|_{L^\infty}\right)$  and for any  $t \in [0, 1]$ . By (3.31) and (3.32) we hence obtain that

$$\|\tilde{u}_\epsilon(\cdot, t)\|_{L^2} \rightarrow \|u(\cdot, s)\|_{L^2},$$

as  $\epsilon \rightarrow 0$ , uniformly in  $[0, 1]$ , and in particular, by (3.22), there exists  $0 < \epsilon_0 = \epsilon_0(\|u(\cdot, 0)\|_{L^2}, N_1)$  such that

$$(3.33) \quad \|\tilde{u}_\epsilon(\cdot, t)\|_{L^2} \geq \frac{\|u(\cdot, 0)\|_{L^2}}{2N_1},$$

for any  $t \in [0, 1]$  and any  $\epsilon \in (0, \epsilon_0)$ . By (3.30) and (3.33) we finally obtain

$$(3.34) \quad M_2, \epsilon := \sup_{t \in [0, 1]} \frac{\left\| e^{\gamma_\epsilon |\cdot|^2} \tilde{F}_\epsilon(\cdot, t) \right\|_{L^2}}{\|\tilde{u}_\epsilon(\cdot, t)\|_{L^2}} \leq 2N_1^2 \frac{\beta}{\alpha} e^{\epsilon \|V_1\|_{L^\infty}} \sup_{s \in [0, 1]} \left\| e^{\frac{|\cdot|^2}{(\alpha s + \beta(1-s))^2}} V_2(\cdot, s) \right\|_{L^\infty} < \infty,$$

by assumptions (1.12), (1.13), for  $\epsilon > 0$  small enough; this ensures the validity of the second condition (2.44) (the first condition in (2.44) is quite immediate, thanks to (3.17) and (1.12)).

We now pass to condition (2.45). By (3.16), writing  $g_\epsilon(t) := \sqrt{\alpha_\epsilon \beta_\epsilon} / (\alpha_\epsilon(1-t) + \beta_\epsilon t)$  we explicitly compute

$$(3.35) \quad \partial_t \tilde{A}_\epsilon(x, t) = g'_\epsilon(t) [A(xg_\epsilon(t)) + g_\epsilon(t)x^t DA(xg_\epsilon(t))]$$

$$(3.36) \quad x^t \tilde{B}_\epsilon(x, t) := x^t (D\tilde{A}_\epsilon - D\tilde{A}_\epsilon^t)(x, t) = g_\epsilon^2(t)x^t B(xg_\epsilon(t)).$$

Writing

$$g'_\epsilon(t) = \frac{(\alpha_\epsilon - \beta_\epsilon)}{\sqrt{\alpha_\epsilon \beta_\epsilon}} g_\epsilon^2(t)$$

and estimating

$$\sup_{t \in [0, 1]} g_\epsilon^2(t) \leq \frac{\beta_\epsilon}{\alpha_\epsilon},$$

we easily obtain, using the above identities, (1.11) and (3.4),

$$(3.37) \quad \begin{aligned} M_{\tilde{A}, \epsilon} &:= \frac{1}{\gamma_\epsilon} \sup_{t \in [0, 1]} \left\| \partial_t \tilde{A}_\epsilon(\cdot, t) \right\|_{L^\infty}^2 + 4\gamma_\epsilon(\epsilon^2 + 1) \sup_{t \in [0, 1]} \left\| x^t \cdot \tilde{B}(\cdot, t) \right\|_{L^\infty}^2 \\ &\leq \frac{2(\alpha_\epsilon^2 + \beta_\epsilon^2)\beta_\epsilon^2}{\alpha_\epsilon^2} (\|A\|_{L^\infty}^2 + \|x^t DA\|_{L^\infty}^2) + \frac{4(\epsilon^2 + 1)}{\alpha_\epsilon^2} \|x^t B\|_{L^\infty}^2 \\ &\leq \frac{4}{\alpha_\epsilon^2} [(\alpha_\epsilon^2 + \beta_\epsilon^2)\beta_\epsilon^2 + \epsilon^2 + 1] M_A < \infty, \end{aligned}$$

$M_A$  being the constant in (1.11).

Finally, notice that from (3.1) and (3.16), and from (3.3) and (3.35) it follows that

$$x \cdot \tilde{A}_\epsilon \equiv 0, \quad \text{and} \quad x \cdot \partial_t \tilde{A}_\epsilon \equiv 0,$$

respectively.

The above argument shows that we can apply the results in Lemmata 2.14 and 2.16 to obtain

(3.38)

$$\begin{aligned} & \left\| e^{\gamma_\epsilon |\cdot|^2} \tilde{u}_\epsilon(\cdot, t) \right\|_{L^2} \\ & \leq e^{N_1 [M_{\tilde{A}, \epsilon} + \sqrt{\epsilon^2 + 1} (M_{1, \epsilon} + M_{2, \epsilon}) + (\epsilon^2 + 1) (M_{1, \epsilon}^2 + M_{2, \epsilon}^2)]} \left\| e^{\gamma_\epsilon |\cdot|^2} \tilde{u}_\epsilon(\cdot, 0) \right\|_{L^2}^{1-t} \left\| e^{\gamma_\epsilon |\cdot|^2} \tilde{u}_\epsilon(\cdot, 1) \right\|_{L^2}^t \end{aligned}$$

(3.39)

$$\begin{aligned} & \left\| \sqrt{t(1-t)} e^{\gamma_\epsilon |x|^2} \nabla_{\tilde{A}_\epsilon} \tilde{u}_\epsilon(x, t) \right\|_{L^2(\mathbb{R}^n \times [0, 1])} + \gamma_\epsilon \left\| \sqrt{t(1-t)} e^{\gamma_\epsilon |x|^2} |x| \tilde{u}_\epsilon(x, t) \right\|_{L^2(\mathbb{R}^n \times [0, 1])} \\ & \leq N_{2, \epsilon} \left[ (M_{1, \epsilon} + 1) \sup_{t \in [0, 1]} \left\| e^{\gamma_\epsilon |\cdot|^2} \tilde{u}_\epsilon(\cdot, t) \right\|_{L^2} + \sup_{t \in [0, 1]} \left\| e^{\gamma_\epsilon |\cdot|^2} \tilde{F}_\epsilon(\cdot, t) \right\|_{L^2} \right], \end{aligned}$$

with  $N_1$  an universal constant and  $N_{2, \epsilon} = N_{2, \epsilon}(\epsilon, \gamma_\epsilon) > 0$ .

**3.4. Step IV: conclusion of the proof.** It is now simple to conclude the proof of Theorem 1.5. Indeed, it is sufficient to rewrite estimates (3.38) and (3.39) in terms of the function  $u_\epsilon(t)$ , using Corollary 2.8; finally, (1.15) and (1.16) follow by taking the limit as  $\epsilon$  tends to 0. We omit further details.

**Remark 3.3.** In the case  $\alpha = \beta$  the same proof as above holds, in a much simpler version. Indeed, in this case it is useless to apply the Appell transformation and the proof can be directly performed on the function  $u_\epsilon$ , by means of Lemmata 2.14 and 2.16.

#### 4. PROOF OF THEOREM 1.1

**Lemma 4.1** (Carleman estimate). *Let  $n \geq 3$ ,  $A = A(x, t) : \mathbb{R}^{n+1} \rightarrow \mathbb{R}^n$ , denote by  $B = DA - DA^t$  and assume that  $x^t B \in L^\infty$ . In addition, assume that*

$$(4.1) \quad x \cdot A_t(x) \equiv 0, \quad v \cdot A_t(x) \equiv 0, \quad \text{and} \quad v^t B(x) \equiv 0,$$

for any  $x \in \mathbb{R}^n$  and some unit vector  $v \in \mathcal{S}^{n-1}$ . Then, for any  $\epsilon > 0$ ,  $\mu > 0$ ,  $g = g(x, t) \in C_0^\infty(\mathbb{R}^{n+1})$ , and  $R > 8\mu\epsilon^{-\frac{1}{2}} \|x^t B\|_{L^\infty}$ , the following inequality holds:

$$(4.2) \quad \begin{aligned} & \frac{R}{4} \sqrt{\frac{\epsilon}{\mu}} \left\| e^{\mu|x+Rt(1-t)v|^2 - \frac{(1+\epsilon)R^2 t(1-t)}{16\mu}} g(x, t) \right\|_{L^2(\mathbb{R}^{n+1})} \\ & \leq \left\| e^{\mu|x+Rt(1-t)v|^2 - \frac{(1+\epsilon)R^2 t(1-t)}{16\mu}} (\partial_t - i\Delta_A) g(x, t) \right\|_{L^2(\mathbb{R}^{n+1})}. \end{aligned}$$

*Proof.* For simplicity, we can assume without loss of generality that  $v = e_1 = (1, 0, \dots, 0)$ . Let

$$f(x, t) := e^{\mu|x+Rt(1-t)e_1|^2 - \frac{(1+\epsilon)R^2 t(1-t)}{16\mu}} g(x, t).$$

Then we have

$$(4.3) \quad e^{\mu|x+Rt(1-t)e_1|^2 - \frac{(1+\epsilon)R^2 t(1-t)}{16\mu}} (\partial_t - i\Delta_A) g = (\partial_t - \mathcal{S} - \mathcal{A}) f,$$

$\mathcal{S}$  and  $\mathcal{A}$  being the ones in (2.23) and (2.24), respectively, with  $a = 0$  and  $b = 1$ .

Following the usual method to prove Carleman estimates (see [11]), we now write

$$\begin{aligned}
(4.4) \quad & \|(\partial_t - \mathcal{S} - \mathcal{A})f\|_{L^2(\mathbb{R}^{n+1})}^2 \\
&= \|(\partial_t - \mathcal{A})f\|_{L^2(\mathbb{R}^{n+1})}^2 + \|\mathcal{S}f\|_{L^2(\mathbb{R}^{n+1})}^2 - 2\Re \iint \mathcal{S}f \overline{(\partial_t - \mathcal{A})f} dx dt \\
&\geq \iint (\mathcal{S}_t + [\mathcal{S}, \mathcal{A}])f \bar{f} dx dt.
\end{aligned}$$

Applying now (2.25) and (2.26) with the choices  $a = 0$ ,  $b = 1$ , and

$$\varphi(x, t) = \mu |x + Rt(1-t)e_1|^2 - \frac{(1+\epsilon)R^2t(1-t)}{16\mu},$$

noticing that  $\nabla\varphi \cdot A_t \equiv 0$  by the first two conditions in (4.1), an easy computation involving the completion of two squares leads to

$$\begin{aligned}
(4.5) \quad & \iint (\mathcal{S}_t + [\mathcal{S}, \mathcal{A}])f \bar{f} dx dt \\
&= 32\mu^3 \iint |f|^2 \left| x + Rt(1-t)e_1 - \frac{R}{16\mu^2}e_1 \right|^2 + \frac{\epsilon R^2}{8\mu} \iint |f|^2 + 8\mu \iint |\nabla_{A,x'}f|^2 \\
&\quad + 8\mu \iint \left| \partial_A^1 f + i \frac{R(1-2t)}{2} f \right|^2 + 8\mu \Im \iint f (x + Rt(1-t)e_1)^t B \cdot \overline{\nabla_A f},
\end{aligned}$$

where  $\nabla_A = \nabla - iA := (\partial_A^1, \dots, \partial_A^n)$ ,  $\nabla_{A,x'} := (0, \partial_A^2, \dots, \partial_A^n)$ . Notice that, since  $e_1^t B = 0$  and  $B$  is anti-symmetric, we can write

$$\begin{aligned}
(4.6) \quad & f (x + Rt(1-t)e_1)^t B \cdot \overline{\nabla_A f} = f x^t B \cdot \overline{\nabla_A f} = f x^t B \cdot \left( \overline{\nabla_A f} + i \frac{R(1-2t)}{2} e_1 f \right) \\
(4.7) \quad & = f x^t B \cdot \overline{\nabla_{A,x'} f} + f x^t B \cdot \left( \partial_A^1 f + i \frac{R(1-2t)}{2} f \right) e_1.
\end{aligned}$$

Therefore, by Cauchy-Schwartz and the elementary inequality  $ab \leq \delta a^2 + \frac{1}{4\delta} b^2$ , with the choice  $\delta := 8\mu$ , we can estimate

$$\begin{aligned}
(4.8) \quad & \left| 8\mu \Im \iint f (x + Rt(1-t)e_1)^t B \cdot \overline{\nabla_A f} \right| \\
&\leq 4\mu \|x^t B\|_{L^\infty}^2 \iint |f|^2 + 8\mu \iint \left| \partial_A^1 f + i \frac{R(1-2t)}{2} f \right|^2 + 8\mu \iint |\nabla_{A,x'} f|^2.
\end{aligned}$$

In conclusion, by (4.5) and (4.8), neglecting the term with cubic growth in  $\mu$  we get

$$\iint (\mathcal{S}_t + [\mathcal{S}, \mathcal{A}])f \bar{f} \geq \left[ \frac{\epsilon R^2}{8\mu} - 4\mu \|x^t B\|_{L^\infty}^2 \right] \iint |f|^2.$$

The last inequality, together with (4.3), (4.4) and the condition  $R > 8\mu\epsilon^{-\frac{1}{2}} \|x^t B\|_{L^\infty}$ , completes the proof of (4.2).  $\square$

*Proof of Theorem 1.1.* With the tools introduced up to now, the proof of Theorem 1.1 is now reduced to a typical argument in the Carleman's spirit.



Let  $u \in \mathcal{C}([0, 1]; L^2(\mathbb{R}^n))$  be the solution to (1.2). As in the first step of the previous section, we first reduce to the Cronström gauge, passing from  $A$  to  $\tilde{A}$  by means of Lemma 2.2. It is hence sufficient to prove that  $\tilde{u} = e^{i\varphi}u \equiv 0$ , where  $\varphi$  is given by (2.4). From now on, by abuse of notation, we keep calling  $u$  the gauged function  $\tilde{u}$  and by  $A$  the transformed potential  $\tilde{A}$ , which satisfy identities (2.5), (2.6), (2.7).

Now apply the Appell transformation (Lemma 2.7) with  $a = 0$  and  $b = 1$ , to obtain the new function  $\tilde{u}$  in (2.11), satisfying

$$\partial_t \tilde{u} = i \left( \Delta_{\tilde{A}} u + \tilde{V} \tilde{u} \right),$$

where  $\tilde{A}$  and  $\tilde{V}$  are defined by (2.13) and (2.14), respectively, and  $V := V_1 + V_2$ . Assumption (1.8) then gives that  $\|e^{\gamma|x|^2} \tilde{u}(0)\|_{L^2} + \|e^{\gamma|x|^2} \tilde{u}(1)\|_{L^2} < \infty$ , for any  $\gamma > \frac{1}{2}$ .

In addition, by estimates (3.38) and (3.39), in the limit as  $\epsilon$  tends to 0, we have

$$(4.9) \quad \sup_{t \in [0, 1]} \left\| e^{\gamma|\cdot|^2} \tilde{u}(\cdot, t) \right\|_{L^2} + \left\| \sqrt{t(1-t)} e^{\gamma|\cdot|^2} \nabla_{\tilde{A}} \tilde{u}(\cdot, t) \right\|_{L^2(\mathbb{R}^n \times [0, 1])} =: N_\gamma < \infty.$$

Now, let  $R > 8\mu\epsilon^{-\frac{1}{2}} \|x^t B\|_{L^\infty}$ , as in the statement of Lemma 4.1, and let  $M > 0$ , to be chosen later. Then, localize the function  $\tilde{u}$  as follows: let  $\theta_M(x), \eta_R(t)$  be two smooth functions such that

$$\theta_M \equiv 1 \text{ if } |x| \leq M \quad \theta_M \equiv 0 \text{ if } |x| \geq 2M$$

$$\eta_R(t) \equiv 1 \text{ if } t \in \left[ \frac{1}{R}, 1 - \frac{1}{R} \right] \quad \eta_R(t) \equiv 0 \text{ if } t \in \left[ 0, \frac{1}{2R} \right] \cup \left[ 1 - \frac{1}{2R}, 1 \right],$$

and define

$$g(x, t) = \theta_M(x) \eta_R(t) \tilde{u}(x, t).$$

It turns out that  $g$  solves

$$(4.10) \quad (\partial_t - i\Delta_{\tilde{A}}) g = i\tilde{V}g + \theta_M \eta'_R \tilde{u} - i(2\nabla\theta_M \cdot \nabla_{\tilde{A}} \tilde{u} + \tilde{u}\Delta\theta_M) \eta_R.$$

Assume without loss of generality that the magnetic field  $B$  satisfies the condition (1.4) with  $v = e_1$ .

Now choose

$$(4.11) \quad \mu \leq \frac{\gamma}{1 + \epsilon},$$

for some fixed small  $\epsilon > 0$ . Notice that, in the support of the second term of the right-hand side of (4.10), we have

$$(4.12) \quad \mu |x + Rt(1-t)e_1|^2 - \frac{(1+\epsilon)R^2 t(1-t)}{16\mu} \leq \gamma |x|^2 + \frac{\gamma}{\epsilon};$$

analogously, in the support of the last term of the right-hand side of (4.10) we have

$$(4.13) \quad \mu |x + Rt(1-t)e_1|^2 - \frac{(1+\epsilon)R^2 t(1-t)}{16\mu} \leq \gamma |x|^2 + \frac{\gamma R^2}{\epsilon}.$$

By condition (1.4) with  $v = e_1$ , (2.6), (2.7), identity (3.35) with  $\epsilon = 0$  and the fact that  $B$  is anti-symmetric, we get  $x \cdot \partial_t \tilde{A} \equiv 0 \equiv e_1 \cdot \partial_t \tilde{A}$ . Hence, applying (4.2) to  $g$ ,

by (4.10), (4.12), (4.13) and the bounds for  $\theta_M, \eta_R$  and their derivatives we easily get

$$(4.14) \quad \begin{aligned} & R \left\| e^{\mu|x+Rt(1-t)v|^2 - \frac{(1+\epsilon)R^2t(1-t)}{16\mu}} g \right\|_{L^2(\mathbb{R}^n \times [0,1])} \\ & \leq N_{\epsilon,\mu} \left\| \tilde{V} \right\|_{L^\infty(\mathbb{R}^n \times [0,1])} \left\| e^{\mu|x+Rt(1-t)v|^2 - \frac{(1+\epsilon)R^2t(1-t)}{16\mu}} g \right\|_{L^2(\mathbb{R}^n \times [0,1])} \\ & + N_{\epsilon,\mu} R e^{\frac{\gamma}{2}} \sup_{[0,1]} \left\| e^{\gamma|\cdot|^2} \tilde{u}(\cdot, t) \right\|_{L^2} + N_\epsilon M^{-1} e^{\frac{\gamma R^2}{\epsilon}} \left\| e^{\gamma|x|^2} (|\tilde{u}| + |\nabla_{\tilde{A}} \tilde{u}|) \right\|_{L^2(\mathbb{R}^n \times [\frac{1}{2R}, 1 - \frac{1}{2R}])}, \end{aligned}$$

with  $N_{\epsilon,\mu} = 4\sqrt{\mu/\epsilon}$ . Notice that, choosing  $R \geq 2N_\epsilon \left\| \tilde{V} \right\|_{L^\infty(\mathbb{R}^n \times [0,1])}$ , the first term in the right-hand side of the last inequality can be hidden in the left-hand side. Moreover, by (4.9), we have that

$$(4.15) \quad \lim_{M \rightarrow \infty} N_\epsilon M^{-1} e^{\frac{\gamma R^2}{\epsilon}} \left\| e^{\gamma|x|^2} (|\tilde{u}| + |\nabla_{\tilde{A}} \tilde{u}|) \right\|_{L^2(\mathbb{R}^n \times [\frac{1}{2R}, 1 - \frac{1}{2R}])} = 0,$$

for any fixed  $R$ . Finally, choose

$$M := f(\epsilon) \frac{R}{8},$$

for some positive function  $f(\epsilon)$  such that  $f(\epsilon) < 1 - \epsilon^2$  and  $f(\epsilon) \rightarrow 0$  as  $\epsilon$  tends to 0. Notice that  $g \equiv \tilde{u}$  in  $B_{f(\epsilon)R/8 \times [(1-\epsilon)/2, (1+\epsilon)/2]}$ ; in this set, one can easily estimate

$$\begin{aligned} & \mu |x + Rt(1-t)v|^2 - \frac{(1+\epsilon)R^2t(1-t)}{16\mu} \\ & \geq \frac{R^2}{16\mu} \left\{ \mu^2 [(1-\epsilon^2)^2 - (1-\epsilon^2)f(\epsilon)] - \frac{1}{4}(1+\epsilon) \right\}. \end{aligned}$$

Consequently, choosing

$$(4.16) \quad \mu^2 > \frac{1}{4} \cdot \frac{1+\epsilon}{(1-\epsilon^2)^2 - (1-\epsilon^2)f(\epsilon)}$$

one obtains that

$$\mu |x + Rt(1-t)v|^2 - \frac{(1+\epsilon)R^2t(1-t)}{16\mu} \geq 0$$

in  $B_{f(\epsilon)R/8 \times [(1-\epsilon)/2, (1+\epsilon)/2]}$ , in which we also have  $g \equiv \tilde{u}$ . Comparing (4.11) and (4.16), we see that they are compatible if and only if  $\gamma > \frac{1}{2}$ , i.e.  $\alpha\beta > 2$ , as required in the statement of Theorem 1.1.

Therefore, by (4.14), and the above considerations, there exist  $C(\gamma, \epsilon), N_{\gamma,\epsilon} > 0$  such that

$$(4.17) \quad R e^{C(\gamma,\epsilon)R^2} \left\| \tilde{u}(x, t) \right\|_{L^2\left(B_{\frac{R}{8}} \times [\frac{1-\epsilon}{2}, \frac{1+\epsilon}{2}]\right)} \leq N_{\gamma,\epsilon} R,$$

for any  $R > \max\{8\mu\epsilon^{-\frac{1}{2}} \|x^t B\|_{L^\infty}, 2N_\epsilon \|\tilde{V}\|_{L^\infty(\mathbb{R}^n \times [0,1])}\}$ . By (3.22) in Lemma 3.2, (4.9) and (4.17) we now conclude that there exists a constant  $N = N(\gamma, \epsilon, V)$  depending on  $N_\gamma, \epsilon$  and  $\sup_{[0,1]} \|V\|_{L^\infty}$  such that

$$e^{C(\gamma,\epsilon)R^2} \left\| \tilde{u}(\cdot, 0) \right\|_{L^2} \leq N(\gamma, \epsilon, V).$$

Letting  $R$  tend to infinity, this implies that  $\tilde{u} \equiv u \equiv 0$ .  $\square$

## REFERENCES

- [1] Bonami, A., and Demange, B., A survey on uncertainty principles related to quadratic forms. *Collect. Math.* 2006, Vol. Extra, 1–36.
- [2] COWLING, M., ESCAURIAZA, L., KENIG, C., PONCE, G., AND VEGA, L., The Hardy Uncertainty Principle Revisited, *Indiana U. Math. J.* **59** (2010), no. 6, 2007–2026.
- [3] ESCAURIAZA, L., FANELLI, L., AND VEGA, L., Carleman estimates and necessary conditions for the existence of waveguides, to appear on *Indiana Univ. Math. J.*
- [4] ESCAURIAZA, L., KENIG, C., PONCE, G., AND VEGA, L., On Uniqueness Properties of Solutions of Schrödinger Equations, *Comm. PDE.* **31** (2006), no. 12, 1811–1823.
- [5] ESCAURIAZA, L., KENIG, C., PONCE, G., AND VEGA, L., Convexity properties of solutions to the free Schrödinger equation with Gaussian decay, *Math. Res. Lett.* **15** (2008), no. 5, 957–971.
- [6] ESCAURIAZA, L., KENIG, C., PONCE, G., AND VEGA, L., Hardy’s uncertainty principle, convexity and Schrödinger evolutions, *J. Eur. Math. Soc. (JEMS)* **10** (2008), no. 4, 883–907.
- [7] ESCAURIAZA, L., KENIG, C., PONCE, G., AND VEGA, L., The sharp Hardy uncertainty principle for Schrödinger evolutions, *Duke Math. J.* **155** (2010), no. 1, 163–187.
- [8] ESCAURIAZA, L., KENIG, C., PONCE, G., AND VEGA, L., Uncertainty principle of Morgan type and Schrödinger evolutions, *J. Lond. Math. Soc. (2)* **83** (2011), no. 1, 187–207.
- [9] ESCAURIAZA, L., KENIG, C., PONCE, G., AND VEGA, L., Unique continuation for Schrödinger evolutions, with applications to profiles of concentration and traveling waves, *Comm. Math. Phys.* **305** (2011), no. 2, 487–512.
- [10] FANELLI, L., AND VEGA, L., Magnetic virial identities, weak dispersion and Strichartz inequalities, *Math. Ann.* **344** (2009), no. 2, 249–278.
- [11] HÖRMANDER, L., *Linear partial differential operators*, Springer, Berlin, (1969).
- [12] IWATSUKA, A., Spectral representation for Schrödinger operators with magnetic vector potentials, *J. Math. Kyoto Univ.* **22**, (1982), no. 2, 223–242.
- [13] PAZY, A., *Semigroups of linear operators with application to partial differential equations*, Springer-Verlag, Berlin, New York (1983).
- [14] REED, M., AND SIMON, B., *Methods of Modern Mathematical Physics vol. I: Functional Analysis*. Academic Press, New York, San Francisco, London 1980.
- [15] SIMON, B., Schrödinger semigroups, *Bull. AMS* **7** (1982), 447–526.
- [16] SITARAM, A., SUNDARI, M., AND THANGAVELU, S., *Uncertainty principles on certain Lie groups*, *Proc., Indian Acad. Sci. Math. Sci.* **105** (1995), 135–151.
- [17] STEIN, E.M., AND SHAKARCHI, R., *Princeton Lecture in Analysis II. Complex Analysis*, Princeton University Press.

JUAN ANTONIO BARCELÓ: ETSI DE CAMINOS, UNIVERSIDAD POLITÉCNICA DE MADRID, 28040 MADRID, SPAIN

*E-mail address:* [juanantonio.barcelo@upm.es](mailto:juanantonio.barcelo@upm.es)

LUCA FANELLI: SAPIENZA UNIVERSITÀ DI ROMA, DIPARTIMENTO DI MATEMATICA, P.LE A. MORO 5, 00185-ROMA, ITALY

*E-mail address:* [fanelli@mat.uniroma1.it](mailto:fanelli@mat.uniroma1.it)

SUSANA GUTIÉRREZ: SCHOOL OF MATHEMATICS, THE WATSON BUILDING, UNIVERSITY OF BIRMINGHAM, EDGBASTON, BIRMINGHAM, B15 2TT, ENGLAND

*E-mail address:* [S.Gutierrez@bham.ac.uk](mailto:S.Gutierrez@bham.ac.uk)

ALBERTO RUIZ: DEPARTAMENTO DE MATEMÁTICAS, UNIVERSIDAD AUTÓNOMA DE MADRID, 28049, MADRID, SPAIN

*E-mail address:* [alberto.ruiz@uam.es](mailto:alberto.ruiz@uam.es)

MARI CRUZ VILELA: UNIVERSIDAD DE VALLADOLID, DEPARTAMENTO DE MATEMÁTICA APLICADA, E.U. DE INFORMÁTICA, PLZA. DE SANTA EULALIA 9 Y 11, 40005 SEGOVIA, SPAIN.

*E-mail address:* [maricruz@eii.uva.es](mailto:maricruz@eii.uva.es)