

# Conflicts with transcription make early replication late

Petermann, Eva

DOI:

[10.1016/j.molcel.2022.08.026](https://doi.org/10.1016/j.molcel.2022.08.026)

License:

Creative Commons: Attribution-NonCommercial-NoDerivs (CC BY-NC-ND)

*Document Version*

Peer reviewed version

*Citation for published version (Harvard):*

Petermann, E 2022, 'Conflicts with transcription make early replication late', *Molecular Cell*, vol. 82, no. 18, pp. 3315-3317. <https://doi.org/10.1016/j.molcel.2022.08.026>

[Link to publication on Research at Birmingham portal](#)

## General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

## Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact [UBIRA@lists.bham.ac.uk](mailto:UBIRA@lists.bham.ac.uk) providing details and we will remove access to the work immediately and investigate.

## **Conflicts with transcription make early replication late**

Eva Petermann

Institute of Cancer and Genomic Sciences, University of Birmingham, Birmingham B15 2TT, United Kingdom

Birmingham Centre for Genome Biology, University of Birmingham, Birmingham B15 2TT, United Kingdom

**By sequencing sites of mitotic DNA synthesis in cells lacking homologous recombination, Groelly et al. (2022) and Bhowmick et al. (2022) show how conflicts between transcription and replication in early S phase can cause under-replicated DNA to persist into mitosis.**

Genomic instability is a potential driving force in cancer development, and DNA replication stress has been implicated in contributing to genomic instability in tumour cells. Replication stress can result from oncogene activation or loss of tumour suppressors. In this issue of *Molecular Cell*, two reports by Groelly, Bhowmick and colleagues shed new light on the genomic features and molecular mechanisms of replication stress in cells lacking the tumour suppressor pathway homologous recombination (HR) (Bhowmick et al., 2022; Groelly et al., 2022).

Replication stress occurs when replication forks slow down, stall or even speed up. This can lead to subsequent problems during mitosis, which exacerbate genomic instability. Common fragile sites (CFS) are difficult-to-replicate, late-replicating genomic regions that are sensitive to mild replication stress (Debatisse et al., 2012). The high frequency of genomic rearrangements at CFS in cancer samples provides some of the strongest evidence for the importance of replication stress in cancer (Dereli-Oz et al., 2011). Under conditions of mild replication inhibition, CFS enter mitosis before completing replication. This results in mitotic DNA synthesis (MiDAS), a specialised break-induced replication pathway using RAD52 that helps prevent genomic instability (Minocherhomji et al., 2015). The phenomenon of MiDAS further emphasises the connection between late-replicating genomic regions and the mitotic effects of replication stress. However, early-replicating genomic regions can also be especially susceptible to damage, especially under high levels of replication stress (Barlow et al., 2013).

The new studies by Groelly and Bhowmick use cutting-edge sequencing approaches to provide evidence that early replicating genomic regions can be hotspots of replication stress phenotypes manifesting in mitosis. They exploit MiDAS to understand how replication stress contributes to genomic instability in cells lacking critical HR proteins. HR, catalysed by the recombinase RAD51, can rescue and repair stalled replication forks (Wassing and Esashi, 2021). Several HR factors also prevent and resolve conflicts between transcription and replication (TRCs). The tumour suppressor BRCA2, involved in RAD51 loading onto DNA, can counteract the accumulation of replication-blocking RNA-DNA hybrids called R-loops (Bhatia et al., 2014). Groelly et al. use human H1299 non-small cell lung carcinoma cells shRNA-depleted for BRCA2, while Bhowmick et al. siRNA-deplete RAD51 in human U2OS osteosarcoma cells.

Both studies use previously established cell synchronisation and EdU-labelling protocols to measure DNA synthesis during mitosis in cells that have experienced replication stress in the preceding S phase, either due to low-dose Aphidicolin treatment or BRCA2 or RAD51 depletion (Figure 1). They then employ a newly developed method, MiDAS-seq, where the EdU-labelled genomic areas are isolated and sequenced for high resolution mapping (Macheret et al., 2020). Intelligent combination with existing datasets allows comparison of the genomic features of MiDAS sites in HR-deficient cells and canonical MiDAS sites, which are induced by Aphidicolin.

The most important finding is that MiDAS regions in HR-deficient cells, which the authors call “atypical” MiDAS sites (Bhowmick et al., 2022), are vastly different from the canonical MiDAS sites. They are in early replicating regions that are also transcribed in early S phase, and are characterised by open chromatin, high density of replication origins, the presence of short- to average-length genes and increased R-loop formation. These features, combined with known roles of HR factors, point towards TRCs as a potential cause of MiDAS. Indeed, both studies provide evidence for such conflicts. Groelly et al. use existing datasets from DNA-RNA immunoprecipitation sequencing (DRIP-seq) as support for atypical MiDAS events mapping to regions of R-loop formation, with the caveat that the datasets were obtained in different cell lines. They then rescue MiDAS in BRCA2-deficient cells with RNase H1 overexpression. Bhowmick et al. quantify both nascent and steady-state transcription levels by combining the high throughput sequencing approaches SLAM-seq and QUANT-seq. This reveals that genes up-regulated in early S-phase are over-represented among MiDAS sites in HR-deficient cells. Both studies show that these MiDAS events require transcription in early S phase. Other lines of evidence include increased co-localisation of replication and transcription factors in early S phase combined with transcription-dependent RAD51 recruitment in S-phase. Therefore, the data suggest that both BRCA2 and RAD51 are required to prevent conflicts between transcription and replication at early replicating genomic regions.

CFS are origin-poor and have been associated with a lack of backup replication origins as well as TRCs (Debatisse et al., 2012). The present studies agree with previous suggestions that TRCs are the main source of stress in early replicating genomic regions (Barlow et al., 2013).

New questions arise from this work. Why do HR-deficient cells display in no canonical MiDAS at CFS, even after Aphidicolin treatment (Bhowmick et al., 2022)? The authors suggest that in absence of RAD51, the late-replicating nature of CFS may allow for collapsed replication forks to be rescued by RAD52-dependent break-induced replication in G2 (Figure 1). Groelly et al. show that transcription and replication are predominantly co-directional at MiDAS sites in HR-deficient cells, and that the replication stress in these cells involves ATM signalling and 53BP1 foci formation, consistent with DNA double-strand break (DSB) formation at co-directional TRCs. Interestingly, DSB formation and RAD51-independent, RAD52-mediated fork restart have been proposed to occur at TRCs involving R-loops

(Chappidi et al., 2020). Similar mechanisms could occur, possibly at a higher rate, in BRCA2- and RAD51-deficient cells, with replication timing affecting whether this leads to MiDAS (Figure 1). In addition, Bhowmick et al. report that RAD51 recruitment in early S phase can be induced by overexpressing oncogenes MYC and Cyclin E. The authors speculate that “atypical” MiDAS sites in early replicating regions may be prone to amplification in cancer, and both studies provide evidence for such amplifications in the genomes of BRCA2-mutated breast cancers. These findings suggest new directions for investigating the mechanisms around TRCs at early replicating regions, and the role of HR in preventing and resolving these conflicts.

To conclude, in cells lacking tumour suppressors, unresolved TRCs in early replicating regions can persist through S phase and into mitosis, resulting in MiDAS. The findings draw exciting new connections between transcription, replication, DNA repair and the cell cycle, with potential implications for genomic instability in cancer.

### Figure legend

**A:** Summary of the experimental approach. Cells are synchronised and EdU labelled, followed by next generation sequencing of EdU-containing DNA. Hydroxyurea (HU) and Aphidicolin treatment protocols differ between the studies.

**B:** In the canonical MiDAS model, insufficient time to resolve replication stress, such as transcription-replication conflicts (TRCs), in late-replicating regions leads to DNA synthesis in mitosis. In the new model for MiDAS in homologous recombination (HR)-deficient cells, unresolved TRCs in early S phase persist and lead to mitotic DNA synthesis, while stress in late-replicating regions might be resolved by RAD52-dependent repair (dotted lines).

### Declaration of interests

The author declares no competing interests.

### Acknowledgements

E.P. is funded by Cancer Research UK (G25526/A28275) and Medical Research Council (MR/S021310/1, MR/W00190X/1, MR/W031442/1).

### References

Barlow, J.H., Faryabi, R.B., Callén, E., Wong, N., Malhowski, A., Chen, H.T., Gutierrez-Cruz, G., Sun, H.-W., McKinnon, P., Wright, G., *et al.* (2013). Identification of early replicating fragile sites that contribute to genome instability. *Cell* 152, 620-632.

Bhatia, V., Barroso, S.I., Garcia-Rubio, M.L., Tumini, E., Herrera-Moyano, E., and Aguilera, A. (2014). BRCA2 prevents R-loop accumulation and associates with TREX-2 mRNA export factor PCID2. *Nature* 511, 362-365.

Bhowmick, R., Lerdrup, M., Gadi, S.A., Rossetti, G.G., Singh, M.I., Liu, Y., Halazonetis, T.D., and Hickson, I.D. (2022). Rad51 protects human cells from transcription-replication conflicts. *Molecular cell*.

Chappidi, N., Nascakova, Z., Boleslavskaja, B., Zellweger, R., Isik, E., Andrs, M., Menon, S., Dobrovolna, J., Balbo Pogliano, C., Matos, J., *et al.* (2020). Fork Cleavage-Religation Cycle and Active Transcription Mediate Replication Restart after Fork Stalling at Co-transcriptional R-Loops. *Mol Cell* 77, 528-541 e528.

Debatisse, M., Le Tallec, B., Letessier, A., Dutrillaux, B., and Brison, O. (2012). Common fragile sites: mechanisms of instability revisited. *Trends Genet* 28, 22-32.


Dereli-Oz, A., Versini, G., and Halazonetis, T.D. (2011). Studies of genomic copy number changes in human cancers reveal signatures of DNA replication stress. *Mol Oncol* 5, 308-314.

Groelly, F.J., Dagg, R.A., Petropoulos, M., Rossetti, G.G., Prasad, B., Panagopoulos, A., Paulsen, T., Karamichali, A., Jones, S.E., Ochs, F., *et al.* (2022). Mitotic DNA Synthesis Is caused by transcription-replication conflicts in BRCA2-deficient cells. *Molecular cell*.

Macheret, M., Bhowmick, R., Sobkowiak, K., Padayachy, L., Mailler, J., Hickson, I.D., and Halazonetis, T.D. (2020). High-resolution mapping of mitotic DNA synthesis regions and common fragile sites in the human genome through direct sequencing. *Cell Res* 30, 997-1008.


Minocherhomji, S., Ying, S., Bjerregaard, V.A., Bursomanno, S., Aleliunaite, A., Wu, W., Mankouri, H.W., Shen, H., Liu, Y., and Hickson, I.D. (2015). Replication stress activates DNA repair synthesis in mitosis. *Nature* 528, 286-290.

Wassing, I.E., and Esashi, F. (2021). RAD51: Beyond the break. *Semin Cell Dev Biol* 113, 38-46.

**A**

EdU-sequencing

**Canonical MiDAS sites**      **MiDAS sites in HR-deficient cells**



**Common fragile sites**  
Late-replicating  
Long genes  
Few origins

**Rich in R-loops**  
Early-replicating  
Short genes  
Rich in origins



**B****Canonical MiDAS****MiDAS in HR-deficient cells**

Figure 1