

On the Strichartz estimates for the kinetic transport equation

Bennett, Jonathan; Bez, Neal; Gutierrez, Susana; Lee, Sanghyuk

DOI:

[10.1080/03605302.2013.850880](https://doi.org/10.1080/03605302.2013.850880)

License:

Creative Commons: Attribution (CC BY)

Document Version

Publisher's PDF, also known as Version of record

Citation for published version (Harvard):

Bennett, J, Bez, N, Gutierrez, S & Lee, S 2014, 'On the Strichartz estimates for the kinetic transport equation', *Communications in Partial Differential Equations*, vol. 39, no. 10, pp. 1821-1826.
<https://doi.org/10.1080/03605302.2013.850880>

[Link to publication on Research at Birmingham portal](#)

Publisher Rights Statement:

Eligibility for repository : checked 12/09/2014

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.


If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

This article was downloaded by: [University of Birmingham]

On: 12 September 2014, At: 06:41

Publisher: Taylor & Francis

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Communications in Partial Differential Equations

Publication details, including instructions for authors and subscription information:

<http://www.tandfonline.com/loi/lpde20>

On the Strichartz Estimates for the Kinetic Transport Equation

Jonathan Bennett^a, Neal Bez^b, Susana Gutiérrez^a & Sanghyuk Lee^c

^a School of Mathematics, University of Birmingham, Birmingham, England

^b Department of Mathematics, Saitama University, Saitama, Japan

^c Department of Mathematical Sciences, Seoul National University, Seoul, Korea

Published online: 19 Aug 2014.

To cite this article: Jonathan Bennett, Neal Bez, Susana Gutiérrez & Sanghyuk Lee (2014) On the Strichartz Estimates for the Kinetic Transport Equation, Communications in Partial Differential Equations, 39:10, 1821-1826, DOI: [10.1080/03605302.2013.850880](https://doi.org/10.1080/03605302.2013.850880)

To link to this article: <http://dx.doi.org/10.1080/03605302.2013.850880>

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Versions of published Taylor & Francis and Routledge Open articles and Taylor & Francis and Routledge Open Select articles posted to institutional or subject repositories or any other third-party website are without warranty from Taylor & Francis of any kind, either expressed or implied, including, but not limited to, warranties of merchantability, fitness for a particular purpose, or non-infringement. Any opinions and views expressed in this article are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor & Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Terms & Conditions of access and use can be found at <http://www.tandfonline.com/page/terms-and-conditions>

It is essential that you check the license status of any given Open and Open Select article to confirm conditions of access and use.

On the Strichartz Estimates for the Kinetic Transport Equation

JONATHAN BENNETT¹, NEAL BEZ², SUSANA GUTIÉRREZ¹, AND SANGHYUK LEE³

¹School of Mathematics, University of Birmingham, Birmingham, England

²Department of Mathematics, Saitama University, Saitama, Japan

³Department of Mathematical Sciences, Seoul National University, Seoul, Korea

We show that the endpoint Strichartz estimate for the kinetic transport equation is false in all dimensions. We also present an alternative approach to proving the non-endpoint cases using multilinear analysis.

Keywords Endpoint estimates; Kinetic transport equations; Multilinear analysis; Strichartz estimates.

Mathematics Subject Classification 42B37.

1. Introduction

The solution of the kinetic transport equation

$$\partial_t f(t, x, v) + v \cdot \nabla_x f(t, x, v) = 0, \quad f(0, x, v) = f^0(x, v)$$

for $(t, x, v) \in \mathbb{R} \times \mathbb{R}^d \times \mathbb{R}^d$, satisfies the Strichartz estimates

$$\|f\|_{L_t^q L_x^r L_v^r} \lesssim \|f^0\|_{L_{x,v}^a}, \quad (1.1)$$

where

$$\frac{2}{q} = d \left(\frac{1}{r} - \frac{1}{p} \right), \quad \frac{1}{a} = \frac{1}{2} \left(\frac{1}{r} + \frac{1}{p} \right), \quad q > a, \quad p \geq a. \quad (1.2)$$

(We write $X \lesssim Y$ and $Y \gtrsim X$ if $X \leq CY$ for some finite constant C depending at most on the parameters (q, p, r, a, d) , and $X \sim Y$ if $X \lesssim Y$ and $X \gtrsim Y$.) With

© Jonathan Bennett, Neal Bez, Susana Gutiérrez and Sanghyuk Lee

Received August 28, 2013; Accepted September 23, 2013

Address correspondence to Neal Bez, Department of Mathematics, Graduate School of Science and Engineering, Saitama University, Saitama 338-8570, Japan; E-mail: nealbez@mail.saitama-u.ac.jp

(q, p, r, a) satisfying (1.2), but with the further condition $q > 2 \geq a$, this was proved by Castella and Perthame [1], and it was observed by Keel and Tao [5] that this latter condition can be relaxed to $q > a$ and hence (1.2) suffices. In [5] it is tentatively conjectured that the Strichartz estimate (1.1) holds at the endpoint $q = a$, at least for $d > 1$. Using the invariance under the transformations

$$f^0 \leftrightarrow (f^0)^\lambda, \quad f \leftrightarrow f^\lambda, \quad (q, p, r, a) \leftrightarrow \left(\frac{q}{\lambda}, \frac{p}{\lambda}, \frac{r}{\lambda}, \frac{a}{\lambda}\right) \quad (1.3)$$

this conjectured endpoint can be (and usually is) stated for initial data in $L^2_{x,v}$ as

$$\|f\|_{L_t^2 L_x^{\frac{2d}{d-1}} L_v^{\frac{2d}{d+1}}} \lesssim \|f^0\|_{L^2_{x,v}}. \quad (1.4)$$

The main purpose of this paper is to disprove this conjecture.

Theorem 1. *The endpoint Strichartz estimate (1.4) for the kinetic transport equation fails for all $d \geq 1$.*

The case $d = 1$ of Theorem 1 was proved by Guo and Peng [4] and later by Ovcharov [7] using different arguments (where the norm in x on the left-hand side is L^∞_x). The argument we present uses key ideas from recent work of Frank et al. [3].

We remark that for the free Schrödinger propagator, the endpoint Strichartz estimate fails in the case $d = 2$ (see [6]) but is true for all $d > 2$, as shown in the landmark paper of Keel and Tao [5]. Thus, Theorem 1 highlights a fundamental difference in the Strichartz estimates for these related equations.

In the next section, we prove Theorem 1. In the final section, we provide an alternative proof of the Strichartz estimates (1.1) in all non-endpoint cases using multilinear analysis.

2. Proof of Theorem 1

Using the invariance under the transformation (1.3) with $\lambda = \frac{2d}{d+1}$, estimate (1.4) is equivalent to

$$\|f\|_{L_t^{\frac{d+1}{d}} L_x^{\frac{d+1}{d-1}} L_v^1} \lesssim \|f^0\|_{L^{\frac{d+1}{d}}_{x,v}}. \quad (2.1)$$

Since $f(t, x, v) = f^0(x - tv, v)$, it is clear that (2.1) implies

$$\|\rho(f^0)\|_{L_t^{\frac{d+1}{d}} L_x^{\frac{d+1}{d-1}}} \lesssim \|f^0\|_{L^{\frac{d+1}{d}}_{x,v}}, \quad (2.2)$$

where $\rho(f^0)$ is the macroscopic density defined by the linear mapping

$$\rho(f^0)(t, x) = \int_{\mathbb{R}^d} f^0(x - tv, v) dv.$$

Hence, by duality, (2.2) implies

$$\|\rho^* g\|_{L^{d+1}_{x,v}} \lesssim \|g\|_{L_t^{d+1} L_x^{\frac{d+1}{2}}}, \quad (2.3)$$

where the adjoint ρ^* is given by

$$\rho^* g(x, v) = \int_{\mathbb{R}} g(t, x + tv) dt.$$

From here, the argument strongly uses ideas from the paper of Frank et al. [3] concerning refined Strichartz estimates for the free Schrödinger propagator associated with orthonormal initial data.

Suppose $g \in \mathcal{S}(\mathbb{R} \times \mathbb{R}^d) \setminus \{0\}$ is nonnegative and such that $\widehat{g} \in C_c^\infty(\mathbb{R} \times \mathbb{R}^d)$ is also nonnegative. Here, we use \widehat{g} to denote the space-time Fourier transform of g given by

$$\widehat{g}(\tau, \xi) = \int_{\mathbb{R} \times \mathbb{R}^d} g(t, x) e^{-i(t\tau + x \cdot \xi)} dt dx.$$

In this proof, we shall also use c to denote a constant depending on at most d , which may change from line to line.

Proceeding formally, using Fourier inversion we get

$$\begin{aligned} \|\rho^* g\|_{L_{x,v}^{d+1}}^{d+1} &= \int \prod_{j=1}^{d+1} g(t_j, x + t_j v) d\vec{t} dx dv \\ &= c \int \prod_{j=1}^{d+1} \widehat{g}(\tau_j, \xi_j) \prod_{k=1}^{d+1} e^{it_k(\tau_k + v \cdot \xi_k)} e^{ix \cdot \sum_{\ell=1}^{d+1} \xi_\ell} d\vec{t} d\vec{\xi} d\vec{t} dx dv \\ &= c \int \prod_{j=1}^{d+1} \widehat{g}(\tau_j, \xi_j) \prod_{k=1}^{d+1} \delta(\tau_k + v \cdot \xi_k) \delta\left(\sum_{\ell=1}^{d+1} \xi_\ell\right) d\vec{t} d\vec{\xi} dv \\ &= c \int \prod_{j=1}^{d+1} \widehat{g}(-v \cdot \xi_j, \xi_j) \delta\left(\sum_{k=1}^{d+1} \xi_k\right) d\vec{\xi} dv \end{aligned}$$

and hence

$$\|\rho^* g\|_{L_{x,v}^{d+1}}^{d+1} = c \int \prod_{j=1}^d \widehat{g}(-v \cdot \xi_j, \xi_j) \widehat{g}\left(v \cdot \sum_{k=1}^d \xi_k, -\sum_{\ell=1}^d \xi_\ell\right) d\vec{\xi} dv. \quad (2.4)$$

We remark that by appropriately truncating the integrals in the above identities and limiting arguments, (2.4) makes sense in $[0, \infty]$ for the class of g under consideration.

Define K to be the d by d matrix whose consecutive rows are $-\xi_1, \dots, -\xi_d$. Using the change of variables $w = Kv$, so that $w_j = -\xi_j \cdot v$ for each $1 \leq j \leq d$, we obtain

$$\|\rho^* g\|_{L_{x,v}^{d+1}}^{d+1} = c \int \prod_{j=1}^d \widehat{g}(w_j, \xi_j) \widehat{g}\left(-\sum_{k=1}^d w_k, -\sum_{\ell=1}^d \xi_\ell\right) \frac{1}{|\det K|} d\vec{w} d\vec{\xi}.$$

Writing each $\xi_j = r_j \theta_j$ in polar coordinates, we have

$$|\det K| = \left(\prod_{j=1}^d r_j\right) |\det(\theta_1 \cdots \theta_d)|.$$

Since $\widehat{g}(0, 0) > 0$ and \widehat{g} is continuous, it follows that

$$\|\rho^* g\|_{L_{x,v}^{d+1}}^{d+1} \gtrsim \int_{|r| \lesssim 1} \int_{(\mathbb{S}^{d-1})^d} \left(\prod_{j=1}^d r_j^{d-2} \right) \frac{1}{|\det(\theta_1 \cdots \theta_d)|} d\vec{r} d\vec{\theta}. \quad (2.5)$$

For $d = 1$ the radial integral is infinite, and for $d \geq 2$,

$$\int_{(\mathbb{S}^{d-1})^d} \frac{1}{|\det(\theta_1 \cdots \theta_d)|} d\vec{\theta} = \infty,$$

so the angular integral is infinite. Hence, for all $d \geq 1$ we have shown that (2.3), and consequently (1.4), cannot hold.

Remark. The above argument shows that the endpoint estimate (2.3) fails rather generically. For example, the space-velocity norm $\|\rho^* g\|_{d+1}$ is infinite whenever $g \in \mathcal{S}(\mathbb{R} \times \mathbb{R}^d) \setminus \{0\}$ is nonnegative and such that $\widehat{g} \in C_c^\infty(\mathbb{R} \times \mathbb{R}^d)$ is also nonnegative.

3. A Multilinear Approach to the Non-Endpoint Cases

Fix $\sigma > 1$. In this section, the notation \lesssim allows, in addition, the implicit constant to depend on σ .

We shall prove

$$\|\rho^* g\|_{L_{x,v}^{\sigma(d+1)}} \lesssim \|g\|_{L_t^{q(\sigma)} L_x^{\frac{(d+1)\sigma}{2}}}, \quad (3.1)$$

for all $g \in L_t^{q(\sigma)} L_x^{\frac{(d+1)\sigma}{2}}$, where the exponent $q(\sigma)$ satisfies

$$\frac{1}{q(\sigma)} + \frac{d}{(d+1)\sigma} = 1.$$

Our argument is a multilinear variant of that in [1] and [9] (see also [8]), which deals with all non-endpoint cases directly. Similar multilinear arguments have been used recently in [3].

Using the invariance under transformations in (1.3), to prove the full range of non-endpoint Strichartz estimates, it suffices to consider (q, p, a) satisfying

$$q > a, \quad p \geq a, \quad \frac{2}{q} = d \left(1 - \frac{1}{p} \right), \quad \frac{1}{a} = \frac{1}{2} \left(1 + \frac{1}{p} \right) \quad (3.2)$$

and show that (1.1) holds with $r = 1$, or equivalently, that

$$\|\rho(f^0)\|_{L_t^q L_x^p} \lesssim \|f^0\|_{L_{x,v}^a}$$

holds for all $f^0 \in L_{x,v}^a$. By duality, this is equivalent to

$$\|\rho^* g\|_{L_{x,v}^{a'}} \lesssim \|g\|_{L_t^{q'} L_x^{p'}} \quad (3.3)$$

for all $g \in L_t^{q'} L_x^{p'}$. Note that (3.2) implies that $a' = 2p'$ and $\frac{1}{q'} + \frac{d}{a'} = 1$, in which case (3.3) reads

$$\|\rho^* g\|_{L_{x,v}^{a'}} \lesssim \|g\|_{L_t^{q'} L_x^{\frac{a'}{2}}},$$

and the condition $q > a$ is equivalent to $a' > d + 1$. Therefore, (3.1) with $\sigma = \frac{a'}{d+1} > 1$ implies the full range of non-endpoint Strichartz estimates (1.1).

Proof of (3.1). Without loss of generality, suppose g is nonnegative. By multiplying out and using Minkowski's integral inequality, we get

$$\begin{aligned} \|\rho^* g\|_{L_{x,v}^{\sigma(d+1)}}^{d+1} &= \left(\int \left(\int \prod_{j=1}^{d+1} g(t_j, x + t_j v) d\vec{t} \right)^\sigma dx dv \right)^{1/\sigma} \\ &\leq \int \left(\int \prod_{j=1}^{d+1} g(t_j, x + t_j v)^\sigma dx dv \right)^{1/\sigma} d\vec{t}. \end{aligned}$$

Now fix t_1, \dots, t_{d+1} and consider the multilinear form

$$\int \prod_{j=1}^{d+1} g_j(t_j, x + t_j v) dx dv.$$

A straightforward estimate via the change of variables $(x, v) \mapsto (x + t_i v, x + t_j v)$ gives

$$\int \prod_{j=1}^{d+1} g_j(t_j, x + t_j v) dx dv \lesssim \frac{1}{|t_i - t_j|^d} \|g_i(t_i, \cdot)\|_{L_x^1} \|g_j(t_j, \cdot)\|_{L_x^1} \prod_{k \neq i, j} \|g_k(t_k, \cdot)\|_{L_x^\infty}$$

for each $1 \leq i < j \leq d$. A multilinear interpolation argument yields

$$\int \prod_{j=1}^{d+1} g_j(t_j, x + t_j v) dx dv \lesssim \prod_{1 \leq i < j \leq d} |t_i - t_j|^{-\frac{2}{d+1}} \prod_{k=1}^{d+1} \|g_k(t_k, \cdot)\|_{L_x^{\frac{d+1}{2}}}.$$

Applying this with g^σ for each g_j we get

$$\begin{aligned} \|\rho^* g\|_{L_{x,v}^{\sigma(d+1)}}^{d+1} &\lesssim \int_{\mathbb{R}^{d+1}} \prod_{1 \leq i < j \leq d} |t_i - t_j|^{-\frac{2}{(d+1)\sigma}} \prod_{k=1}^{d+1} \|g(t_k, \cdot)\|_{L_x^{\frac{(d+1)\sigma}{2}}} d\vec{t} \\ &\lesssim \|g\|_{L_t^{q(\sigma)} L_x^{\frac{(d+1)\sigma}{2}}}^{d+1}. \end{aligned}$$

The last inequality is a consequence of the multilinear Hardy–Littlewood–Sobolev inequality due to Christ [2]. \square

Remark. It seems natural to conjecture that ρ^* satisfies the weak-type estimate

$$\|\rho^* g\|_{L_{x,v}^{d+1, \infty}} \lesssim \|g\|_{L_t^{d+1} L_x^{\frac{d+1}{2}}}.$$

A similar comment was made in an operator-theoretic context in [3].

Certain replacements for the endpoint are already known. For example, with $q = a = 2$, in [5], Keel and Tao obtain a substitute for (1.4) with the $L_x^p L_v^r$ norm (where $p = \frac{2d}{d-1}$ and $r = \frac{2d}{d+1}$) replaced by that of a certain real interpolation space which is between $L_x^{p,1} L_v^{r,1}$ and $L_x^{p,\infty} L_v^{r,\infty}$. See also work of Ovcharov [8] where a different substitute bound was given for velocities v belonging to a bounded subset of \mathbb{R}^d .

Acknowledgments

We would like to thank José Cañizo for his contributions during the early stages of this project.

Funding

This work was supported by the European Research Council (grant number 307617; Bennett); the Engineering and Physical Sciences Research Council (grant numbers EP/J021490/1, EP/J01155X/1; Bez, Gutiérrez); National Research Foundation of Korea (grant number 2012008373; Lee).

References

- [1] Castella, F., Perthame, B. (1996). Estimations de Strichartz pour les équations de transport cinétique [Strichartz estimates for kinetic transport equations]. *C. R. Acad. Sci. Paris Sér. I Math.* 332:535–540.
- [2] Christ, M. (1985). On the restriction of the Fourier transform to curves: endpoint results and the degenerate case. *Trans. Amer. Math. Soc.* 287:223–238.
- [3] Frank, R., Lewin, M., Lieb, E.H., Seiringer, R. A Strichartz inequality for orthonormal functions. *J. Euro. Math. Soc.*, to appear.
- [4] Guo, Z., Peng, L. (2007). Endpoint Strichartz estimate for the kinetic transport equation in one dimension. *C. R. Math. Acad. Sci. Paris* 345:253–256.
- [5] Keel, M., Tao, T. (1998). Endpoint Strichartz estimates. *Amer. J. Math.* 120:955–980.
- [6] Montgomery-Smith, S.J. (1998). Time decay for the bounded mean oscillation of solutions of the Schrödinger and wave equations. *Duke Math. J.* 91:393–408.
- [7] Ovcharov, E. (2011). Counterexamples to Strichartz estimates for the kinetic transport equation based on Besicovitch sets. *Nonlinear Anal.* 74:2515–2522.
- [8] Ovcharov, E. (2011). Strichartz estimates for the kinetic transport equation. *SIAM J. Math. Anal.* 43:1282–1310.
- [9] Perthame, B. (2004). Mathematical tools for kinetic equations. *Bull. Amer. Math. Soc.* 41:205–244.