UNIVERSITY^{OF} BIRMINGHAM University of Birmingham Research at Birmingham

Pseudo-heterolepticity in low-symmetry metalorganic cages

Lewis, Jamie

DOI: 10.1002/anie.202212392

License: Creative Commons: Attribution (CC BY)

Document Version Publisher's PDF, also known as Version of record

Citation for published version (Harvard):

Lewis, J 2022, 'Pseudo-heterolepticity in low-symmetry metal-organic cages', *Angewandte Chemie International Edition*, vol. 61, no. 44, e202212392. https://doi.org/10.1002/anie.202212392

Link to publication on Research at Birmingham portal

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

•Users may freely distribute the URL that is used to identify this publication.

•Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.

•User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?) •Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

Research Articles

Angewandte Chemie www.angewandte.org

Check for updates

How to cite: German Edition:

International Edition: doi.org/10.1002/anie.202212392 doi.org/10.1002/ange.202212392

Pseudo-heterolepticity in Low-Symmetry Metal-Organic Cages

James E. M. Lewis*

Host-Guest Systems Hot Paper

Abstract: Heteroleptic metal-organic cages, formed through integrative self-assembly of ligand mixtures, are highly attractive as reduced symmetry supramolecular hosts. Ensuring high-fidelity, non-statistical self-assembly, however, presents a significant challenge in molecular engineering due to the inherent difficulty in predicting thermodynamic energy landscapes. In this work, two conceptual strategies are described that circumvent this issue, using ligand design strategies to access structurally sophisticated metal-organic hosts. Using these approaches, it was possible to realise cavity environments described by two inequivalent, unsymmetrical ligand frameworks, representing a significant step forward in the construction of highly anisotropic confined spaces.

Introduction

Metal-organic self-assembly has proven to be a powerful tool to construct functional supramolecular architectures from relatively simple components.^[1] Of particular interest are systems possessing internal cavities^[2] capable of binding ions or small molecules-typically referred to as metalorganic cages (MOCs) or polyhedra (MOPs).^[3] The ability to bind guest species,^[4] such as drugs,^[5] pollutants^[6] and anions,^[7] within these cavities has been exploited for catalysis,^[8] stabilising reactive species^[9] and modulation of photophysical properties,^[10] amongst other applications.

To advance the functionality of these systems, and move towards more structurally sophisticated assemblies, researchers have recently begun to develop strategies to access MOCs of reduced symmetry in order to tailor the shape and functionality of the cavity environment.[11] Two main approaches have shown success in this endeavour. Employing unsymmetrical ligands will inherently generate MOCs of

[*] Dr. J. E. M. Lewis School of Chemistry, University of Birmingham Edgbaston, Birmingham B15 2TT (UK) and Previous address: Department of Chemistry, Molecular Sciences Research Hub, Imperial College London 82 Wood Lane, London W12 0BZ (UK) E-mail: j.e.m.lewis@bham.ac.uk

reduced symmetry (Figure 1b).^[12] Alternatively, the incorporation of more than one ligand type to generate heteroleptic,^[13] or mixed-ligand, cages is also effective (Figure 1c).^[14] The latter approach has the added benefit of being able to introduce different endo-/exohedral functional moieties into a single structure.^[15] For both of these strategies, ensuring high-fidelity self-sorting in the assembly process is essential to avoid formation of mixtures of cage structures, which makes their design challenging. It can further be imagined that combining heteroleptic and lowsymmetry strategies through the incorporation of multiple, unsymmetrical ligands into a single MOC in a defined manner (Figure 1d) would be especially difficult, which likely explains why no examples have been reported to date.

Herein are detailed two strategies to access cavity environments within MOCs that are described by two, unsymmetrical ligand structures. The first employs a covalent-tethering strategy to specifically constrain ligand fragments in a trans orientation within a polytopic framework. The second exploits the geometrically-enforced arrangement of ligands with $C_{\rm s}$ symmetry in the selective assembly of $C_{\rm 2h}$ -symmetry palladium nanocages (Figure 1b) within a previously unreported class of double-cavity Pd₃L₄ cages. With both of these approaches, the cavities generated are heteroleptic environments-i.e. they are defined by two inequivalent, unsymmetrical ligand frameworks. With the cages as a whole assembled from one type of ligand, the term pseudo-heteroleptic is proposed to describe these structures.

The significant advantage of these pseudo-heterolepticity approaches lies in the ability to construct sophisticated, multi-functional internal spaces without the requirement for integrative self-sorting of multiple components, a prerequisite of traditional heteroleptic assemblies. The delineation of these strategies opens the door to develop more sophisticated and functional confined spaces in combination with facile self-assembly.

Results and Discussion

First demonstrated by McMorran and Steel,^[16] ditopic ligands are able to assemble with Pd^{II} ions to form quadruply-stranded, Pd₂L₄-type cage structures.^[17] For ligands with C_{2v} symmetry there is only one possible configurational isomer of these cages, with (pseudo-) D_{4h} symmetry (Figure 1a). Ligands with reduced C_s symmetry, however, can assemble into four possible Pd₂L₄ diastereoisomers (Figure 1b). Unless sufficient directing effects are employed in the design of the ligand, a statistical mixture of these isomers will be generated.^[18] For example, Chand and

Angew. Chem. Int. Ed. 2022, e202212392 (1 of 7)

© 2022 The Authors. Angewandte Chemie International Edition published by Wiley-VCH GmbH

C 2022 The Authors. Angewandte Chemie International Edition published by Wiley-VCH GmbH. This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.



Figure 1. a) Self-assembly of C_{2v} -symmetric ditopic ligands with square-planar Pd^{II} ions leads to formation of Pd₂L₄ cages with (pseudo-)D_{4h} symmetry. b) Ditopic ligands lacking bilateral symmetry (C_2 rotation axis) can assemble into four possible Pd₂L₄ cage diastereoisomers. c) Integrative self-assembly of ligand mixtures leading to selective formation of heteroleptic structures. d) Integrative self-assembly of unsymmetrical ligands into heteroleptic cages has not been previously reported.

co-workers have previously reported the self-assembly of ditopic ligand **L** (Figure 2a), which is unsymmetrical by virtue of the methylene ester backbone that separates the two coordinating pyridine units. Upon self-assembly with Pd^{II} , a statistical mixture of the four possible Pd_2L_4 cage isomers was observed to form both in solution and the solid state.^[19]

This group,^[20] and others,^[21,22] have previously used geometric design principles within reduced symmetry, ditopic ligands to successfully bias the self-assembly process with Pd^{II} ions towards *cis*-Pd₂L₄ cages with C_{2h} symmetry, due to the favourable antiparallel arrangement of *trans*oriented ligands (Figure 1b). Accessing Pd₂L₄ cage isomers of alternative symmetries represents a significant design challenge. The use of geometric parameters to prepare C_{2v} *trans*-Pd₂L₄ cages, for example, would require enforcement of a parallel relationship between *trans*-oriented ligands, and antiparallel configuration of *cis*-oriented ligands. Recently, an M₂L₄ system with C_{4v} symmetry was able to be realised by Crowley and co-workers, using a sub-component selfassembly strategy in the synthesis of a heteronuclear PdPtL₄ cage.^[23]

The 1,2-bis(pyridin-2-ylethynyl)benzene motif has previously been used as a *trans*-chelating unit for square-planar Pd^{II} ions (Figure 2b),^[24] and has been investigated for its potential as a *trans* analogue of Fujita's *cis*-protection strategy^[25] in metal-organic assembly.^[26] It was envisaged that covalently tethering ligand fragments^[27] via this unit could be used to constrain the relative orientations of ligands held *trans* to each other. In this manner it might be possible to obtain cages of C_{2v} symmetry that have previously proved difficult to access.

trans-Pd2L2 C2v Cage

 L^{P4} was readily synthesised from 1,2-diethynylbenzene and commercially available reagents in 2 steps in 73% overall

in an anti-parallel arrangement around two Pd^{II} ions. Combining L^{P4} with $Pd(NO_3)_2 \cdot 2H_2O$ in d_6 -DMSO at room temperature gradually led to quantitative formation of a single species (C^{P4}), as observed by ¹H NMR (Figure 2d). Formation of an assembly with the anticipated $[Pd_2(L^{P4})_2]^{4+}$ formula was confirmed by electrospray ionisation mass spectrometry (ESI-MS; Fig S19 and S20). Ultimately, despite the poor quality of the crystals, unambiguous confirmation of the structure was achieved via single crystal X-ray diffraction (SCXRD) studies of the host–guest complex with an encapsulated chloride anion, $[Pd_2(L^{P4})_2 \supset CI]^{3+}$ (Figure 2f).^[28]

yield. It was anticipated that two L^{P4} ligands would assemble

Due to the ligand structure and the steric bulk of the diethynylbenzene linker, $\mathbf{C}^{\mathbf{P4}}$ exclusively assembles as a $\mathrm{Pd_2L_2}$ cage with C_{2v} symmetry. Consequently, the cavity of the cage can be described as being surrounded by four iterations of the ligand **L**, reported by Chand, in a (pseudo-)*trans*-Pd_2L_4 configuration (Figure 1b). As with **L**, efficient formation of $\mathbf{C}^{\mathbf{P4}}$ required the presence of a suitable anion template such as NO₃⁻ or Cl⁻, with BF₄⁻ alone resulting in a mixture of products (Figure S22). Cl⁻ appeared to be a better guest than NO₃⁻, evidenced by an increased downfield shift of the endohedral protons H_a and H_f ($\Delta \delta = 0.43$ and 0.41 ppm, respectively).

trans-Pd₂L₂ Pseudo-heteroleptic C₂ Cage

Having successfully demonstrated the viability of this tethering strategy to constrain two dipyridyl ligand moieties in a *trans* arrangement, the thought occurred that the two ditopic units need not be identical. Consequently, \mathbf{L}^{P3Q} (Figure 2c) was prepared incorporating one of the previously described dipyridyl fragments, and a second ditopic ligand with one pyridine and one quinoline coordinating unit. Self-assembly of a stoichiometric mixture of \mathbf{L}^{P3Q} with Pd(NO₃)₂·2H₂O in *d*₆-DMSO at 50 °C for 2 h again yielded a single species,





Figure 2. a) Chand's low-symmetry ditopic ligand, L, and SCXRD structure of the dinuclear Pd^{III} cage structure $[Pd_2L_4 \supset Cl]^{3+}$ (CCDC# 1941617; only C_{4v} isomer shown for clarity); b) 1,2-bis (pyridin-2-ylethynyl)benzene and the SCXRD structure of its mononuclear Pd^{III} complex (CCDC# 185471); c) tetratopic ligands L^{P4} and L^{P3Q} with cartoon diagrams representing the relative orientations of the ligand fragments within C^{P4} and C^{P3Q}; partial ¹H NMR spectra (500 MHz, d_6 -DMSO, 298 K) of d) C^{P4} and e) C^{P3Q}; SCXRD structures of f) $[Pd_2(L^{P3Q})_2 \supset NO_3]^{3+}$.

determined to be $[Pd_2(\mathbf{L}^{P3Q})_2](NO_3)_4$ (\mathbf{C}^{P3Q}) by NMR and ESI-MS. Diastereotopic splitting of the signals assigned to the methylene units was observed (Figure 2e), giving two pairs of doublets (5.74/5.51 ppm and 5.42/5.12 ppm), indicative of inequivalent chemical environments on either face of the ditopic ligand fragments. Vapour diffusion of Et₂O into a solution of the assembly in 1:1 DMSO/CH₃CN yielded Xray quality crystals, with SCXRD revealing the anticipated $[Pd_2(\mathbf{L}^{P4})_2 \supset NO_3]^{3+}$ structure (Figure 2g).^[28] By virtue of the inequivalence of the two ditopic ligand fragments within \mathbf{L}^{P3Q} , the structural framework of \mathbf{C}^{P3Q} is analogous to that of a heteroleptic $Pd_2L_2L'_2$ cage^[29] assembled from two unsymmetrical ligands (L and L'). Within the assembly, pairs

of identical ligands are held cis to each other in an antiparallel arrangement, giving a structure with C_2 symmetry.

Of note, due to the (bis-)*trans*-heterobidentate nature of \mathbf{L}^{P3Q} , each of the Pd^{II} centres in \mathbf{C}^{P3Q} possesses axial chirality.^[30] From the SCXRD structure it could be shown that within \mathbf{C}^{P3Q} both ions are of the same stereochemistry, and the system crystallises as a racemic mixture of both (S,S)- and (R,R)- \mathbf{C}^{P3Q} enantiomers (see Supporting Information section 6).

This covalent tethering strategy was clearly highly effective at constraining the relative orientations of ligand fragments, and even for the incorporation of more than one framework into an assembly. The *bent*, or *V*-shaped, nature of L^{P3Q} gave rise to the C_2 symmetry of C^{P3Q} . To widen the scope of accessible $Pd_2L_2L'_2$ cavity symmetries, there was motivation to investigate alternative modes of tethering that might lead to C_s symmetry structures (Figure 1d). These would be formed from pairs of unsymmetrical ligands with identical ligands arranged *cis* to each other, but now in a parallel orientation. Tethering of ligand fragments in a *linear* fashion appeared to present a solution to accessing these structures.

cis-Pd₂L₄ Cages from Unsymmetrical Ditopic Ligands

Based on principles of geometric complementarity delineated in previous work,^[20] two low-symmetry, ditopic ligands were synthesised with isostructural core frameworks (Figure 3a). The ligands differed only in the identity of the



Figure 3. a) Ligands L1^{Ph} and L1^{xy}; partial ¹H NMR spectra (400 MHz, d_6 -DMSO, 298 K) of b) C1^{Ph} and c) C1^{xy}; d) SCXRD structure of $[Pd_2(L1^{xy})_4 \supset 2BF_4]^{2+}$.

flanking aromatic units—1,4-phenyl ($L1^{Ph}$) and 2,5-*m*-xylyl ($L1^{Xy}$). In each case, self-assembly with Pd^{II} as the tetrafluoroborate salt in a 2:1 ligand/metal ratio resulted in formation of the anticipated *cis*-[Pd₂L₄](BF₄)₄ architectures (C1^{Ph} and C1^{Xy}), as determined by NMR (Figure 3b and c), ESI-MS (Figure S75–77 and S100–102) and, for C1^{Xy}, SCXRD (see below).

The impact of exchanging the flanking phenyl group in $C1^{Ph}$ for a xylyl unit in $C1^{Xy}$ resulted in some spectroscopic differences, noticeably a shift in the resonance of the signal assigned to endohedral proton H_j from 8.31 ppm to 7.71 ppm. The symmetry of the ¹H NMR spectra of $C1^{Ph}$ and $C1^{Xy}$, through-space interactions between *ortho*-pyridyl protons observed by NOESY ($H_a \cdots H_n$, $H_b \cdots H_m$; Figure S73 and S98) and splitting of the H_e (7.46 and 7.31 ppm, Figure 3c) and H_f (2.13 and 1.99 ppm) resonances within $C1^{Xy}$ into two distinct sets of signals, were all congruent with the C_{2h} symmetry associated with selective formation of the *cis* assembly (Figure 1b). This was ultimately further confirmed by the solid-state SCXRD structure of $C1^{Xy}$ (Figure 3d).^[28]

cis-Pd₃L₄ Double-Cavity Cages

There has been some interest in recent years in the synthesis of linearly-elongated, polytopic ligand scaffolds, and their self-assembly to form multi-cavity cage architectures.^[31] In this manner, larger assemblies can be formed that possess several small cavities^[32] instead of a single large one.^[33] The formation of such a species from low-symmetry ligands has, to the best of my knowledge, never been reported.^[34]

To probe whether the geometric complementarity approach would remain effective for the assembly of lowsymmetry, multi-cavity systems, tritopic ligand $L2^{Ph}$ (Figure 4a) was synthesised, incorporating two linearly-fused iterations of the $L1^{Ph}$ scaffold. Pleasingly, self-assembly with [Pd(CH₃CN)₄](BF₄)₂ in a 3:4 metal/ligand ratio in 3:1 d_6 -DMSO/CDCl₃ successfully yielded the desired *cis*-Pd₃L₄ double-cavity cage (C2^{Ph}), determined by NMR (Figure 4b; see below) and ESI-MS (Figure S151–153).

As with previous work on the self-assembly of unsymmetrical ditopic ligands,^[20] the symmetry of the ¹H NMR spectrum and through-space interactions observed by NO-ESY between opposite ends of the ligand framework $(H_a \cdots H_w, H_t \cdots H_m;$ Figure S149) within the trinuclear architecture were consistent with cages of either C_{2v} (*trans*) or C_{2h} (*cis*) symmetry. Disappointingly, the growth of X-ray quality crystals proved elusive. Based, however, on previous work, the dinuclear cages C1^{Ph} and C1^{xy}, and molecular modelling of the *cis*-Pd₃L₄ structure (Figure 4d), it was concluded that C2^{Ph} must be the *cis* assembly. As such, the efficacy of using geometric design parameters to enforce ligands *trans* to each other to arrange in an anti-parallel fashion is retained by the elongated, tritopic ligand.

Although the two cavities of the double-decker assembly $C2^{Ph}$ are chemically equivalent, separate sets of signals in the ¹H NMR spectrum were observed for the protons associated with each half of the ligand framework (Figure 4b), confirming the difference in their chemical environ-



Figure 4. a) Tritopic ligands $L2^{Ph}$ and $L2^{xy}$; partial ¹H NMR spectra (400 MHz, 3:1 d_6 -DMSO/CDCl₃, 298 K) of b) $C2^{Ph}$ and c) $C2^{xy}$; molecular models (PM6) of d) $C2^{Ph}$ and e) $C2^{xy}$ with some chemically inequivalent core ligand fragments shown in green and purple, highlighting the pseudo-heteroleptic nature of the cavities.

ments. This can readily be seen by the different chemical shifts in the ¹H NMR spectrum of the homologous *ortho*-pyridyl protons H_a and H_m (δ =9.89 and 9.94 ppm, respectively) and H_l and H_w (δ =9.60 and 9.51 ppm, respectively), and endohedral protons H_j and H_s (δ =8.42 and 8.34 ppm, respectively).

The consequence of the combination of chemically inequivalent fragments of $L2^{Ph}$ and the symmetry of $C2^{Ph}$ is that each of the identical cage cavities are encompassed by two chemically distinct, unsymmetrical ligand fragments (Figure 4d). As a result, the cavities can be visualised as

(pseudo-)heteroleptic $Pd_2L_2L'_2$ environments of C_s symmetry.

This concept was further demonstrated through the selfassembly of ligand $L2^{xy}$ (Figure 4a) with $[Pd(CH_3CN)_4]$ - $(BF_4)_2$, under the same conditions as $L2^{Ph}$, to give $C2^{xy}$, in which one half of the ligand scaffold possesses a phenyl ring as the flanking aromatic unit, the other a xylyl moiety. As with the dinuclear C1 cages and $C2^{Ph}$, confirmation of successfully forming the targeted *cis*-Pd₃L₄ isomer came from ESI-MS (Figure S168–170), the high symmetry of the NMR spectrum (Figure 4c), NOE interactions only feasible between ligands in an anti-parallel arrangement ($H_a \cdots H_w$, $H_i \cdots H_m$; Figure S165) and, consistent with C1^{xy}, splitting of the aromatic (H_n ; $\delta = 7.47$ and 7.28 ppm) and methyl (H_o ; $\delta =$ 2.03 and 1.92 ppm) protons of the xylyl unit in C2^{xy} into two distinct signals.

Within the trinuclear $C2^{xy}$, two pairs of ligands surround each individual cage cavity (Figure 4e)—one with phenyl units as the flanking aromatic moieties (as per $C1^{Ph}$), the other with *m*-xylyl (as per $C1^{xy}$), the close proximity of which was demonstrated by NOE interactions observed between the two ($H_e \cdots H_o$; Figure S165 and S166). Again, the difference in the ligand environments was readily shown by ¹H NMR (Figure 4c) in the chemical shifts of homologous endohedral protons (e.g. H_j and H_s —8.53 and 7.62 ppm, respectively).

The geometric parameters that drive selective formation of the cis-[Pd₃(L2)₄]⁶⁺ cage structures mean that ligands positioned *trans* from each other across the Pd^{II} ions are in an anti-parallel relative orientation. The result is that each cavity is described by two chemically distinct, unsymmetrical ligand environments (Figure 4d and e). Within a single, dinuclear cage structure, such an assembly would be termed heteroleptic. As the cis-[Pd₃(**L2**)₄]⁶⁺ assemblies are homoleptic as a whole, and it is the individual cavities that possess an induced heteroleptic environment, the term pseudoheteroleptic is suggested to describe the cavity environments that result from the self-assembly of the low-symmetry ligands. This term would also seem appropriate for the C^{P3Q} architecture described above, in which two distinct, unsymmetrical ligand fragments were incorporated into a single cage structure in a configurationally defined manner. As such, these pseudo-heterolepticity approaches offer the potential for incorporating multiple functionalities into the cavity environment of the cage that could not otherwise be achieved via integrative self-sorting of multiple ligands, and under facile self-assembly conditions afforded by the formally homoleptic architectures.

Conclusion

With general principles of metal-organic self-assembly well evolved, and growing interest in the various applications of metal-organic cages, the development of strategies to access more structurally sophisticated systems would greatly advance their utility. Constructing MOCs with more than one ligand framework offers the potential to precision engineer both the shape and functionality of the cavity environment.

Angew. Chem. Int. Ed. 2022, e202212392 (5 of 7)

Previous approaches to the assembly of heteroleptic metalorganic architectures rely on integrative self-assembly of multiple ligands, either resulting from geometric complementarity, or coordination sphere engineering strategies. Whilst these have proven effective to a certain extent, there are inherent limitations to these approaches.

In this work, strategies have been outlined towards the development of *pseudo*-heteroleptic MOCs-cages that, although they are assembled from single ligands, possess cavity spaces described by multiple ligand environments. These rely on covalent tethering of ligand fragments to constrain their relative orientations. By controlling the geometry of the tethering-either *bent* or *linearly fused*-MOCs have been realised that incorporate two inequivalent, unsymmetrical ligand scaffolds in a defined manner, generating cavities of C_2 and C_s symmetry, respectively—a feat that has so far not been achieved via integrative self-assembly of ligand mixtures.

The delicate balance of entropic and enthalpic factors can be easily disturbed to shift thermodynamic minima away from desired products. Removing the reliance on integrative self-sorting greatly simplifies the self-assembly process, whilst enhancing the structural and functional complexity of the cavity space. As such, pseudo-heteroleptic design strategies will allow fine-tuning of the cavity shape and functionality of discrete, porous MOCs, under facile conditions, to an extent not previously possible.

Acknowledgements

Dr. Andrew J. P. White (Imperial College London) is thanked for collection and analysis of SCXRD data. The Imperial College Research Fellowship programme is acknowledged for funding. Professor Matthew J. Fuchter is thanked for discussions and access to resources.

Conflict of Interest

The authors declare no conflict of interest.

Data Availability Statement

The data that support the findings of this study are available from the corresponding author upon reasonable request.

Keywords: Cage • Heteroleptic • Low-Symmetry • Metallosupramolecular • Self-Assembly

- a) M. Fujita, Chem. Soc. Rev. 1998, 27, 417–425; b) B.J.
 Holliday, C. A. Mirkin, Angew. Chem. Int. Ed. 2001, 40, 2022–2043; Angew. Chem. 2001, 113, 2076–2097; c) T. R. Cook, P. J.
 Stang, Chem. Rev. 2015, 115, 7001–7045.
- [2] R. S. Forgan, G. O. Lloyd, *Reactivity in confined spaces*, Royal Society of Chemistry, London, 2021.

 \bigcirc 2022 The Authors. Angewandte Chemie International Edition published by Wiley-VCH GmbH

Angewandte

- [3] a) M. M. J. Smulders, I. A. Riddell, C. Browne, J. R. Nitschke, *Chem. Soc. Rev.* 2013, 42, 1728–1754; b) K. Harris, D. Fujita, M. Fujita, *Chem. Commun.* 2013, 49, 6703–6712; c) N. B. Debata, D. Tripathy, H. S. Sahoo, *Coord. Chem. Rev.* 2019, 387, 273–298; d) A. E. Martín Díaz, J. E. M. Lewis, *Front. Chem.* 2021, 9, 706462; e) T. Tateishi, M. Yoshimura, S. Tokuda, F. Matsuda, D. Fujita, S. Furukawa, *Coord. Chem. Rev.* 2022, 467, 214612; f) A. J. McConnell, *Chem. Soc. Rev.* 2022, 51, 2957–2971.
- [4] a) V. Maurizot, M. Yoshizawa, M. Kawano, M. Fujita, *Dalton Trans.* **2006**, 2750–2756; b) C. T. McTernan, J. A. Davies, J. R. Nitschke, *Chem. Rev.* **2022**, *122*, 10393–10437.
- [5] a) S. K. Samanta, L. Isaacs, *Coord. Chem. Rev.* 2020, 410, 213181. For selected examples, see: b) B. Therrien, G. Süss-Fink, P. Govindaswamy, A. K. Renfrew, P. J. Dyson, *Angew. Chem. Int. Ed.* 2008, 47, 3773–3776; *Angew. Chem.* 2008, 120, 3833–3836; c) J. E. M. Lewis, E. L. Gavey, S. A. Cameron, J. D. Crowley, *Chem. Sci.* 2012, *3*, 778–784; d) F. Schmitt, J. Freudenreich, N. P. E. Barry, L. Juillerat-Jeanneret, G. Süss-Fink, B. Therrien, *J. Am. Chem. Soc.* 2012, 134, 754–757; e) Y.-R. Zheng, K. Suntharalingam, T. C. Johnstone, S. J. Lippard, *Chem. Sci.* 2015, *6*, 1189–1193.
- [6] a) E. G. Percástegui, Chem. Commun. 2022, 58, 5055–5071. For selected examples, see: b) I. A. Riddell, M. M. J. Smulders, J. K. Clegg, J. R. Nitschke, Chem. Commun. 2011, 47, 457–459; c) D. Preston, K. F. White, J. E. M. Lewis, R. A. S. Vasdev, B. F. Abrahams, J. D. Crowley, Chem. Eur. J. 2017, 23, 10559–10567; d) J. S. Wright, A. J. Metherell, W. M. Cullen, J. R. Piper, R. Dawson, M. D. Ward, Chem. Commun. 2017, 53, 4398–4401; e) D. Zhang, T. K. Ronson, R. Lavendomme, J. R. Nitschke, J. Am. Chem. Soc. 2019, 141, 18949–18953; f) S. de Jesús Valencia-Loza, A. López-Olvera, E. Martínez-Ahumada, D. Martínez-Otero, I. A. Ibarra, V. Jancik, E. G. Percástegui, ACS Appl. Mater. Interfaces 2021, 13, 18658–18665.
- [7] a) R. Custelcean, Chem. Soc. Rev. 2014, 43, 1813–1824. For selected examples, see: b) D. Zhang, T. K. Ronson, J. Mosquera, A. Martinez, J. R. Nitschke, Angew. Chem. Int. Ed. 2018, 57, 3717–3721; Angew. Chem. 2018, 130, 3779–3783; c) A. J. Plajer, E. G. Percástegui, M. Santella, F. J. Rizzuto, Q. Gan, B. W. Laursen, J. R. Nitschke R, Angew. Chem. Int. Ed. 2019, 58, 4200–4204; Angew. Chem. 2019, 131, 4244–4248; d) R. Andrews, S. Begum, C. J. Clemett, R. A. Faulkner, M. L. Ginger, J. Harmer, M. Molinari, G. M. B. Parkes, Z. M. H. Qureshi, C. R. Rice, M. D. Ward, H. M. Williams, P. B. Wilson, Angew. Chem. Int. Ed. 2020, 59, 20480–20484; Angew. Chem. 2020, 132, 20660–20664; e) H. Lee, D. Kim, H. Oh, O.-S. Jung, Chem. Commun. 2020, 56, 2841–2844; f) D. Preston, K. M. Patil, A. T. O'Neil, R. A. S. Vasdev, J. A. Kitchen, P. E. Kruger, Inorg. Chem. Front. 2020, 7, 2990–3001.
- [8] a) Y. Ueda, H. Ito, D. Fujita, M. Fujita, J. Am. Chem. Soc. 2017, 139, 6090–6093; b) W. Cullen, A. J. Metherell, A. B. Wragg, C. G. P. Taylor, N. H. Williams, M. D. Ward, J. Am. Chem. Soc. 2018, 140, 2821–2828; c) V. Martí-Centelles, A. L. Lawrence, P. J. Lusby, J. Am. Chem. Soc. 2018, 140, 2862–2868; d) C. Ngai, C. M. Sanchez-Marsetti, W. H. Harman, R. J. Hooley, Angew. Chem. Int. Ed. 2020, 59, 23505–23509; Angew. Chem. 2020, 132, 23711–23715; e) J. Guo, Y.-Z. Fan, Y.-L. Lu, S.-P. Zheng, C.-Y. Su, Angew. Chem. Int. Ed. 2020, 59, 8661–8669; Angew. Chem. 2020, 132, 8739–8747; f) A. Paul, M. A. Shipman, D. Y. Onabule, S. Sproules, M. D. Symes, Chem. Sci. 2021, 12, 5082–5090; g) S. M. Bierschenk, J. Y. Pan, N. S. Settineri, U. Warzok, R. G. Bergman, K. N. Raymond, F. D. Toste, J. Am. Chem. Soc. 2022, 144, 11425–11433.
- [9] a) P. Mal, B. Breiner, K. Rissanen, J. R. Nitschke, *Science* 2009, 324, 1697–1699; b) M. Yamashina, Y. Sei, M. Akita, M. Yoshizawa, *Nat. Commun.* 2014, 5, 4662; c) S. Hasegawa, S. L.

Meichsner, J. J. Holstein, A. Baksi, M. Kasanmascheff, G. H. Clever, *J. Am. Chem. Soc.* **2021**, *143*, 9718–9723; d) R. Sumida, Y. Tanaka, K. Niki, Y. Sei, S. Toyota, M. Yoshizawa, *Chem. Sci.* **2021**, *12*, 9946–9951.

- [10] a) J. Gemen, J. Ahrens, L. J. W. Shimon, R. Klajn, *J. Am. Chem. Soc.* 2020, *142*, 17721–17729; b) X. Tang, H. Jiang, Y. Si, N. Rampal, W. Gong, C. Cheng, X. Kang, D. Fairen-Jimenez, Y. Cui, Y. Liu, *Chem* 2021, *7*, 2771–2786.
- [11] S. Pullen, J. Tessarolo, G. H. Clever, *Chem. Sci.* 2021, *12*, 7269– 7293.
- [12] a) J. E. M. Lewis, J. D. Crowley, *ChemPlusChem* 2020, 85, 815–827; b) D. Tripathy, N. B. Debata, K. C. Naik, H. S. Sahoo, *Coord. Chem. Rev.* 2022, 456, 214396.
- [13] a) W. M. Bloch, G. H. Clever, *Chem. Commun.* 2017, *53*, 8506–8516; b) S. Pullen, G. H. Clever, *Acc. Chem. Res.* 2018, *51*, 3052–3064; c) D. Bardhan, D. K. Chand, *Chem. Eur. J.* 2019, 25, 12241–12269.
- [14] For some recent examples of heteroleptic metal-organic cages, see: a) S. Sudan, R.-J. Li, S. M. Jansze, A. Platzek, R. Rudolf, G. H. Clever, F. Fadaei-Tirani, R. Scopelliti, K. Severin, J. Am. Chem. Soc. 2021, 143, 1773–1778; b) R.-J. Li, F. Fadaei-Tirani, R. Scopelliti, K. Severin, Chem. Eur. J. 2021, 27, 9439–9445; c) J. A. Findlay, K. M. Patil, M. G. Gardiner, H. I. MacDermott-Opeskin, M. L. O'Mara, P. E. Kruger, D. Preston, Chem. Asian J. 2022, 17, e202200093; d) S.-C. Li, L.-X. Cai, M. Hong, Q. Chen, Q.-F. Sun, Angew. Chem. Int. Ed. 2022, 61, e202204732; Angew. Chem. 2022, 134, e202204732.
- [15] K. Wu, J. Tessarolo, A. Baksi, G. H. Clever, Angew. Chem. Int. Ed. 2022, 61, e202205725; Angew. Chem. 2022, 134, e202205725.
- [16] a) D. A. McMorran, P. J. Steel, Angew. Chem. Int. Ed. 1998, 37, 3295–3297; Angew. Chem. 1998, 110, 3495–3497; b) P. J. Steel, D. A. McMorran, Chem. Asian J. 2019, 14, 1098–1101.
- [17] a) A. Schmidt, A. Casini, F. E. Kühn, *Coord. Chem. Rev.* 2014, 275, 19–36. For some recent examples of homoleptic Pd₂L₄ cages, see: b) T. Tsutsui, L. Catti, K. Yozac, M. Yoshizawa, *Chem. Sci.* 2020, *11*, 8145–8150; c) A. P. Birvé, H. D. Patel, J. R. Price, W. M. Bloch, T. Fallon, *Angew. Chem. Int. Ed.* 2022, *61*, e202115468; *Angew. Chem.* 2022, *134*, e202115468; d) H. Lee, J. Tessarolo, D. Langbehn, A. Baksi, R. Herges, G. H. Clever, *J. Am. Chem. Soc.* 2022, *144*, 3099–3105.
- [18] R. A. S. Vasdev, D. Preston, C. A. Casey-Stevens, V. Martí-Centelles, P. J. Lusby, A. L. Garden, J. D. Crowley, *Inorg. Chem.* 2022, https://doi.org/10.1021/acs.inorgchem.2c00937.
- [19] S. Samantray, S. Krishnaswamy, D. K. Chand, Nat. Commun. 2020, 11, 880.
- [20] a) J. E. M. Lewis, A. Tarzia, A. J. P. White, K. E. Jelfs, *Chem. Sci.* 2020, *11*, 677–683; b) J. E. M. Lewis, *Chem. Eur. J.* 2021, 27, 4454–4460; c) A. Tarzia, J. E. M. Lewis, K. E. Jelfs, *Angew. Chem. Int. Ed.* 2021, *60*, 20879–20887; *Angew. Chem.* 2021, *133*, 21047–21055.
- [21] a) D. Ogata, J. Yuasa, Angew. Chem. Int. Ed. 2019, 58, 18424– 18428; Angew. Chem. 2019, 131, 18595–18599; b) S. S. Mishra, S. V. K. Kompella, S. Krishnaswamy, S. Balasubramanian, D. K. Chand, Inorg. Chem. 2020, 59, 12884–12894.
- [22] Higher nuclearity Pd^{II} assemblies with unsymmetrical ditopic ligands have also been reported: a) R.-J. Li, A. Marcus, F. Fadaei-Tirani, K. Severin, *Chem. Commun.* 2021, 57, 10023– 10026; b) S. S. Mishra, D. K. Chand, *Dalton Trans.* 2022, 51, 11650–11657.
- [23] L. S. Lisboa, J. A. Findlay, L. J. Wright, C. G. Hartinger, J. D. Crowley, Angew. Chem. Int. Ed. 2020, 59, 11101–11107; Angew. Chem. 2020, 132, 11194–11200.
- [24] a) E. Bosch, C. L. Barnes, *Inorg. Chem.* 2001, 40, 3097–3100;
 b) Y.-Z. Hu, C. Chamchoumis, J. S. Grebowicz, R. P. Thummel, *Inorg. Chem.* 2002, 41, 2296–2300.

Angew. Chem. Int. Ed. 2022, e202212392 (6 of 7)

© 2022 The Authors. Angewandte Chemie International Edition published by Wiley-VCH GmbH

- [25] M. Fujita, M. Tominaga, A. Hori, B. Therrien, Acc. Chem. Res. 2005, 38, 369–378.
- [26] F. A. Pereira, T. Fallows, M. Frank, A. Chen, G. H. Clever, Z. Anorg. Allg. Chem. 2013, 639, 1598–1605.
- [27] a) S. Tashiro, M. Tominaga, T. Kusukawa, M. Kawano, S. Sakamoto, K. Yamaguchi, M. Fujita, Angew. Chem. Int. Ed. 2003, 42, 3267–3270; Angew. Chem. 2003, 115, 3389–3392;
 b) Q.-F. Sun, T. Murase, S. Sato, M. Fujita, Angew. Chem. Int. Ed. 2011, 50, 10318–10321; Angew. Chem. 2011, 123, 10502–10505; c) I. A. Bhat, D. Samanta, P. S. Mukherjee, J. Am. Chem. Soc. 2015, 137, 9497–9502; d) L. Escobar, E. C. Escuedro-Adán, P. Ballester, Angew. Chem. Int. Ed. 2019, 58, 16105–16109; Angew. Chem. 2019, 131, 16251–16255; e) K. Wu, B. Zhang, C. Drechsler, J. J. Holstein, G. H. Clever, Angew. Chem. Int. Ed. 2021, 60, 6403–6407; Angew. Chem. 2021, 133, 6473–6478.
- [28] Deposition numbers 2202605 (\mathbb{C}^{P4}), 2202606 ($\mathbb{C1}^{xy}$), and 2202607 (\mathbb{C}^{P3Q}) contain the supplementary crystallographic data for this paper. These data are provided free of charge by the joint Cambridge Crystallographic Data Centre and Fachinformationszentrum Karlsruhe Access Structures service.
- [29] a) W. M. Bloch, Y. Abe, J. J. Holstein, C. M. Wandtke, B. Dittrich, G. H. Clever, J. Am. Chem. Soc. 2016, 138, 13750–13755; b) W. M. Bloch, J. J. Holstein, W. Hiller, G. H. Clever, Angew. Chem. Int. Ed. 2017, 56, 8285–8289; Angew. Chem. 2017, 129, 8399–8404; c) D. Preston, J. E. Barnsley, K. C. Gordon, J. D. Crowley, J. Am. Chem. Soc. 2016, 138, 10578–10585; d) R. Zhu, W. M. Bloch, J. J. Holstein, S. Mandal, L. V. Schäfer, G. H. Clever, Chem. Eur. J. 2018, 24, 12976–12982; e) B. Chen, J. J. Holstein, A. Platzek, L. Schneider, K. Wu, G. H. Clever, Chem. Sci. 2022, 13, 1829–1834.
- [30] T. R. Schulte, J. J. Holstein, L. Krause, R. Michel, D. Stalke, E. Sakuda, K. Umakoshi, G. Longhi, S. Abbate, G. H. Clever, J. Am. Chem. Soc. 2017, 139, 6863–6866.

- [31] R. A. S. Vasdev, D. Preston, J. D. Crowley, *Chem. Asian J.* 2017, *12*, 2513–2523.
- [32] a) S. Bandi, A. K. Pal, G. S. Hanan, D. K. Chand, *Chem. Eur. J.* 2014, 20, 13122–13126; b) M. D. Johnstone, E. K. Schwarze, G. H. Clever, F. M. Pfeffer, *Chem. Eur. J.* 2015, 21, 3948–3955; c) D. Preston, J. E. M. Lewis, J. D. Crowley, *J. Am. Chem. Soc.* 2017, 139, 2379–2386; d) K. Yazaki, M. Akita, S. Prusty, D. K. Chand, T. Kikuchi, H. Sato, M. Yoshizawa, *Nat. Commun.* 2017, 8, 15914; e) R. Zhu, I. Regeni, J. J. Holstein, B. Dittrich, M. Simon, S. Prévost, M. Gradzielski, G. H. Clever, *Angew. Chem. Int. Ed.* 2018, 57, 13652–13656; *Angew. Chem.* 2018, 130, 13840–13844.
- [33] a) M. Tominaga, K. Suzuki, M. Kawano, T. Kusukawa, T. Ozeki, S. Sakamoto, K. Yamaguchi, M. Fujita, *Angew. Chem. Int. Ed.* 2004, *43*, 5621–5625; *Angew. Chem.* 2004, *116*, 5739–5743; b) Q.-F. Sun, J. Iwasa, D. Ogawa, Y. Ishido, S. Sato, T. Ozeki, Y. Sei, K. Yamaguchi, M. Fujita, *Science* 2010, *328*, 1144–1147; c) D. Fujita, Y. Ueda, S. Sato, N. Mizuno, T. Kumasaka, M. Fujita, *Nature* 2016, *540*, 563–567.
- [34] Crowley and co-workers have recently reported a heteronuclear, double-cavity [Pd₂PtL₄] cage using a sub-component self-assembly strategy. The unsymmetrical ligand is formed in situ using dynamic covalent chemistry, distinguishing it from the systems investigated in this work: L. S. Lisboa, D. Preston, C. J. McAdam, L. J. Wright, C. G. Hartinger, J. D. Crowley, *Angew. Chem. Int. Ed.* 2022, *61*, e202201700; *Angew. Chem.* 2022, *134*, e202201700.

Manuscript received: August 22, 2022

Accepted manuscript online: September 8, 2022 Version of record online: **••**, **••**



Research Articles

Research Articles

Host-Guest Systems

J. E. M. Lewis* _____ e202212392

Pseudo-heterolepticity in Low-Symmetry Metal-Organic Cages **PSEUDO-HETEROLEPTICITY**



HETEROLEPTIC CAVITY ENVIRONMENT UNSYMMETRICAL LIGAND FRAMEWORKS FACILE HOMOLEPTIC SELF-ASSEMBLY Heteroleptic metal-organic cages are prized for their structural complexity and multi-functionality. Controlling integrative self-sorting processes necessary to achieve defined self-assembly of these supramolecular hosts is a significant design challenge. In this work, ligandtethering strategies are described that allow access to metal-organic cages with highly anisotropic cavities described by two inequivalent, unsymmetrical ligand scaffolds.