

Panel unit-root tests with structural breaks

Chen, Pengyu; Karavias, Yiannis; Tzavalis, Elias

DOI:

[10.1177/1536867X221124541](https://doi.org/10.1177/1536867X221124541)

License:

Creative Commons: Attribution (CC BY)

Document Version

Publisher's PDF, also known as Version of record

Citation for published version (Harvard):

Chen, P, Karavias, Y & Tzavalis, E 2022, 'Panel unit-root tests with structural breaks', *Stata Journal*, vol. 22, no. 3, pp. 664-678. <https://doi.org/10.1177/1536867X221124541>

[Link to publication on Research at Birmingham portal](#)

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.



Panel unit-root tests with structural breaks

Pengyu Chen
University of Birmingham
Birmingham, U.K.
cpy1416@outlook.com

Yiannis Karavias
University of Birmingham
Birmingham, U.K.
i.karavias@bham.ac.uk

Elias Tzavalis
Athens University of Economics and Business
Athens, Greece
e.tzavalis@aueb.gr

Abstract. In this article, we introduce a new community-contributed command called `xtbunitroot`, which implements the panel-data unit-root tests developed by Karavias and Tzavalis (2014, *Computational Statistics and Data Analysis* 76: 391–407). These tests allow for one or two structural breaks in deterministic components of the series and can be seen as panel-data counterparts of the tests by Zivot and Andrews (1992, *Journal of Business and Economic Statistics* 10: 251–270) and Lumsdaine and Papell (1997, *Review of Economics and Statistics* 79: 212–218). The dates of the breaks can be known or unknown. The tests allow for intercepts and linear trends, nonnormal errors, and cross-section heteroskedasticity and dependence. They have power against homogeneous and heterogeneous alternatives and can be applied to panels with small or large time-series dimensions.

Keywords: `st0687`, `xtbunitroot`, panel data, unit root, structural break, banking, COVID-19

1 Introduction

A fundamental property of time-series data is stationarity. If a time series is stationary, then all the standard tools of statistical analysis can be applied; the series can be used in regression models, its mean and variance estimated, and its future forecasted. Frequently, though, time series are nonstationary, or unit-root processes as they are called, and in this case their statistical analysis is different. If nonstationarity is not accounted for, many problems arise for inference and prediction, the most notable of which is that of spurious regression; see, for example, Granger and Newbold (1974). Spurious regression leads to high R^2 and statistically significant coefficients, when in fact there might be no relationship between the series of interest. Therefore, determining whether a series is stationary should be one of the first steps in any time-series and panel-data (which consist of multiple time series) analysis. Unit-root tests are statistical hypothesis tests used to infer whether a series is a unit root or a stationary process; see, for example, Dickey and Fuller (1979).

In a seminal article, Perron (1989) demonstrated that structural breaks can adversely impact the behavior of unit-root tests. Structural breaks are shocks that are exogenous to the model but have a lasting effect because they change the model parameters. Structural breaks occur when a system is hit by large-scale phenomena; in economics, for example, such events include wars, policy changes, a pandemic like COVID-19, or a financial crisis. These shocks can make stationary series look as if they are nonstationary and therefore mislead the unit-root tests to accept the null hypothesis of nonstationarity when it is not true. To deal with this problem, Perron (1989) proposed new unit-root tests that allow for a structural break in the constant and trend of the series. Perron's approach, however, assumed that the date of the structural break is known to the researcher. Zivot and Andrews (1992) and Banerjee, Lumsdaine, and Stock (1992) extended Perron (1989) by allowing the date of the break to be endogenously determined by the data, and Lumsdaine and Papell (1997) further extended Zivot and Andrews (1992) to the case of two structural breaks at unknown dates.

Structural breaks can affect panel (longitudinal) data unit-root tests in the same way. In response, Karavias and Tzavalis (2014) proposed panel-data unit-root tests that allow for structural breaks in the intercepts of the series or in both the intercepts and linear trends. The break dates are assumed to be common for all series, but the magnitude of the break can differ across series. The null hypothesis is the same as in Zivot and Andrews (1992) and Lumsdaine and Papell (1997); under the null, the panel series are assumed to constitute unit-root processes without breaks, while under the alternative, they are stationary around breaking means or breaking means and trends.

The Karavias and Tzavalis (2014) tests are widely applicable and possess some unique optimality properties, as has been shown in Karavias and Tzavalis (2017, 2019). In terms of applicability, they can be used in both small- and large- T settings, where T is the number of time-series observations. They allow for multiple common breaks, and the dates of the breaks can be known or unknown. In the latter case, they can be endogenously determined from the data. The errors can be nonnormal and have cross-sectional heteroskedasticity and dependence. Finally, the autoregressive coefficients under the alternative can be homogeneous or heterogeneous, which means that either they can be the same for all units or they can differ between units. In terms of their optimality properties, the tests are invariant under the null to the initial condition, which means that no assumptions on the first observations are necessary, as is the case in other fixed- T tests. Furthermore, the tests are invariant to the coefficients of the deterministic components, and are powerful in the presence of linear trends.

This article introduces `xtbunitroot`, a new community-contributed command that implements the panel-data unit-root tests of Karavias and Tzavalis (2014). The command includes the options of one or two breaks at known or unknown dates. This is the first command that allows for panel unit-root tests with structural breaks, and therefore it is a complement to the official `xtunitroot` command and the community-contributed `multipurt` (Eberhardt 2011), `xtfisher` (Merryman 2005), and `pescadf` (Lewandowski 2007) commands.

The breaks can affect the intercepts only or the intercepts and the linear trends of the series. If the dates of the breaks are unknown, then a bootstrap procedure described in Karavias and Tzavalis (2019) is used to derive the critical value and the p -value of the test. Other options include the allowance of cross-section heteroskedasticity, cross-section dependence as in O’Connell (1998), and normal errors. Unbalanced panels are also supported.

The `xtbunitroot` command is applied to examine stationarity in four fundamental bank balance sheet variables frequently used in the banking literature: returns on assets, returns on equity, total assets, and noninterest income. Their stationarity properties affect model building and economic evaluation; see, for example, Kripfganz and Sarafidis (2021) and Delis and Karavias (2015), and in panel forecasting, see, for example, Liu, Moon, and Schorfheide (2020). We examine a sample of 500 randomly selected U.S. banks for the period 2018Q3 to 2020Q4. This period contains the shock of the COVID-19 pandemic and the lockdown, which could have caused a structural break in the series. We find that returns on assets, returns on equity, and noninterest income are stationary, but total assets are nonstationary.

The remainder of the article is organized as follows. In section 2, we present the panel unit-root tests as developed by Karavias and Tzavalis (2014) and their extension to two breaks along the lines of Karavias and Tzavalis (2019). This section also provides some Monte Carlo simulations on the behavior of the tests with two breaks. The simulations have been done using the `xtbunitroot` command. Section 3 describes the syntax of `xtbunitroot`. Section 4 illustrates the command by analyzing the four banking variables. Section 5 concludes the article.

2 Panel unit-root tests with structural breaks

2.1 One break

For panels with N cross-section units, T time-series observations, and one common break, Karavias and Tzavalis (2014) give two models. The first model can be used to test the null hypothesis of a random walk against the alternative hypothesis of a stationary series with a break in the intercepts (means) of the series,

$$\begin{aligned} H_0: & \quad y_{i,t} = y_{i,t-1} + u_{i,t} \\ H_1: & \quad y_{i,t} = \varphi y_{i,t-1} + (1 - \varphi) \{a_{1,i}I(t \leq b) + a_{2,i}I(t > b)\} + u_{i,t} \end{aligned} \quad (1)$$

where $i = 1, \dots, N$ and $t = 1, \dots, T$. In the above model, φ is the autoregressive parameter, and $a_{1,i}$ and $a_{2,i}$ are the fixed effects before and after the break, which happens on date b . The notation $I(\cdot)$ denotes the indicator function.

The second model tests the null hypothesis of a random walk with drift against the alternative of a trend-stationary panel process with a break in the intercepts and linear trends at time b :

$$H_0: \quad y_{i,t} = y_{i,t-1} + \beta_i + u_{i,t}$$

and

$$\begin{aligned}
 H_1: \quad y_{i,t} = & \varphi y_{i,t-1} + \varphi \{ \beta_{1,i} I(t \leq b) + \beta_{2,i} I(t > b) \} \\
 & + (1 - \varphi) \{ a_{1,i} I(t \leq b) + a_{2,i} I(t > b) \} \\
 & + (1 - \varphi) \{ \beta_{1,i} t I(t \leq b) + \beta_{2,i} t I(t > b) \} + u_{i,t}
 \end{aligned} \tag{2}$$

In the above formulation, β_i is the drift under the null hypothesis, while $\beta_{1,i}$ and $\beta_{2,i}$ are the trend coefficients under the alternative hypothesis.

We will henceforth denote by M1 the model with intercepts (1) and by M2 the model with both intercepts and trends (2). For M1, the break is allowed to be in $I_1 = \{1, 2, \dots, T-1\}$, and for M2 the break is allowed to be in $I_2 = \{2, \dots, T-2\}$.¹

The alternative hypothesis is homogeneous across different individuals, but Karavias and Tzavalis (2016) have shown that the test has power against heterogeneous alternatives as well, when $\varphi_i \neq \varphi_j$ and $\varphi_i, \varphi_j < 1$ for $i, j = 1, \dots, N$ and $i \neq j$. Furthermore, Juodis, Karavias, and Sarafidis (2021) argue that pooled estimators can lead to power gains as opposed to mean-group-type estimators like those of Im, Pesaran, and Shin (2003).

Karavias and Tzavalis (2014) propose estimating the autoregressive parameter φ with the following pooled least-squares estimator,

$$\hat{\varphi} = \left(\sum_{i=1}^N \mathbf{y}'_{i,-1} \mathbf{Q}_m^b \mathbf{y}_{i,-1} \right)^{-1} \left(\sum_{i=1}^N \mathbf{y}'_{i,-1} \mathbf{Q}_m^b \mathbf{y}_i \right), \quad m = \{\text{M1}, \text{M2}\}$$

where $\mathbf{y}_i = (y_{i,1}, \dots, y_{i,T})'$ and $\mathbf{y}_{i,-1} = (y_{i,0}, \dots, y_{i,T-1})'$ are $T \times 1$ vectors. The orthogonal projection matrix \mathbf{Q}_m^b is defined as $\mathbf{Q}_m^b = \mathbf{I}_T - \mathbf{X}_m^b (\mathbf{X}_m^{b'} \mathbf{X}_m^b)^{-1} \mathbf{X}_m^{b'}$, where \mathbf{I}_T is the $T \times T$ identity matrix and $\mathbf{X}_{\text{M1}}^b = (\mathbf{e}_1, \mathbf{e}_2)$ and $\mathbf{X}_{\text{M2}}^b = (\mathbf{e}_1, \mathbf{e}_2, \boldsymbol{\tau}_1, \boldsymbol{\tau}_2)$ are $T \times 2$ and $T \times 4$ matrices, respectively, where

$$\mathbf{e}_{1,t} = \begin{cases} 1 & \text{if } t \leq b \\ 0 & \text{elsewhere} \end{cases}, \quad \mathbf{e}_{2,t} = \begin{cases} 1 & \text{if } t > b \\ 0 & \text{elsewhere} \end{cases}$$

and

$$\boldsymbol{\tau}_{1,t} = \begin{cases} t & \text{if } t \leq b \\ 0 & \text{elsewhere} \end{cases}, \quad \boldsymbol{\tau}_{2,t} = \begin{cases} t & \text{if } t > b \\ 0 & \text{elsewhere} \end{cases}$$

The superscripts b in \mathbf{Q}_m^b and \mathbf{X}_m^b denote dependence on the break date.

The estimator $\hat{\varphi}$ is asymptotically inconsistent; therefore, it must be modified. If the date of the break b is known, the following statistic is asymptotically normally distributed as $N \rightarrow \infty$,

$$Z(b) = \sqrt{N} \{C^b(k_u, \sigma_u^2)\}^{-1/2} (\hat{\varphi} - B^b - 1) \xrightarrow{L} N(0, 1)$$

1. Notice that in the above setup, we have a total of $T+1$ observations to simplify notation. In the presentation of the `xtbunitroot` command and in the example below, we assume that the total number of time-series observations is T , and therefore, I_1 and I_2 become $I_1 = \{2, \dots, T-1\}$ and $I_2 = \{3, \dots, T-2\}$.

where B^b is the bias correction and C^b is the variance of the bias-corrected (modified) estimator. The parameters k_u and σ_u^2 are error moments to be estimated, but if the errors are normally distributed, they are removed from C^b , which becomes a simple function of the break date and the number of time-series observations. Normality also removes any nuisance parameters related to cross-section heteroskedasticity. The assumptions for the above result are mild, with the strongest being that there is no residual serial correlation in the errors $u_{i,t}$.

To deal with cross-sectional dependence in error terms, one popular approach is demeaning the series across i by subtracting the cross-sectional averages for all t before conducting the test (see, for example, O'Connell [1998]),

$$\tilde{y}_{i,t} = y_{i,t} - \bar{y}_t$$

where

$$\bar{y}_t = \frac{1}{N} \sum_{i=1}^N y_{i,t}$$

If the date of the break is unknown, Karavias and Tzavalis (2014) follow Zivot and Andrews (1992) and propose the following statistic to test for unit roots:

$$\min \mathcal{Z} = \min_{b \in I_m} Z(b) \text{ for } m = \{M_1, M_2\}$$

The limiting distribution of this statistic is shown to depend on the time dimension T . Following Karavias and Tzavalis (2019), the `xtbunitroot` command implements a bootstrap algorithm to derive the critical values and p -values of $\min \mathcal{Z}$. The asymptotic distribution is valid for T fixed and $N \rightarrow \infty$. The `xtbunitroot` command also reports the date with the most evidence against the null hypothesis $\hat{b} = \operatorname{argmin}_{b \in I_m} Z(b)$. Notice, however, that the break date estimator is not consistent. For consistent break date estimation in small or large T , one may use the `xtbreak` command of Ditzen, Karavias, and Westerlund (2021) (see also Karavias, Narayan, and Westerlund [Forthcoming]).

The `xtbunitroot` command supports unbalanced panels. In this case, the missing values in the formulas are zeroed out because this method maximizes the power of tests; see, for example, Karavias, Tzavalis, and Zhang (2022).

2.2 Extension to two-break case

The alternative hypotheses in M_1 and M_2 allow for two breaks. In this case, (1) and (2) become

$$\begin{aligned} H_1: \quad y_{i,t} = & \varphi y_{i,t-1} \\ & + (1 - \varphi) \{a_{1,i}I(t \leq b_1) + a_{2,i}I(b_1 < t \leq b_2) + a_{3,i}I(t > b_2)\} + u_{i,t} \end{aligned} \quad (3)$$

and

$$\begin{aligned}
 H_1: \quad y_{i,t} = & \varphi y_{i,t-1} + \varphi \{ \beta_{1,i} I(t \leq b_1) + \beta_{2,i} I(b_1 < t \leq b_2) + \beta_{3,i} I(t > b_2) \} \\
 & + (1 - \varphi) \{ a_{1,i} I(t \leq b_1) + a_{2,i} I(b_1 < t \leq b_2) + a_{3,i} I(t > b_2) \} \\
 & + (1 - \varphi) \{ \beta_{1,i} t I(t \leq b_1) + \beta_{2,i} t I(b_1 < t \leq b_2) + \beta_{3,i} t I(t > b_2) \} + u_{i,t} \quad (4)
 \end{aligned}$$

The pooled least-squares estimator is based on the $\mathbf{Q}_m^{b_1, b_2}$ orthogonal projection matrix, where $\mathbf{Q}_m^{b_1, b_2} = \mathbf{I}_T - \mathbf{X}_m^{b_1, b_2} (\mathbf{X}_m^{b_1, b_2'} \mathbf{X}_m^{b_1, b_2})^{-1} \mathbf{X}_m^{b_1, b_2'}$ and where $\mathbf{X}_{M1}^{b_1, b_2} = (\mathbf{e}_1, \mathbf{e}_2, \mathbf{e}_3)$ and $\mathbf{X}_{M2}^{b_1, b_2} = (\mathbf{e}_1, \mathbf{e}_2, \mathbf{e}_3, \boldsymbol{\tau}_1, \boldsymbol{\tau}_2, \boldsymbol{\tau}_3)$ are $T \times 3$ and $T \times 6$ matrices, respectively, with

$$e_{1,t} = \begin{cases} 1 & \text{if } t \leq b_1 \\ 0 & \text{elsewhere} \end{cases}, \quad e_{2,t} = \begin{cases} 1 & \text{if } b_1 < t \leq b_2 \\ 0 & \text{elsewhere} \end{cases}, \quad e_{3,t} = \begin{cases} 1 & \text{if } t > b_2 \\ 0 & \text{elsewhere} \end{cases}$$

The vector $\boldsymbol{\tau}_t$ is defined as

$$\tau_{1,t} = \begin{cases} t & \text{if } t \leq b_1 \\ 0 & \text{elsewhere} \end{cases}, \quad \tau_{2,t} = \begin{cases} t & \text{if } b_1 < t \leq b_2 \\ 0 & \text{elsewhere} \end{cases}, \quad \tau_{3,t} = \begin{cases} t & \text{if } t > b_2 \\ 0 & \text{elsewhere} \end{cases}$$

The distributions of the test statistics are as before; $Z(b_1, b_2)$ is standard normal when b_1 and b_2 are known, and

$$\min \mathcal{Z} = \min_{b_1, b_2 \in I_m} Z(b_1, b_2) \text{ for } m = \{M_1, M_2\}$$

can be used when the dates of the breaks are unknown. Its limiting distribution can be calculated using the bootstrap. Notice that b_1 and b_2 cannot be in two consecutive dates for M2, which is the same situation as in a single time series; see, for example, Lumsdaine and Papell (1997). The `xtbunitroot` command implements a bootstrap algorithm to derive the critical values and p -values. This asymptotic distribution is valid for T fixed and $N \rightarrow \infty$, and the combination of dates with the most evidence against the null hypothesis $\{\hat{b}_1, \hat{b}_2\} = \operatorname{argmin}_{b_1, b_2 \in I_m} Z(b)$ is reported.

To provide some evidence on the behavior of the test statistic with two breaks, we conduct a small Monte Carlo experiment. Consider the data-generation process in (1) and (2) under the null and (3) and (4) under the alternative. We assume that the error term $u_{i,t} \sim$ independent and identically distributed $N(0, 1)$. The panel-data series $y_{i,t}$ is generated with autoregressive coefficient $\varphi = 1$ to investigate the size at 5% significance level and $\varphi = \{0.90, 0.95\}$ to investigate the power of the test. As we set φ very close to 1, it will be hard to reject the null under $H_1: \varphi < 1$. The initial values $y_{i,0}$ are generated as $y_{i,0} \sim$ independent and identically distributed $N(0, 1)$. For the M1 and M2 models, we generate the intercept and trend coefficients as follows: $a_{1,i} \sim U(-0.5, 0)$, $a_{2,i} \sim U(0, 0.5)$, and $a_{3,i} \sim U(0.5, 1)$. The slope coefficients for the linear trends are generated as $\beta_{1,i} \sim U(0, 0.025)$, $\beta_{2,i} \sim U(0.025, 0.05)$, and $\beta_{3,i} \sim U(0.05, 0.075)$. These parameter choices follow Karavias and Tzavalis (2014) and are made to make it harder to reject the null $H_0: \phi = 1$. We set break dates to the following fractions of the sample, $\lambda_1 = 0.25, \lambda_2 = 0.75$. The number of bootstrap replications is set to 100, which is the default in `xtbunitroot`.

The results of the exercise appear in table 1. We can see that for both known and unknown breaks, the size of the tests is always very close to its nominal 5% level, and the power is satisfactory and increasing with both N and T .

Table 1. Size and power of the tests for two known and unknown breaks

		Panel A: Known breaks								
		$\varphi = 1$			$\varphi = 0.95$			$\varphi = 0.90$		
	(N, T)	10	25	50	10	25	50	10	25	50
M1	25	0.060	0.057	0.071	0.180	0.359	0.719	0.311	0.684	0.990
	50	0.055	0.062	0.053	0.226	0.566	0.915	0.441	0.918	1.000
	100	0.050	0.054	0.061	0.355	0.809	0.996	0.687	0.995	1.000
M2	25	0.056	0.059	0.070	0.062	0.087	0.184	0.077	0.170	0.607
	50	0.052	0.051	0.060	0.059	0.093	0.258	0.077	0.240	0.840
	100	0.055	0.061	0.057	0.064	0.117	0.384	0.095	0.375	0.985
		Panel B: Unknown breaks								
		$\varphi = 1$			$\varphi = 0.95$			$\varphi = 0.90$		
	(N, T)	10	15	25	10	15	25	10	15	25
M1	25	0.030	0.030	0.030	0.070	0.250	0.410	0.270	0.580	0.770
	50	0.020	0.050	0.070	0.300	0.440	0.700	0.580	0.810	0.970
	100	0.020	0.070	0.040	0.400	0.720	0.930	0.810	0.990	1.000
M2	25	0.080	0.050	0.060	0.080	0.070	0.060	0.100	0.100	0.130
	50	0.060	0.080	0.050	0.070	0.060	0.050	0.070	0.100	0.080
	100	0.060	0.040	0.010	0.050	0.040	0.040	0.050	0.080	0.200

NOTES: The reported values are rejection probabilities. For $\varphi = 1$, the reported rejection rates give the size of the test, while for $\varphi < 1$, they give the power of the test. The M1 model includes intercepts, while the M2 model includes both intercepts and linear trends. The dates of the breaks are $b_1 = \lfloor 0.25T \rfloor$, $b_2 = \lfloor 0.75T \rfloor$.

3 The xtbunitroot command

3.1 Syntax

```
xtbunitroot varname [if] [in] [, trend known(#1 [#2]) unknown(#3 [#4])
normal csd het nobootstrap showindex level(#) seed(#)]
```

where *varname* is the variable to be tested for nonstationarity. You must `xtset` your data before using `xtbunitroot`. If no option is specified, the default will be the model

with a single break in the intercepts, at an unknown date: `xtbunitroot varname, unknown(1 100)`. *varname* can contain time-series operators.

3.2 Options

trend specifies that the deterministic component of the model include individual intercepts (means) and individual linear trends. The common breaks affect both intercepts and trends. Breaks in consecutive dates are not allowed in this model. In this model, the breaks can be in dates from 3 to $T - 2$, while in the model with intercepts, the breaks can take place from 2 up to $T - 1$.

known($\#_1$ [$\#_2$]) specifies the number and places of breaks. This option implements the case when the dates of the breaks are known. The $\#_1$ input specifies the location of the first break, and the [$\#_2$] input specifies the location of the second break. If only one break is assumed, **known**($\#_1$) can be used. The inputs should be in terms of time ordering (from 1 to T), instead of using dates, that is, 1995 or 2020. The break dates should be in order. For example, **known**(5) means the break occurs in period 5, and **known**(3 7) means the first break occurs in period 3 and the second break occurs in period 7.

unknown($\#_3$ [$\#_4$]) specifies the number of breaks and the number of bootstrap replications. This option is used when the dates of the breaks are unknown to the researcher. The number of bootstrap replications, [$\#_4$], can be omitted. The option **unknown**(2) states that there are two unknown breaks and the critical and p -values will be calculated based on the default number of bootstrap replications, which is set to 100.

normal specifies that the errors be normally distributed.

csd subtracts the cross-section averages for each time period and applies the tests in the demeaned series.

het specifies that the errors be cross-sectionally heteroskedastic. If both **het** and **normal** are specified, the results will be the same as in the case that only **normal** is used, because if the errors are normal, then heteroskedastic variances drop out.

nobootstrap prevents the command from running the bootstrap. The bootstrap is necessary for calculating critical values and p -values for the test when the dates of the breaks are unknown. However, if the dataset is very large, then the bootstrap can be time consuming. This option stops the bootstrap but returns the $\min Z$ statistic and the estimated break dates. The $\min Z$ statistic can then be compared with the approximate critical values that appear in table 1 of Karavias and Tzavalis (2014), which is reported in the command output. This option can be used only with the option **normal** because the available critical values are for normally distributed errors.

`showindex` specifies that the estimated break date be reported as a time index. For example, an estimated break point at the 9th observation will be reported as 9 instead of 2021q4.

`level(#)` specifies the level of the test used. The default is `level(5)`, and all integers between 1 and 99 are applicable. For example, `level(10)` sets the level of the one-sided null hypothesis to 10%.

`seed(#)` specifies the seed used in the bootstrap process for the case of unknown breaks. The default is `seed(123)`. Seed is important for reproducing bootstrap-based results.

3.3 Stored results

`xtbunitroot` stores the following in `r()`:

Scalars

<code>r(N)</code>	number of total observations
<code>r(N_g)</code>	number of cross-sectional units
<code>r(T)</code>	number of time periods
<code>r(breaks)</code>	number of breaks
<code>r(break1)</code>	time index of the first break
<code>r(break2)</code>	time index of the second break
<code>r(seed)</code>	seed
<code>r(Z)</code>	Z or $\min Z$ statistic
<code>r(pvalue)</code>	p -value for the Z or $\min Z$ statistic
<code>r(cv)</code>	asymptotic or bootstrap critical value
<code>r(boot)</code>	number of bootstrap replications
<code>r(fihat)</code>	estimate of autoregressive parameter
<code>r(khat)</code>	estimate of k
<code>r(shat)</code>	estimate of error variance

Macros

<code>r(idvar)</code>	name of panel ID variable
<code>r(tvar)</code>	name of panel time variable
<code>r(varname)</code>	name of tested variable
<code>r(model)</code>	type of model: <code>constant</code> or <code>trend</code>
<code>r(date1)</code>	date of the first break
<code>r(date2)</code>	date of the second break
<code>r(avert)</code>	average number of time periods

Matrices

<code>r(kui)</code>	individual k for heteroskedastic errors
<code>r(sigmai)</code>	individual variances for heteroskedastic errors

4 Example

4.1 Unit-root tests for bank balance sheet variables

Unit-root processes were originally important for macroeconomic variables. However, as Holtz-Eakin, Newey, and Rosen (1988) argue, such dynamic relationships can appear in other economic variables for which long time series may not be available. In this case, the panel dimension of the data can be used for inference.

In this section, we focus on bank balance sheet variables taken from the call reports of the Federal Deposit Insurance Corporation. This is a very popular dataset; see, for example, Kripfganz and Sarafidis (2021) and Juodis, Karavias, and Sarafidis (2021) for some recent applications. However, while stationarity of the bank balance sheet variables is frequently assumed, it is almost never tested. The contribution of this section is to examine the stationarity of four variables of interest, namely, returns on assets, returns on equity, total assets, and noninterest income. The returns on assets (**roa**) are defined as net income after taxes and extraordinary items and is given as a percent of average total assets. The returns on equity (**roe**) are defined as net income over average total equity; total assets (**tassets**) are the year-to-date average of total assets; and noninterest income (**nii**) is defined as income derived from bank services and sources other than interest bearing assets, over average total assets.

We collect data for a random sample of 500 banks from the third quarter of 2018 to the fourth quarter of 2020. This period includes the COVID-19 pandemic, which may have caused breaks in the intercepts and trends of the series. The short dimension of the data is chosen so that our sample of banks does not suffer from survivorship bias. The data are publicly available, and they have been downloaded from the Federal Deposit Insurance Corporation website.²

In the following, we perform panel unit-root tests allowing for structural breaks in the aforementioned variables. We assume that our sample is large enough so that the idiosyncratic errors $u_{i,t}$ are normally distributed. The errors can be cross-sectionally heteroskedastic, but under the normality assumption, the tests are invariant to heteroskedasticity. Finally, we assume error cross-section dependence, caused possibly by the monetary policy rate.

We start the analysis with the **roa** series. First, we assume that the date of the break is known to be the first quarter of 2020. Second, we allow the break to be unknown and determined from the data. In the latter case, the critical values are taken using 100 bootstrap samples, which is the default. The results are given below:

2. See <https://www.fdic.gov/>. The sample used is available at <https://sites.google.com/site/yianniskaravias/> and the Boston College Statistical Software Components Archive.

```
. use xtbunitroot_example
. xtset fed_rssd time
Panel variable: fed_rssd (strongly balanced)
Time variable: time, 1 to 10
Delta: 1 unit

. xtbunitroot roa, known(7) normal csd
Karavias and Tzavalis (2014) panel unit root test for roa
```

H0: All panel time series are unit root processes
H1: Some or all of the panel time series are stationary processes

Number of panels:	500	Avrge number of periods:	10.00
Number of breaks:	1		
Cross-section dependence:	Yes	Linear time trend:	No
Cross-section heteroskedasticity:	No	Normal errors:	Yes

	Statistic	Asymtotic critical-value	p-value
Z-statistic	-15.5212	-1.6449	0.0000

Result: the null is rejected
Known break date(s): 7
Significance level of test: .05

The output above indicates that the $Z(b)$ statistic is equal to -15.521 , which is far less than the critical value of -1.645 ; therefore, we can reject the null hypothesis of nonstationarity. The break under the alternative takes place in observation 7, which corresponds to the first quarter of 2020. The output also reports the result “the null is rejected”, which is the outcome at the 5% significance level. The following output presents the results for *roa*, when the date of the break is unknown.

```
. xtbunitroot roa, unknown(1) normal csd
Karavias and Tzavalis (2014) panel unit root test for roa
```

H0: All panel time series are unit root processes
H1: Some or all of the panel time series are stationary processes

Number of panels:	500	Avrge number of periods:	10.00
Number of breaks:	1	Bootstrap replications:	100
Cross-section dependence:	Yes	Linear time trend:	No
Cross-section heteroskedasticity:	No	Normal errors:	Yes

	Statistic	Bootstrap critical-value	p-value
minZ-statistic	-25.1619	8.5546	0.0000

Result: the null is rejected
Estimated break date(s): 6
Significance level of test: .05

We can see from the above output that the $\min Z$ statistic is equal to -25.162 , which is much smaller than the bootstrap critical value of 8.555 ; therefore, we once more reject the null hypothesis. The p -value is also reported and implies that the null hypothesis is rejected at the 1% significance level. The date with the most evidence against the null is observation 6, which corresponds to the fourth quarter of 2019, although this estimator is not expected to provide a consistent estimate of the break date, as mentioned earlier.

For the remaining three variables, we will implement the tests being agnostic about the date of the break. For `tassets` and `nii`, we also include linear trends.

```
. xtunitroot roe, unknown(1) normal csd
Karavias and Tzavalis (2014) panel unit root test for roe
```

```
H0: All panel time series are unit root processes
H1: Some or all of the panel time series are stationary processes
```

Number of panels:	500	Avrge number of periods:	10.00
Number of breaks:	1	Bootstrap replications:	100
Cross-section dependence:	Yes	Linear time trend:	No
Cross-section heteroskedasticity:	No	Normal errors:	Yes

	Statistic	Bootstrap critical-value	p-value
minZ-statistic	-20.9965	-19.4133	0.0100

```
Result: the null is rejected
Estimated break date(s):      8
Significance level of test: .05

. xtunitroot tassets, unknown(1) normal csd trend
Karavias and Tzavalis (2014) panel unit root test for tassets
```

```
H0: All panel time series are unit root processes
H1: Some or all of the panel time series are stationary processes
```

Number of panels:	500	Avrge number of periods:	10.00
Number of breaks:	1	Bootstrap replications:	100
Cross-section dependence:	Yes	Linear time trend:	Yes
Cross-section heteroskedasticity:	No	Normal errors:	Yes

	Statistic	Bootstrap critical-value	p-value
minZ-statistic	-8.3437	-69.5491	0.6200

```
Result: the null is not rejected
Significance level of test: .05
```

```
. xtbunitroot nii, unknown(1) normal csd trend
Karavias and Tzavalis (2014) panel unit root test for nii
```

```
H0: All panel time series are unit root processes
H1: Some or all of the panel time series are stationary processes
```

Number of panels:	500	Avrge number of periods:	10.00
Number of breaks:	1	Bootstrap replications:	100
Cross-section dependence:	Yes	Linear time trend:	Yes
Cross-section heteroskedasticity:	No	Normal errors:	Yes

	Statistic	Bootstrap critical-value	p-value
minZ-statistic	-26.7284	-18.3424	0.0000

```
Result: the null is rejected
Estimated break date(s):      6
Significance level of test: .05
```

As we can see from the above outputs, `roe` and `nii` are stationary, while `tassets` is not. When the null hypothesis is not rejected at the 5% level, the output does not report the break date; however, this date can be recovered from `r(break1)`. In smaller datasets, one may wish to employ higher levels of significance using the `level()` option. For `tassets`, the date with the most evidence against the null is the first quarter of 2019; however, the *p*-value is high, and we cannot reject the null hypothesis.

5 Concluding remarks

This article introduced a new community-contributed command, `xtbunitroot`, that implements the panel-data unit-root tests with structural breaks, developed by Karavias and Tzavalis (2014). This is the first command that allows for panel unit-root tests with structural breaks; it allows for one or two breaks under an alternative that can be known or unknown. It also allows for linear trends, cross-sectional heteroskedasticity and dependence, and nonnormal errors. The `xtbunitroot` command was applied to four bank balance sheet variables, and it was found that returns on assets, returns on equity, and noninterest income are stationary variables, but total assets are nonstationary.

6 Programs and supplemental materials

To install a snapshot of the corresponding software files as they existed at the time of publication of this article, type

```
. net sj 22-3
. net install st0687      (to install program files, if available)
. net get st0687          (to install ancillary files, if available)
```

7 References

- Banerjee, A., R. L. Lumsdaine, and J. H. Stock. 1992. Recursive and sequential tests of the unit-root and trend-break hypotheses: Theory and international evidence. *Journal of Business and Economic Statistics* 10: 271–287. <https://doi.org/10.2307/1391542>.
- Delis, M. D., and Y. Karavias. 2015. Optimal versus realized bank credit risk and monetary policy. *Journal of Financial Stability* 16: 13–30. <https://doi.org/10.1016/j.jfs.2014.11.004>.
- Dickey, D. A., and W. A. Fuller. 1979. Distribution of the estimators for autoregressive time series with a unit root. *Journal of the American Statistical Association* 74: 427–431. <https://doi.org/10.2307/2286348>.
- Ditzen, J., Y. Karavias, and J. Westerlund. 2021. Testing and estimating structural breaks in time series and panel data in Stata. ArXiv Working Paper No. arXiv:2110.14550. <https://arxiv.org/abs/2110.14550>.
- Eberhardt, M. 2011. multiptur: Stata module to run 1st and 2nd generation panel unit root tests for multiple variables and lags. Statistical Software Components S457239, Department of Economics, Boston College. <https://ideas.repec.org/c/boc/bocode/s457239.html>.
- Granger, C. W. J., and P. Newbold. 1974. Spurious regressions in econometrics. *Journal of Econometrics* 2: 111–120. [https://doi.org/10.1016/0304-4076\(74\)90034-7](https://doi.org/10.1016/0304-4076(74)90034-7).
- Holtz-Eakin, D., W. Newey, and H. S. Rosen. 1988. Estimating vector autoregressions with panel data. *Econometrica* 56: 1371–1395. <https://doi.org/10.2307/1913103>.
- Im, K. S., M. H. Pesaran, and Y. Shin. 2003. Testing for unit roots in heterogeneous panels. *Journal of Econometrics* 115: 53–74. [https://doi.org/10.1016/S0304-4076\(03\)00092-7](https://doi.org/10.1016/S0304-4076(03)00092-7).
- Juodis, A., Y. Karavias, and V. Sarafidis. 2021. A homogeneous approach to testing for Granger non-causality in heterogeneous panels. *Empirical Economics* 60: 93–112. <https://doi.org/10.1007/s00181-020-01970-9>.
- Karavias, Y., P. K. Narayan, and J. Westerlund. Forthcoming. Structural breaks in interactive effects panels and the stock market reaction to COVID-19. *Journal of Business and Economic Statistics*. <https://doi.org/10.1080/07350015.2022.2053690>.
- Karavias, Y., and E. Tzavalis. 2014. Testing for unit roots in short panels allowing for a structural break. *Computational Statistics and Data Analysis* 76: 391–407. <https://doi.org/10.1016/j.csda.2012.10.014>.
- . 2016. Local power of fixed-T panel unit root tests with serially correlated errors and incidental trends. *Journal of Time Series Analysis* 37: 222–239. <https://doi.org/10.1111/jtsa.12144>.

- . 2017. Local power of panel unit root tests allowing for structural breaks. *Econometric Reviews* 36: 1123–1156. <https://doi.org/10.1080/07474938.2015.1059722>.
- . 2019. Generalized fixed-T panel unit root tests. *Scandinavian Journal of Statistics* 46: 1227–1251. <https://doi.org/10.1111/sjos.12392>.
- Karavias, Y., E. Tzavalis, and H. Zhang. 2022. Missing values in panel data unit root tests. *Econometrics* 10: 12. <https://doi.org/10.3390/econometrics10010012>.
- Kripfganz, S., and V. Sarafidis. 2021. Instrumental-variable estimation of large-T panel-data models with common factors. *Stata Journal* 21: 659–686. <https://doi.org/10.1177/1536867X211045558>.
- Lewandowski, P. 2007. pescadf: Stata module to perform Pesaran's CADF panel unit root test in presence of cross section dependence. Statistical Software Components S456732, Department of Economics, Boston College. <https://ideas.repec.org/c/boc/bocode/s456732.html>.
- Liu, L., H. R. Moon, and F. Schorfheide. 2020. Forecasting with dynamic panel data models. *Econometrica* 88: 171–201. <https://doi.org/10.3982/ECTA14952>.
- Lumsdaine, R. L., and D. H. Papell. 1997. Multiple trend breaks and the unit-root hypothesis. *Review of Economics and Statistics* 79: 212–218. <https://doi.org/10.1162/003465397556791>.
- Merryman, S. 2005. xtfisher: Stata module to compute Fisher type unit root test for panel data. Statistical Software Components S448201, Department of Economics, Boston College. <https://ideas.repec.org/c/boc/bocode/s448201.html>.
- O'Connell, P. G. J. 1998. The overvaluation of purchasing power parity. *Journal of International Economics* 44: 1–19. [https://doi.org/10.1016/S0022-1996\(97\)00017-2](https://doi.org/10.1016/S0022-1996(97)00017-2).
- Perron, P. 1989. The great crash, the oil price shock, and the unit root hypothesis. *Econometrica* 57: 1361–1401. <https://doi.org/10.2307/1913712>.
- Zivot, E., and D. W. K. Andrews. 1992. Further evidence on the Great Crash, the oil price shock, and the unit-root hypothesis. *Journal of Business and Economic Statistics* 10: 251–270. <https://doi.org/10.2307/1391541>.

About the authors

Pengyu Chen is a full-time student in the MSc in economics program at the London School of Economics and Political Science.

Yiannis Karavias is an assistant professor in financial economics at the Birmingham Business School in the University of Birmingham.

Elias Tzavalis is a professor of economics at the Department of Economics at the Athens University of Economics and Business.