

Toxicity of ceria nanoparticles to the regeneration of freshwater planarian *Dugesia japonica*

Xie, Changjian; Li, Xiaowei; Hei, Lisha; Chen, Yiqing; Dong, Yuling; Zhang, Shujing; Ma, Shan; Xu, Jianing; Pang, Qiuxiang; Lynch, Iseult; Guo, Zhiling; Zhang, Peng

DOI:

[10.1016/j.scitotenv.2022.159590](https://doi.org/10.1016/j.scitotenv.2022.159590)

License:

Creative Commons: Attribution-NonCommercial-NoDerivs (CC BY-NC-ND)

Document Version

Peer reviewed version

Citation for published version (Harvard):

Xie, C, Li, X, Hei, L, Chen, Y, Dong, Y, Zhang, S, Ma, S, Xu, J, Pang, Q, Lynch, I, Guo, Z & Zhang, P 2023, 'Toxicity of ceria nanoparticles to the regeneration of freshwater planarian *Dugesia japonica*: The role of biotransformation', *Science of the Total Environment*, vol. 857, 159590. <https://doi.org/10.1016/j.scitotenv.2022.159590>

[Link to publication on Research at Birmingham portal](#)

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

***Toxicity of ceria nanoparticles to the regeneration of freshwater
planarian Dugesia japonica: the role of biotransformation***

Changjian Xie ^{a,*}, Xiaowei Li ^a, Lisha Hei ^a, Yiqing Chen ^a, Yuling Dong ^a, Shujing Zhang ^a, Shan Ma ^d, Jianing Xu ^a, Qiuxiang Pang ^{a,*}, Iseult Lynch ^b, Zhiling Guo ^{b,*}, and Peng Zhang ^{b, c}

^a School of life Sciences and medicine, Shandong University of Technology, Zibo 255000, Shandong, China

^b School of Geography, Earth & Environmental Sciences, University of Birmingham, Edgbaston, Birmingham B15 2TT, United Kingdom

^c Department of Environmental Science and Engineering, University of Science and Technology of China, Hefei 230026, China

^d Zibo Environment Monitoring Center, Zibo25500, Shandong, China

* Corresponding authors:

E-mail addresses: xiecj@sdut.edu.cn (Changjian Xie), pangqiuxiang@sdut.edu.cn (Qiuxiang Pang), z.guo@bham.ac.uk (Zhiling Guo).

Abstract

Cerium oxide nanoparticles (n-CeO₂) have wide applications ranging from industrial to consumer products, which would inevitably lead to their release into the environment. Despite the toxicity of n-CeO₂ on aquatic organisms has been largely reported, research on developing organisms is still lacking. In this study, we investigate the toxic effects of n-CeO₂ on the stem cells, tissue- and neuro-regeneration, using freshwater planarian *Dugesia japonica* as a model. Effects of bulk sized (μ -) CeO₂ and ionic Ce (Ce³⁺) were compared with that of n-CeO₂ to explore the origin of the toxic effects of n-CeO₂. No overt toxicity was observed in μ -CeO₂ treatment. n-CeO₂ not only impaired the homeostasis of normal planarians, but also inhibited the regeneration processes of regenerated planarians, demonstrated by the inhibited blastema growth, disturbed antioxidant defense system at molecular levels, elevated DNA-damage and decreased stem cell proliferation. Regenerating organisms are more susceptible to n-CeO₂ than the normal ones. Ce³⁺ exhibited significantly higher toxicity than n-CeO₂, even though the total Ce uptake is 0.2% less in Ce³⁺ than in n-CeO₂ treated in planarian. X-ray absorption near edge spectroscopy (XANES) analysis revealed that 12.8% of n-CeO₂ (5.95 mg/kg Ce per planarian) was transformed to Ce³⁺ after interaction with planarian, suggesting that biotransformation at the nano-bio interface might play an important role in the observed toxicity. Since the biotransformation of n-CeO₂ is a slow process, it may cause long-term chronic toxicity to planarians due to the slow whilst sustained release of toxic Ce³⁺ ions.

Keywords: n-CeO₂, toxicity, planarian, XANES; Regeneration

1. Introduction

The rapid development of nanotechnology promotes the applications of nanomaterials in various fields. Worldwide, about 10, 000 metric tons of cerium oxide nanoparticles (n-CeO₂) are produced per year¹. n-CeO₂ are used in numerous applications such as biomedical fields, fuel additives, gas sensors, polishing agent, and metal-oxide semiconductor devices^{2, 3}. The increasing production and widespread application would inevitably lead to the release of n-CeO₂ into the environment, causing concerns for the potential toxicological risks to aquatic organisms as well as human health⁴. Developing organisms and tissues are often more susceptible to environmental pollutants including nanoparticles⁵.

The bio-effects of n-CeO₂ remain controversial. On one hand, n-CeO₂ is reported to have

inherent or acquired ability to scavenge reactive oxygen species (ROS)⁶⁻⁸, which was equivalent to the antioxidant enzymes in biological systems^{7,9}. For example, n-CeO₂ can induce the proliferation of stem cells of mouse¹⁰ and human¹¹ by regulating the intracellular ROS and the relative expression of genes responsible for the stem cell proliferation, migration and antioxidant defense system. Singh, et al.¹² also confirmed that n-CeO₂ can scavenge ROS accumulation and regulate the status of cell redox and thereby accelerating the growth in human keratinocytes (HaCaT cells). On the contrary, studies also showed that n-CeO₂ could exhibit adverse effects both *in vivo* and *in vitro*¹³⁻¹⁵. For instance, n-CeO₂ induced ROS accumulation and caused oxidative stress in *Caenorhabditis elegans* at environmental relevant concentration (1 nM), and ultimately led to a decreased lifespan¹³. The toxicity of n-CeO₂ depends on the species, dose, and the physiochemical property of n-CeO₂^{14, 16-20}. The biological fates of n-CeO₂ are still unclear. Some studies suggest that the primary bioavailable form of n-CeO₂ in tissue is particulate form rather than the ionic (Ce³⁺)²¹ as n-CeO₂ was considered as highly insoluble under environmental conditions²². Studies based on synchrotron radiation techniques have revealed the biotransformation of n-CeO₂ in plants and bacteria²³⁻²⁵. Is the transformed n-CeO₂ responsible for the toxicity? This issue needs to be urgent studied.

Developing organisms and tissues are often more susceptible to environmental pollutants than mature organisms^{26, 27}. Stem cells, which undergo rapid cell division, are highly vulnerable to environmental chemicals, impairing their differentiation and development²⁸. However, vertebrate experimental protocols need to be more prepared in terms of licensing and ethics. In fact, planarian as an alternative invertebrate organism has been proven to be an excellent stem cell model for more than 20 years²⁹. Planarian has large numbers of pluripotent stem cells (20-30%) and can regenerate any body part including the whole head and the entire central nervous system³⁰, which makes them a suitable model for studying developmental (neuro-) toxicity³¹. In addition, planarian can provide early warning of harmful substances even at very low concentrations, thus, it is widely used as an indicator for water quality^{5, 32}. As a benthic animal in the water body, planarians use their motile cilia to glide over sediment surfaces³³ and interacted with the deposited nanoparticles (NPs), which may make them an excellent model for investigating the effects of NPs on aquatic animals.

Planarians have been used for assessing nanotoxicity of Ag nanoparticles⁵, polystyrene microplastics^{34, 35}, and Fe₃O₄ nanoparticles³⁶. For instance, Kustov, et al.³⁷ found that Ag NPs was moderately toxic to the planarian with a dose-dependent effect. Compared with regeneration

planarians, the behavioral changes such as side-lying and screw-like movements, was only happened at the highest concentration (50 mg/L) in the homeostatic planarians, while the stem cell proliferation was not affected⁵. In a previous study, Fe₃O₄ NPs had not affected the stem cell population dynamics, nor did they induce significant changes in either homeostatic or regenerative planarians³⁶. Instead, Ermakov, et al.³⁸ found that CeO₂ NPs (an average size of 1-2 nm) may play an inorganic mitogens role in stimulating *Schmidtea mediterranea* (planarian) regeneration at 10⁻¹¹ M colloid solution, and CeF₃ NPs (F⁻ can stabilize Ce³⁺ that makes Ce³⁺ less prone to oxidation, $K_{sp}=8 \times 10^{-16}$ of CeF₃) also did not show any toxic to the homeostatic and regeneration planarians³⁹. In fact, the toxicity of n-CeO₂ is related to its physicochemical properties, such as size, shapes, concentrations, Ce³⁺ released and species-specific. However, only little is known about the effects of n-CeO₂ sizes, doses, and biotransformation on the (neuro-) development toxicity of the aquatic organisms. In this study, we studied the toxicity of n-CeO₂ to freshwater planarian *Dugesia japonica* and compared their toxic effects to bulk-CeO₂ (μ -CeO₂), ionic Ce (Ce³⁺) and control treatment (CT) on stem cells, tissue regeneration and development *in vivo*. We also used X-ray absorption near-edge spectroscopy (XANES) to identify the chemical species to examine the role of biotransformation in the observed biological effects.

2. Materials and Methods

2.1 Chemical and animals. n-CeO₂ (nominal size < 25nm, determined by nitrogen sorption, purity 99.9%, manufacturer's data) were purchased from Sigma-Aldrich (St. Louis, MO, USA). As previously reported⁴⁰, the particles were 100% cubic ceria with diameters ranging from 6.2 nm up to 48 nm, with the center of the distribution at 16.3 nm. The diameters of μ -CeO₂ (nominal size < 5 μ m, 99.9% trace metal basis, Sigma-Aldrich) varied from 237.5 nm to 1320 nm, and the distribution center was 473.5 nm (**Fig. S1**). Hydrodynamic size of n-CeO₂ and μ -CeO₂ in DI water and MW medium were 138.2 ± 31.8 nm and 2703.4 ± 147.6 nm, 162.3 ± 42.5 nm and 2863.2 ± 258.7 nm respectively. The zeta potential were 13.8 ± 4.0 mV and 11.5 ± 5.7 mV in DI water and 11.3 ± 2.9 mV and 9.8 ± 3.2 mV in MW medium for n-CeO₂ and μ -CeO₂ respectively. Cerium (III) chloride (CAS no. 7790-86-5, > 99.99%) was also purchased from Sigma-Aldrich.

Planarians *D. japonica* used in the study were collected from a spring in Yiyuan (Zibo, China), and acclimated in mineral water (MW, Lushan fountain water, Zibo, China) in a 12h/12h day/night

cycle at 20 °C for about 2 months to allow reproduce asexually in the laboratory. Then the planarians were maintained in our lab under the same cultural conditions for further studies. The planarians used in this study have been maintained in the lab for more than 1 year. Before exposure, the planarians were fed twice a week with bovine liver homogenate and the water was changed regularly (every two days) to ensure a clean environment. Planarians with lengths of 8 ~10 mm were selected for experiments and starved for at least a week before use.

2.2 Exposure experiment. Both homeostatic and regenerating planarians were used to examine the toxicity of n-CeO₂. Regeneration was induced right before exposure by transversally cutting homeostatic planarians anterior to the pharynx (mouth part) using an ethanol-sterilized razor blade, creating a head-regeneration part. Tissue regeneration, such as new blastema formation was monitored over time, indicated as days post amputation (dpa). Stock of n-CeO₂ (10 mg/mL), μ -CeO₂ (20 mg/mL) and Ce³⁺ (1000 mg/L) were prepared. *D. japonica* was exposed to different treatments for 7 days. MW medium was used as a control treatment. In brief, thirty planarians were exposed to 10 mL of each Ce-based material suspension with 3 replicates for each treatment. A series of n-CeO₂ suspensions (0.01, 0.1, 1, 10, 50, 100, and 200 mg/L), μ -CeO₂ suspensions (1, 10, 50, 100, 200, 500, and 1000 mg/L), and Ce³⁺ ions suspensions (0.01, 0.1, 0.5, 1, 5, 10, and 50 mg/L) were prepared in MW medium (the truth concentration was also detected by ICP-MS, **Table. S1**). The suspensions were sonicated in the MW medium for 15 mins before exposure. All experiments were performed in a 12h/12h day/night rhythm at 20 °C. No food was given during the experiment. For homeostatic and regeneration planarians, the number of live planarians was recorded every day. Besides, based on acute toxicity tests, 40 mg/L n-CeO₂, 40 mg/L μ -CeO₂, and 3 mg/L Ce³⁺ were chosen for the follow-up studies as these concentrations were close to LC₅₀ based on our preliminary experiment (**Fig. S2**).

2.3 Phenotype assay For the regenerating planarians, the growth of the regeneration bud (blastema) was evaluated by measuring the size using a computer morphometry as described previously⁴¹. The planarians were photographed with a Nikon SMZ 1500 stereomicroscope (Nikon, Japan) at 7 dpa. The area of the blastema and the total area of the body were determined using Image J software (Version 1.8.0, Wayne Rasband, National Institutes of Health, Bethesda, MD, USA). The quantitative measure of blastema growth was calculated as the values of new blastema size divided by the sizes of the whole body were used as. Data from 30 animals were averaged. All experiments at each time

point were repeated in triplicate. Each planarian was photographed at the same exposition and magnification. Details of the experiments are shown in the SI.

2.3 Post-exposure locomotion velocity (*pLMV*) assay. At 7 dpa, thirty planarians per treatment were individually placed in the center of a clear acrylic box (50.0 cm × 50.0 cm) positioned over graph paper (grid lines spaced 0.5 cm apart) with mineral water covering the bottom of the paper. The observation started 30 s after placement of the planarians at the center of the grid and lasted for 8 min. *pLMV* was measured as the number of grid lines that each planarian crossed or re-crossed per min over an 8 min observation period.

2.4 Whole-mount *in situ* hybridization (WISH). After 2 days of exposure to Ce-based suspensions, the regenerated planarians (2 dpa) were used for *in situ* hybridizations. The experiments were performed to characterize the expression level of *Djpiwi-A*, which is the main stem cell marker gene⁴² following a previously described method⁴³. In brief, *Djpiwi-A* probe primers (F: 5'-AAGAGAGATAGGAAGACTGCG-3' and R: 5'-GATCACTAATACGACTCACTATAGGGAAGAGAGATAGGAAGACTGCG-3') were designed, synthesized and labeled with digoxigenin (DIG) using an *in vitro* labeling kit (Roche, Switzerland). Planarians were then analyzed using a stereomicroscope (Nikon, SMZ1500, Japan). Each planarian was photographed at the same exposition and magnification. Details of the experiments are shown in the SI.

2.5 Whole-mount immunofluorescence. At 2 dpa, 30 planarians were collected for immunochemistry following a previously described method⁴⁴. Anti-phospho-histone-3 (H₃P, Millipore, Billerica, MA, USA) was used at a 1:500 dilution to investigate the effects on the stem cell proliferation (*in vivo* mitotic activity of stem cells). Phospho-histone-3-positive cells were counted under an Olympus DP80 inverted microscope (Olympus, Tokyo, Japan). To analyze differences in the regeneration of the center nervous system (CNS), immunostaining was carried out using anti-SYNORF1 (3C11, mouse monoclonal antibody specific for synapsin, 1:500 dilution) to investigate the effects on the neurodevelopmental process, and nerve fiber density of SYNORF1-immunopositive neurons were measured at 10 dpa. The detailed method is shown in the SI.

2.6 ICP-MS and XANES measurement. To quantify the concentration of Ce in the regenerating planarians, the samples were collected at 2 dpa after a thorough wash using DI water. For ICP-MS

analysis, these samples were freeze-dried, ground into fine powders and digested with a 3:1 (v:v) mixture of HNO₃ (75%) and H₂O₂ (25%) on a heating plate (80 °C for 1 h, 120 °C for 3 h, and 160 °C for 0.5 h). Ce concentrations in the digestion solution were then analyzed by inductively coupled plasma-mass spectrometry (ICP-MS, 7900, Agilent Technologies). Ce standard solutions (0.5-50 mg/L) were used for calibration. The recovery rate of Ce was tested to be 99.75%.

XANES was used to quantify the Ce³⁺ fraction in the regenerating planarians. At 2 dpa, the planarians were collected, freeze-dried, motor-homogenized, and pressed into circular slices with a diameter of 10 mm and a thickness of 1 mm. Fluorescence mode was applied for the collection of Ce *L_{III}*-edge XANES spectra of the samples and reference standards. Athena software was used to perform linear combination fitting (LCF analysis) of XANES spectra⁴⁵.

2.7 Oxidative Stress responses. At 2 dpa, 30 planarians were collected and homogenized in PBS on ice using a tissue grinder (Tiangen, OSE-Y20, Beijing, China) with a pestle (Tiangen, OSE-Y001, Beijing, China). The homogenates were centrifuged at 12,000 g at 4 °C for 10 min. The supernatant was collected for determination of activities of superoxide dismutase (SOD), glutathione peroxidase (GPx), and glutathione S-transferase (GST) and contents of malondialdehyde (MDA), glutathione (GSH), and glutathione disulfide (GSSG) following the manufacturer's instructions. Details of the analyses are provided in the **SI**.

At 2 dpa, 10 planarians were collected for real-time quantitative PCR analysis (qPCR). Total RNA was isolated using TRIzol (Invitrogen, USA). cDNAs were generated from total RNA with the reverse transcription system (Thermo, USA). The expression of oxidative stress-associated genes (*gpx*, *gst*, *p53* and *nak*) in planarians were quantified (LightCycler 480 II Real-Time PCR System (Roche, Basel, Switzerland)). The primer sequences are provided in **Table S3**. The detailed method is shown in the **SI**.

2.8 ELISA for Caspase-3 and 8-OHdG activities. At 2 dpa, 5-6 planarians were collected for ELISA. Cleaved cysteine-aspartic proteases-3 (caspase-3), an apoptotic marker, was quantified in planarian's tissue using a commercially available ELISA assay (Jiangsu Jingmei Biotechnology Co., Ltd, Jiangsu, China). The content of 8-hydroxy-2 deoxyguanosine (8-OHdG) was measured to assess the oxidative damage to DNA using a DNA Damage ELISA kit (Jiangsu Jingmei Biotechnology Co., Ltd, Jiangsu, China). Details of sample preparation and the analyses are shown in the **SI**.

2.9 Statistical analysis. All tests were repeated at least three times. Data were expressed as mean ±

SD (standard deviation). Statistical significance was analyzed by ANOVA or student's t-test. $p < 0.05$ was considered as the level of significance.

3. Results and discussion

3.1 Effect of n-CeO₂ exposure on tissue regeneration and mobility of regenerating planarian

A pilot experiment was carried out to identify the exposure concentration (**Fig. 1a**). As shown in **Fig. S2** and **Table. S4-S5**, the LC₅₀ values after 7-day n-CeO₂ exposure for homeostatic and regeneration planarians were 51.2 mg/L and 46.3 mg/L, respectively. The LC₅₀ of Ce³⁺ for the homeostatic and regenerating groups were 3.5 mg/L and 3.2 mg/L, respectively (**Table. S4-S5**). Therefore, we chose 40 mg/L n-CeO₂, 40 mg/L μ -CeO₂, and 3 mg/L Ce³⁺ which were close to the LC₅₀ values for the following studies.

After an injury, the stem cells (neoblasts cells) in the planarian would proliferate and migrate to the wound site to form a protective mass of new cells which called blastema⁴⁶. We examined the influences on the tissue regeneration by measuring the size of newly growing blastema after decapitation. As shown in **Fig. 1b** and **1c**, the average anterior new blastema sizes in the n-CeO₂ and Ce³⁺-exposed planarians were significantly smaller than CT and μ -CeO₂ groups. In the CT and μ -CeO₂ groups, 100% planarians developed eyes at the 7 dpa, whereas only 12.5% and 6.7% of planarians reformed eyes under n-CeO₂ and Ce³⁺ ions exposure, respectively (**Fig. 1d**). These results indicated that the n-CeO₂ and Ce³⁺ treatment delayed planarian's regeneration. Locomotion velocity test result showed that n-CeO₂ and Ce³⁺ reduced the mobility of the regenerating planarians (**Fig. 1e**). The data were plotted as the cumulative means of each group over an 8-min observation period at 7 dpa. The slope of the plot represents the speed of movement across the gridlines. The planarians in the CT group displayed an average *pLMV* of about 18.3-24.3 gridlines per minute (the slope value is 20.1), similar to that of the μ -CeO₂ group (the slope value is 17.8). n-CeO₂ and Ce³⁺ ions reduced the slope of the *pLMV* to 8.9 and 3.8, respectively.




Fig. 1 n-CeO₂ and Ce³⁺ delay planarian regeneration and affect the mobility. **a)** Experimental time points of Ce-based suspensions (40 mg/L n-CeO₂, 40 mg/L μ-CeO₂, and 3 mg/L Ce³⁺) exposure, amputation, and morphological observation. **b)** Blastema reform at 7 dpa in different treatments (**Black Arrow**: eyespots; **Red Arrow**: blastema). **c)** Average size of regenerated blastema in proportion to the body area of the planarian. **d)** Percentage of planarians with eyespots regeneration in different treatments. **e)** Effect of Ce-based suspensions on the regenerating planarians' motility at 7 dpa. PLMV was quantified as the number of gridlines crossed or re-crossed over an 8-min. This was plotted as cumulative crosses vs. time. Thirty animals per experimental condition were used, and each experiment was repeated three times. The results are presented as the mean ± SD, one-way ANOVA.

3.2 n-CeO₂ impaired stem cell dynamics and regeneration of the nervous system.

The stem cell proliferation promotes the tissue regeneration. Planarian has large number of pluripotent stem cells for rapid cell division and regeneration. The decreased size of anterior new blastema may arise from the disrupted cell division and regeneration. Stem cells are highly vulnerable to environmental chemicals, which can lead to adverse effects on differentiation processes and development⁴⁷. The detailed experimental design was described in **Fig. 2a**. *DjpiwiA*, a main stem cell gene marker⁴⁴, was quantified using the WISH method. As shown in **Fig. 2b**, at 2 dpa, *DjpiwiA* were extensively expressed (shown as the blue color) in the whole body except for the pharynx in CT and μ-CeO₂ group, and the expression were reduced by 29.2% and 38.4% in n-CeO₂ and Ce³⁺ group, respectively.

During development, mitotic activity (cell proliferation) can be regarded as one of the primary

events of tissue remodeling and repair⁴⁸. Anti-phospho-Histone H3 antibody can be used to label the nuclei of stem cells to examine the *in vivo* mitotic activity⁴⁹. Compared to CT group, the number of labeled nuclei (green dots) significantly decreased by 63.1% and 65.2% after exposure to n-CeO₂ and Ce³⁺, respectively (**Fig. 2c**), while μ -CeO₂ group did not show significant difference with CT. This suggested that n-CeO₂ and Ce³⁺ treatment impaired the proliferation of stem cell.

A recent study found that Ag NPs can affect the locomotion of freshwater planarians *Schmidtea mediterranea* by inducing neurotoxicity and disorder in neurodevelopment⁵. These suggested that n-CeO₂ induced locomotive inhibition may be attributed to the neurotoxicity. Planarian is a unique in the animal kingdom as they can regenerate its brain, including a fully functional CNS⁵⁰. To evaluate whether the reduced stem cell proliferation further impeded the regeneration of the CNS, the neuron morphologies were observed at 10 dpa using immunohistochemistry method. The anti-SYNORF1 antibody was used as a biomarker, which allows visualization of the planarian CNS. The normal planarian brain has an inverted U-shaped structure comprised of a cortex of neural cells and a core of axons, which displays rich branches⁵¹. Based on the behavioral changes and regeneration defects, we observed that planarian could not regenerate its complete CNS. As shown in **Fig. 2d**, n-CeO₂ and Ce³⁺ treatment significantly reduced the number of lateral branches, indicating that n-CeO₂ and Ce³⁺ induced neural regeneration defects in the treated planarians. μ -CeO₂ groups showed no significant effect. Overall, these results suggest that n-CeO₂ and Ce³⁺ exposure impaired the stem cell activity and neurodevelopment, and thus the health and regeneration capacity of the CNS in the exposed planarians.




Fig. 2 n-CeO₂ and Ce³⁺ affects stem cell proliferation. **(a)** Experimental time points of Ce-based suspensions (40 mg/L n-CeO₂, 40 mg/L μ-CeO₂, and 3 mg/L Ce³⁺ ions) exposure to regeneration planarians and detect biomarkers, such as oxidative stress marker MDA and SOD, WISH and Immunochemistry. **(b) Left**, *in situ* hybridization to detect *DjpiwiA* transcripts in planarians. **Right**, the gene expression of *DjpiwiA* in the regenerating planarians in different treatment. **(c)** Immunohistochemistry using the mitotic marker Anti-H3P antibody. The **left** panel shows a view of the mitotic cells in regenerating planarian in different treatments. **Right**, the number of mitotic events in Ce-based suspensions treatment and control (CT) groups. The results are presented as the mean ± SD of a minimum of 10 biological replicates per condition (1 planarian per point, asterisks indicate the level of significant differences between treatment and CT groups of the planarian, one-way ANOVA, **p* < 0.05 ***p* < 0.01). **(d)** Whole-mount immunostaining of the nervous system with anti-SYNORF1 (3C11) in regeneration planarians after exposure to Ce-based suspensions for 10 days. Scale bars represent 2 mm or 100 μm.

3.3 Chemical origin of the toxicity: role of bioaccumulation and biotransformation

Biotransformation and bioaccumulation determine the ultimate fate and toxicity of nanomaterials in living organisms^{52, 53}, such as *Lactuca plants*³⁹, *Bacillus subtilis*²⁴, and *Chlorella pyrenoidosa*¹⁴. *In vivo* studies rarely consider physicochemical characteristics and actual exposure concentrations when linking them to uptake, distribution patterns, and potential adverse effects. In the present study, the actual size of the n-CeO₂ and μ-CeO₂ were considerably larger than their nominal reported size (**Fig. S3**). In fact, n-CeO₂ and μ-CeO₂ changes in both concentration and particle size were observed during the exposure concentration (**Fig. S4**, the hydrodynamic diameter from 162 nm and 2863 nm at the beginning up to 312 nm and 3312 nm at 48 h by n-CeO₂ and μ-CeO₂ interacted with planarian, respectively; **Table S6**). Sedimentation also depends on the surrounding environment⁵, such as the production of organic material mucus, in which particles agglomerate (**Fig.S4-S5**). This way, the bioavailability in the aquatic column is reduced, while increasing the risk for benthic animals.

Planarians inhabit the lowest water layer, and are able to take up NPs as they glide over the substrate⁵. Then we used ICP-MS to detect the Ce-based materials and found that Ce was significantly accumulated in the regeneration planarian after 2 days of exposure to different Ce treatments, and n-CeO₂ group showed the highest accumulation, followed by μ -CeO₂ and Ce³⁺ group (**Fig. 2b**). For the 40 mg/L μ -CeO₂, n-CeO₂, and 3 mg/L Ce³⁺ ions, Ce accumulations in the planarians were 27.6 ± 4.7 , 46.5 ± 7.8 , and 2.8 ± 0.4 mg Ce kg⁻¹, respectively. The percentages of the accumulation from the total applied Ce are 0.31%, 0.52% and 0.32% for μ -CeO₂, n-CeO₂ and Ce³⁺, respectively. Many factors would influence the bioaccumulation and bioavailable of Ce in planarian, such as sizes and culture environment. On the one hand, n-CeO₂ with smaller size is easier to move into the organism thus accumulate more than μ -CeO₂. For example, neither μ -CeO₂ nor n-CeO₂ are ingested by planarian, however, the latter with a smaller size allows them to stay a longer time in body during this ingestion process. In addition, planarians use their motile cilia to glide across the sediment surface and come into contact with the deposited materials, which make them adsorb μ -CeO₂ and n-CeO₂ in their body. Besides, planarian will secrete mucus after being stimulated, as shown in **Fig. S5**, large numbers of mucus proteins were present in the n-CeO₂ treatment group while a small amount these proteins were present in the μ -CeO₂ group, which may promote n-CeO₂ adsorption by planarian. Once n-CeO₂ penetrated into glial and neurons cells, it would be directed to the lysosomes or persist in the cytoplasm and subsequently interact with other organelles⁵⁴. Although μ -CeO₂ also accumulated significant amount of Ce (equal to 60% of that in n-CeO₂ group), it did not show significant effects on locomotion, regeneration and neurodevelopment of planarian. Therefore, both the higher accumulation of Ce and higher reactivity of n-CeO₂ may contribute to its higher toxicity than μ -CeO₂. However, n-CeO₂ exposure resulted in 16.6-fold higher Ce accumulation but led to less toxicity than Ce³⁺, further demonstrating that Ce³⁺ is more toxic than n-CeO₂. It is well known that nanomaterials in environment and biota may undergo transformation either through chemical or biological processes, and the transformation process may determine their toxicity^{24, 55}. We therefore hypothesize that the biotransformation of n-CeO₂ might contribute to the significant toxicity.

We firstly examined the release of Ce³⁺ from μ -CeO₂ and n-CeO₂ in the culture media. Only less than 26.5 μ g/L of Ce were detected, suggesting toxicity is unlikely caused by the Ce³⁺ that were initially in the exposure solution. However, biotransformation is a continuous process once the nanomaterials contact the biota⁵⁶. Biotransformation of n-CeO₂ may occur on or within the planarian

during the exposure. Therefore, we examined the real amount of transformation in the organism using synchrotron radiation based XANES technique (**Fig. 3b**, **Table. S7**). The spectrum of Ce in the planarians under treatments of μ -CeO₂ at 40 mg/L shows a small feature of Ce(III) oxidation state ($2.9 \pm 0.2\%$), suggesting a small extent of transformation. However, the feature of Ce(III) under n-CeO₂ (40 mg/L) account for $12.8 \pm 2.3\%$ of the total Ce. This is equivalent to a release of 5.96 mg/kg ($46.5 \times 12.8\%$) Ce³⁺ in planarian, which demonstrated our hypothesis that biotransformation contributed to the toxicity observed for n-CeO₂. It should be also noted that the concentration of Ce(III) is twice of that in Ce³⁺ treated group (2.8 mg/kg), while the toxicity is lower than Ce³⁺. This can be attributed to the fact that the biotransformation of nanomaterials is a time-dependent process. For example, it can take up to 7 days to show 20% of transformation of n-CeO₂ in plant shoot⁵⁷. Depending on different species, the transformation rate can be lower or higher. Unlike Ce³⁺ which instantly release all the Ce³⁺ to organism and thus cause acute toxicity, n-CeO₂ released Ce³⁺ slowly by continuously interact with the planarian thus the toxicity is more chronic rather than acute. As with all chemicals and bulk materials, the biological and/or toxicological effects of NPs will be affected by both their physicochemical properties and the biological environment around the organism. NPs interactions with biological environments result in the formation of a protein corona^{58, 59}. A corona-NPs complex may feature NPs undergoing different changes as a result of their adsorption behavior, agglomeration, accumulation, distribution, biotransformation, and dissolution. A protein corona complex and NPs represent a new entity in every biological environment which will inevitably define NPs toxicology, pharmacokinetics, and dynamics^{60, 61}. In the present study, we found that n-CeO₂ induced a more mucus secreted by planarian and the higher biotransformation occurred with a higher toxicity in the homeostatic and regeneration planarians. In fact, n-CeO₂ would gain a negative charge due to the formation of a protein-corona-NPs complex and cause planarian absorbed a more n-CeO₂, which may promote NPs' biotransformation and dissolution (**Fig. S7**, **Table. S4**). Thus, we hypothesized that the protein corona affects not only n-CeO₂'s distribution, but also its toxicity. In summary, n-CeO₂ induced toxicity to planarian via both its nano-specific effect as well as the ions induced effects.




Fig. 3 a) Cerium contents in the regeneration planarians at 2 dpa in different treatments. Data are expressed as mean \pm SD (n=6) an average of six replicates. Different letters stand for statistical differences between treatments at $p < 0.05$. **b)** XANES normalized Ce LIII-edge spectra of reference compounds (CePO₄ and CeO₂) and samples. μ -CeO₂, n-CeO₂ and Ce³⁺ ions exposure to regeneration planarians for 2 days and the samples were detected by XANES. Ce speciation in Ce³⁺ treatment sample could not be detected, as the Ce signal was too low to obtain a reliable spectrum.

3.4 n-CeO₂ disrupts the antioxidant system and damage DNA integrity


In previous studies, Ce³⁺/Ce⁴⁺ fractions (two reversible states) on n-CeO₂ particle surfaces were found to determine the particle's SOD mimicking activity⁶². Compared with the smaller n-CeO₂, n-CeO₂ with dimensions bigger than 5 nm exhibit a lower Ce³⁺/Ce⁴⁺ ratio, and thus exhibit negligible activity⁶³. Once n-CeO₂ undergoes biotransformation and releases Ce³⁺ ions, which plays a role of pro-oxidative effects, and the toxicity would be exhibited²⁴. In addition, a Fenton-like reaction could be catalyzed by the released Ce³⁺ ions with hydrogen peroxide, producing damaging oxygen radicals, which may cause oxidative stress happened⁶⁴. Oxidative stress has been considered a major contributor to the toxicity of environmental pollutants including nanomaterials⁶⁵⁻⁶⁷. Changes in activities of antioxidant enzymes and contents of MDA are the primary indicators of oxidative stress and cell damage. The antioxidant enzymes such as SOD, GPX, and GST, are involved in the protective mechanisms adapted by animals to scavenge ROS⁶⁸. To examine whether the disorder of redox balance underlies the toxicity of n-CeO₂, we evaluated the expression levels of different categories of antioxidative in regeneration planarians. In this study, the antioxidant defense system including enzymatic (SOD, GST and GPx) and non-enzymatic (GSH, GSSG) and MDA levels were

determined in the regenerating planarians. As shown in **Fig. 4a-c**, both n-CeO₂ and Ce³⁺ treatments induced significant changes in the activities of SOD, GST, and GPx in planarians. n-CeO₂ and Ce³⁺ decreased the activity of SOD by 28.6% and 34.4%, respectively, compared with CT. The activity of GST in planarians in n-CeO₂ and Ce³⁺ treatment was significantly decreased as compared to that of CT. n-CeO₂ and Ce³⁺ significantly decreased the GPx activity by 38.8% and 37.5%, respectively. The results of the present study indicate that both n-CeO₂ and Ce³⁺ treatments disturbed the homeostasis of antioxidant defense system in regeneration planarians. A similar pattern of impacts on the non-enzymatic antioxidant system was also observed (**Fig. 4d** and **4e**). The GSH contents at 2 dpa decreased by 26.8% and 32.1 % after exposure to n-CeO₂ and Ce³⁺, respectively. Correspondingly, a significant increase of the GSSG content, an oxidized form of GSH, was observed in planarians after exposure to n-CeO₂ and Ce³⁺. n-CeO₂ and Ce³⁺ decreased activities of SOD, GST, GPx and GSH in the regeneration planarians compared to control with a higher level of MDA and GSSG which suggests the treatments provoked lipid peroxidation.

We further examined whether the antioxidant system was impaired at the genetic level by examining the mRNA expression of key genes responsible for antioxidant production (*gpx*, *gst*), ATP synthesis (*nak*), and cell apoptosis (*p53*) (**Fig. 5**). Consistent with the above result, *gpx*, *gst*, and *nak* expression were significantly down-regulated after 2 days of exposure to n-CeO₂ and Ce³⁺. μ -CeO₂ did not show significant difference with CT in all cases. *Gpx* and *gr* are key antioxidant defense genes involved in the glutathione metabolism. In planarians, *p53* has a wider range of functions than known tumor suppressor⁶⁹. Pearson and Alvarado⁷⁰ found that *p53* involved in tumor inhibition and regulation of stem cell self-renewal in planarians. In this work, *p53* expression was significantly up-regulated under n-CeO₂ and Ce³⁺ exposure when compared with CT. The apoptosis was affected in regenerating planarians, which could be due to the ability of Ce³⁺ to promote programmed cell death.

Oxidative stress can cause tissue alternations, trigger apoptosis and induce cell death. So we studied whether n-CeO₂ and Ce³⁺ exposure can induce cell apoptosis (**Fig. 4g**) and DNA damage at the molecular levels (**Fig. 4h**). 8-OHdG, a maker for oxidative DNA damage, was increased after n-CeO₂ and Ce³⁺ exposure (**Fig. 4h**), indicating n-CeO₂ and Ce³⁺ exposure can cause genotoxic effects. Similarly, Arslan, et al.⁷¹ found 0.78 mg/L n-CeO₂ induced genotoxic effect *in vitro* using micronucleus and chromosome aberration tests.

407 The overall oxidative responses of planarian to n-CeO₂ and Ce³⁺ are similar and to similar
 408 extents on molecular level (**Fig. 5**), although the phenotypic effects caused by the two are different.
 409 Since the phenotypic changes are essentially originated from the changes at molecular or genetic
 410 levels and given that n-CeO₂ can release toxic Ce³⁺ in a slow and sustained way, these findings give
 411 implications that n-CeO₂ may cause longer-term adverse impacts to planarian, which might be worse
 412 than that caused by acute exposure to Ce³⁺.



413
 414 **Fig. 4** n-CeO₂ regulates planarian's antioxidant defense system and cell damage. SOD (**a**), GST (**b**), GPx (**c**), GSH
 415 (**d**), GSSG (**e**), and MDA levels (**f**) in regeneration planarians after exposure to Ce-based suspensions for 48 h.
 416 Contents of caspase-3 (**g**) and 8-OHdG (**h**) in planarians affected by Ce-based suspensions treatments. Different
 417 letters stand for statistical differences between treatments at $p < 0.05$.




Fig. 5 The effect of Ce-based suspensions on relative mRNA expression levels of gene *gpx* (a), *gst* (b), *nak* (c), and *p53* (d) measured in regeneration planarians. Different letters stand for statistical differences between treatments at $p < 0.05$.

4. Conclusion

In this study we compared the toxicity effects of n-CeO₂, μ -CeO₂, and ionic Ce (CeCl₃) in planarian *D. japonica*. We not only showed that n-CeO₂ induced nano-specific toxicity to planarian compared with μ -CeO₂, but also demonstrated that the biotransformation of n-CeO₂ contributed significantly to the observed toxicity. Interestingly, although n-CeO₂ induced more Ce accumulation in the planarians than Ce³⁺, the toxicity is actually lower which is attributed to that the release of Ce³⁺ from n-CeO₂ is a slow process. However, this in turn implicates that n-CeO₂ may cause long-term adverse effects to planarian, especially to the regenerating ones which are more sensitive to the n-CeO₂ toxicity. This knowledge gap should be addressed in the future. On the other hand, interests in exploring the beneficial effects of n-CeO₂ to enhance regeneration of neural systems have increasingly growing due to the ROS capturing activity of n-CeO₂. However, such effects are only reported at very low concentrations and particles with extremely small size (usually less than 10 nm). Our results suggest that a threshold between an inhibitor and an enhancer should be established, and caution should be made for this type of application.

CRedit authorship contribution statement

Changjian Xie: Conceptualization, Methodology, Investigation, Formal analysis, Writing-original draft, Funding acquisition. **Xiaowei Li:** Methodology, Resources. **Lisha Hei:** Resources, review & editing. **Yiqing Chen:** Resources, review & editing. **Yuling Dong:** Investigation, Funding acquisition. **Shujing Zhang:** Resources. **Shan Ma:** Investigation. **Jianing Xu:** Formal analysis, Funding acquisition. **Qiuxiang Pang:** Conceptualization, Methodology, Formal analysis, Writing-review & editing, Resources. **Iseult Lynch:** Funding acquisition, Writing-review & editing. **Zhiling Guo:** Conceptualization, Methodology, Formal analysis, Writing-review & editing, Resources. **Peng Zhang:** Writing-review & editing, Resources.

Declaration of competing interest

The authors declare that they have no competing interests.

Acknowledgments

This work was supported by the National Natural Science Foundation of China (Grant Nos. 12105163, 32001987) and the Natural Science Foundation of Shandong Province (Grant Nos. ZR2020QD133, ZR2020QC241, ZR2020MC142). Additional support from H2020 project NanoCommons (Grant Agreement No. 731032) is acknowledged.

References

1. Keller, A. A.; McFerran, S.; Lazareva, A.; Suh, S. J. J. o. n. r., Global life cycle releases of engineered nanomaterials. **2013**, *15*, (6), 1-17.
2. Dahle, J. T.; Arai, Y. J. I. j. o. e. r.; health, p., Environmental geochemistry of cerium: applications and toxicology of cerium oxide nanoparticles. **2015**, *12*, (2), 1253-1278.
3. Das, S.; Dowding, J. M.; Klump, K. E.; McGinnis, J. F.; Self, W.; Seal, S. J. N., Cerium oxide nanoparticles: applications and prospects in nanomedicine. **2013**, *8*, (9), 1483-1508.
4. Hoecke, K. V.; Quik, J. T.; Mankiewicz-Boczek, J.; Schamphelaere, K. A. D.; Elsaesser, A.; Meeren, P. V. d.; Barnes, C.; McKerr, G.; Howard, C. V.; Meent, D. V. D. J. E. s.; technology, Fate and effects of CeO₂ nanoparticles in aquatic ecotoxicity tests. **2009**, *43*, (12), 4537-4546.
5. Leynen, N.; Van Belleghem, F. G.; Wouters, A.; Bove, H.; Ploem, J.-P.; Thijssen, E.; Langie, S. A.; Carleer, R.; Ameloot, M.; Artois, T. J. N., In vivo toxicity assessment of silver nanoparticles in homeostatic versus regenerating planarians. **2019**, *13*, (4), 476-491.
6. Chen, J.; Patil, S.; Seal, S.; McGinnis, J. F. J. N. n., Rare earth nanoparticles prevent retinal degeneration induced by intracellular peroxides. **2006**, *1*, (2), 142-150.
7. Li, Y.; He, X.; Yin, J. J.; Ma, Y.; Zhang, P.; Li, J.; Ding, Y.; Zhang, J.; Zhao, Y.; Chai, Z. J. A. C., Acquired superoxide - scavenging ability of ceria nanoparticles. **2015**, *127*, (6), 1852-1855.

- 469 8. Lee, S. S.; Song, W.; Cho, M.; Puppala, H. L.; Nguyen, P.; Zhu, H.; Segatori, L.; Colvin, V. L. J. A. n., Antioxidant
470 properties of cerium oxide nanocrystals as a function of nanocrystal diameter and surface coating. **2013**, 7, (11),
471 9693-9703.
- 472 9. Pirmohamed, T.; Dowding, J. M.; Singh, S.; Wasserman, B.; Heckert, E.; Karakoti, A. S.; King, J. E.; Seal, S.; Self, W. T. J.
473 C. c., Nanoceria exhibit redox state-dependent catalase mimetic activity. **2010**, 46, (16), 2736-2738.
- 474 10. Popov, A. L.; Popova, N. R.; Selezneva, I. I.; Akkizov, A. Y.; Ivanov, V. K. J. M. S.; C, E., Cerium oxide nanoparticles
475 stimulate proliferation of primary mouse embryonic fibroblasts in vitro. **2016**, 68, 406-413.
- 476 11. Popov, A.; Ermakov, A.; Savintseva, I.; Selezneva, I.; Poltavtseva, R.; Zaisky, E.; Poltavtsev, A.; Stepanov, A.; Ivanov,
477 V.; Sukhikh, G., Citrate-stabilized nanoparticles of ceo 2 stimulate proliferation of human mesenchymal stem cells in
478 vitro. *Nanoscience and Technology: An International Journal* **2016**, 7, (3).
- 479 12. Singh, R.; Karakoti, A. S.; Self, W.; Seal, S.; Singh, S. J. L., Redox-sensitive cerium oxide nanoparticles protect human
480 keratinocytes from oxidative stress induced by glutathione depletion. **2016**, 32, (46), 12202-12211.
- 481 13. Zhang, H.; He, X.; Zhang, Z.; Zhang, P.; Li, Y.; Ma, Y.; Kuang, Y.; Zhao, Y.; Chai, Z. J. E. s.; technology, Nano-CeO₂
482 exhibits adverse effects at environmental relevant concentrations. **2011**, 45, (8), 3725-3730.
- 483 14. Xie, C.; Ma, Y.; Zhang, P.; Zhang, J.; Li, X.; Zheng, K.; Li, A.; Wu, W.; Pang, Q.; He, X., Elucidating the origin of the
484 toxicity of nano-CeO₂ to *Chlorella pyrenoidosa*: the role of specific surface area and chemical composition.
485 *Environmental Science: Nano* **2021**, 8, 1701-1712.
- 486 15. Rosbach, L. M.; Brede, D. A.; Nuyts, G.; Cagno, S.; Olsson, R. M. S.; Oughton, D. H.; Falkenberg, G.; Janssens, K.;
487 Lind, O. C., Synchrotron XRF Analysis Identifies Cerium Accumulation Colocalized with Pharyngeal Deformities in CeO₂
488 NP-Exposed *Caenorhabditis elegans*. *Environmental Science & Technology* **2022**, 56, (8), 5081-5089.
- 489 16. Kuang, Y.; He, X.; Zhang, Z.; Li, Y.; Zhang, H.; Ma, Y.; Wu, Z.; Chai, Z. J. J. o. n.; nanotechnology, Comparison study on
490 the antibacterial activity of nano-or bulk-cerium oxide. **2011**, 11, (5), 4103-4108.
- 491 17. Manier, N.; Bado-Nilles, A.; Delalain, P.; Aguerre-Chariol, O.; Pandard, P. J. E. p., Ecotoxicity of non-aged and aged
492 CeO₂ nanomaterials towards freshwater microalgae. **2013**, 180, 63-70.
- 493 18. Sendra, M.; Yeste, P.; Moreno-Garrido, I.; Gatica, J. M.; Blasco, J. J. S. o. T. T. E., CeO₂ NPs, toxic or protective to
494 phytoplankton? Charge of nanoparticles and cell wall as factors which cause changes in cell complexity. **2017**, 590,
495 304-315.
- 496 19. Alabresm, A.; Decho, A. W.; Lead, J. J. N., A novel method to estimate cellular internalization of nanoparticles into
497 gram-negative bacteria: Non-lytic removal of outer membrane and cell wall. **2021**, 21, 100283.
- 498 20. Bour, A.; Mouchet, F.; Verneuil, L.; Evariste, L.; Silvestre, J.; Pinelli, E.; Gauthier, L. J. C., Toxicity of CeO₂
499 nanoparticles at different trophic levels—effects on diatoms, chironomids and amphibians. **2015**, 120, 230-236.
- 500 21. Yang, X.; Pan, H.; Wang, P.; Zhao, F.-J. J. J. o. H. M., Particle-specific toxicity and bioavailability of cerium oxide
501 (CeO₂) nanoparticles to *Arabidopsis thaliana*. **2017**, 322, 292-300.
- 502 22. Bandyopadhyay, S.; Peralta-Videa, J. R.; Plascencia-Villa, G.; José-Yacamán, M.; Gardea-Torresdey, J. L. J. J. o. h. m.,
503 Comparative toxicity assessment of CeO₂ and ZnO nanoparticles towards *Sinorhizobium meliloti*, a symbiotic alfalfa
504 associated bacterium: use of advanced microscopic and spectroscopic techniques. **2012**, 241, 379-386.
- 505 23. Zhang, P.; Ma, Y.; Zhang, Z.; He, X.; Zhang, J.; Guo, Z.; Tai, R.; Zhao, Y.; Chai, Z., Biotransformation of ceria
506 nanoparticles in cucumber plants. *ACS nano* **2012**, 6, (11), 9943-9950.
- 507 24. Xie, C.; Zhang, J.; Ma, Y.; Ding, Y.; Zhang, P.; Zheng, L.; Chai, Z.; Zhao, Y.; Zhang, Z.; He, X., *Bacillus subtilis* causes
508 dissolution of ceria nanoparticles at the nano–bio interface. *Environmental Science: Nano* **2019**, 6, (1), 216-223.
- 509 25. Thill, A.; Zeyons, O.; Spalla, O.; Chauvat, F.; Rose, J.; Auffan, M.; Flank, A. M. J. E. s.; technology, Cytotoxicity of
510 CeO₂ nanoparticles for *Escherichia coli*. Physico-chemical insight of the cytotoxicity mechanism. **2006**, 40, (19),
511 6151-6156.
- 512 26. Leynen, N.; Van Belleghem, F. G.; Wouters, A.; Bove, H.; Ploem, J.-P.; Thijssen, E.; Langie, S. A.; Carleer, R.; Ameloot,

513 M.; Artois, T., In vivo toxicity assessment of silver nanoparticles in homeostatic versus regenerating planarians.
514 *Nanotoxicology* **2019**, *13*, (4), 476-491.

515 27. Kluever, A. N.; Aungst, J.; Gu, Y.; Hatwell, K.; Shackelford, M., Infant toxicology: State of the science and
516 considerations in evaluation of safety. *Food. Chem. Toxicol* **2014**, *70*, (2), 68-83.

517 28. Perera, F.; Herbstman, J., Prenatal environmental exposures, epigenetics, and disease. *Reprod. Toxicol* **2011**, *31*, (3),
518 363-373.

519 29. Zeng, A.; Li, H.; Guo, L.; Gao, X.; McKinney, S.; Wang, Y.; Yu, Z.; Park, J.; Semerad, C.; Ross, E. J. C., Prospectively
520 isolated tetraspanin+ neoblasts are adult pluripotent stem cells underlying planaria regeneration. **2018**, *173*, (7),
521 1593-1608. e20.

522 30. Dong, Z.; Huo, J.; Liang, A.; Chen, J.; Chen, G.; Liu, D., Gamma-Secretase Inhibitor (DAPT), a potential therapeutic
523 target drug, caused neurotoxicity in planarian regeneration by inhibiting Notch signaling pathway. *Science of The Total*
524 *Environment* **2021**, *781*, 146735.

525 31. Morris, J.; Bealer, E. J.; Souza, I. D.; Repmann, L.; Bonelli, H.; Stanzione III, J. F.; Staehle, M. M. J. T. S., Chemical
526 Exposure-Induced Developmental Neurotoxicity in Head-Regenerating Schmidtea mediterranea. **2022**, *185*, (2),
527 220-231.

528 32. Zhang; Jianyong; Bin; Wang; Bosheng; Zhao; Yanqing; Li; Xiuyun; Pollution, Z. J. E., Blueberry anthocyanin alleviate
529 perfluorooctanoic acid-induced toxicity in planarian (*Dugesia japonica*) by regulating oxidative stress biomarkers, ATP
530 contents, DNA methylation and mRNA expression. **2019**.

531 33. Rompolas, P.; Patel-King, R. S.; King, S. M., *Schmidtea mediterranea*: a model system for analysis of motile cilia. In
532 *Methods in cell biology*, Elsevier: 2009; Vol. 93, pp 81-98.

533 34. Gao, T.; Sun, B.; Xu, Z.; Chen, Q.; Yang, M.; Wan, Q.; Song, L.; Chen, G.; Jing, C.; Zeng, E. Y. J. J. o. H. M., Exposure to
534 polystyrene microplastics reduces regeneration and growth in planarians. **2022**, *432*, 128673.

535 35. Gambino, G.; Falleni, A.; Nigro, M.; Salvetti, A.; Cecchetti, A.; Ippolito, C.; Guidi, P.; Rossi, L. J. A. T., Dynamics of
536 interaction and effects of microplastics on planarian tissue regeneration and cellular homeostasis. **2020**, *218*, 105354.

537 36. Tran, T. A.; Hesler, M.; Moriones, O. H.; Jimeno-Romero, A.; Fischer, B.; Bastús, N. G.; Puentes, V.; Wagner, S.; Kohl, Y.
538 L.; Gentile, L. J. A.-A. t. a. e., Assessment of iron oxide nanoparticle ecotoxicity on regeneration and homeostasis in the
539 replacement model system *Schmidtea mediterranea*. **2019**, *36*, (4), 583-596.

540 37. Kustov, L.; Tiras, K.; Al-Abed, S.; Golovina, N.; Ananyan, M. J. A. t. L. A., Estimation of the toxicity of silver
541 nanoparticles by using planarian flatworms. **2014**, *42*, (1), 51-58.

542 38. Ermakov, A.; Popov, A.; Ermakova, O.; Ivanova, O.; Baranchikov, A.; Kamenskikh, K.; Shekunova, T.; Shcherbakov, A.;
543 Popova, N.; Ivanov, V., The first inorganic mitogens: Cerium oxide and cerium fluoride nanoparticles stimulate planarian
544 regeneration via neoblastic activation. *Mater. Sci. Eng. C* **2019**, *104*, 109924.

545 39. Zhang, P.; Ma, Y.; Zhang, Z.; He, X.; Li, Y.; Zhang, J.; Zheng, L.; Zhao, Y. J. N., Species-specific toxicity of ceria
546 nanoparticles to *Lactuca plants*. **2015**, *9*, (1), 1-8.

547 40. Rui, Y.; Zhang, P.; Zhang, Y.; Ma, Y.; He, X.; Gui, X.; Li, Y.; Zhang, J.; Zheng, L.; Chu, S. J. E. P., Transformation of ceria
548 nanoparticles in cucumber plants is influenced by phosphate. **2015**, *198*, 8-14.

549 41. Ermakov, A.; Popov, A.; Ermakova, O.; Ivanova, O.; Baranchikov, A.; Kamenskikh, K.; Shekunova, T.; Shcherbakov, A.;
550 Popova, N.; Ivanov, V., The first inorganic mitogens: Cerium oxide and cerium fluoride nanoparticles stimulate planarian
551 regeneration via neoblastic activation. *Materials Science and Engineering: C* **2019**, *104*, 109924.

552 42. Shibata, N.; Rouhana, L.; Agata, K. J. D., growth; differentiation, Cellular and molecular dissection of pluripotent
553 adult somatic stem cells in planarians. **2010**, *52*, (1), 27-41.

554 43. Gao, L.; Li, A.; Lv, Y.; Huang, M.; Liu, X.; Deng, H.; Liu, D.; Zhao, B.; Liu, B.; Pang, Q., Planarian
555 gamma-interferon-inducible lysosomal thiol reductase (GILT) is required for gram-negative bacterial clearance.
556 *Developmental & Comparative Immunology* **2021**, *116*, 103914.

557 44. Dong, Z.; Cheng, F.; Yang, Y.; Zhang, F.; Chen, G.; Liu, D., Expression and functional analysis of flotillins in *Dugesia*
558 *japonica*. *Experimental cell research* **2019**, *374*, (1), 76-84.

559 45. Collin, B.; Oostveen, E.; Tsyusko, O. V.; Unrine, J. M. J. E. s.; technology, Influence of natural organic matter and
560 surface charge on the toxicity and bioaccumulation of functionalized ceria nanoparticles in *Caenorhabditis elegans*.
561 **2014**, *48*, (2), 1280-1289.

562 46. Lobo, D.; Beane, W. S.; Levin, M. J. P. c. b., Modeling planarian regeneration: a primer for reverse-engineering the
563 worm. **2012**, *8*, (4), e1002481.

564 47. Perera, F.; Herbstman, J. J. R. t., Prenatal environmental exposures, epigenetics, and disease. **2011**, *31*, (3),
565 363-373.

566 48. Matson, J. P.; Cook, J. G. J. T. F. j., Cell cycle proliferation decisions: the impact of single cell analyses. **2017**, *284*, (3),
567 362-375.

568 49. González-Estévez, C.; Felix, D. A.; Aboobaker, A. A.; Saló, E. J. P. o. t. N. A. o. S., Gtdap-1 promotes autophagy and is
569 required for planarian remodeling during regeneration and starvation. **2007**, *104*, (33), 13373-13378.

570 50. Brown, D. D.; Pearson, B. J. J. F. i. n., A brain unfixed: unlimited neurogenesis and regeneration of the adult
571 planarian nervous system. **2017**, *11*, 289.

572 51. Umesono, Y.; Watanabe, K.; Agata, K., Distinct structural domains in the planarian brain defined by the expression
573 of evolutionarily conserved homeobox genes. *Development genes and evolution* **1999**, *209*, (1), 31-39.

574 52. Cui, D.; Zhang, P.; Ma, Y.; He, X.; Li, Y.; Zhang, J.; Zhao, Y.; Zhang, Z. J. E. S. N., Effect of cerium oxide nanoparticles on
575 asparagus lettuce cultured in an agar medium. **2014**, *1*, (5), 459-465.

576 53. Ding, X.; Pu, Y.; Tang, M.; Zhang, T. J. N. T., Environmental and health effects of graphene-family nanomaterials:
577 potential release pathways, transformation, environmental fate and health risks. **2022**, *42*, 101379.

578 54. Ding, L.; Liu, Z.; Okweesi Aggrey, M.; Li, C.; Chen, J.; Tong, L., Nanotoxicity: the toxicity research progress of metal
579 and metal-containing nanoparticles. *Mini reviews in medicinal chemistry* **2015**, *15*, (7), 529-542.

580 55. Guo, Z.; Zhang, P.; Chakraborty, S.; Chetwynd, A. J.; Abdolahpur Monikh, F.; Stark, C.; Ali-Boucetta, H.; Wilson, S.;
581 Lynch, I.; Valsami-Jones, E., Biotransformation modulates the penetration of metallic nanomaterials across an artificial
582 blood–brain barrier model. *Proceedings of the National Academy of Sciences* **2021**, *118*, (28), e2105245118.

583 56. Zhang, P.; Guo, Z.; Zhang, Z.; Fu, H.; White, J. C.; Lynch, I., Nanomaterial transformation in the soil–plant system:
584 implications for food safety and application in agriculture. *Small* **2020**, *16*, (21), 2000705.

585 57. Ma, Y.; Zhang, P.; Zhang, Z.; He, X.; Zhang, J.; Ding, Y.; Zhang, J.; Zheng, L.; Guo, Z.; Zhang, L. J. E. s.; technology,
586 Where does the transformation of precipitated ceria nanoparticles in hydroponic plants take place? **2015**, *49*, (17),
587 10667-10674.

588 58. Docter, D.; Westmeier, D.; Markiewicz, M.; Stolte, S.; Knauer, S.; Stauber, R. J. C. S. R., The nanoparticle biomolecule
589 corona: lessons learned—challenge accepted? **2015**, *44*, (17), 6094-6121.

590 59. Lane, L. A.; Qian, X.; Smith, A. M.; Nie, S. J. A. r. o. p. c., Physical Chemistry of Nanomedicine: Understanding the
591 Complex Behaviors of Nanoparticles In-Vivo. **2015**, *66*, 521.

592 60. Fleischer, C. C.; Payne, C. K. J. A. o. c. r., Nanoparticle–cell interactions: molecular structure of the protein corona
593 and cellular outcomes. **2014**, *47*, (8), 2651-2659.

594 61. Walczyk, D.; Bombelli, F. B.; Monopoli, M. P.; Lynch, I.; Dawson, K. A. J. J. o. t. A. C. S., What the cell “sees” in
595 bionanoscience. **2010**, *132*, (16), 5761-5768.

596 62. Wei, H.; Wang, E. J. C. S. R., Nanomaterials with enzyme-like characteristics (nanozymes): next-generation artificial
597 enzymes. **2013**, *42*, (14), 6060-6093.

598 63. Dowding, J. M.; Das, S.; Kumar, A.; Dosani, T.; McCormack, R.; Gupta, A.; Sayle, T. X.; Sayle, D. C.; von Kalm, L.; Seal,
599 S. J. A. n., Cellular interaction and toxicity depend on physicochemical properties and surface modification of
600 redox-active nanomaterials. **2013**, *7*, (6), 4855-4868.

64. Rico, C. M.; Johnson, M. G.; Marcus, M. A. J. E. S. N., Cerium oxide nanoparticles transformation at the root–soil interface of barley (*Hordeum vulgare* L.). **2018**, *5*, (8), 1807-1812.
65. Limbach, L. K.; Wick, P.; Manser, P.; Grass, R. N.; Bruinink, A.; Stark, W. J. J. E. s.; technology, Exposure of engineered nanoparticles to human lung epithelial cells: influence of chemical composition and catalytic activity on oxidative stress. **2007**, *41*, (11), 4158-4163.
66. Gong, X.; Huang, D.; Liu, Y.; Zeng, G.; Wang, R.; Wan, J.; Zhang, C.; Cheng, M.; Qin, X.; Xue, W. J. E. s.; technology, Stabilized nanoscale zerovalent iron mediated cadmium accumulation and oxidative damage of *Boehmeria nivea* (L.) Gaudich cultivated in cadmium contaminated sediments. **2017**, *51*, (19), 11308-11316.
67. Ma, Y.; Xie, C.; He, X.; Zhang, B.; Yang, J.; Sun, M.; Luo, W.; Feng, S.; Zhang, J.; Wang, G., Effects of Ceria Nanoparticles and CeCl₃ on Plant Growth, Biological and Physiological Parameters, and Nutritional Value of Soil Grown Common Bean (*Phaseolus vulgaris*). *Small* **2020**, 1907435.
68. Yang, J.; Cao, W.; Rui, Y., Interactions between nanoparticles and plants: phytotoxicity and defense mechanisms. *Journal of Plant Interactions* **2017**, *12*, (1), 158-169.
69. Van Roten, A.; Barakat, A. Z. A.-Z.; Wouters, A.; Tran, T. A.; Mouton, S.; Noben, J.-P.; Gentile, L.; Smeets, K., A carcinogenic trigger to study the function of tumor suppressor genes in *Schmidtea mediterranea*. *Disease models & mechanisms* **2018**, *11*, (9), dmm032573.
70. Pearson, B. J.; Alvarado, A. S. J. D., A planarian p53 homolog regulates proliferation and self-renewal in adult stem cell lineages. **2010**, *137*, (2), 213-221.
71. Arslan, K.; Akbaba, G. B. J. T.; Health, I., In vitro genotoxicity assessment and comparison of cerium (IV) oxide micro-and nanoparticles. **2020**, *36*, (2), 76-83.