

InSAR application for the detection of precursors on the Achoma landslide, Peru

Dini, Benedetta; Lacroix, Pascal; Doin, Marie-Pierre

DOI:

[10.5194/egusphere-egu22-12583](https://doi.org/10.5194/egusphere-egu22-12583)

License:

Creative Commons: Attribution (CC BY)

Document Version

Publisher's PDF, also known as Version of record

Citation for published version (Harvard):

Dini, B, Lacroix, P & Doin, M-P 2022, InSAR application for the detection of precursors on the Achoma landslide, Peru. in *EGU General Assembly 2022.*, EGU22-12583, European Geosciences Union, EGU General Assembly 2022, Vienna, Austria, 23/05/22. <https://doi.org/10.5194/egusphere-egu22-12583>

[Link to publication on Research at Birmingham portal](#)

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

EGU22-12583

<https://doi.org/10.5194/egusphere-egu22-12583>

EGU General Assembly 2022

© Author(s) 2023. This work is distributed under the Creative Commons Attribution 4.0 License.



InSAR application for the detection of precursors on the Achoma landslide, Peru

Benedetta Dini, Pascal Lacroix, and Marie-Pierre Doin

University of Grenoble, ISTerre, France (b.dini@univ-grenoble-alpes.fr)

In the last few decades, InSAR has been used to identify ground deformation related to slope instability and to retrieve time series of landslide displacements. In some cases, retrospective retrieval of time series revealed acceleration patterns precursory to failure. This suggests that, the higher temporal sampling of new generation satellites, may indeed offer the opportunity to detect motion precursory to failure with viable lead time.

However, the possibility to retrieve continuous time series over landslides is often impaired by factors such as unfavourable orientation or landcover and fast movements, which make phase unwrapping difficult if not, in certain cases, impossible.

One way to retrieve precursors of destabilisation for landslides that present characteristics unfavourable to unwrapping and to time series inversion is to analyse in detail changes in successive interferograms in the phase domain in combination with interferometric coherence.

We generated and analysed 102 Sentinel-1 interferograms, covering the period between April 2015 and February 2021, at high spatial resolution (8 and 2 looks in range and azimuth respectively) over the Achoma landslide in the Colca valley, Peru. This large, deep-seated landslide, covering an area of about 40 hectares, previously unidentified, failed on 18th June 2020, damming the Rio Colca and giving origin to a lake.

We developed a method to analyse the changes through time of the unwrapped phase difference between a stable point and points within the landslide. In combination with this, we investigated patterns of coherence loss both within the landslide and in the surrounding area.

We observed that, in the weeks prior to the landslide, there was an increase of the phase difference between a stable reference and points within the landslide, indicating an acceleration of the downslope displacements. In addition to that, seasonal coherence loss is seen both within the landslide and in the surrounding area, in correspondence with wet periods. However, we observed also significant, local coherence loss outlining the scarp and the southeastern flank of the landslide, intermittently in the years before failure, in periods in which coherence was overall higher. Moreover, we observe a sharp decrease in the ratio between the coherence within the landslide and in the surrounding area, roughly six months before the failure.

This type of approach is promising with respect to the extraction of relevant information from

interferometric data when the generation of accurate and continuous time series of displacements is hindered by the nature of landcover or of the landslide studied, such in the case of the Achoma landslide. The combination of key, relevant parameters and their changes through time obtained with this methodology may prove necessary for the identification of precursors over a wider range of landslides than with time series generation alone.