

# Reviving the use of inhibitors of matrix metalloproteases in spinal cord injury

Ahmed, Zubair

DOI:

[10.4103/1673-5374.367837](https://doi.org/10.4103/1673-5374.367837)

License:

Creative Commons: Attribution-NonCommercial-ShareAlike (CC BY-NC-SA)

*Document Version*

Publisher's PDF, also known as Version of record

*Citation for published version (Harvard):*

Ahmed, Z 2023, 'Reviving the use of inhibitors of matrix metalloproteases in spinal cord injury: a case for specificity', *Neural Regeneration Research*, vol. 18, no. 9, pp. 1925-1930. <https://doi.org/10.4103/1673-5374.367837>

[Link to publication on Research at Birmingham portal](#)

## General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

## Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact [UBIRA@lists.bham.ac.uk](mailto:UBIRA@lists.bham.ac.uk) providing details and we will remove access to the work immediately and investigate.

# Reviving the use of inhibitors of matrix metalloproteases in spinal cord injury: a case for specificity

Zubair Ahmed<sup>1,2,\*</sup><https://doi.org/10.4103/1673-5374.367837>

Date of submission: September 19, 2022

Date of decision: December 2, 2022

Date of acceptance: January 12, 2023

Date of web publication: February 6, 2023

## Abstract

At present, there are no restorative therapies in the clinic for spinal cord injury, with current treatments offering only palliative treatment options. The role of matrix metalloproteases is well established in spinal cord injury, however, translation into the clinical space was plagued by early designs of matrix metalloprotease inhibitors that lacked specificity and fears of musculoskeletal syndrome prevented their further development. Newer, much more specific matrix metalloprotease inhibitors have revived the possibility of using these inhibitors in the clinic since they are much more specific to their target matrix metalloproteases. Here, the evidence for use of matrix metalloproteases after spinal cord injury is reviewed and researchers are urged to overcome their old fears regarding matrix metalloprotease inhibition and possible side effects for the field to progress. Recently published work by us shows that inhibition of specific matrix metalloproteases after spinal cord injury holds promise since four key consequences of spinal cord injury could be alleviated by specific, next-generation matrix metalloprotease inhibitors. For example, specific inhibition of matrix metalloprotease-9 and matrix metalloprotease-12 within 24 hours after injury and for 3 days, alleviates spinal cord injury-induced edema, blood-spinal cord barrier breakdown, neuropathic pain and restores sensory and locomotor function. Attempts are now underway to translate this therapy into the clinic.

**Key Words:** axon regeneration; blood-spinal cord barrier; edema; functional recovery; matrix metalloprotease-9; matrix metalloproteases-12; pain; spinal cord injury

## From the Contents

Introduction	1925
Search Strategy and Selection Criteria	1926
Matrix Metalloproteases in Spinal Cord Injury	1926
Matrix Metalloproteases and Their Inhibitors in Spinal Cord Injury	1926
Inflammation and Blood-Spinal Cord Barrier Breakdown	1926
Oxidative Stress and Apoptosis	1927
Matrix Metalloproteases in Glial Scar Formation	1927
Matrix Metalloproteases in Neuropathic Pain	1927
Matrix Metalloproteases in Human Spinal Cord Injury	1927
Matrix Metalloprotease Inhibitors	1928
Inhibition of Matrix Metalloprotease-9 and Matrix Metalloprotease-12 Using AZD1236 to Improve Function after Spinal Cord Injury	1928

## Introduction

Spinal cord injury (SCI) is a devastating condition, often causing permanent loss of function and significant disability (Wilson et al., 2012). At present, there are no fully restorative treatments available for SCI and hence this remains an area of urgent medical need. Treatments such as Lyrica and Medrol, only offer symptomatic relief to patients from neuropathic pain but do not address the underlying mechanisms that cause neuropathic pain or inflammation (Tong et al., 2021). A fully restorative therapy for SCI would therefore promote the protection of neurons and glia from further damage, neutralize inhibitory molecules, inhibit the glial scar, replace lost neurons, and promote axon regeneration and appropriate synapse formation.

Matrix metalloproteases (MMPs) are zinc-dependent enzymes that are involved in a variety of proteolytic events during development, wound healing, and repair processes (Sternlicht and Werb, 2001). For several decades, the role of MMPs has been investigated in the pathogenesis of neurodegenerative diseases and models of central nervous system (CNS)

injury, including SCI and traumatic brain injury (Yong, 2005; Zhang et al., 2010). In SCI, MMPs contribute to the degradation of the blood-spinal cord barrier (BSCB), oxidative stress, demyelination, and a progressive neuroinflammatory response (Noble et al., 2002; Wells et al., 2003; Zhang et al., 2011).

The MMPs are a family of 23 human or 24 murine members and can be subdivided into eight different types based on their domain structure and substrate specificity (Yong, 2005; Page-McCaw et al., 2007; Zhang et al., 2011; **Table 1**). MMPs are initially produced as inactive zymogens and require catalytic activation where the propeptide domain is cleaved and thus exposing the active catalytic site. MMPs are regulated at both the transcriptional and post-translation levels and their activity is regulated by four endogenous inhibitors of metalloproteases, called tissue inhibitors of MMPs (TIMPs) (Sternlicht and Werb, 2001). TIMPs can sequester MMP activity by binding to their catalytic sites and thus precisely control their activity and limit extracellular matrix (ECM) degradation (Sternlicht and Werb, 2001; Crocker et al., 2004).

**Table 1 | Classification of MMPs based on domain structures**

Domain structure	MMP
Simple hemopexin domain-containing MMPs	MMP-1 (collagenase 1), MMP3 (stromelysin 1), MMP-8 (collagenase 2), MMP-10 (stromelysin 2), MMP-12 (matelloelastase), MMP-13 (collagenase 3), MMP-19 (RASI 1), MMP-20 (enammelysin), MMP-27
Gelatinase MMP	MMP-2 (gelatinase A) and MMP-9 (gelatinase B)
Furin-activated secreted MMPs	MMP-11 (stromelysin 3), MMP28 (epilysin)
Minimal domain MMPs	MMP-7 (matrilysin) and MMP-2 (endometase)
GPI-linked MT-MMPs	MMP-17 (MT-MMP), MMP-25 (MT-MMP)
Transmembrane MMPs	MMP1 (MT1-MMP), MMP-15 (MT2-MMP), MMP-1 (MT3-MMP), MMP2 (MT5-MMP)
Cysteine/proline-rich IL-1 receptor-like domain MMP	MMP-23
ADAMs	ADAMs

ADAM: A disintegrin and metalloproteinase; GPI: glycosylphosphatidylinositol; IL-1: interleukine-1; MMP: matrix metalloprotease; MT-MMP: membrane type matrix metalloprotease.

<sup>1</sup>Neuroscience and Ophthalmology, Institute of Inflammation and Ageing, University of Birmingham, Edgbaston, Birmingham, UK; <sup>2</sup>Center for Trauma Sciences Research, University of Birmingham, Edgbaston, Birmingham, UK

\*Correspondence to: Zubair Ahmed, PhD, z.ahmed.1@bham.ac.uk.  
<https://orcid.org/0000-0001-6267-6442> (Zubair Ahmed)

**How to cite this article:** Ahmed Z (2023) Reviving the use of inhibitors of matrix metalloproteases in spinal cord injury: a case for specificity. *Neural Regen Res* 18(9):1925-1930.

## Search Strategy and Selection Criteria

The search strategy and selection criteria were limited to articles published in peer-reviewed journals articles. A literature review of articles was conducted by searching PubMed and Web of Science databases, updated until August 2022, searching for the following topics: matrix metalloproteases and/or MMP and spinal cord injury, matrix metalloproteases and/or MMP and SCI, matrix metalloprotease and/or MMP-9 and spinal cord injury, matrix metalloprotease and/or MMP-12 and spinal cord injury, matrix metalloprotease and/or MMP inhibitors and spinal cord injury, matrix metalloprotease and/or MMP-9 inhibitor and spinal cord injury, matrix metalloprotease and/or MMP-12 inhibitor and spinal cord injury. The selected articles focused on MMP expression and MMP inhibition in spinal cord injury. Human and animal studies were all screened.

## Matrix Metalloproteases in Spinal Cord Injury

A number of studies have profiled the expression of MMPs after SCI. The first MMPs to be studied in SCI were gelatinases MMP-9 and MMP-2. It was reported that MMP-9 activity was observed within 12–24 hours postinjury whilst MMP-2 rose significantly by 5 days after contusion injury in the rat (de Castro et al., 2000). These results were later confirmed in other species and models of SCI, including hemisection and compression in a flurry of studies. For example, zymography and western blot assay showed transient increases in MMP-9 at 24 hours post injury with a gradual increase in MMP-2 (Duchossoy et al., 2001; Xu et al., 2001; Goussev et al., 2003; Hsu et al., 2006; Yu et al., 2008). In a compression model of SCI, mRNAs for MMP-9, MMP-3, MMP-7, MMP-10, MMP-11, MMP13, MMP-19, and MMP-20 were significantly upregulated within 24 hours whereas MMP-2, MMP-12, and MMP-13 upregulation was delayed until 5 days after SCI (Wells et al., 2003; Veeravalli et al., 2009; Table 2).

After SCI, MMP-1 and MMP-9 were both localized in glia and neurons (Xu et al., 2001), whilst MMP-9 was also observed in blood vessels, neutrophils,

and macrophages (Xu et al., 2001; Noble et al., 2002; Yu et al., 2008). MMP-2 not only localizes to neurons and astrocytes but also to microglia and macrophages (Wells et al., 2003; Hsu et al., 2006; Veeravalli et al., 2009). Other MMPs were also localized in the spinal cord after injury but their temporal and cellular localization varied after SCI and related to the type and severity of the injury.

## Matrix Metalloproteases and Their Inhibitors in Spinal Cord Injury

In the acute phase after CNS injury, MMPs largely play a number of detrimental roles such as degradation of the basal lamina, disruption of BSCB, oxidative stress, demyelination, leukocyte infiltration, and progressive neuroinflammation (Noble et al., 2002; Wells et al., 2003; Hsu et al., 2006; Zhang et al., 2011). In the chronic phase, however, MMPs play a beneficiary role in modulating the inhibitory glial scar, promoting cell survival, supporting axonal regeneration, and attenuating mechanical allodynia (Chang and Werb, 2001; Duchossoy et al., 2001; Yong, 2005; Zhao et al., 2006). Therefore, inhibiting MMP activity is a complex process and requires intricate knowledge of the timeline of MMP activities after SCI.

## Inflammation and Blood-Spinal Cord Barrier Breakdown

MMPs facilitate the extravasation of inflammatory cells into the injured spinal cord and contribute to the breakdown of the BSCB. When infiltrating leukocytes migrate across the vascular wall, they secrete MMPs which then degrade tight junction proteins since zonulae occludens-1, VE-cadherin, and occludins are substrates for MMP-2, -3, -7, and -9 (Asahi et al., 2001; Caron et al., 2005; Yang et al., 2007; Buhler et al., 2009). The target of MMPs also included proteins found in the basal lamina such as laminin, fibronectin, and heparan sulfate (Rosenberg and Yang, 2007).

**Table 2 | Summary of MMPs in experimental spinal cord injury models**

MMP	Type of injury	Species	Changes in MMP	Peak	Localization	Studies
MMP-1	Contusion	Rat	Increased activity	24 h	Neurons, glia	Xu et al., 2001; Veeravalli et al., 2009
MMP-2	Contusion	Rat	Increased mRNA	1–21 d	Not studied	de Castro et al., 2000 Goussev et al., 2003; Hsu et al., 2006 Duchossoy et al., 2001 Yu et al., 2008 Kawasaki et al., 2008 Wells et al., 2003 Dang et al., 2008
	Contusion	Mouse		7–14 d	Astrocytes	
	Hemisection	Rat		1 d to 4 wk	Not studied	
	Compression	Rat		3–7 d	Not studied	
	Compression	Mouse	Increased mRNA	5 d	Not studied	
	Contusion	Mouse		5 d	Not studied	
	Contusion	Rat		3–21 d	Neurons	
MMP-9	Contusion	Rat	Increased activity	12–24 h	Not studied	Veeravalli et al., 2009 Noble et al., 2002; Goussev et al., 2003 Xu et al., 2001 Duchossoy et al., 2001 Yu et al., 2008 Wells et al., 2003 Veeravalli et al., 2009
	Contusion	Mouse		24 h	Astrocytes, blood vessels, macrophages	
	Contusion	Rat		24 h	Neurons, glia	
	Hemisection	Rat		24–48 h	Not studied	
	Compression	Rat		24 h	Neutrophil, endothelial cells, neurons	
	Compression	Mouse	Increased mRNA	24 h	Not studied	
	Compression	Mouse	Increased mRNA	5–14 d	Macrophages/microglia	
MMP-12	Contusion	Rat		21 d	Not studied	Wells et al., 2003 Veeravalli et al., 2009
	Compression	Mouse	Increased mRNA	24 h	Not studied	
MMP-3	Contusion	Rat		3 d	Not studied	Wells et al., 2003; Veeravalli et al., 2009
	Compression	Mouse		24 h	Not studied	
MMP-7	Contusion	Rat		3 d		
	Compression	Mouse		24 h		
MMP-10	Contusion	Rat		3–21 d		
	Compression	Mouse		24 h		
MMP-11	Contusion	Rat		3 d		
	Compression	Mouse		5 d		
MMP-13	Contusion	Rat		3 d		
	Compression	Mouse		5 d		
MMP-14	Contusion	Rat		3 d		
				1–21 d		
MMP-16				1–21 d		
MMP-17				24 h		
MMP-19				24 h		
MMP-20				21 d		
MMP-21				1–2 d		
MMP-23	Compression	Mouse	Decreased mRNA	1–2 d		Wells et al., 2003
MMP-24				2–5 d		
MMP-15			Decreased mRNA	1–3 d		Veeravalli et al., 2009
			Increased mRNA	7–21 d		
MMP-8			Decreased mRNA	1 d		
MMP-25			Increased	3–21 d		

MMP: Matrix metalloprotease.

MMP-9 mediates acute SCI pathogenesis in the injured spinal cord, derived primarily from infiltrating leukocytes since depleting leukocytes reduces MMP-9 activity in injured tissues (de Castro et al., 2000). MMP-9 expression (in both rodents and humans) peaks within one day after SCI and is detected in glia, macrophages, neutrophils, and vascular element in the SCI 24 hours after injury (Noble et al., 2002; Goussev et al., 2003; Buss et al., 2007). Excessive activity of MMP-9 in the acute phase of SCI disrupts the BSCB, edema, excitotoxicity, infiltration of leukocytes, mitochondrial dysfunction, apoptosis, demyelination of neurons, increase in inflammatory responses and astrogliosis (Noble et al., 2002). Disruption of BSCB after SCI is maximal 24 hours after injury, coinciding with the peak activity of MMP-9 in the injured spinal cord (Noble et al., 2002). Indeed, BSCB disruption was reduced in MMP-9 KO mice and mice treated with the broad-spectrum MMP inhibitor, GM6001 (Noble et al., 2002). Moreover, treatment with SB-3CT, a gelatinase inhibitor, at 2 hours before SCI, reduced MMP-9 activity and BSCB disruption, ultimately decreasing apoptosis (Yu et al., 2008).

Similar findings were also observed in MMP-12 KO mice, where greater stabilization of the BSCB was observed. Indeed, MMP-12 (macrophage metalloelastase) is required for the migration of blood-borne macrophages across endothelial cell basement membranes and into areas of inflammation (Shipley et al., 1996; Wells et al., 2003). MMP-12 is also expressed by macrophages and increases 182-fold in a compression SCI model, perpetuating the inflammatory response and acute edema and promoting the development of the secondary injury response (Wells et al., 2003, 2005).

## Oxidative Stress and Apoptosis

MMPs are regulated by oxidative stress and reactive oxygen species such as nitric oxide and hypochlorous acid increase MMP-9 and MMP-9 activity and decrease TIMPs (Rajagopalan et al., 1996; Morita-Fujimura et al., 2000). Increased levels of oxidative stress after SCI lead to MMP-9 upregulation and BSCB breakdown as well as apoptosis via neurotoxicity (Yu et al., 2008). In support of this, transgenic rats that overexpress the antioxidant enzyme superoxide dismutase 1, showed that oxidative stress and MMP-9-mediated BSCB breakdown and apoptosis were all attenuated (Morita-Fujimura et al., 2000; Sugawara et al., 2002). MMP-2 contributes to apoptosis and is upregulated after injury, coinciding with glial and neuronal apoptosis. MMP-2 therefore has been suggested to mediate apoptosis after SCI since the suppression of MMP-2/MMP-9 reduced the levels of apoptosis in glia and neurons, which may promote long-term recovery (Dang et al., 2008).

Importantly, these studies demonstrate that MMPs direct disruption of the BSCB, infiltration of leukocytes and apoptosis after SCI. Since early and short-term (for up to 5 days after SCI) inhibition of MMPs stabilized the BSCB, reduced infiltration of leukocytes may confer acute and long-term neuroprotection, suggesting that acute inhibition of MMPs may be a promising strategy for SCI.

## Matrix Metalloproteases in Glial Scar Formation

A glial scar at the lesion site normally forms after SCI, containing axon growth inhibitory molecules, ECM molecules, microglia/macrophages, and reactive astrocytes and ECM molecules such as chondroitin sulphate proteoglycans (CSPG). Increased expression of CSPGs results from the interaction of astrocytes, oligodendrocytes, and macrophages, which in turn inhibits axon regeneration (Bradbury et al., 2002; Jones et al., 2002). MMPs degrade the core protein of a variety of inhibitory molecules including CSPGs, Nogo, and tenascin-C, whilst CSPGs such as tenascin-C, brevican, neurocan, NG2, phosphacan, and versican are degraded by MMP activity (Pizzi and Crowe, 2007). Therefore, MMPs support axonal regeneration in the injured CNS by degrading CSPGs and other inhibitory molecules (Yong, 2005; Pizzi and Crowe, 2007).

MMPs also facilitate the migration of astrocytes since gelatinase activity is correlated with scar formation (Duchosoy et al., 2001). *In vitro* assays demonstrate that astrocytes derived from MMP-9 KO mice or astrocytes treated with an MMP-9 inhibitor both attenuate their migration (Hsu et al., 2008). Consistent with this observation, the glial scar is abrogated after SCI in MMP-9 KO mice along with reduced CSPG immunohistochemistry (Hsu et al., 2008). On the other hand, MMP-2 KO mice showed increased immunoreactivity for CSPG, fewer serotonergic fibers and significantly reduced motor recovery compared to wild-type mice, suggesting reduced axonal sprouting across the lesion site and the important role that MMP-2 plays in promoting functional recovery after SCI (Hsu et al., 2006). MMPs also play a role in axonal dieback since inhibition of MMP-9 prevented macrophage-mediated axon retraction (Busch et al., 2009). These studies suggest that MMPs are critical to the formation of a glial scar but can degrade inhibitory molecules and cleave ECM proteins that sequester growth factors and hence support function recovery (Yong et al., 2001; Pizzi and Crowe, 2007).


## Matrix Metalloproteases in Neuropathic Pain

Neuropathic pain is common in as many as 85% of SCI patients in dermatomes above and below the lesion, manifesting in allodynia and hyperalgesia (Siddall et al., 2003) but the mechanisms causing neuropathic pain remains unanswered. Neuropathic pain is normally studied using the peripheral nerve injury model where neuropathic pain can be modeled in the absence of damage to central pathways. However, studies suggest that the

mechanisms causing neuropathic pain after SCI and peripheral nerve injury are similar (Detloff et al., 2008). In peripheral nerve injury, MMPs both induce and maintain neuropathic pain, primarily regulated by MMP-9 and MMP-2. MMP-9 levels rise rapidly within 24 hours after injury whilst MMP-2 levels are delayed until day 7 but persist for up to 21 days (Kawasaki et al., 2008). The differential time course of MMP-9 and MMP-2 in the development and maintenance of neuropathic pain after injury suggests a role for MMP-9 and MMP-2 but a direct role is yet to be established.

It is thought that mechanical injury to axons causes Schwann cells to secrete MMP-9 which then activates the infiltration of macrophages and stimulates the degradation of myelin proteins such as myelin basic protein (Chandler et al., 1995; Kobayashi et al., 2008). This causes exposure of the bare nerve endings, resulting in increased sodium channel expression and ectopic hyperexcitability of afferents (Devor, 2006). Action potentials now outlast the stimulus and create central sensitization, a common mechanism that is found in the transmission of neuropathic pain.

It has long been established that proinflammatory cytokines and growth factors also regulate neuropathic pain since they mediate the expression of gelatinases in the CNS. For example, tumor necrosis factor- $\alpha$  or interleukin-1 $\beta$  (IL-1 $\beta$ ) induces MMP-9 expression by Schwann cells, leading to neuronal sensitization (Chattopadhyay et al., 2007; Schafers and Sorkin, 2008). Deletion of MMP-9 reduces pain behaviors after peripheral nerve injury, whilst blocking IL-1 $\beta$  signaling with a neutralizing antibody-prevented allodynia (Chattopadhyay et al., 2007; Kawasaki et al., 2008). Conversely, intrathecal injection of MMP-9 increased the levels of allodynia with increased cleavage of IL-1 $\beta$  and microglial activation in the dorsal horn and colocalized with activated p38 MAP kinase (Kawasaki et al., 2008). MMP-2 also cleaves IL-1 $\beta$  and activates spinal astrocytes at later time points, potentially maintaining neuropathic pain (Kawasaki et al., 2008). This suggests that MMP-9 and MMP-2 are intricately related to IL-1 $\beta$ , which may be a potential downstream regulator of neuropathic pain (Figure 1).



**Figure 1 | Schematic to show mechanisms of MMP-induced neuropathic pain.**

Nerve injury activates proinflammatory cytokines such as TNF- $\alpha$  which in turn increase MMP-9 expression in the DRG. MMP-9 is then activated and increases IL-1 $\beta$  cleavage in the DRG and spinal cord which sensitizes primary sensory neurons and enhances synaptic transmission of dorsal horn neurons, leading to pain hypersensitivity. A positive feedback loop between IL-1 $\beta$  and p38 exists and can enhance IL-1 $\beta$  production. Adapted from Kawasaki et al. (2008). DRG: Dorsal root ganglia; IL-1 $\beta$ : interleukin-1 $\beta$ ; MMP: matrix metalloprotease; TNF- $\alpha$ : tumor necrosis factor- $\alpha$ .

Apart from gene deletion (Table 3), short interfering RNA (siRNA) and peptide inhibitors to MMP-9 and MMP-2 have been used and demonstrated to be effective in reducing allodynia in the peripheral nerve injury model. Daily injections of broad-spectrum gelatinase inhibitor, GM6001, resulted in immediate and sustained attenuation of allodynia, preserving myelin basic protein, reducing infiltration of macrophages and reducing glial activation in the dorsal horn (Kobayashi et al., 2008). Likewise, treatment with siRNA to MMP-9 prevented the onset of allodynia whilst MMP-2 siRNA also attenuated allodynia (Kawasaki et al., 2008). Administration of TIMP-1 and TIMP-2 led to a greater reversal of allodynia but the effects were transient and lasted 3 to 24 hours (Kawasaki et al., 2008). Inhibition of MMP-9 using Inhibitor-I delayed the onset of allodynia by 6 days whilst inhibition of MMP-2 using Inhibitor III attenuated allodynia for 10 days (Kawasaki et al., 2008) suggesting that MMP-9 and MMP-2 inhibition can attenuate neuropathic pain.

## Matrix Metalloproteases in Human Spinal Cord Injury

Although the majority of the data on MMPs in SCI is from animal studies, there are a small number of studies in humans that corroborate some of the findings. For example, in post-mortem spinal cord MMP-9 was found at 2 days after injury (which was the earliest time studied) at the lesion site and localized in neutrophils and macrophages (Buss et al., 2007) whilst another study localized MMP-9 in neutrophils with peak expression at 4 hours – 3 days, in agreement with rodent studies (Fleming et al., 2006). However, MMP-12 showed a delayed expression of 24 days, far greater than the 5-day peak expression observed in mouse/rat studies (Buss et al., 2007). This may be due to an artifact arising from the usage of post-mortem spinal cord for experiments or could be a discrepancy arising from mouse and human



**Table 3 | KO animals and their effects in SCI**

MMP KO or Inhibitor	Mechanism	Effects after SCI	Studies
MMP-2 <sup>-/-</sup>	Elimination of MMP-2; compensatory increase in MMP-9	Reduced white matter sparing and fewer serotonergic fibres, widespread reactive astrogliosis, and impairment in locomotor recovery.	Hsu et al., 2006
MMP-9 <sup>-/-</sup>	Elimination of MMP-9	Improved function recovery and reduced barrier disruption, neutrophil infiltration, neuropathic pain and inflammatory pain.	Noble et al., 2002; Kawasaki et al., 2008
MMP-12 <sup>-/-</sup>	Elimination of MMP-12	Improved functional recovery, attenuation of barrier disruption, and reduces microglial/macrophages.	Wells et al., 2003

KO: Knockout; MMP: matrix metalloproteases; SCI: spinal cord injury.

disease models. Although an exploratory study suggested using serum levels of MMP-8 and MMP-9 as early markers for remission, the sample sizes were too low to make definitive conclusions (Moghaddam et al., 2017). Further studies into the temporal expression of MMPs after human SCI are required but are obviously difficult to perform. However, serum and cerebrospinal fluid levels of MMPs will give us an indication of what is happening in terms of MMP activation after SCI in humans.

## Matrix Metalloprotease Inhibitors

The search for MMP inhibitors began over three decades ago (Table 4) but initial clinical trials were disappointing and painted a negative picture of MMPs as therapeutic targets. With an improved understanding of the biology of MMPs, it emerged that clinical trials were conducted prematurely. Better-designed MMP inhibitors had desirable selectivity's and improved pharmacokinetic profiles, resulting in lower toxicity. Selective MMP inhibitors demonstrated that MMP-2, MMP-9, MMP-13, and MT1-MMP were not involved in musculoskeletal syndrome that plagued early, broad-spectrum MMP inhibitors. Musculoskeletal syndrome appeared to be caused by off-target effects of inhibiting not just one MMP but a combination of MMPs and/or possibly other related enzymes (Peterson, 2006).

The first generation of peptidomimetics MMP inhibitors was broad-spectrum, containing a hydroxamate moiety that chelated catalytic zinc and inactivated the protein such as batimastat, marimastat, and ilomastat. The hydroxamate moiety was readily metabolized and non-specifically inhibited ADAMs as well as producing musculoskeletal syndrome (Peterson, 2004; Jacobsen et al., 2010). Other inhibitors include those that contain a mercaptoacyl as a zinc-binding group (Rebimastat), and tanomastat has a zinc-binding carboxylate group. However, as these are zinc chelators they target gelatinases as well as other zinc-dependent enzymes, including other MMPs. Doxycycline and Minocycline are tetracycline-based inhibitors with broad-spectrum MMP inhibitory activity (Romero-Perez et al., 2008), whilst bisphosphonates and carbamoyl phosphobates also inhibit various MMPs (Teronen et al., 1999; Matziari et al., 2004).

SB-3CT was the first mechanism-based inhibitor of gelatinase where the inhibitor-enzyme complex underwent a requisite conformational change that did not readily reverse. SB-3CT is selective for MMP-2 and MMP-9 but does not inhibit other MMPs except for MMP-14 (Toth et al., 2000). SB-3CT showed enhanced efficacy than previous MMP inhibitors but it was rapidly metabolized to a more active gelatinase inhibitor than the parent molecule, limiting its use (Lee et al., 2007). Although these early MMP inhibitors displayed high affinity *in vitro*, the first generation of MMP inhibitors failed in clinical trials due to their low selectivity and nonspecific inhibition of MMPs and non-related proteins that can be affected by the metal chelating groups *in vivo* (Overall and Kleifeld, 2006; Fingleton, 2008; Lopez-Otin et al., 2009).

Better designs of MMP inhibitors soon followed with the targeting of inhibitory sites outside the conserved catalytic cleft, the exosites, which produced selective MMP inhibition *in vivo* and thus revived the field's interest to selectively control the pathophysiological activity of these enzymes (Turk, 2006; Sela-Passwell et al., 2010; Deu et al., 2012). This led to the development of macromolecule inhibitors, which were inhibitory antibodies against various MMPs beyond the enzyme catalytic domain. These antibodies were much more specific and highly selective but still retained the homeostatic activity of other MMP family members. Highly selective MMP-9 inhibitors such as AB0041 and AB0046 were developed either as full-length human MMP-9 or a mouse Pro-Cat construct (Marshall et al., 2015). Other inhibitory antibodies such as SDS3, an MMP-2/-9 inhibitor soon followed (Sela-Passwell et al., 2011) together with the first anti-MMP-9 antibody (REGA-3GL12) (Paemen et al., 1995).

**Table 4 | MMP inhibitors and their use after SCI**

Inhibitor	Mechanism	Effects after SCI	Studies
MMP-2/MMP-9 inhibitor, subcutaneous	Inhibit MMP activity, especially MMP-9 and MMP-2	Reduced apoptosis in neurons and glia.	Dang et al., 2008
SB-3CT	Inhibit MMPs, especially MMP-9 and MMP-2	Reduced MMP activity, barrier disruption, and apoptosis.	Yu et al., 2008
GM-6001	Broad spectrum MMP inhibitor (active against MMP-1, MMP-3, MMP-2, MMP-8, and MMP-9)	Tissue sparing and improved locomotor function when given over the first 3 days post-injury	Noble et al., 2002
		In dogs, GM6001+DMSO or DMSO alone improved functional recovery. NB: Authors concluded functional improvement were due to DMSO and not GM6001.	Levine et al., 2014
Lipitor	Inhibit MMP-9	Prevented endothelial dysfunction, facilitated neuroprotection, and promoted locomotor recovery	Pannu et al., 2007
Fluoxetine	Inhibit MMP-2, MMP-9, and MMP-12	Prevented BSCB disruption, prevented infiltration of leucocytes, reduced expression of inflammatory mediators, attenuated apoptotic cell death and improved locomotor function recovery	Scali et al., 2013
ND-378	Inhibit MMP-2	No changes in long-term neurological outcomes	Gao et al., 2016
Sulforaphane	Inhibit MMP-9	Inhibited microvascular leakage and inflammatory cytokine expression	Benedict et al., 2012

BSCB: Blood-spinal cord barrier; DMSO: dimethyl sulfoxide; MMP: matrix metalloproteases; SCI: spinal cord injury.

There are several clinical trials listed on ClinicalTrials.gov (last accessed 30/05/2022) evaluating MMP inhibitors in a variety of conditions such as cancer, glioblastoma, open-angle glaucoma, asthma, and chronic obstructive pulmonary disease. With the advent of much more specific MMP inhibitors, the time is now right to let go of the dogma of viewing MMP inhibition as intractable.

## Inhibition of Matrix Metalloprotease-9 and Matrix Metalloprotease-12 Using AZD1236 to Improve Function after Spinal Cord Injury

AZD1236 is a potent and reversible inhibitor of MMP-9 and MMP-12, with an IC<sub>50</sub> of 4.5 and 6.1 nM, respectively. It has > 10-fold selectivity to MMP-12 and MMP-13 and > 350-fold selectivity to other members of the MMP family. AZD1236 was originally developed by AstraZeneca for use in chronic obstructive pulmonary disease and has been tested in single doses up to 1500 mg and 500 mg QD for 13 days, whilst COPD patients were dosed for 6 weeks with 75 mg BID (Magnussen et al., 2011; Dahl et al., 2012). We re-purposed AZD1236 in SCI and showed unprecedented benefits (Ahmed et al., 2022).

By critically analyzing the known pathophysiology associated with SCI, it is evident that excessive MMP activity, in particular, MMP-9 and MMP-12, contribute to four of the key pathological drivers of SCI, namely; disruption of the BSCB, edema, neuropathic pain and loss of motor/sensory function. We found that MMP-9 levels and their activity in the spinal cord, serum, and cerebrospinal fluid peaked within the first 24 hours, whilst MMP-12 peaked 5 days post-SCI (Ahmed et al., 2022). Inhibition of MMP-9 and MMP-12 activity using oral or intrathecal delivery of AZD1236, attenuated the SCI-induced rise in water content in the spinal cord by 98%; an effect that was directly related to inhibition of both MMP-9 and MMP-12. Measures of proinflammatory pain markers, BSCB breakdown, and scarring at the lesion site were also suppressed by 80%, 75%, and 80%, respectively. Inhibition of MMP-9 and MMP-12 promoted axon regeneration and protected against SCI-induced sensory and locomotor deficits such that animals behaved as uninjured sham controls by 3 weeks post-injury. Suppressed microglial/macrophage activation and reduced scar-derived inhibitors of axon growth at the lesion site, including Semaphorin-3A and CS-56, were also reduced further contributing to the promotion of axon regeneration after SCI (Bradbury et al., 2002).

Interestingly, the reduction of SCI-induced edema was reliant on the combined suppression of both MMP-9 and MMP-12, since the suppression of either of these individually had a suboptimal impact on spinal cord edema (Ahmed et al., 2022). Indeed, the clinical outcomes after SCI are greatly

affected by spinal cord edema and if left unchecked this can lead to further damage and death, posing a significant challenge to neurosurgeons (Rutges et al., 2017).

Our findings, therefore, demonstrate that selective short-term inhibition of MMP-9 and MMP-12 has the potential to offer previously unparalleled levels of protection against secondary damage to the spinal cord as well as the dramatic potential for recovery. The effects of AZD1236 after SCI show that the beneficial effects of MMP inhibition lie primarily in their selectivity and this plagued early MMP inhibitor designs. However, with advances in protein engineering and new technologies, it is possible to design next-generation MMP inhibitors with unparalleled selectivity and specificity such that old fears regarding MMP inhibition and possible side effects need to be overcome for the field to progress further. Our work in the spinal cord brings hope that MMP inhibition to target specific diseases may become a possible reality for future clinical translation.

**Author contributions:** ZA conceptualized and conducted the literature review, wrote the manuscript and created the Figures and Tables. ZA edited text and figures, revised the whole manuscript, and approved the final version of the manuscript for publication.

**Conflicts of interest:** The author is an inventor on a patent related to MMP-9 and MMP-12 inhibitors in neurological diseases.

**Data availability statement:** No additional data are available.

**Open access statement:** This is an open access journal, and articles are distributed under the terms of the Creative Commons AttributionNonCommercial-ShareAlike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

## References

- Ahmed Z, Alhajlah S, Thompson AM, Fairclough RJ (2022) Clinic-ready inhibitor of MMP-9/-12 restores sensory and functional decline in rodent models of spinal cord injury. *Clin Transl Med* 12:e884.
- Asahi M, Wang X, Mori T, Sumii T, Jung JC, Moskowitz MA, Fini ME, Lo EH (2001) Effects of matrix metalloproteinase-9 gene knock-out on the proteolysis of blood-brain barrier and white matter components after cerebral ischemia. *J Neurosci* 21:7724-7732.
- Benedict AL, Mountney A, Hurtado A, Bryan KE, Schnaar RL, Dinkova-Kostova AT, Talalay P (2012) Neuroprotective effects of sulforaphane after contusive spinal cord injury. *J Neurotrauma* 29:2576-2586.
- Bradbury EJ, Moon LD, Popat RJ, King VR, Bennett GS, Patel PN, Fawcett JW, McMahon SB (2002) Chondroitinase ABC promotes functional recovery after spinal cord injury. *Nature* 416:636-640.
- Buhler LA, Samara R, Guzman E, Wilson CL, Krizanac-Bengez L, Janigro D, Ethell DW (2009) Matrix metalloproteinase-7 facilitates immune access to the CNS in experimental autoimmune encephalomyelitis. *BMC Neurosci* 10:17.
- Busch SA, Horn KP, Silver DJ, Silver J (2009) Overcoming macrophage-mediated axonal dieback following CNS injury. *J Neurosci* 29:9967-9976.
- Buss A, Pech K, Kakulas BA, Martin D, Schoenen J, Noth J, Brook GA (2007) Matrix metalloproteinases and their inhibitors in human traumatic spinal cord injury. *BMC Neurol* 7:17.
- Caron A, Desrosiers RR, Beliveau R (2005) Ischemia injury alters endothelial cell properties of kidney cortex: stimulation of MMP-9. *Exp Cell Res* 310:105-116.
- Chandler S, Coates R, Gearing A, Lury J, Wells G, Bone E (1995) Matrix metalloproteinases degrade myelin basic protein. *Neurosci Lett* 201:223-226.
- Chang C, Werb Z (2001) The many faces of metalloproteases: cell growth, invasion, angiogenesis and metastasis. *Trends Cell Biol* 11:537-43.
- Chattopadhyay S, Myers RR, Janes J, Shubayev V (2007) Cytokine regulation of MMP-9 in peripheral glia: implications for pathological processes and pain in injured nerve. *Brain Behav Immun* 21:561-568.
- Crocker SJ, Pagenstecher A, Campbell IL (2004) The TIMPs tango with MMPs and more in the central nervous system. *J Neurosci Res* 75:1-11.
- Dahl R, Titlestad I, Lindqvist A, Wielders P, Wray H, Wang M, Samuelsson V, Mo J, Holt A (2012) Effects of an oral MMP-9 and -12 inhibitor, AZD1236, on biomarkers in moderate/severe COPD: a randomised controlled trial. *Pulm Pharmacol Ther* 25:169-177.
- Dang AB, Tay BK, Kim HT, Nauth A, Alfonso-Jaume MA, Lovett DH (2008) Inhibition of MMP2/MMP9 after spinal cord trauma reduces apoptosis. *Spine (Phila Pa 1976)* 33:E576-579.
- de Castro RC, Jr., Burns CL, McAdoo DJ, Romanic AM (2000) Metalloproteinase increases in the injured rat spinal cord. *Neuroreport* 11:3551-3554.
- Detloff MR, Fisher LC, McGaughy V, Longbrake EE, Popovich PG, Basso DM (2008) Remote activation of microglia and pro-inflammatory cytokines predict the onset and severity of below-level neuropathic pain after spinal cord injury in rats. *Exp Neurol* 212:337-347.
- Deu E, Verdoes M, Bogoy M (2012) New approaches for dissecting protease functions to improve probe development and drug discovery. *Nat Struct Mol Biol* 19:9-16.
- Devor M (2006) Centralization, central sensitization and neuropathic pain. Focus on "sciatic chronic constriction injury produces cell-type-specific changes in the electrophysiological properties of rat substantia gelatinosa neurons". *J Neurophysiol* 96:522-523.
- Duchossay Y, Horvat JC, Stettler O (2001) MMP-related gelatinase activity is strongly induced in scar tissue of injured adult spinal cord and forms pathways for ingrowing neurites. *Mol Cell Neurosci* 17:945-956.
- Fingleton B (2008) MMPs as therapeutic targets--still a viable option? *Semin Cell Dev Biol* 19:61-68.
- Fleming JC, Norenberg MD, Ramsay DA, Dekaban GA, Marcillo AE, Saenz AD, Pasquale-Styles M, Dietrich WD, Weaver LC (2006) The cellular inflammatory response in human spinal cords after injury. *Brain* 129:3249-3269.
- Gao M, Zhang H, Trivedi A, Mahasenan KV, Schroeder VA, Wolter WR, Suckow MA, Mobashery S, Noble-Haeusslein LJ, Chang M (2016) Selective inhibition of MMP-2 does not alter neurological recovery after spinal cord injury. *ACS Chem Neurosci* 7:1482-1487.
- Goussev S, Hsu JY, Lin Y, Tjoa T, Maida N, Werb Z, Noble-Haeusslein LJ (2003) Differential temporal expression of matrix metalloproteinases after spinal cord injury: relationship to revascularization and wound healing. *J Neurosurg* 99:188-197.
- Hsu JY, McKeon R, Goussev S, Werb Z, Lee JU, Trivedi A, Noble-Haeusslein LJ (2006) Matrix metalloproteinase-2 facilitates wound healing events that promote functional recovery after spinal cord injury. *J Neurosci* 26:9841-9850.
- Hsu JY, Bourguignon LY, Adams CM, Peyrollier K, Zhang H, Fandel T, Cun CL, Werb Z, Noble-Haeusslein LJ (2008) Matrix metalloproteinase-9 facilitates glial scar formation in the injured spinal cord. *J Neurosci* 28:13467-13477.
- Jacobsen JA, Major Jourden JL, Miller MT, Cohen SM (2010) To bind zinc or not to bind zinc: an examination of innovative approaches to improved metalloproteinase inhibition. *Biochim Biophys Acta* 1803:72-94.
- Jones LL, Yamaguchi Y, Stallcup WB, Tuszynski MH (2002) NG2 is a major chondroitin sulfate proteoglycan produced after spinal cord injury and is expressed by macrophages and oligodendrocyte progenitors. *J Neurosci* 22:2792-2803.
- Kawasaki Y, Xu ZZ, Wang X, Park JY, Zhuang ZY, Tan PH, Gao YJ, Roy K, Corfas G, Lo EH, Ji RR (2008) Distinct roles of matrix metalloproteases in the early- and late-phase development of neuropathic pain. *Nat Med* 14:331-336.
- Kobayashi H, Chattopadhyay S, Kato K, Dolkas J, Kikuchi S, Myers RR, Shubayev VI (2008) MMPs initiate Schwann cell-mediated MBP degradation and mechanical nociception after nerve damage. *Mol Cell Neurosci* 39:619-627.
- Lee M, Villegas-Estrada A, Celenza G, Boggess B, Toth M, Kreitinger G, Forbes C, Fridman R, Mobashery S, Chang M (2007) Metabolism of a highly selective gelatinase inhibitor generates active metabolite. *Chem Biol Drug Des* 70:371-382.
- Levine JM, Cohen ND, Heller M, Fajt VR, Levine GJ, Kerwin SC, Trivedi AA, Fandel TM, Werb Z, Modestino A, Noble-Haeusslein LJ (2014) Efficacy of a metalloproteinase inhibitor in spinal cord injured dogs. *PLoS One* 9:e96408.
- Lopez-Otin C, Palavalli LH, Samuels Y (2009) Protective roles of matrix metalloproteinases: from mouse models to human cancer. *Cell Cycle* 8:3657-3662.
- Magnussen H, Watz H, Kirsten A, Wang M, Wray H, Samuelsson V, Mo J, Kay R (2011) Safety and tolerability of an oral MMP-9 and -12 inhibitor, AZD1236, in patients with moderate-to-severe COPD: a randomised controlled 6-week trial. *Pulm Pharmacol Ther* 24:563-570.
- Marshall DC, Lyman SK, McCauley S, Kovalenko M, Spangler R, Liu C, Lee M, O'Sullivan C, Barry-Hamilton V, Ghermazien H, Mikels-Vigdal A, Garcia CA, Jorgensen B, Velayo AC, Wang R, Adamkewicz JI, Smith V (2015) Selective allosteric inhibition of MMP9 is efficacious in preclinical models of ulcerative colitis and colorectal cancer. *PLoS One* 10:e0127063.

- Matziari M, Beau F, Cuniassé P, Dive V, Yiotakis A (2004) Evaluation of P1'-diversified phosphinic peptides leads to the development of highly selective inhibitors of MMP-11. *J Med Chem* 47:325-336.
- Moghaddam A, Heller R, Daniel V, Swing T, Akbar M, Gerner HJ, Biglari B (2017) Exploratory study to suggest the possibility of MMP-8 and MMP-9 serum levels as early markers for remission after traumatic spinal cord injury. *Spinal Cord* 55:8-15.
- Morita-Fujimura Y, Fujimura M, Gasche Y, Copin JC, Chan PH (2000) Overexpression of copper and zinc superoxide dismutase in transgenic mice prevents the induction and activation of matrix metalloproteinases after cold injury-induced brain trauma. *J Cereb Blood Flow Metab* 20:130-138.
- Noble LJ, Donovan F, Igarashi T, Goussev S, Werb Z (2002) Matrix metalloproteinases limit functional recovery after spinal cord injury by modulation of early vascular events. *J Neurosci* 22:7526-7535.
- Overall CM, Kleinfeld O (2006) Tumour microenvironment- opinion: validating matrix metalloproteinases as drug targets and anti-targets for cancer therapy. *Nat Rev Cancer* 6:227-239.
- Paemen L, Martens E, Masure S, Opendakker G (1995) Monoclonal antibodies specific for natural human neutrophil gelatinase B used for affinity purification, quantitation by two-site ELISA and inhibition of enzymatic activity. *Eur J Biochem* 234:759-765.
- Page-McCaw A, Ewald AJ, Werb Z (2007) Matrix metalloproteinases and the regulation of tissue remodelling. *Nat Rev Mol Cell Biol* 8:221-233.
- Pannu R, Christie DK, Barbosa E, Singh I, Singh AK (2007) Post-trauma Lipitor treatment prevents endothelial dysfunction, facilitates neuroprotection, and promotes locomotor recovery following spinal cord injury. *J Neurochem* 101:182-200.
- Peterson JT (2004) Matrix metalloproteinase inhibitor development and the remodeling of drug discovery. *Heart Fail Rev* 9:63-79.
- Peterson JT (2006) The importance of estimating the therapeutic index in the development of matrix metalloproteinase inhibitors. *Cardiovasc Res* 69:677-687.
- Pizzi MA, Crowe MJ (2007) Matrix metalloproteinases and proteoglycans in axonal regeneration. *Exp Neurol* 204:496-511.
- Rajagopalan S, Meng XP, Ramasamy S, Harrison DG, Galis ZS (1996) Reactive oxygen species produced by macrophage-derived foam cells regulate the activity of vascular matrix metalloproteinases in vitro. Implications for atherosclerotic plaque stability. *J Clin Invest* 98:2572-2579.
- Romero-Perez D, Fricovsky E, Yamasaki KG, Griffin M, Barraza-Hidalgo M, Dillmann W, Villarreal F (2008) Cardiac uptake of minocycline and mechanisms for in vivo cardioprotection. *J Am Coll Cardiol* 52:1086-1094.
- Rosenberg GA, Yang Y (2007) Vasogenic edema due to tight junction disruption by matrix metalloproteinases in cerebral ischemia. *Neurosurg Focus* 22:E4.
- Rutges J, Kwon BK, Heran M, Ailon T, Street JT, Dvorak MF (2017) A prospective serial MRI study following acute traumatic cervical spinal cord injury. *Eur Spine J* 26:2324-2332.
- Scali M, Begenisic T, Mainardi M, Milanese M, Bonifacino T, Bonanno G, Sale A, Maffei L (2013) Fluoxetine treatment promotes functional recovery in a rat model of cervical spinal cord injury. *Sci Rep* 3:2217.
- Schafers M, Sorkin L (2008) Effect of cytokines on neuronal excitability. *Neurosci Lett* 437:188-193.
- Sela-Passwell N, Rosenblum G, Shoham T, Sagi I (2010) Structural and functional bases for allosteric control of MMP activities: can it pave the path for selective inhibition? *Biochim Biophys Acta* 1803:29-38.
- Sela-Passwell N, Kikkeri R, Dym O, Rozenberg H, Margalit R, Arad-Yellin R, Eisenstein M, Brenner O, Shoham T, Danon T, Shanzer A, Sagi I (2011) Antibodies targeting the catalytic zinc complex of activated matrix metalloproteinases show therapeutic potential. *Nat Med* 18:143-147.
- Shipley JM, Wesselschmidt RL, Kobayashi DK, Ley TJ, Shapiro SD (1996) Metalloelastase is required for macrophage-mediated proteolysis and matrix invasion in mice. *Proc Natl Acad Sci U S A* 93:3942-3946.
- Siddall PJ, McClelland JM, Rutkowski SB, Cousins MJ (2003) A longitudinal study of the prevalence and characteristics of pain in the first 5 years following spinal cord injury. *Pain* 103:249-257.
- Sternlicht MD, Werb Z (2001) How matrix metalloproteinases regulate cell behavior. *Annu Rev Cell Dev Biol* 17:463-516.
- Sugawara T, Lewen A, Gasche Y, Yu F, Chan PH (2002) Overexpression of SOD1 protects vulnerable motor neurons after spinal cord injury by attenuating mitochondrial cytochrome c release. *FASEB J* 16:1997-1999.
- Teronen O, Heikkilä P, Kontinen YT, Laitinen M, Salo T, Hanemaaijer R, Teronen A, Maisi P, Sorsa T (1999) MMP inhibition and downregulation by bisphosphonates. *Ann N Y Acad Sci* 878:453-465.
- Tong C, Zhengyao Z, Mei L, Dongpo S, Qian H, Fengqun M (2021) Pregabalin and gabapentin in patients with spinal cord injury-related neuropathic pain: a network meta-analysis. *Pain Ther* 10:1497-1509.
- Toth M, Bernardo MM, Gervasi DC, Soloway PD, Wang Z, Bigg HF, Overall CM, DeClerck YA, Tschesche H, Cher ML, Brown S, Mobashery S, Fridman R (2000) Tissue inhibitor of metalloproteinase (TIMP)-2 acts synergistically with synthetic matrix metalloproteinase (MMP) inhibitors but not with TIMP-4 to enhance the (Membrane type 1)-MMP-dependent activation of pro-MMP-2. *J Biol Chem* 275:41415-41423.
- Türk B (2006) Targeting proteases: successes, failures and future prospects. *Nat Rev Drug Discov* 5:785-799.
- Veeravalli KK, Dasari VR, Tsung AJ, Dinh DH, Gujrati M, Fassett D, Rao JS (2009) Human umbilical cord blood stem cells upregulate matrix metalloproteinase-2 in rats after spinal cord injury. *Neurobiol Dis* 36:200-212.
- Wells JE, Rice TK, Nuttall RK, Edwards DR, Zekki H, Rivest S, Yong VW (2003) An adverse role for matrix metalloproteinase 12 after spinal cord injury in mice. *J Neurosci* 23:10107-10115.
- Wells JE, Biernaskie J, Szymanska A, Larsen PH, Yong VW, Corbett D (2005) Matrix metalloproteinase (MMP)-12 expression has a negative impact on sensorimotor function following intracerebral haemorrhage in mice. *Eur J Neurosci* 21:187-196.
- Wilson JR, Cadotte DW, Fehlings MG (2012) Clinical predictors of neurological outcome, functional status, and survival after traumatic spinal cord injury: a systematic review. *J Neurosurg Spine* 17:11-26.
- Xu J, Kim GM, Ahmed SH, Xu J, Yan P, Xu XM, Hsu CY (2001) Glucocorticoid receptor-mediated suppression of activator protein-1 activation and matrix metalloproteinase expression after spinal cord injury. *J Neurosci* 21:92-97.
- Yang Y, Estrada EY, Thompson JF, Liu W, Rosenberg GA (2007) Matrix metalloproteinase-mediated disruption of tight junction proteins in cerebral vessels is reversed by synthetic matrix metalloproteinase inhibitor in focal ischemia in rat. *J Cereb Blood Flow Metab* 27:697-709.
- Yong VW, Power C, Forsyth P, Edwards DR (2001) Metalloproteinases in biology and pathology of the nervous system. *Nat Rev Neurosci* 2:502-511.
- Yong VW (2005) Metalloproteinases: mediators of pathology and regeneration in the CNS. *Nat Rev Neurosci* 6:931-944.
- Yu F, Kamada H, Niizuma K, Endo H, Chan PH (2008) Induction of mmp-9 expression and endothelial injury by oxidative stress after spinal cord injury. *J Neurotrauma* 25:184-195.
- Zhang H, Adwanikar H, Werb Z, Noble-Haeusslein LJ (2010) Matrix metalloproteinases and neurotrauma: evolving roles in injury and reparative processes. *Neuroscientist* 16:156-170.
- Zhang H, Chang M, Hansen CN, Basso DM, Noble-Haeusslein LJ (2011) Role of matrix metalloproteinases and therapeutic benefits of their inhibition in spinal cord injury. *Neurotherapeutics* 8:206-220.
- Zhao BQ, Wang S, Kim HY, Storrie H, Rosen BR, Mooney DJ, Wang X, Lo EH (2006) Role of matrix metalloproteinases in delayed cortical responses after stroke. *Nat Med* 12:441-445.

C-Editors: Zhao M, Liu WJ, Qiu Y; T-Editor: Jia Y