

Brownian Motion in an N-scale periodic Potential

Duncan, A. B.; Duong, Manh Hong; Pavliotis, G. A.

DOI:

[10.1007/s10955-023-03092-9](https://doi.org/10.1007/s10955-023-03092-9)

License:

Creative Commons: Attribution (CC BY)

Document Version

Publisher's PDF, also known as Version of record

Citation for published version (Harvard):

Duncan, AB, Duong, MH & Pavliotis, GA 2023, 'Brownian Motion in an N-scale periodic Potential', *Journal of Statistical Physics*, vol. 190, no. 4, 82 . <https://doi.org/10.1007/s10955-023-03092-9>

[Link to publication on Research at Birmingham portal](#)

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.



Brownian Motion in an N -Scale Periodic Potential

A. B. Duncan¹ · M. H. Duong² · G. A. Pavliotis³

Received: 16 June 2022 / Accepted: 3 March 2023
© The Author(s) 2023

Abstract

We study the problem of Brownian motion in a multiscale potential. The potential is assumed to have $N + 1$ scales (i.e. N small scales and one macroscale) and to depend periodically on all the small scales. We show that for nonseparable potentials, i.e. potentials in which the microscales and the macroscale are fully coupled, the homogenized equation is an overdamped Langevin equation with multiplicative noise driven by the free energy, for which the detailed balance condition still holds. This means, in particular, that homogenized dynamics is reversible and that the coarse-grained Fokker–Planck equation is still a Wasserstein gradient flow with respect to the coarse-grained free energy. The calculation of the effective diffusion tensor requires the solution of a system of N coupled Poisson equations.

Keywords Brownian dynamics · Multiscale analysis · Reiterated homogenization · Reversible diffusions · Free energy

Mathematics Subject Classification 35B27 · 35Q82 · 60H30

1 Introduction

The evolution of complex systems arising in chemistry and biology often involve dynamic phenomena occurring at a wide range of time and length scales. Many such systems are characterised by the presence of a hierarchy of barriers in the underlying energy landscape, giving rise to a complex network of metastable regions in configuration space. Such energy

Communicated by Eric A. Carlen.

✉ M. H. Duong
h.duong@bham.ac.uk
A. B. Duncan
a.duncan@imperial.ac.uk
G. A. Pavliotis
g.pavliotis@imperial.ac.uk

- ¹ Department of Mathematics, Imperial College London, London, UK
- ² School of Mathematics, University of Birmingham, Birmingham, UK
- ³ Department of Mathematics, Imperial College London, London, UK

landscapes occur naturally in macromolecular models of solvated systems, in particular protein dynamics. In such cases the rugged energy landscape is due to the many competing interactions in the energy function [11], giving rise to frustration, in a manner analogous to spin glass models [10, 40]. Although the large scale structure will determine the minimum energy configurations of the system, the small scale fluctuations of the energy landscape will still have a significant influence on the dynamics of the protein, in particular the behaviour at equilibrium, the most likely pathways for binding and folding, as well as the stability of the conformational states. Rugged energy landscapes arise in various other contexts, for example nucleation at a phase transition and solid transport in condensed matter.

To study the influence of small scale potential energy fluctuations on the system dynamics, a number of simple mathematical models have been proposed which capture the essential features of such systems. In one such model, originally proposed by Zwanzig [56], the dynamics are modelled as an overdamped Langevin diffusion in a rugged two-scale potential V^ϵ ,

$$dX_t^\epsilon = -\nabla V^\epsilon(X_t) dt + \sqrt{2\sigma} dW_t, \quad \sigma = \beta^{-1} = k_B T, \quad (1)$$

where T is the temperature and k_B is Boltzmann's constant. The function $V^\epsilon(x) = V(x, x/\epsilon)$ is a smooth potential which has been perturbed by a rapidly fluctuating function with wave number controlled by the small scale parameter $\epsilon > 0$. See Fig. 1 for an illustration. Zwanzig's analysis was based on an effective medium approximation of the mean first passage time, from which the standard Lifson–Jackson formula [33] for the effective diffusion coefficient was recovered. In the context of protein dynamics, phenomenological models based on (1) are widespread in the literature, including but not limited to [3, 28, 37, 53]. Theoretical aspects of such models have also been previously studied. In [13] the authors study diffusion in a strongly correlated quenched random potential constructed from a periodically-extended path of a fractional Brownian motion. Numerical study of the effective diffusivity of diffusion in a potential obtained from a realisation of a stationary isotropic Gaussian random field is performed in [6]. More recent works include [22, 23] where the authors study systems of weakly interacting diffusions moving in a multiwell potential energy landscape, coupled via a Curie–Weiss type (quadratic) interaction potential and [34] in which the authors consider enhanced diffusion for Brownian motion in a tilted periodic potential expressing the effective diffusion in terms of the eigenvalue band structure. It is also worth mentioning a series of works [4, 19, 48, 54] studying multiscale behaviour of diffusion processes with multiple-well potentials in which the limiting process is a chemical reactions instead of a diffusion. We also mention [14], where the combined mean field/homogenization limit for diffusions interacting via a periodic potential is considered. The main result of this paper is that, in the presence of phase transitions, the mean field and homogenization limits do not commute.

For the case where (1) possesses one characteristic lengthscale controlled by $\epsilon > 0$, the convergence of X_t^ϵ to a coarse-grained process X_t^0 in the limit $\epsilon \rightarrow 0$ over a finite time interval is well-known. When the rapid oscillations are periodic, under a diffusive rescaling this problem can be recast as a periodic homogenization problem, for which it can be shown that the process X_t^ϵ converges weakly to a Brownian motion with constant effective diffusion tensor D (covariance matrix) which can be calculated by solving an appropriate Poisson equation posed on the unit torus, see for example [8, 46]. The analogous case where the rapid fluctuations arise from a stationary ergodic random field has been studied in [31, Chap. 9]. The case where the potential V^ϵ possesses periodic fluctuations with two or three well-separated characteristic timescales, i.e. $V^\epsilon(x) = V(x, x/\epsilon, x/\epsilon^2)$ follow from the results in [8, Chap. 3.7], in which case the dynamics of the coarse-grained model in the $\epsilon \rightarrow 0$ limit are characterised by an Itô SDE whose coefficients can be calculated in terms of the solution




Fig. 1 Example of a multiscale potential. The left panel shows the isolines of the Mueller potential [39, 49]. The right panel shows the corresponding rugged energy landscape where the Mueller potential is perturbed by high frequency periodic fluctuations

of an associated Poisson equation. A generalization of these results to diffusion processes having N -well separated scales was explored in Sect. 3.11.3 of the same text, but no proof of convergence is offered in this case. Similar diffusion approximations for systems with one fast scale and one slow scale, where the fast dynamics is not periodic have been studied in [43].

A model for Brownian dynamics in a potential V possessing infinitely many characteristic lengthscales was studied in [7]. In particular, the authors studied the large-scale diffusive behaviour of the overdamped Langevin dynamics in potentials of the form

$$V^n(x) = \sum_{k=0}^n U_k \left(\frac{x}{R_k} \right), \quad (2)$$

obtained as a superposition of Hölder continuous periodic functions with period 1. It was shown in [7] that the effective diffusion coefficient decays exponentially fast with the number of scales, provided that the scale ratios R_{k+1}/R_k are bounded from above and below, which includes cases where there is no scale separation. From this the authors were able to show that the effective dynamics exhibits subdiffusive behaviour, in the limit of infinitely many scales. See also the analytical calculation presented in [15] for a piecewise linear periodic potential; in the limit of infinitely many scales, the homogenized diffusion coefficient converges to zero, signaling that, in this limit, the coarse-grained dynamics is characterized by subdiffusive behaviour.

In this paper we study the dynamics of diffusion in a rugged potential possessing N well-separated lengthscales. More specifically, we study the dynamics of (1) where the multiscale potential is chosen to have the form

$$V^\epsilon(x) = V\left(x, x/\epsilon, x/\epsilon^2, \dots, x/\epsilon^N\right), \tag{3}$$

where V is a smooth function, which is periodic with period 1 in all but the first argument. Clearly, V can always be written in the form

$$V(x_0, x_1, \dots, x_N) = V_0(x_0) + V_1(x_0, x_1, \dots, x_N), \tag{4}$$

where $(x_0, x_1, \dots, x_N) \in \mathbb{R}^d \times (\mathbb{T}^d)^N$. We will assume that the large scale component of the potential V_0 is smooth and confining in \mathbb{R}^d , and that the perturbation V_1 is a smooth bounded function which is periodic in all but the first variable. Unlike [7], we work under the assumption of explicit scale separation, however we also permit more general potentials than those of the form (2), allowing possibly nonlinear interactions between the different scales, and even full coupling between scales.¹ To emphasize the fact that the potential (4) leads to a fully coupled system across scales, we introduce the auxiliary processes $X_t^{(j)} = X_t/\epsilon^j$, $j = 0, \dots, N$. The SDE (1) can then be written as a fully coupled system of SDEs driven by the same Brownian motion W_t ,

$$dX_t^{(0)} = - \sum_{i=0}^N \epsilon^{-i} \nabla_{x_i} V\left(X_t^{(0)}, X_t^{(1)}, \dots, X_t^{(N)}\right) dt + \sqrt{2\sigma} dW_t, \tag{5a}$$

$$dX_t^{(1)} = - \sum_{i=0}^N \epsilon^{-(i+1)} \nabla_{x_i} V\left(X_t^{(0)}, X_t^{(1)}, \dots, X_t^{(N)}\right) dt + \sqrt{\frac{2\sigma}{\epsilon^2}} dW_t, \tag{5b}$$

⋮

$$dX_t^{(N)} = - \sum_{i=0}^N \epsilon^{-(i+N)} \nabla_{x_i} V\left(X_t^{(0)}, X_t^{(1)}, \dots, X_t^{(N)}\right) dt + \sqrt{\frac{2\sigma}{\epsilon^{2N}}} dW_t, \tag{5c}$$

in which case $X_t^{(0)}$ is considered to be a ‘‘slow’’ variable, while $X_t^{(1)}, \dots, X_t^{(N)}$ are ‘‘fast’’ variables. In this paper, we first provide an explicit proof of the convergence of the solution of (1), X_t^ϵ to a coarse-grained (homogenized) diffusion process X_t^0 given by the unique solution of the following Itô SDE:

$$dX_t^0 = -\mathcal{M}(X_t^0) \nabla \Psi(X_t^0) dt + \sigma \nabla \cdot \mathcal{M}(X_t^0) dt + \sqrt{2\sigma \mathcal{M}(X_t^0)} dW_t, \tag{6}$$

where

$$\Psi(x) = -\sigma \log Z(x),$$

denotes the free energy, for

$$Z(x) = \int_{\mathbb{T}^d} \dots \int_{\mathbb{T}^d} e^{-V_1(x, y_1, \dots, y_N)/\sigma} dy_1 \dots dy_N,$$

and where $\mathcal{M}(x)$ is a symmetric uniformly positive definite tensor which is independent of ϵ . The formula of the effective diffusion tensor is given in Sect. 2.

¹ We will refer to potentials of the form $V^\epsilon(x) = V_0(x) + V_1(x/\epsilon, \dots, x/\epsilon^N)$ where V_1 is periodic in all variables as separable.

Our assumptions on the potential V^ϵ in (4) guarantee that the full dynamics (1) is reversible with respect to the Gibbs measure μ^ϵ by construction. It is important to note that the coarse-grained dynamics (6) is also reversible with respect to the equilibrium Gibbs measure

$$\mu^0(x) = Z(x)/\bar{Z}.$$

Indeed, the natural interpretation of $\Psi(x) = -\sigma \log Z(x)$ is as the free energy corresponding to the coarse-grained variable X_t^0 . The weak convergence of X_t^ϵ to X_t^0 implies in particular that the distribution of X_t^ϵ will converge weakly to that of X_t^0 , uniformly over finite time intervals $[0, T]$, which does not say anything about the convergence of the respective stationary distributions μ^ϵ to μ^0 . In Sect. 4 we study the equilibrium behaviour of X_t^ϵ and X_t^0 and show that the long-time limit $t \rightarrow \infty$ and the coarse-graining limit $\epsilon \rightarrow 0$ commute, and in particular that the equilibrium measure μ^ϵ of X_t^ϵ converges in the weak sense to μ^0 . We also study the rate of convergence to equilibrium for both processes, and we obtain bounds relating the two rates. This question is naturally related to the study of the Poincaré constants for the full and coarse-grained potentials [24, 41].

We can summarize the above discussion as follows: the (Wasserstein) gradient structure, reversibility and detailed balance property of the dynamics (the three properties are equivalent) are preserved under the homogenization/coarse-graining process: the reversibility of X_t^ϵ with respect to μ^ϵ is preserved under the homogenization procedure. Indeed, any general diffusion process that is reversible with respect to $\mu^0(x)$ will have the form (18), see [45, Sect. 4.7]. It is not necessarily always the case that the gradient structure is preserved under coarse-graining, as has been shown recently [47]. The creation of non-gradient/nonreversible effects due to the multiscale structure of the dynamics is a very interesting problem that we will return to in future work.

We also remark that the homogenized SDE corresponds to the kinetic/Klimontovich interpretation of the stochastic integral [27], i.e. it can be written in the form

$$dX_t^0 = -\mathcal{M}(X_t^0)\nabla\Psi(X_t^0)dt + \sqrt{2\sigma\mathcal{M}(X_t^0)} \circ^{\text{Klim}} dW_t, \quad (7)$$

where we use the notation \circ^{Klim} to denote the Klimontovich stochastic differential/integral. The Klimontovich interpretation of the stochastic integral leads to a thermodynamically consistent Langevin dynamics, in the sense that it is reversible with respect to the coarse-grained Gibbs measure.

The multiplicative noise is due to the full coupling between the macroscopic and the N microscopic scales.² For one-dimensional potentials, we are able to obtain an explicit expression for $\mathcal{M}(x)$, regardless of the number of scales involved. In higher dimensions, $\mathcal{M}(x)$ will be expressed in terms of the solution of a recursive family of Poisson equations which can be solved only numerically. We also obtain a variational characterization of the effective diffusion tensor, analogous to the standard variational characterisations for the effective conductivity tensor for multiscale conductivity problems, see for example [29]. Using this variational characterisation, we are able to derive tight bounds on the effective diffusion tensor, and in particular, show that as $N \rightarrow \infty$, the eigenvalues of the effective diffusion tensor will converge to zero, suggesting that diffusion in potentials with infinitely many scales will exhibit anomalous diffusion. The focus of this paper is the rigorous analysis of the homogenization problem for (1) with V^ϵ given by (4). More precisely, we are interested in establishing the convergence of both the dynamics (over finite time domain) and of the equilibrium measure of (1) as ϵ tends to zero.

² For additive potentials of the form (2), i.e. when there is no interaction between the macroscale and the microscales, the noise in the homogenized equation is additive.

Our proof of the homogenization theorem, Theorem 1 is based on the well known martingale approach to proving limit theorems [8, 42, 43]. The main technical difficulty in applying such well known techniques is the construction of the corrector field/compensator and the analysis of the obtained Poisson equations. This turns out to be a challenging task, since we consider the case where all scales, the macroscale and the N -microscales, are fully coupled. For recent applications of the techniques, we refer the reader to [32, 50] where the authors study metastable behaviour of multiscale diffusion processes.

The rest of the paper is organized as follows. In Sect. 2 we state the assumptions on the structure of the multiscale potential and state the main results of this paper. In Sect. 3 we study properties of the effective dynamics, providing expressions for the diffusion tensor in terms of a variational formula, and derive various bounds. In Sect. 4 we study properties of the effective potential, and prove convergence of the equilibrium distribution of X_t^ϵ to the coarse-grained equilibrium distribution μ^0 . The proof of the main theorem, Theorem 1, is presented in Sect. 5. Finally, in Sect. 6 we provide further discussion and outlook.

2 Setup and Statement of Main Results

In this section we provide conditions on the multiscale potential which are required to obtain a well-defined homogenization limit. In particular, we shall highlight assumptions necessary for the ergodicity of the full model as well as the coarse-grained dynamics.

We will consider the overdamped Langevin dynamics

$$dX_t^\epsilon = -\nabla V^\epsilon(X_t^\epsilon) dt + \sqrt{2\sigma} dW_t, \tag{8}$$

where $V^\epsilon(x)$ is of the form (3). The multiscale potentials we consider in this paper can be viewed as a smooth confining potential perturbed by smooth, bounded fluctuations which become increasingly rapid as $\epsilon \rightarrow 0$, see Fig. 1 for an illustration. More specifically, we will assume that the multiscale potential V satisfies the following assumptions.³

Assumption 1 The potential V is given by

$$V(x_0, x_1, \dots, x_N) = V_0(x_0) + V_1(x_0, x_1, \dots, x_N), \tag{9}$$

where $(x_0, x_1, \dots, x_N) \in \mathbb{R}^d \times (\mathbb{T}^d)^N$, and

1. V_0 is a smooth confining potential, i.e. $e^{-V_0(x)} \in L^1(\mathbb{R}^d)$ and $V_0(x) \rightarrow \infty$ as $|x| \rightarrow \infty$.
2. The perturbation $V_1(x_0, x_1, \dots, x_N)$ is smooth and bounded uniformly in x_0 .
3. There exists $C > 0$ such that $\|\nabla^2 V_0\|_{L^\infty(\mathbb{R}^d)} \leq C$.

Remark 1 We note that Assumption 1 is quite stringent, since it implies that V_0 is quadratic to leading order. This assumption is also made in [43]. In cases where the process $X_0^\epsilon \sim \mu^\epsilon$, i.e. the process is stationary, this condition can be relaxed considerably.

The infinitesimal generator \mathcal{L}^ϵ of X_t^ϵ is the selfadjoint extension of

$$\mathcal{L}^\epsilon f(x) = -\nabla V^\epsilon(x) \cdot \nabla f(x) + \sigma \Delta f(x), \quad f \in C_c^\infty(\mathbb{R}^d). \tag{10}$$

It follows from the assumption on V_0 that the corresponding overdamped Langevin equation

$$dY_t = -\nabla V_0(Y_t) dt + \sqrt{2\sigma} dW_t, \tag{11}$$

³ We remark that we can always write (4) in the form (9) where $V_0(x) = \int_{\mathbb{T}^d} \dots \int_{\mathbb{T}^d} V(x, x_1, \dots, x_N) dx_1 \dots dx_N$.

is ergodic with the unique stationary distribution

$$\mu_{ref}(x) = \frac{1}{Z_{ref}} \exp(-V_0(x)/\sigma), \quad Z_{ref} = \int_{\mathbb{R}^d} e^{-V_0(x)/\sigma} dx.$$

Since V_1 is bounded uniformly, by Assumption 1, it follows that the potential V^ϵ is also confining, and therefore X_t^ϵ is ergodic, possessing a unique invariant distribution given by $\mu^\epsilon(x) = \frac{e^{-V^\epsilon(x)/\sigma}}{Z^\epsilon}$, where $Z^\epsilon = \int_{\mathbb{R}^d} e^{-V^\epsilon(x)/\sigma}$. Moreover, noting that the generator \mathcal{L}^ϵ of X_t^ϵ can be written as

$$\mathcal{L}^\epsilon f(x) = \sigma e^{V^\epsilon(x)/\sigma} \nabla \cdot \left(e^{-V^\epsilon(x)/\sigma} \nabla f(x) \right), \quad f \in C_c^2(\mathbb{R}^d).$$

It follows that μ^ϵ is reversible with respect to the dynamics X_t^ϵ , c.f. [20, 45].

Our main objective in this paper is to study the dynamics (8) in the limit of infinite scale separation $\epsilon \rightarrow 0$. Having introduced the model and the assumptions we can now present the main result of the paper.

Theorem 1 (Weak convergence of X_t^ϵ to X_t^0) *Suppose that Assumption 1 holds and let $T > 0$, and the initial condition X_0 is distributed according to some probability distribution ν on \mathbb{R}^d . Then as $\epsilon \rightarrow 0$, the process X_t^ϵ converges weakly in $(C[0, T]; \mathbb{R}^d)$ to the diffusion process X_t^0 with generator defined by*

$$\mathcal{L}^0 f(x) = \frac{\sigma}{Z(x)} \nabla_x \cdot (Z(x) \mathcal{M}(x) \nabla_x f(x)), \quad f \in C_c^2(\mathbb{R}^d), \tag{12}$$

and where

$$Z(x) = \int_{\mathbb{T}^d} \dots \int_{\mathbb{T}^d} e^{-V_1(x, x_1, \dots, x_N)/\sigma} dx_N \dots dx_1, \tag{13}$$

and

$$\mathcal{M}(x) = \frac{1}{Z(x)} \int_{\mathbb{T}^d} \dots \int_{\mathbb{T}^d} (1 + \nabla_{x_N} \theta_N) \dots (1 + \nabla_{x_1} \theta_1) e^{-V_1(x, x_1, \dots, x_N)/\sigma} dx_N \dots dx_1. \tag{14}$$

The correctors are defined recursively as follows: define $\theta_{N-k} = \left(\theta_{N-k}^1, \dots, \theta_{N-k}^d \right)$ to be the weak vector-valued solution of the PDE

$$\nabla_{x_{N-k}} \cdot (\mathcal{K}_{N-k}(x_0, \dots, x_{N-k}) (\nabla_{x_{N-k}} \theta_{N-k}(x_0, \dots, x_{N-k}) + I)) = 0, \tag{15}$$

where $\theta_{N-k}(x_0, \dots, x_{N-k-1}, \cdot) \in H^1(\mathbb{T}^d; \mathbb{R}^d)$, with the notation $[\nabla_{x_n} \theta_n]_{\cdot, j} = \nabla_{x_n} \theta_n^j$, for $j = 1, \dots, d$ and $n = 1, \dots, N$ and where

$$\begin{aligned} &\mathcal{K}_{N-k}(x_0, \dots, x_{N-k}) \\ &= \int_{\mathbb{T}^d} \dots \int_{\mathbb{T}^d} (I + \nabla_{x_N} \theta_N) \dots (I + \nabla_{x_{N-k+1}} \theta_{N-k+1}) e^{-V_1/\sigma} dx_N \dots dx_{N-k+1}, \end{aligned} \tag{16}$$

for $k = 1, \dots, N - 1$, and

$$\mathcal{K}_N(x, x_1, \dots, x_N) = e^{-V_1(x, x_1, \dots, x_N)/\sigma} I, \tag{17}$$

where I denotes the identity matrix in $\mathbb{R}^{d \times d}$. Provided that Assumption 1 hold, Proposition 5 guarantees the existence and uniqueness (up to a constant) of solutions to the coupled Poisson

equations (15). Furthermore, the solutions will depend smoothly on the slow variable x_0 as well as the fast variables x_1, \dots, x_N . The process X_t^0 is the unique solution to the Itô SDE

$$dX_t^0 = -\mathcal{M}(X_t^0)\nabla\Psi(X_t^0) dt + \sigma\nabla \cdot \mathcal{M}(X_t^0) dt + \sqrt{2\sigma\mathcal{M}(X_t^0)} dW_t, \tag{18}$$

where

$$\Psi(x) = -\sigma \log Z(x) = -\sigma \log \left(\int_{\mathbb{T}^d} \dots \int_{\mathbb{T}^d} e^{-V_1(x, y_1, \dots, y_N)/\sigma} dy_1 \dots dy_N \right).$$

The proof, which closely follows that of [43] is postponed to Sect. 5. Theorem 1 confirms the intuition that the coarse-grained dynamics is driven by the coarse-grained free energy. On the other hand, the corresponding SDE has multiplicative noise given by a space dependent diffusion tensor $\mathcal{M}(x)$. We can show that the homogenized process (18) is ergodic with unique invariant distribution

$$\mu^0(x) = \frac{Z(x)}{\bar{Z}} = \frac{1}{\bar{Z}} e^{-\Psi(x)/\sigma}, \quad \text{where } \bar{Z} = \int_{\mathbb{R}^d} Z(x) dx.$$

Other qualitative properties of the solution to the homogenized equation (6), including noise-induced transitions and noise-induced hysteresis behaviour has been studied in [15]. It is also important to note that the reversibility of X_t^ϵ with respect to μ^ϵ is preserved under the homogenization procedure. Indeed, any general diffusion process that is reversible with respect to $\mu^0(x)$ will have the form (18), see [45, Sect. 4.7]. See Sect. 6 for further discussion on this point.

As is characteristic with homogenization problems, when $d = 1$ we can obtain, up to quadratures, an explicit expression for the homogenized SDE. In this case, we obtain explicit expressions for the correctors $\theta_1, \dots, \theta_N$, so that the intermediary coefficients $\mathcal{K}_1, \dots, \mathcal{K}_N$ can be expressed as (see also [15])

$$\mathcal{K}_i(x_0, x_1, \dots, x_i) = \left(\int e^{V_1(x_0, x_1, \dots, x_i, x_{i+1}, \dots, x_N)/\sigma} dx_{i+1} \dots dx_N \right)^{-1}, \quad i = 1, \dots, N.$$

Thus we obtain the following result.

Proposition 1 (Effective Dynamics in one dimension) *When $d = 1$, the effective diffusion coefficient $\mathcal{M}(x)$ in (18) is given by*

$$\mathcal{M}(x) = \frac{1}{Z_1(x)\widehat{Z}_1(x)}, \tag{19}$$

where

$$Z_1(x) = \int \dots \int e^{-V_1(x, x_1, \dots, x_N)/\sigma} dx_1 \dots dx_N,$$

and

$$\widehat{Z}_1(x) = \int \dots \int e^{V_1(x, x_1, \dots, x_N)/\sigma} dx_1 \dots dx_N.$$

Equation (19) generalises the expression for the effective diffusion coefficient for a two-scale potential that was derived in [56] without any appeal to homogenization theory. In higher dimensions we will not be able to obtain an explicit expression for $\mathcal{M}(x)$, however we are able to obtain bounds on the eigenvalues of $\mathcal{M}(x)$. In particular, we are able to show that (19) acts as a lower bound for the eigenvalues of $\mathcal{M}(x)$.

Proposition 2 *The effective diffusion tensor \mathcal{M} is uniformly positive definite over \mathbb{R}^d . In particular,*

$$0 < e^{-osc(V_1)/\sigma} \leq \frac{1}{Z_1(x)\widehat{Z}_1(x)} \leq e \cdot \mathcal{M}(x)e \leq 1, \quad x \in \mathbb{R}^d, \tag{20}$$

for all $e \in \mathbb{R}^d$ such that $|e| = 1$, where

$$osc(V_1) = \sup_{\substack{x \in \mathbb{R}^d, \\ y_1, \dots, y_N \in \mathbb{T}^d}} V_1(x, y_1, \dots, y_N) - \inf_{\substack{x \in \mathbb{R}^d, \\ y_1, \dots, y_N \in \mathbb{T}^d}} V_1(x, y_1, \dots, y_N).$$

This result follows immediately from Lemmas 1 and 2 which are proved in Sect. 3.

Remark 2 The bounds in (20) highlight the two extreme possibilities for fluctuations occurring in the potential V^ϵ . The equality $\frac{1}{Z_1(x)\widehat{Z}_1(x)} = e \cdot \mathcal{M}(x)e$ is attained when the multiscale fluctuations $V_1(x_0, \dots, x_N)$ are constant in all but one dimension (e.g. the analogue of a layered composite material, [12, Sect. 5.4], [46, Sect. 12.6.2]). In the other extreme, the inequality $e \cdot \mathcal{M}(x)e = 1$ is attained in the absence of fluctuations, i.e. when $V_1 = 0$.

Remark 3 Clearly, the lower bound in (20) becomes exponentially small in the limit as $\sigma \rightarrow 0$.

While Theorem 1 guarantees weak convergence of X_t^ϵ to X_t^0 in $C([0, T]; \mathbb{R}^d)$ for fixed T , it makes no claims regarding the convergence at infinity, i.e. of μ^ϵ to μ^0 . However, under the conditions of Assumption 1 we can show that μ^ϵ converges weakly to μ^0 , so that the $T \rightarrow \infty$ and $\epsilon \rightarrow 0$ limits commute, in the sense that:

$$\lim_{\epsilon \rightarrow 0} \lim_{T \rightarrow \infty} \mathbb{E}[f(X_T^\epsilon)] = \lim_{T \rightarrow \infty} \lim_{\epsilon \rightarrow 0} \mathbb{E}[f(X_T^\epsilon)],$$

for all $f \in L^2(\mu_{ref})$.

Proposition 3 (Weak convergence of μ^ϵ to μ^0) *Suppose that Assumption 1 holds. Then for all $f \in L^2(\mu_{ref})$,*

$$\int_{\mathbb{R}^d} f(x) \mu^\epsilon(dx) \rightarrow \int_{\mathbb{R}^d} f(x) \mu^0(dx), \tag{21}$$

as $\epsilon \rightarrow 0$.

If Assumption 1 holds, then for every $\epsilon > 0$, the potential V^ϵ is confining, so that the process X_t^ϵ is ergodic. If the ‘‘unperturbed’’ process defined by (11) converges to equilibrium exponentially fast in $L^2(\mu_{ref})$, then so will X_t^ϵ and X_t^0 . Moreover, we can relate the rates of convergence of the three processes. We will use the notation $\text{Var}_\mu(f) = \mathbb{E}_\mu(f - \mathbb{E}_\mu f)^2$ to denote the variance with respect to a measure μ .

Proposition 4 *Suppose that Assumption 1 holds and let P_t be the semigroup associated with the dynamics (11) and suppose that $\mu_{ref}(x) = \frac{1}{Z_0} e^{-V_0(x)/\sigma}$ satisfies Poincaré’s inequality with constant ρ/σ , i.e.*

$$\text{Var}_{\mu_{ref}}(f) \leq \frac{\sigma}{\rho} \int |\nabla f(x)|^2 \mu_{ref}(dx), \quad f \in H^1(\mu_{ref}), \tag{22}$$

or equivalently⁴

$$\text{Var}_{\mu_{ref}}(P_t f) \leq e^{-2\rho t/\sigma} \text{Var}_{\mu_{ref}}(f), \quad f \in L^2(\mu_{ref}), \tag{23}$$

⁴ The equivalence between (22) and (23) follows since P_t is a reversible Markov semigroup with respect to the measure μ_{ref} . See [5].

for all $t \geq 0$. Let P_t^ϵ and P_t^0 denote the semigroups associated with the full dynamics (8) and homogenized dynamics (18), respectively. Then for all $f \in L^2(\mu_{ref})$,

$$\text{Var}_{\mu^\epsilon}(P_t^\epsilon f) \leq e^{-2\gamma t/\sigma} \text{Var}_{\mu^\epsilon}(f), \tag{24}$$

and

$$\text{Var}_{\mu^0}(P_t^0 f) \leq e^{-2\tilde{\gamma} t/\sigma} \text{Var}_{\mu^0}(f). \tag{25}$$

for $\gamma = \rho e^{-osc(V_1)/\sigma}$ and $\tilde{\gamma} = \rho e^{-2osc(V_1)/\sigma}$.

The proof of Propositions 3 and 4 can be found in Sect. 4.

3 Properties of the Coarse-Grained Process

In this section we study the properties of the coefficients of the homogenized SDE (18) and its dynamics.

3.1 Separable Potentials

Consider the special case where the potential V^ϵ is *separable*, in the sense that the fast scale fluctuations do not depend on the slow scale variable, i.e.

$$V(x_0, x_1, \dots, x_N) = V_0(x_0) + V_1(x_1, x_2, \dots, x_N).$$

Then, it is clear from the construction of the effective diffusion tensor (14) that $\mathcal{M}(x)$ will not depend on $x \in \mathbb{R}^d$. Moreover, since

$$Z(x) = \int_{\mathbb{T}^d} \cdots \int_{\mathbb{T}^d} e^{-\frac{V_0(x) + V_1(y_1, \dots, y_N)}{\sigma}} dy_1 \dots dy_N = \frac{1}{K} e^{-V_0(x)/\sigma},$$

where $K = \int_{\mathbb{T}^d} \cdots \int_{\mathbb{T}^d} \exp(-V_1(y_1, \dots, y_N)/\sigma) dy_1 \dots dy_N$, then it follows that the coarse-grained stationary distribution μ^0 equals the stationary distribution $\mu_{ref} \propto \exp(-V_0(x)/\sigma)$ of the process (11). For general multiscale potentials however, μ^0 will be different from μ_{ref} . Indeed, introducing multiscale fluctuations can dramatically alter the qualitative equilibrium behaviour of the process, including noise-induced transitions and noise induced hysteresis, as has been studied for various examples in [15].

3.2 Variational Bounds on $\mathcal{M}(x)$

A first essential property is that the constructed matrices $\mathcal{K}_N, \dots, \mathcal{K}_1$ are positive definite over all parameters. For convenience, we shall introduce the following notation

$$\mathbb{X}_k = \mathbb{R}^d \times \times_{i=1}^k \mathbb{T}^d, \tag{26}$$

for $k = 1, \dots, N$, and set $\mathbb{X}_0 = \mathbb{R}^d$ for consistency. First we require the following existence and regularity result for a uniformly elliptic Poisson equation on \mathbb{T}^d .

Lemma 1 For $k = 1, \dots, N$, for x_0, \dots, x_{k-1} fixed, the tensor $\mathcal{K}_k(x_0, \dots, x_{k-1}, \cdot)$ is uniformly positive definite and in particular satisfies, for all unit vectors $e \in \mathbb{R}^d$,

$$\frac{1}{\widehat{Z}_k(x_0, x_1, \dots, x_{k-1})} \leq e \cdot \mathcal{K}_k(x_0, x_1, \dots, x_{k-1}, x_k) e, \quad x_k \in \mathbb{T}^d, \tag{27}$$

where

$$\widehat{Z}_k(x_0, x_1, \dots, x_{k-1}) = \int \dots \int e^{V(x_0, x_1, \dots, x_{k-1}, x_k, \dots, x_N)/\sigma} dx_N dx_{N-1} \dots dx_k,$$

which is independent of x_k .

Proof We prove the result by induction on k starting from $k = N$. For $k = N$ the tensor \mathcal{K}_N is clearly uniformly positive definite for fixed $x_0, \dots, x_{N-1} \in \mathbb{X}_{N-1}$. By [8, Thms III.3.2, III.3.3] there exists a unique (up to a constant) solution such that $\theta_N(x, x_1, \dots, x_{N-1}, \cdot) \in H^2(\mathbb{T}^d; \mathbb{R}^d)$ of (15). In particular,

$$\int_{\mathbb{T}^d} |\nabla_{x_N} \theta_N(x_0, x_1, \dots, x_{N-1}, x_N)|_F^2 dx_N < \infty,$$

where $|\cdot|_F$ denotes the Frobenius norm, so that \mathcal{K}_{N-1} is well defined. Fix $(x_0, \dots, x_{N-2}) \in \mathbb{X}_{N-2}$. To show that $\mathcal{K}_{N-1}(x_0, \dots, x_{N-2}, \cdot)$ is uniformly positive definite on \mathbb{T}^d we first note that

$$\begin{aligned} & \int_{\mathbb{T}^d} (I + \nabla_{x_N} \theta_N)^\top (I + \nabla_{x_N} \theta_N) e^{-V/\sigma} dx_N \\ &= \int_{\mathbb{T}^d} \left(I + \nabla_{x_N} \theta_N + \nabla_{x_N} \theta_N^\top + \nabla_{x_N} \theta_N^\top \nabla_{x_N} \theta_N \right) e^{-V/\sigma} dx_N, \end{aligned} \tag{28}$$

where $V = V(x_0, x_1, \dots, x_N)$ and \top denotes the transpose. From the Poisson equation for θ_N we have

$$\int \theta_N \otimes \nabla_{x_N}^\top (e^{-V/\sigma} (\nabla_{x_N} \theta_N + I)) dx_N = \mathbf{0},$$

from which we obtain, after integrating by parts:

$$\int_{\mathbb{T}^d} \nabla_{x_N} \theta_N^\top (\nabla_{x_N} \theta_N + I) e^{-V/\sigma} dx_N = 0. \tag{29}$$

From (28) and (29) we deduce that

$$\begin{aligned} \mathcal{K}_{N-1} &= \int_{\mathbb{T}^d} (I + \nabla_{x_N} \theta_N) e^{-V/\sigma} dx_N \\ &= \int_{\mathbb{T}^d} \left[I + \nabla_{x_N} \theta_N + \nabla_{x_N} \theta_N^\top (\nabla_{x_N} \theta_N + I) \right] e^{-V/\sigma} dx_N \\ &= \int_{\mathbb{T}^d} (I + \nabla_{x_N} \theta_N)^\top (I + \nabla_{x_N} \theta_N) e^{-V/\sigma} dx_N. \end{aligned}$$

Thus \mathcal{K}_{N-1} is well-defined and symmetric. We note that

$$\int_{\mathbb{T}^d} (I + \nabla_{x_N} \theta_N) dx_N = I,$$

therefore, it follows by Hölder’s inequality that

$$|v|^2 = \left| v^\top \int_{\mathbb{T}^d} (I + \nabla_N \theta_N) dx_N \right|^2 \leq v^\top (\mathcal{K}_{N-1}) v \left(\int_{\mathbb{T}^d} e^{V/\sigma} dx_N \right),$$

so that

$$\frac{|v|^2}{\widehat{Z}_N(x_0, \dots, x_{N-1})} \leq v^\top \mathcal{K}_{N-1}(x_0, \dots, x_{N-1}) v, \quad \forall (x_0, x_1, \dots, x_{N-1}).$$

Since \widehat{Z}_N is uniformly bounded for (x_0, \dots, x_{N-1}) it follows $\mathcal{K}_{N-1}(x_0, \dots, x_{N-2}, \cdot)$ is uniformly positive definite, and arguing as above we establish existence of a unique θ_{N-1} , up to a constant, solving (15) for $k = 2$.

Now, assume that the corrector θ_{N-k+1} has been constructed, and so \mathcal{K}_{N-k+1} is well defined. By multiplying the cell equation for θ_{N-k+1}

$$\nabla_{x_{N-k+1}} \cdot \left[\mathcal{K}_{N-k+1}(\nabla_{x_{N-k+1}} \theta_{N-k+1} + I) \right] = 0,$$

by θ_{N-k+1} then integrating with respect to x_{N-k+1} and using integration by parts as well as the symmetry of \mathcal{K}_{N-k+1} from the inductive hypothesis we obtain

$$\int \nabla_{x_{N-k+1}} \theta_{N-k+1}^\top \mathcal{K}_{N-k+1} (I + \nabla_{x_{N-k+1}} \theta_{N-k+1}) dx_{N-k+1} = \mathbf{0}.$$

Therefore, we have

$$\begin{aligned} \mathcal{K}_{N-k} &= \int_{\mathbb{T}^d} \mathcal{K}_{N-k+1}(I + \nabla_{x_{N-k+1}} \theta_{N-k+1}) dx_{N-k+1} \\ &= \int_{\mathbb{T}^d} \left[\mathcal{K}_{N-k+1}(I + \nabla_{x_{N-k+1}} \theta_{N-k+1}) + \nabla_{x_{N-k+1}} \theta_{N-k+1}^\top \mathcal{K}_{N-k+1} \right. \\ &\quad \left. \times (I + \nabla_{x_{N-k+1}} \theta_{N-k+1}) \right] dx_{N-k+1} \\ &= \int_{\mathbb{T}^d} (I + \nabla_{x_{N-k+1}} \theta_{N-k+1})^\top \mathcal{K}_{N-k+1} (I + \nabla_{x_{N-k+1}} \theta_{N-k+1}) dx_{N-k+1}. \end{aligned}$$

Thus \mathcal{K}_{N-k} is also well-defined and symmetric. To show (27) we note that

$$\int \dots \int (I + \nabla_{x_N} \theta_N) \dots (I + \nabla_{x_{N-k}} \theta_{N-k}) dx_N \dots dx_{N-k} = I.$$

Therefore, for any vector $v \in \mathbb{R}^d$:

$$\begin{aligned} |v|^2 &= \left| v^\top \left(\int \dots \int (I + \nabla_{x_N} \theta_N) \dots (I + \nabla_{x_{N-k}} \theta_{N-k}) dx_N \dots dx_{N-k} \right) \right|^2 \\ &\leq v^\top \left(\int \dots \int (I + \nabla_{x_{N-k}} \theta_{N-k})^\top \dots (I + \nabla_{x_{N-k}} \theta_{N-k}) e^{-V/\sigma} dx_N \dots dx_{N-k} \right) \\ &\quad \times v \int e^{V/\sigma} dx_N \dots dx_{N-k} \\ &= \left(v^\top \mathcal{K}_{N-k}(x_1, \dots, x_{N-k}) v \right) \widehat{Z}(x_1, \dots, x_{N-k}). \end{aligned}$$

The fact that we have strict positivity then follows immediately. □

To obtain upper bounds for the effective diffusion coefficient, we will express the intermediary diffusion tensors \mathcal{K}_i as solutions of a quadratic variational problem. This variational formulation of the diffusion tensors can be considered as a generalisation of the analogous representation for the effective conductivity coefficient of a two-scale composite material, see for example [8, 29, 36].

Lemma 2 For $i = 1, \dots, N$, the tensor \mathcal{K}_i satisfies

$$\begin{aligned}
 e \cdot \mathcal{K}_i(x_0, \dots, x_i)e &= \inf_{\substack{v_{i+1} \in C(\mathbb{X}_i; H^1(\mathbb{T}^d)) \\ \vdots \\ v_N \in C(\mathbb{X}_{N-1}; H^1(\mathbb{T}^d))}} \int_{(\mathbb{T}^d)^N} |e + \nabla v_{i+1}(x_0, \dots, x_{i+1}) + \dots + \nabla v_N(x_0, \dots, x_N)|^2 \\
 &\times e^{-V(x_0, \dots, x_N)/\sigma} dx_N \dots, dx_{i+1},
 \end{aligned} \tag{30}$$

for all $e \in \mathbb{R}^d$.

Proof For $i = 1, \dots, N$, from the proof of Lemma 1 we can express the intermediary diffusion tensor \mathcal{K}_i in the following recursive manner,

$$\begin{aligned}
 \mathcal{K}_i(x_0, \dots, x_i) &= \int_{\mathbb{T}^d} (I + \nabla_{x_{i+1}} \theta_{i+1}(x_0, \dots, x_i, x_{i+1}))^\top \\
 &\times \mathcal{K}_{i+1}(x_0, \dots, x_{i+1})(I + \nabla_{x_{i+1}} \theta_{i+1}(x_0, \dots, x_{i+1})) dx_{i+1}.
 \end{aligned}$$

Consider the tensor $\tilde{\mathcal{K}}_i$ defined by the following symmetric minimization problem

$$\begin{aligned}
 e \cdot \tilde{\mathcal{K}}_i(x_0, \dots, x_i)e &= \inf_{v \in C(\mathbb{X}_i; H^1(\mathbb{T}^d))} \int_{\mathbb{T}^d} (e + \nabla v(x_0, \dots, x_{i+1})) \cdot \mathcal{K}_{i+1}(x_0, \dots, x_{i+1}) \\
 &\times (e + \nabla v(x_0, \dots, x_{i+1})) dx_{i+1}.
 \end{aligned} \tag{31}$$

Since \mathcal{K}_{i+1} is a symmetric tensor, the corresponding Euler–Lagrange equation for the minimiser is given by

$$\nabla_{x_{i+1}} \cdot (\mathcal{K}_{i+1}(x_0, \dots, x_{i+1})(\nabla_{x_{i+1}} \chi(x_0, \dots, x_{i+1}) + e)) = 0, \quad x_{i+1} \in \mathbb{T}^d,$$

with periodic boundary conditions. This equation has a unique mean zero solution given by $\chi(x_0, \dots, x_{i+1}) = \theta_i(x_0, \dots, x_{i+1})^\top e$, where θ_i is the unique mean-zero solution of (15). It thus follows that $e^\top \mathcal{K}_i e = e^\top \tilde{\mathcal{K}}_i e$, where $\tilde{\mathcal{K}}_i$ is given by (31). Consider now the minimisation problem

$$\begin{aligned}
 \inf_{\substack{v_2 \in C(\mathbb{X}_i; H^1(\mathbb{T}^d)) \\ v_1 \in C(\mathbb{X}_{i+1}; H^1(\mathbb{T}^d))}} \int_{\mathbb{T}^d} \int_{\mathbb{T}^d} (e + \nabla_{x_{i+2}} v_1(x_0, \dots, x_{i+2}) + \nabla_{x_{i+1}} v_2(x_0, \dots, x_{i+1}))^\top \\
 \times \mathcal{K}_{i+2}(x_0, \dots, x_{i+2})(e + \nabla_{x_{i+2}} v_1(x_0, \dots, x_{i+2}) + \nabla_{x_{i+1}} v_2(x_0, \dots, x_{i+1})) \\
 \times dx_{i+2} dx_{i+1}.
 \end{aligned}$$

Optimising over v_1 for v_2 fixed it follows that $v_1 = (e + \nabla_{x_{i+1}} v_2)^\top \theta_{i+2}$, where θ_{i+2} is the unique mean-zero solution of (15). Thus, the above minimisation can be written as

$$\begin{aligned}
 \inf_{v_2 \in C(\mathbb{X}_i; H^1(\mathbb{T}^d))} \int_{\mathbb{T}^d} \int_{\mathbb{T}^d} (e + \nabla_{x_{i+1}} v_2(x_0, \dots, x_{i+1}))^\top (I + \nabla_{x_{i+2}} \theta_{i+2})^\top \\
 \times \mathcal{K}_{i+2}(x_0, \dots, x_{i+2})(I + \nabla_{x_{i+2}} \theta_{i+2}) \\
 \times (e + \nabla_{x_{i+1}} v_2(x_0, \dots, x_{i+1})) dx_{i+2} dx_{i+1} \\
 = \inf_{v_2 \in C(\mathbb{X}_{i-1}; H^1(\mathbb{T}^d))} \int_{\mathbb{T}^d} (e + \nabla_{x_{i+1}} v_2(x_0, \dots, x_{i+1}))^\top \mathcal{K}_{i+1}(x_0, \dots, x_{i+1}) \\
 \times (e + \nabla_{x_{i+1}} v_2(x_0, \dots, x_{i+1})) dx_{i+2} dx_{i+1} \\
 = e^\top \mathcal{K}_i e.
 \end{aligned}$$

Proceeding recursively, we arrive at the advertised result (30). □

4 Properties of the Equilibrium Distributions

In this section we study in more detail the properties of the equilibrium distributions μ^ϵ and μ^0 of the full (8) and homogenized dynamics (18), respectively. We first provide a proof of Proposition 3. The approach we follow in this proof is based on properties of periodic functions, in a manner similar to [12, Chap. 2].

Proof of Proposition 3 Let $f \in L^2(\mu_{ref})$ and $\delta > 0$. Clearly $C_c^\infty(\mathbb{R}^d)$ is dense in $L^2(\mu_{ref})$ and so, by Assumptions 1 there exists $f_\delta \in C_c^\infty(\mathbb{R}^d)$ such that

$$\left| \int_{\mathbb{R}^d} f(x)e^{-V^\epsilon(x)/\sigma} dx - \int_{\mathbb{R}^d} f_\delta(x)e^{-V^\epsilon(x)/\sigma} dx \right| \leq \frac{\delta}{3}, \tag{32}$$

and

$$\left| \int_{\mathbb{R}^d} \int_{\mathbb{T}^d} \dots \int_{\mathbb{T}^d} (f_\delta(x) - f(x))e^{-V(x,y_1,\dots,y_N)/\sigma} dy_N \dots dy_1 dx \right| \leq \frac{\delta}{3}, \tag{33}$$

uniformly with respect to ϵ . Now, we partition \mathbb{R}^d into pairwise disjoint translations of $[0, 1]^d$ as $\mathbb{R}^d = \cup_{k \in \mathbb{N}} Y_k$, where

$$Y_k = \epsilon^N x_k + \epsilon^N [0, 1]^d,$$

for $\{x_k\}_{k \geq 0} = \mathbb{Z}^d$. With this decomposition we obtain

$$\begin{aligned} \int_{\mathbb{R}^d} f_\delta(x)e^{-V^\epsilon(x)/\sigma} dx &= \sum_{k \in \mathbb{N}} \int_{Y_k} f_\delta(x)e^{-V^\epsilon(x)/\sigma} dx \\ &= \epsilon^{Nd} \sum_{k \in \mathbb{N}} \int_{[0,1]^d} f_\delta(\epsilon^N(x_k + y))e^{-V(\epsilon^N(x_k + y), \dots, \epsilon^N(x_k + y), y)/\sigma} dy, \end{aligned}$$

where in the last equality we use the periodicity of V with respect to the last variable. Since the integrand is smooth with compact support, we can Taylor expand around $\epsilon^N x_k$ to obtain

$$\int_{\mathbb{R}^d} f_\delta(x)e^{-V^\epsilon(x)/\sigma} dx = \epsilon^{Nd} \sum_{k \in \mathbb{N}} \int_{[0,1]^d} f_\delta(\epsilon^N x_k)e^{-V(\epsilon^N x_k, \dots, \epsilon^N x_k, y)/\sigma} dy + C\epsilon,$$

where C is a constant depending on the derivatives of V with respect to the first N variables, and the volume of the support of f_δ .

Noting that the above sum is a Riemann sum approximation, we can write

$$\begin{aligned} &\epsilon^{Nd} \sum_{k \in \mathbb{N}} \int_{[0,1]^d} f_\delta(\epsilon^N x_k)e^{-V(\epsilon^N x_k, \dots, \epsilon^N x_k, y)/\sigma} dy \\ &= \epsilon^{Nd} \sum_{k \in \mathbb{N}} \int_{[0,1]^d} \int_{[0,1]^d} f_\delta(\epsilon^N(x_k + y'))e^{-V(\epsilon^N(x_k + y'), \dots, \epsilon^N(x_k + y'), y)/\sigma} dy dy' + C_1\epsilon \\ &= \int_{\mathbb{R}^d} \int_{[0,1]^d} f_\delta(x)e^{-V(x, \dots, x/\epsilon^{N-1}, y)/\sigma} dy dx + C_1\epsilon, \end{aligned}$$

where C_1 is a constant. Repeating the above process $N - 1$ times, we obtain that

$$\int_{\mathbb{R}^d} f_\delta(x)e^{-V^\epsilon(x)/\sigma} dx = \int_{\mathbb{R}^d} \int_{\mathbb{T}^d} \dots \int_{\mathbb{T}^d} f_\delta(x)e^{-V(x,y_1,\dots,y_N)/\sigma} dy_N \dots dy_1 dx + C_N\epsilon, \tag{34}$$

where $C_N > 0$ is a constant depending on the support of f_δ and derivatives of V with respect to the first N variable. Thus, choosing $\epsilon < \delta/(3C_N)$ and combining (32), (33) and (34) we obtain

$$\left| \int_{\mathbb{R}^d} f(x)e^{-V^\epsilon(x)/\sigma} dx - \int_{\mathbb{R}^d} \int_{\mathbb{T}^d} \dots \int_{\mathbb{T}^d} f(x)e^{-V(x,y_1,\dots,y_N)/\sigma} dy_N \dots dy_1 dx \right| \leq \delta, \tag{35}$$

Choosing $f \equiv 1$ we obtain immediately that

$$Z^\epsilon = \int_{\mathbb{R}^d} e^{-V^\epsilon(x)/\sigma} dx \rightarrow Z^0 = \int_{\mathbb{R}^d} \int_{\mathbb{T}^d} \dots \int_{\mathbb{T}^d} e^{-V(x,y_1,\dots,y_N)} dy_N \dots dy_1 dx,$$

and so for $f \in L^2(\mu_{ref})$ we obtain

$$\int f(x)\mu^\epsilon(x) dx \rightarrow \int f(x)\mu^0(x) dx,$$

as $\epsilon \rightarrow 0$, as required. □

Proof of Proposition 4 Since V_1 is bounded uniformly by Assumption 1, it is straightforward to check that

$$\mu_{ref}(x)e^{-osc(V_1)/\sigma} \leq \mu^\epsilon(x) \leq \mu_{ref}(x)e^{osc(V_1)/\sigma}. \tag{36}$$

It follows from the discussion following [5, Prop 4.2.7], that μ^ϵ satisfies Poincaré’s inequality with constant

$$\gamma = \frac{\rho}{\sigma} e^{-osc(V_1)/\sigma},$$

which implies (24). An identical argument follows for the coarse-grained density $\mu^0(x)$. Finally, by (20) of Proposition 2 we have $|v|^2 e^{-osc(V_1)/\sigma} \leq v \cdot \mathcal{M}(x)v$, for all $v \in \mathbb{R}^d$, and so

$$\begin{aligned} \text{Var}_{\mu^0}(f) &\leq \frac{\sigma}{\rho} e^{osc(V_1)/\sigma} \int_{\mathbb{R}^d} |\nabla f(x)|^2 \mu^0(x) dx \\ &\leq \frac{\sigma}{\rho} e^{2osc(V_1)/\sigma} \int \nabla f(x) \cdot \mathcal{M}(x) \nabla f(x) \mu^0(x) dx, \end{aligned}$$

from which (25) follows. □

Remark 4 Note that one can similarly relate the constants in the logarithmic Sobolev inequalities for the measures μ_{ref} , μ^ϵ and μ^0 in an almost identical manner, based on the Holley-Stroock criterion [26].

Remark 5 Proposition 4 requires the assumption that the multiscale perturbation V_1 is bounded uniformly. If this is relaxed, then it is no longer guaranteed that μ^ϵ will satisfy a Poincaré inequality, even though μ_{ref} does. Consider, for example, the following one dimensional potential

$$V^\epsilon(x) = x^2(1 + \alpha \cos(x/\epsilon)),$$

then the corresponding Gibbs distribution $\mu^\epsilon(x)$ will not satisfy Poincaré’s inequality for any $\epsilon > 0$. Following [25, Appendix A] we demonstrate this by checking that this choice of

μ^ϵ does not satisfy the Muckenhoupt criterion [2, 38] which is necessary and sufficient for the Poincaré inequality to hold, namely that $\sup_{r \in \mathbb{R}} B_\pm(r) < \infty$, where

$$B_\pm(r) = \left(\int_r^{\pm\infty} \mu^\epsilon(x) dx \right)^{\frac{1}{2}} \left(\int_{[0, \pm r]} \frac{1}{\mu^\epsilon(x)} dx \right)^{\frac{1}{2}}.$$

Given $n \in \mathbb{N}$, we set $r/\epsilon = 2\pi n + \pi/2$. Then we have that

$$\begin{aligned} B_+(r) &\geq \left(\int_{\epsilon(2\pi n+2\pi/3)}^{\epsilon(2\pi n+4\pi/3)} e^{-|x|^2(1-\alpha/2)/\sigma} dx \right)^{1/2} \left(\int_{\epsilon(2\pi n-\pi/3)}^{\epsilon(2\pi n+\pi/3)} e^{|x|^2(1+\alpha/2)/\sigma} dx \right)^{1/2} \\ &\geq \left(\frac{2\pi\epsilon}{3} \right) \exp \left(-\frac{|\pi\epsilon(2n+4/3)|^2}{2\sigma} \left(1 - \frac{\alpha}{2}\right) + \frac{|\pi\epsilon(2n-1/3)|^2}{2\sigma} \left(1 + \frac{\alpha}{2}\right) \right) \\ &= \left(\frac{2\pi\epsilon}{3} \right) \exp \left(-\frac{|2\pi\epsilon n|^2 \left(1 + \frac{2}{3n}\right)^2}{2\sigma} \left(1 - \frac{\alpha}{2}\right) + \frac{|2\pi\epsilon n|^2 \left(1 - \frac{1}{6n}\right)^2}{2\sigma} \left(1 + \frac{\alpha}{2}\right) \right) \\ &\approx \left(\frac{2\pi\epsilon}{3} \right) \exp \left(\frac{|2\pi\epsilon n|^2}{2\sigma} (\alpha + o(n^{-1})) \right) \rightarrow \infty, \quad \text{as } n \rightarrow \infty, \end{aligned}$$

so that Poincaré’s inequality does not hold for μ^ϵ .

A natural question to ask is whether the weak convergence of μ^ϵ to μ^0 holds true in a stronger notion of distance such as total variation. The following simple one-dimensional example demonstrates that the convergence cannot be strengthened to total variation.

Example 1 Consider the one dimensional Gibbs distribution

$$\mu^\epsilon(x) = \frac{1}{Z^\epsilon} e^{-V^\epsilon(x)/\sigma},$$

where

$$V^\epsilon(x) = \frac{x^2}{2} + \alpha \cos\left(\frac{x}{\epsilon}\right),$$

and where Z^ϵ is the normalization constant and $\alpha \neq 0$. Then the measure μ^ϵ converges weakly to μ^0 given by

$$\mu^0(x) = \frac{1}{\sqrt{2\pi\sigma}} e^{-x^2/2\sigma}.$$

From the plots of the stationary distributions in Fig. 2a it becomes clear that the density of μ^ϵ exhibits rapid fluctuations which do not appear in μ^0 , thus we do not expect to be able to obtain convergence in a stronger metric. First we consider the distance between μ^ϵ and μ^0 in total variation⁵

$$\|\mu^\epsilon - \mu^0\|_{TV} = \int_{\mathbb{R}} |\mu^\epsilon(x) - \mu^0(x)| dx = \int_{\mathbb{R}} \frac{e^{-x^2/2\sigma}}{\sqrt{2\sigma}} \left| 1 - \frac{e^{-\frac{\alpha}{\sigma} \cos(2\pi x/\epsilon)}}{K^\epsilon} \right| dx,$$

where $K^\epsilon = Z^\epsilon/\sqrt{2\pi\sigma}$. It follows that

$$\|\mu^\epsilon - \mu^0\|_{TV} \geq \sum_{n \geq 0} \int_{\epsilon(2\pi n-\pi/3)}^{\epsilon(2\pi n+\pi/3)} \frac{e^{-x^2/2\sigma}}{\sqrt{2\pi\sigma}} dx \left| 1 - \frac{e^{-\frac{\alpha}{2\sigma}}}{K^\epsilon} \right|$$

⁵ We are using the same notation for the measure and for its density with respect to the Lebesgue measure on \mathbb{R} .

$$\begin{aligned} &\geq \sum_{n \geq 0} \frac{2\epsilon\pi}{3} \frac{e^{-\epsilon^2(2n\pi + \pi/3)^2/2\sigma}}{\sqrt{2\pi\sigma}} \left| 1 - \frac{e^{-\frac{\alpha}{2\sigma}}}{K^\epsilon} \right| \\ &\geq \int_0^\infty \frac{2\pi}{3} \frac{e^{-2\pi^2(x+\epsilon/6)^2/\sigma}}{\sqrt{2\pi\sigma}} \left| 1 - \frac{e^{-\frac{\alpha}{2\sigma}}}{K^\epsilon} \right|, \end{aligned}$$

where we use the fact that $e^{-\alpha/2\sigma}/K^\epsilon \leq 1$ for ϵ sufficiently small. In the limit $\epsilon \rightarrow 0$, we have $K^\epsilon \rightarrow I_0(\alpha/\sigma)$, where $I_n(\cdot)$ is the modified Bessel function of the first kind of order n . Therefore, as $\epsilon \rightarrow 0$,

$$\|\mu^\epsilon - \mu^0\|_{TV} \geq \int_0^\infty \frac{2\pi}{3} \frac{e^{-2\pi^2(x+\epsilon/6)^2/\sigma}}{\sqrt{2\pi\sigma}} \left| 1 - \frac{e^{-\frac{\alpha}{2\sigma}}}{K^\epsilon} \right| = \frac{1}{6} \left| 1 - \frac{e^{-\frac{\alpha}{2\sigma}}}{I_0(\alpha/\sigma)} \right|, \quad (37)$$

which converges to $\frac{1}{6}$ as $\frac{\alpha}{\sigma} \rightarrow \infty$. Since relative entropy controls total variation distance by Pinsker’s theorem, it follows that μ^ϵ does not converge to μ^0 in relative entropy, either. Nonetheless, we shall compute the distance in relative entropy between μ^ϵ and μ^0 to understand the influence of the parameters σ and α . Since both μ^0 and μ^ϵ have strictly positive densities with respect to the Lebesgue measure on \mathbb{R} , we have that

$$\frac{d\mu^\epsilon}{d\mu^0}(x) = \frac{\sqrt{2\pi\sigma}}{Z^\epsilon} e^{-\frac{V^\epsilon(x)}{\sigma} + \frac{x^2}{2\sigma}}.$$

Then, for $Z^0 = \sqrt{2\pi\sigma} I_0(1/\sigma)$,

$$\begin{aligned} H(\mu^\epsilon | \mu^0) &= \frac{1}{Z^\epsilon} \int \left(\frac{1}{2} \log(2\pi\sigma) - \log Z^\epsilon \right) e^{-V^\epsilon(x)/\sigma} dx \\ &\quad + \frac{1}{Z^\epsilon} \int (-V^\epsilon(x)/\sigma + x^2/2\sigma) e^{-V^\epsilon(x)/\sigma} dx \\ &\xrightarrow{\epsilon \rightarrow 0} -\log I_0(\alpha/\sigma) - \frac{\alpha}{\sigma Z^0} \lim_{\epsilon \rightarrow 0} \int \cos(2\pi x/\epsilon) e^{-x^2/2\sigma - \alpha \cos(2\pi x/\epsilon)/\sigma} dx \\ &= -\log I_0(\alpha/\sigma) - \frac{\alpha}{\sigma} \frac{I_1(\alpha/\sigma)}{I_0(\alpha/\sigma)} =: K(\alpha/\sigma), \end{aligned}$$

and it is straightforward to check that $K(s) > 0$, and moreover

$$K(s) \rightarrow \begin{cases} 0 & \text{as } s \rightarrow 0 \\ +\infty & \text{as } s \rightarrow \infty \end{cases}.$$

In Fig. 2b we plot the value of $K(s)$ as a function of s . From this result, we see that for fixed $\alpha > 0$, the measure μ^ϵ will converge in relative entropy only in the limit as $\sigma \rightarrow \infty$, while the measures will become increasingly mutually singular as $\sigma \rightarrow 0$.

5 Proof of Weak Convergence

In this section we show that over finite time intervals $[0, T]$, the process X_t^ϵ converges weakly to a process X_t^0 which is uniquely identified as the weak solution of a coarse-grained SDE. The approach we adopt is based on the classical martingale methodology of [8, Sect. 3]. The proof of the homogenization result is split into three steps.




Fig. 2 Error between $\mu^\epsilon(x) \propto \exp(-V^\epsilon(x)/\sigma)$ and effective distribution μ^0

1. We construct an appropriate test function which is used to decompose the fluctuations of the process X_t^ϵ into a martingale part and a term which goes to zero as $\epsilon \rightarrow 0$.
2. Using this test function, we demonstrate that the path measure \mathbb{P}^ϵ corresponding to the family $\left\{ (X_t^\epsilon)_{t \in [0, T]} \right\}_{0 < \epsilon \leq 1}$ is tight on $C([0, T]; \mathbb{R}^d)$.
3. Finally, we show that any limit point of the family of measures must solve a well-posed martingale problem, and is thus unique.

The test functions will be constructed by solving a recursively defined sequence of Poisson equations on \mathbb{R}^d . We first provide a general well-posedness result for this class of equations.

Proposition 5 *Let $\mathbb{X}_k, k = 0, 1, \dots, N$ be the space defined in Sect. 3.2. For fixed $(x_0, \dots, x_{k-1}) \in \mathbb{X}_{k-1}$, let \mathcal{S}_k be the operator given by*

$$\mathcal{S}_k u = \frac{1}{\rho(x_0, \dots, x_k)} \nabla_{x_k} \cdot (\rho(x_0, \dots, x_k) D(x_0, \dots, x_k) \nabla_{x_k} u(x_0, \dots, x_k)), \quad (38)$$

for $u \in C^2(\mathbb{T}^d)$, where ρ is a smooth and uniformly positive and bounded function, and D is a smooth and uniformly positive definite tensor on \mathbb{X}_k . Let h be a smooth function with bounded derivatives, such that for each $(x_0, \dots, x_{k-1}) \in \mathbb{X}_{k-1}$:

$$\int_{\mathbb{T}^d} h(x_0, \dots, x_k) \rho(x_0, \dots, x_k) dx_k = 0. \quad (39)$$

Then there exists a unique solution $u \in C(\mathbb{X}_{k-1}; H^1(\mathbb{T}^d))$ to the Poisson equation on \mathbb{T}^d given by

$$\mathcal{S}_k u(x_0, \dots, x_k) = h(x_0, \dots, x_k), \quad \int_{\mathbb{T}^d} u(x_0, \dots, x_k) \rho(x_0, \dots, x_k) dx_k = 0. \quad (40)$$

Moreover u is smooth and bounded with respect to the variable $x_k \in \mathbb{T}^d$ as well as the parameters $x_0, \dots, x_{k-1} \in \mathbb{X}_{k-1}$.

Proof Since ρ and D are strictly positive, for fixed values of x_0, \dots, x_{k-1} , the operator \mathcal{S}_k is uniformly elliptic, and since \mathbb{T}^d is compact, \mathcal{S}_k has compact resolvent in $L^2(\mathbb{T}^d)$, see [18, Chap. 6] and [46, Chap. 7]. The nullspace of the adjoint \mathcal{S}^* is spanned by a single function $\rho(x_0, \dots, x_{k-1}, \cdot)$. By the Fredholm alternative, a necessary and sufficient condition for the existence of u is (39) which is assumed to hold. Thus, there exists a unique solution

$u(x_0, \dots, x_{k-1}, \cdot) \in H^1(\mathbb{T}^d)$ having mean zero with respect to $\rho(x_0, \dots, x_k)$. By elliptic estimates and Poincaré’s inequality, it follows that there exists $C > 0$ satisfying

$$\|u(x_0, \dots, x_{k-1}, \cdot)\|_{H^1(\mathbb{T}^d)} \leq C \|h(x_0, \dots, x_{k-1}, \cdot)\|_{L^2(\mathbb{T}^d)},$$

for all $(x_0, \dots, x_{k-1}) \in \mathbb{X}_{k-1}$. Since the components of D and ρ are smooth with respect to x_k , standard interior regularity results [21] ensure that, for fixed $x_0, \dots, x_{k-1} \in \mathbb{X}_{k-1}$, the function $u(x_0, \dots, x_{k-1}, \cdot)$ is smooth. To prove the smoothness and boundedness with respect to the other parameters x_0, \dots, x_{k-1} , we can apply an approach either similar to [8], by showing that the finite differences approximation of the derivatives of u with respect to the parameters has a limit, or otherwise, by directly differentiating the transition density of the semigroup associated with the generator \mathcal{S}_k , see for example [43, 44, 55] as well as [21, Sec 8.4]. \square

Remark 6 Suppose that the function h in Proposition 5 can be expressed as

$$h(x_0, \dots, x_k) = a(x_0, x_1, \dots, x_k) \cdot \nabla \phi_0(x_0)$$

where a is smooth with all derivatives bounded. Then the mean-zero solution of (40) can be written as

$$u(x_0, x_1, \dots, x_k) = \chi(x_0, x_1, \dots, x_k) \cdot \nabla \phi_0(x_i), \tag{41}$$

where χ is the classical mean-zero solution to the following Poisson equation

$$\mathcal{S}_k \chi(x_0, \dots, x_k) = a(x_0, \dots, x_k), \quad (x_0, \dots, x_k) \in \mathbb{X}_k. \tag{42}$$

This can be seen by checking directly that u given in (41) with χ satisfying (42) solves (40), which implies it is the unique solution of (40) due to the uniqueness of a solution. In particular, χ is smooth and bounded over x_0, \dots, x_k , so that given a multi-index $\alpha = (\alpha_0, \dots, \alpha_k)$ on the indices $(0, \dots, k)$, there exists $C_\alpha > 0$ such that

$$|\nabla^\alpha u(x_0, \dots, x_k)|_F \leq C_\alpha \sum_{k=0}^{\alpha_0} |\nabla^{k+1} \phi_0(x_0)|_F, \quad \forall x_0, x_1, \dots, x_k,$$

where $|\cdot|_F$ denotes the Frobenius norm. A similar decomposition is possible for

$$g(x_0, \dots, x_k) = A(x_0, x_1, \dots, x_k) : \nabla^2 \phi_0(x_0),$$

where ∇^2 denotes the Hessian.

5.1 Constructing the Test Functions

It is clear that we can rewrite (8) as

$$dX_t^\epsilon = - \sum_{i=0}^N \epsilon^{-i} \nabla_{x_i} V(x, x/\epsilon, \dots, x/\epsilon^N) dt + \sqrt{2\sigma} dW_t. \tag{43}$$

The generator of X_t^ϵ denoted by \mathcal{L}^ϵ can be decomposed into powers of ϵ as follows

$$(\mathcal{L}^\epsilon f)(x) = - \sum_{i=0}^N \epsilon^{-i} \nabla_{x_i} V(x, x/\epsilon, \dots, x/\epsilon^N) \cdot \nabla f(x) + \sigma \Delta f(x).$$

For functions of the form $f^\epsilon(x) = f(x, x/\epsilon, \dots, x/\epsilon^N)$, we have

$$\begin{aligned}
 (\mathcal{L}^\epsilon f^\epsilon)(x) &= \sum_{i=0}^N \epsilon^{-i} \nabla_{x_i} V(x, x/\epsilon, \dots, x/\epsilon^N) \cdot \left(\sum_{j=0}^N \epsilon^{-j} \nabla_{x_j} f(x, x/\epsilon, \dots, x/\epsilon^N) \right) \\
 &\quad + \sigma \sum_{i,j=0}^k \epsilon^{-(i+j)} \nabla_{x_i x_j}^2 f(x, x/\epsilon, \dots, x/\epsilon^N) \\
 &= \sum_{i,j=0}^N \epsilon^{-(i+j)} \left[e^{V/\sigma} \nabla_{x_i} \cdot \left(\sigma e^{-V/\sigma} \nabla_{x_j} f \right) \right] (x, x/\epsilon, \dots, x/\epsilon^N) \\
 &= \sum_{n=0}^{2N} \epsilon^{-n} (\mathcal{L}_n f)(x, x/\epsilon, \dots, x/\epsilon^N), \tag{44}
 \end{aligned}$$

where for $n = 0, \dots, 2N$

$$(\mathcal{L}_n f)(x, x/\epsilon, \dots, x/\epsilon^N) = \left[e^{V/\sigma} \sum_{\substack{i,j \in \{0, \dots, N\} \\ i+j=n}} \nabla_{x_i} \cdot \left(\sigma e^{-V/\sigma} \nabla_{x_j} f \right) \right] (x, x/\epsilon, \dots, x/\epsilon^N).$$

Given a function ϕ_0 , which will be specified later, our objective is to construct a test function ϕ^ϵ of the form

$$\begin{aligned}
 \phi^\epsilon(x) &= \phi_0(x) + \epsilon \phi_1(x, x/\epsilon) + \dots + \epsilon^N \phi_N(x, x/\epsilon, \dots, x/\epsilon^N) \\
 &\quad + \epsilon^{N+1} \phi_{N+1}(x, x/\epsilon, \dots, x/\epsilon^N) + \dots + \epsilon^{2N} \phi_{2N}(x, x/\epsilon, \dots, x/\epsilon^N),
 \end{aligned}$$

such that

$$(\mathcal{L}^\epsilon \phi^\epsilon)(x) = F(x) + O(\epsilon), \tag{45}$$

for some function F which is independent of ϵ . The above form for the test function is suggested by the calculation (44). Using (44) we compute

$$\begin{aligned}
 (\mathcal{L}^\epsilon \phi^\epsilon)(x) &= \sum_{k=0}^{2N} \epsilon^k (\mathcal{L} \phi_k)(x, x/\epsilon, \dots, x/\epsilon^N) \\
 &= \sum_{k=0}^{2N} \epsilon^k \left(\sum_{n=0}^{2N} \epsilon^{-n} (\mathcal{L}_n \phi_k)(x, x/\epsilon, \dots, x/\epsilon^N) \right) \\
 &= \sum_{k,n=0}^{2N} \epsilon^{k-n} (\mathcal{L}_n \phi_k)(x, x/\epsilon, \dots, x/\epsilon^N),
 \end{aligned}$$

where

$$(\mathcal{L}_n \phi_k)(x, x/\epsilon, \dots, x/\epsilon^N) = \left[e^{V/\sigma} \sum_{\substack{i,j \in \{0, \dots, N\} \\ i+j=n}} \nabla_{x_i} \cdot \left(\sigma e^{-V/\sigma} \nabla_{x_j} \phi_k \right) \right] (x, x/\epsilon, \dots, x/\epsilon^N).$$

Note that $\nabla_{x_j} \phi_k = 0$ for $j > k$. By equating powers of ϵ , from $O(\epsilon^{-N})$ to $O(1)$ respectively, in both sides of (45), we obtain the following sequence of $N + 1$ equations

$$\mathcal{L}_{2N}\phi_N + \mathcal{L}_{2N-1}\phi_{N-1} + \dots + \mathcal{L}_N\phi_0 = 0, \tag{46a}$$

$$\mathcal{L}_{2N}\phi_{N+1} + \mathcal{L}_{2N-1}\phi_N + \dots + \mathcal{L}_{N-1}\phi_0 = 0, \tag{46b}$$

\vdots

$$\mathcal{L}_{2N}\phi_{2N-1} + \dots + \mathcal{L}_1\phi_0 = 0, \tag{46c}$$

$$\mathcal{L}_{2N}\phi_{2N} + \dots + \mathcal{L}_0\phi_0 = F. \tag{46d}$$

This system generalizes the system written for three scales in [8, III–11.3]. We note that each nonzero term in (46a), (46b) to (46c) has the form

$$\sigma e^{V(x_0, \dots, x_N)/\sigma} \nabla_{x_i} \cdot \left(e^{-V(x_0, \dots, x_N)/\sigma} \nabla_{x_j} \phi_k \right),$$

where $1 \leq i + j - k \leq N$. Furthermore, all the terms appearing in (46a), (46b) to (46c) must satisfy $i > 0$. Indeed $i = 0$ would imply $j \geq k + 1 > k$ and so $\nabla_{x_j} \phi_k = 0$ by construction of the test function. Since

$$V(x_0, \dots, x_N) = V_0(x_0) + V_1(x_0, \dots, x_N),$$

all the terms $\mathcal{L}_n\phi_k$ appearing (46a), (46b) to (46c) can be simplified as

$$\begin{aligned} \mathcal{L}_n\phi_k &= e^{(V_0+V_1)/\sigma} \sum_{\substack{i \in \{1, \dots, N\} \\ j \in \{0, \dots, N\} \\ i+j=n}} \nabla_{x_i} \cdot \left(\sigma e^{-(V_0+V_1)/\sigma} \nabla_{x_j} \phi_k \right) \\ &= e^{V_1/\sigma} \sum_{\substack{i \in \{1, \dots, N\} \\ j \in \{0, \dots, N\} \\ i+j=n}} \nabla_{x_i} \cdot \left(\sigma e^{-V_1/\sigma} \nabla_{x_j} \phi_k \right), \end{aligned}$$

where we have used the fact that V_0 is independent of x_i for $i \in \{1, \dots, N\}$ to pull the term e^{V_0} out from the divergence operator. Thus, we can rewrite the first N equations as

$$\mathcal{A}_{2N}\phi_N + \mathcal{A}_{2N-1}\phi_{N-1} + \dots \mathcal{A}_N\phi_0 = 0, \tag{47a}$$

$$\mathcal{A}_{2N}\phi_{N+1} + \mathcal{A}_{2N-1}\phi_N + \dots \mathcal{A}_{N-1}\phi_0 = 0, \tag{47b}$$

\vdots

$$\mathcal{A}_{2N}\phi_{2N-1} + \dots + \mathcal{A}_1\phi_0 = 0, \tag{47c}$$

where

$$\mathcal{A}_n f = \sigma e^{V_1(x_0, \dots, x_N)/\sigma} \sum_{\substack{i \in \{1, \dots, N\} \\ j \in \{0, \dots, N\} \\ i+j=n}} \nabla_{x_i} \cdot \left(e^{-V_1(x_0, \dots, x_N)/\sigma} \nabla_{x_j} f \right).$$

Before constructing the test functions, we first introduce the sequence of spaces on which the sequence of correctors will be constructed. Define \mathcal{H} to be the space of functions on the extended state space, i.e. $\mathcal{H} = L^2(\mathbb{X}_N)$, where \mathbb{X}_N is defined by (26). We construct the following sequence of subspaces of \mathcal{H} . Let

$$\mathcal{H}_N = \left\{ f \in \mathcal{H} : \int f(x_0, \dots, x_N) e^{-V_1/\sigma} dx_N = 0 \right\},$$

Then clearly $\mathcal{H} = \mathcal{H}_N \oplus \mathcal{H}_N^\perp$. Suppose we have defined \mathcal{H}_{N-k+1} then we can define \mathcal{H}_{N-k} inductively by

$$\mathcal{H}_{N-k} = \left\{ f \in \mathcal{H}_{N-k+1} : \int f(x_0, \dots, x_{N-k}) Z_{N-k}(x_0, \dots, x_{N-k}) dx_{N-k} = 0 \right\},$$

where $Z_i(x_0, \dots, x_i) = \int \dots \int e^{-V_1(x_0, \dots, x_N)/\sigma} dx_{i+1} dx_{i+2} \dots dx_N$. Clearly, we have that $\mathcal{H}_1 \oplus \mathcal{H}_1^\perp \oplus \dots \oplus \mathcal{H}_N^\perp = \mathcal{H}$.

Applying Proposition 5 we can now construct the series of test functions ϕ_1, \dots, ϕ_{2N} that solve (47).

Proposition 6 *Given $\phi_0 \in C^\infty(\mathbb{R}^d)$, there exist smooth functions ϕ_i for $i = 1, \dots, 2N - 1$ such that Eqs (47a)–(47c) are satisfied, and moreover we have the following pointwise estimates, which hold uniformly on $x_0, \dots, x_k \in \mathbb{X}_k$:*

$$\|\nabla^\alpha \phi_i(x_0, \dots, x_k)\|_F \leq C \sum_{l=1}^{\alpha_0+2} \|\nabla_{x_0}^l \phi_0(x_0)\|_F, \tag{48}$$

for some constant $C > 0$, and all multiindices α on $(0, \dots, k)$, and all $0 \leq k \leq i \leq 2N - 1$. Finally, Eq. (46d) is satisfied with

$$F(x) = \frac{1}{Z(x)} \nabla_{x_0} \cdot (\mathcal{K}_1(x) \nabla_{x_0} \phi_0(x)). \tag{49}$$

Proof Guideline of the proof. Given ϕ_0 as in the hypothesis of the proposition, we will find the test functions $\phi_i, i = 1, \dots, 2N$ from the system (47). This system consists of N equations. The other N equations come from solvability (compatibility) conditions, which are applications of the Fredholm alternative [46, Theorem 7.9]. More specially, the solvability condition for the $O(\epsilon^{-(N-k)})$ -equation in (47), viewing as an equation for ϕ_{N+k} in terms of $\phi_0, \dots, \phi_{N+k-1}$, will give rise to an equation for ϕ_{N-k} in term of $\phi_0, \dots, \phi_{N-k-1}$, for $k = 1, \dots, N$. The latter is an elliptic equation of the form (38) with $\rho = 1$ and $D = \mathcal{K}_{N-k}$. According to Lemma 1, \mathcal{K}_{N-k} is uniformly positive definite. Hence, the existence of ϕ_{N-k} follows from Proposition 5. Therefore, the solvability condition for ϕ_{N+k} is fulfilled guaranteeing the existence of ϕ_{N+k} . By inductively repeating this process for all $k = 1, \dots, N$, we can construct the test functions ϕ_1, \dots, ϕ_{2N} satisfying the system (47). Finally, the function F is then determined from (46d).

Now we implement this strategy in details. We start from Equation (47a), which can be viewed as an equation for ϕ_N in term of $\phi_0, \dots, \phi_{N-1}$

$$\mathcal{A}_{2N} \phi_N = -(\mathcal{A}_{2N-1} \phi_{N-1} + \dots + \mathcal{A}_0 \phi_0), \quad \mathcal{A}_{2N} f = \sigma e^{V_1/\sigma} \nabla_{x_N} \cdot (e^{-V_1/\sigma} \nabla_{x_N} f). \tag{50}$$

Since the operator \mathcal{A}_{2N} has a compact resolvent in $L^2(\mathbb{T}^d)$, by the Fredholm alternative a necessary and sufficient condition for (47a) to have a solution is that the following compatibility condition holds

$$\int (\mathcal{A}_{2N-1} \phi_{N-1} + \mathcal{A}_{2N-2} \phi_{N-2} + \dots + \mathcal{A}_N \phi_0) e^{-V_1/\sigma} dx_N = 0. \tag{51}$$

Note that every term in this summation is of the form

$$\mathcal{A}_{2N-k} \phi_{N-k} = \sigma \sum_{\substack{0 \leq i, j \leq N \\ i+j=2N-k}} e^{V_1/\sigma} \nabla_{x_j} \cdot (e^{-V_1/\sigma}(x) \nabla_{x_i} \phi_{N-k}). \tag{52}$$

For $\nabla_{x_i} \phi_{N-k}$ to be non-zero it is necessary that $i \leq N - k$. To enforce the condition $i + j = 2N - k$ it must be that $i = N - k$ and $j = N$, and thus the only non-zero terms in the above summation are:

$$\mathcal{A}_{2N-k} \phi_{N-k} = \sigma e^{V_1/\sigma} \nabla_{x_N} \cdot \left(e^{-V_1/\sigma} \nabla_{x_{N-k}} \phi_{N-k} \right), \tag{53}$$

for $k = 1, \dots, N$. It follows that the compatibility condition (51) holds, by the periodicity of the domain. Therefore (47a) has a solution. In addition, it can be written as

$$\begin{aligned} \mathcal{A}_{2N} \phi_N &= - \sum_{k=1}^N \mathcal{A}_{2N-k} \phi_{N-k} \\ &= - \sum_{k=1}^N \sigma e^{V_1/\sigma} \nabla_{x_N} \cdot \left(e^{-V_1/\sigma} \nabla_{x_{N-k}} \phi_{N-k} \right) \\ &= \left(\sigma e^{V_1/\sigma} \nabla_{x_N} \cdot (e^{-V_1/\sigma} I) \right) \cdot \left(\sum_{k=1}^N \nabla_{x_{N-k}} \phi_{N-k} \right). \end{aligned}$$

Note that for $k = 0$, the Poisson equation (15) can be expressed as

$$\mathcal{A}_{2N} \theta_N = \sigma e^{V_1/\sigma} \nabla_{x_N} \cdot (e^{-V_1/\sigma} I),$$

which has unique mean-zero solution θ_N . According to Remark 6, the test function ϕ_N can be written as

$$\phi_N = \theta_N \cdot (\nabla_{x_{N-1}} \phi_{N-1} + \dots + \nabla_{x_0} \phi_0) + r_N^{(1)}(x_0, \dots, x_{N-1}), \tag{54}$$

where

$$\theta_N \cdot (\nabla_{x_{N-1}} \phi_{N-1} + \dots + \nabla_{x_0} \phi_0) \in \mathcal{H}_N,$$

and for some $r_N^{(1)} \in \mathcal{H}_N^\perp$, which will be specified later. Next we consider the $O(\epsilon^{-(N-1)})$ -equation, that is (47b) viewing as an equation for ϕ_{N+1} in terms of ϕ_N, \dots, ϕ_0 :

$$\mathcal{A}_{2N} \phi_{N+1} = -(\mathcal{A}_{2N-1} \phi_N + \dots + \mathcal{A}_{N-1} \phi_0), \tag{55}$$

where \mathcal{A}_{2N} is given in (50). According to the Fredholm alternative, a necessary and sufficient condition for the above equation to have a solution is

$$\int (\mathcal{A}_{2N-1} \phi_N + \dots + \mathcal{A}_{N-2} \phi_1 + \mathcal{A}_{N-1} \phi_0) e^{-V_1/\sigma} dx_N = 0. \tag{56}$$

Similarly as in (53), for $k = 1, \dots, N + 1$, we have

$$\begin{aligned} \mathcal{A}_{2N-k} \phi_{N-k+1} &= \sigma e^{V_1/\sigma} \left[\nabla_{x_{N-1}} \cdot \left(e^{-V_1/\sigma} \nabla_{x_{N-k+1}} \phi_{N-k+1} \right) \right. \\ &\quad \left. + \nabla_{x_N} \cdot (e^{-V_1/\sigma} \nabla_{x_{N-k}} \phi_{N-k+1}) \right]. \end{aligned}$$

Substituting this into (55) we obtain

$$\begin{aligned} 0 &= \int \nabla_{x_{N-1}} \cdot \left[e^{-V_1/\sigma} (\nabla_{x_N} \phi_N + \nabla_{x_{N-1}} \phi_{N-1} + \dots + \nabla_{x_0} \phi_0) \right] dx_N \\ &= \nabla_{x_{N-1}} \cdot \left(\int e^{-V_1/\sigma} \nabla_{x_N} \theta_N (\nabla_{x_{N-1}} \phi_{N-1} \dots + \nabla_{x_0} \phi_0) dx_N \right) \end{aligned}$$

$$+ \nabla_{x_{N-1}} \cdot \left(\int e^{-V_1/\sigma} (\nabla_{x_{N-1}} \phi_{N-1} + \dots + \nabla_{x_0} \phi_0) \right) dx_N,$$

where in the last equality we use the fact that $r_N^{(1)}$ is independent of x_N . Thus we obtain the following equation for ϕ_{N-1} :

$$\nabla_{x_{N-1}} \cdot (\mathcal{K}_{N-1} \nabla_{x_{N-1}} \phi_{N-1}) = -\nabla_{x_{N-1}} \cdot (\mathcal{K}_{N-1} (\nabla_{x_{N-2}} \phi_{N-2} + \dots + \nabla_{x_0} \phi_0)), \tag{57}$$

where

$$\mathcal{K}_{N-1}(x_0, x_1, \dots, x_{N-1}) = \int (I + \nabla_{x_N} \theta_N) e^{-V_1/\sigma} dx_N.$$

By Lemma 1, for fixed x_0, x_1, \dots, x_{N-1} the tensor \mathcal{K}_{N-1} is uniformly positive definite over $x_{N-1} \in \mathbb{T}^d$. As a consequence, the operator defined in (57) is uniformly elliptic, with adjoint nullspace spanned by $Z_N(x_0, x_1, \dots, x_{N-1})$. Since the right hand side has mean zero, this implies that a solution ϕ_{N-1} exists. We recall that the corrector θ_{N-1} satisfies equation (15) with $k = 1$, that is

$$\nabla_{x_{N-1}} \cdot [\mathcal{K}_{N-1} (\nabla_{x_{N-1}} \theta_{N-1} + I)] = 0.$$

According to Remark 6, we can write ϕ_{N-1} as

$$\phi_{N-1} = \theta_{N-1} \cdot (\nabla_{x_{N-2}} \phi_{N-2} + \dots + \nabla_{x_0} \phi_0) + r_{N-1}^{(1)}(x_0, \dots, x_{N-2}),$$

for some $r_{N-1}^{(1)} \in \mathcal{H}_{N-1}^\perp$. Since (56) has been satisfied, it follows from Proposition 5 that there exists a unique decomposition of ϕ_{N+1} into

$$\phi_{N+1}(x_0, x_1, \dots, x_N) = \tilde{\phi}_{N+1}(x_0, x_1, \dots, x_N) + r_{N+1}^{(1)}(x_0, x_1, \dots, x_{N-1}),$$

where $\tilde{\phi}_{N+1} \in \mathcal{H}_N$ and for some $r_{N+1}^{(1)} \in \mathcal{H}_N^\perp$. For the sake of illustration we now consider the $O(\epsilon^{-(N-2)})$ equation in (47)

$$\mathcal{A}_{2N} \phi_{N+2} = - \sum_{k=0}^{N+1} \mathcal{A}_{N+k-2} \phi_k,$$

which, again by the Fredholm alternative, has a solution if and only if

$$\int (\mathcal{A}_{2N-1} \phi_{N+1} + \mathcal{A}_{2N-2} \phi_N + \dots + \mathcal{A}_{N-2} \phi_0) e^{-V/\sigma} dx_N = 0. \tag{58}$$

For $k = 1, \dots, N + 2$, we have

$$\begin{aligned} \mathcal{A}_{2N-k} \phi_{N-k+2} &= \sigma e^{V_1/\sigma} \left[\nabla_{x_{N-2}} \cdot (e^{-V_1/\sigma} \nabla_{x_{N-k+2}} \phi_{N-k+2}) + \nabla_{x_{N-1}} \cdot (e^{-V_1/\sigma} \nabla_{x_{N-k+1}} \phi_{N-k+2}) \right. \\ &\quad \left. + \nabla_{x_N} \cdot (e^{-V_1/\sigma} \nabla_{x_{N-k}} \phi_{N-k+2}) \right]. \end{aligned}$$

Fixing the variables x_0, \dots, x_{N-2} , we can rewrite (58) as an equation for $r_N^{(1)} = r_N^{(1)}(x_0, \dots, x_{N-1})$

$$\tilde{\mathcal{A}}_{2N-2} r_N^{(1)} := \nabla_{x_{N-1}} \cdot (Z_{N-1} \nabla_{x_{N-1}} r_N^{(1)}) = -RHS, \tag{59}$$

where

$$Z_{N-1} = \int e^{-V_1(x)/\sigma} dx_N,$$

and the *RHS* contains all the remaining terms. We note that all the functions of x_{N-1} in the *RHS* are known, so that all the remaining undetermined terms can be viewed as constants for fixed $x_0, \dots, x_{N-2} \in \mathbb{X}_{N-2}$. By the Fredholm alternative, a necessary and sufficient condition for a unique mean zero solution to exist to (59) is that the *RHS* has integral zero with respect to x_{N-1} , which is equivalent to:

$$\nabla_{N-2} \cdot \left(\int \int (\nabla_{x_N} \phi_N + \nabla_{x_{N-1}} \phi_{x_{N-1}} + \dots + \nabla_{x_0} \phi_0) e^{-V/\sigma} dx_N dx_{N-1} \right) = 0,$$

or equivalently:

$$\nabla_{x_{N-2}} \cdot (\mathcal{K}_{N-2} \nabla_{x_{N-2}} \phi_{N-2}) = -\nabla_{x_{N-2}} \cdot (\mathcal{K}_{N-2} (\nabla_{x_{N-3}} \phi_{N-3} + \dots + \nabla_{x_0} \phi_0)).$$

Once again, this implies that

$$\phi_{N-2} = \theta_{N-2} \cdot (\nabla_{x_{N-3}} \phi_{N-3} + \dots + \nabla_{x_0} \phi_0) + r_{N-2}^{(1)}(x_0, \dots, x_{N-3}),$$

where $r_{N-2}^{(1)} \in \mathcal{H}_{N-2}^\perp$ is unspecified. Since the compatibility condition holds, by Proposition 5, Eq. (59) has a solution, so that we can write

$$r_N^{(1)}(x_0, \dots, x_{N-1}) = \tilde{r}_N^{(1)}(x_0, \dots, x_{N-1}) + r_N^{(2)}(x_0, \dots, x_{N-2}),$$

where $\tilde{r}_N^{(1)} \in \mathcal{H}_{N-1}$ is the unique smooth solution of (59) and for some $r_N^{(2)} \in \mathcal{H}_{N-1}^\perp$.

We continue the proof by induction. Suppose that for some $k < N$, the functions $\phi_N, \dots, \phi_{N \pm (k-1)}$ have all been determined. We shall consider the case when k is even, noting that the k odd case follows *mutatis mutandis*.

From the previous steps, each term in

$$\phi_{N+k-2}, \phi_{N+k-4}, \dots, \phi_{N-k-2},$$

admits a decomposition such that in each case we can write:

$$\phi_{N+k-2i} = \tilde{\phi}_{N+k-2i} + r_{N+k-2i}^{(k/2-i)},$$

where

$$\tilde{\phi}_{N+k-2i} \in \mathcal{H}_{k/2-i},$$

has been uniquely specified, and the remainder term

$$r_{N+k-2i}^{(k/2-i)} \in \mathcal{H}_{k/2-i}^\perp,$$

remains to be determined. The $O(\epsilon^{N-k})$ equation is given by

$$\mathcal{A}_{2N} \phi_{N+k} + \mathcal{A}_{2N-1} \phi_{N+k-1} + \dots + \mathcal{A}_{N-k} \phi_0 = 0. \tag{60}$$

Following the example of the $O(\epsilon^{N-2})$ step, in descending order we successively apply the compatibility conditions which must be satisfied for the equations involving $r_{N+k}^{(1)}, \dots, r_{N-k-2}^{(k-1)}$ of the form

$$\tilde{\mathcal{A}}_{2N-2k-2i} r_{N+k-2i}^{(k/2-i)} = RHS, \tag{61}$$

where in (61), all terms dependent on the variable $x_{k/2-i}$ have been specified uniquely and where

$$\tilde{\mathcal{A}}_{2N-2k-2i} u = \nabla_{x_{N-k-i}} \cdot (Z_{N-k-i} \nabla_{x_{N-k-i}} u).$$

This results in (60) being integrated with respect to the variables $N, \dots, N - k + 1$. In particular, all terms $\mathcal{A}_{2N-j}\phi_{N+k-j}$ for $j = 0, \dots, k - 1$ will have integral zero, and thus vanish. The resulting equation is then

$$\int \dots \int (\mathcal{A}_{2N-k}\phi_N + \dots + \mathcal{A}_{N-k}\phi_0) e^{-V_1/\sigma} dx_N \dots dx_{N-k+1} = 0. \tag{62}$$

Moreover, since the function ϕ_{N-i} depends only on the variables x_0, \dots, x_{N-i} , then (62) must be of the form

$$\nabla_{x_{N-k}} \cdot \left(\int \dots \int (\nabla_{x_N}\phi_N + \dots + \nabla_{x_{N-1}}\phi_{N-1} + \dots + \nabla_{x_0}\phi_0) e^{-V/\sigma} dx_N \dots dx_{N-k+1} \right) = 0.$$

We now apply the inductive hypothesis to see that (to shorten the notations, we denote $dx_{N,\dots,N-k+1} := dx_N \dots dx_{N-k+1}$ etc)

$$\begin{aligned} & \int (\nabla_{x_N}\phi_N + \dots + \nabla_{x_0}\phi_0) e^{-V_1/\sigma} dx_{N,\dots,N-k+1} \\ &= \int \int (\nabla_{x_N}\theta_N + I) dx_N (\nabla_{x_{N-1}}\phi_{N-1} + \dots + \nabla_{x_0}\phi_0) e^{-V_1/\sigma} dx_{N-1,\dots,N-k+1} \\ &= \int \int \int (\nabla_{x_N}\theta_N + I) dx_N (\nabla_{x_{N-1}}\theta_{N-1} + I) dx_{N-1} \\ & \quad \times (\nabla_{x_{N-2}}\phi_{N-2} + \dots + \nabla_{x_0}\phi_0) e^{-V_1/\sigma} dx_{N-2,\dots,N-k+1} \\ & \quad \vdots \\ &= \mathcal{K}_{N-k+1} (\nabla_{x_{N-k}}\phi_{N-k} + \dots + \nabla_{x_0}\phi_0). \end{aligned}$$

Thus, the compatibility condition for the $O(\epsilon^{N-k})$ equation reduces to the elliptic PDE

$$\nabla_{x_{N-k}} \cdot (\mathcal{K}_{N-k}\nabla_{x_{N-k}}\phi_{N-k}) = -\nabla_{x_{N-k}} \cdot (\mathcal{K}_{N-k}(\nabla_{x_{N-k-1}}\phi_{N-k-1} + \dots + \nabla_{x_0}\phi_0)) = 0,$$

so that ϕ_{N-k} can be written as

$$\phi_{N-k} = \theta_{N-k} (\nabla_{x_{N-k-1}}\phi_{N-k-1} + \dots + \nabla_{x_0}\phi_0) + r_{N-k}^{(1)}, \tag{63}$$

where $r_{N-k}^{(1)}$ is an element of \mathcal{H}_{N-k}^\perp , which is yet to be determined. Moreover, each remainder term $r_{N+k-2i}^{(k/2-i)}$ can be further decomposed as

$$r_{N+k-2i}^{(k/2-i)} = \tilde{r}_{N+k-2i}^{(k/2-i)} + r_{N+k-2i}^{(k/2-i+1)},$$

where

$$\tilde{r}_{N+k-2i}^{(k/2-i)} \in \mathcal{H}_{k/2-i+1},$$

is uniquely determined and

$$r_{N+k-2i}^{(k/2-i+1)} \in \mathcal{H}_{k/2-i+1}^\perp,$$

is still unspecified. Continuing the above procedure inductively, starting from a smooth function ϕ_0 we construct a series of correctors $\phi_1, \dots, \phi_{2N-1}$.

We now consider the final Eq. (46d). Arguing as before, we note that we can rewrite (46d) as

$$\mathcal{A}_{2N}\phi_{2N} + \dots + \mathcal{A}_{N+1}\phi_{N+1} = F(x) - \sum_{i=1}^N \mathcal{L}_i\phi_i. \tag{64}$$

A necessary and sufficient condition for ϕ_{2N} to have a solution is that

$$\begin{aligned} & \int_{\mathbb{T}^d} (\mathcal{A}_{2N-1}\phi_{2N-1} + \dots + \mathcal{A}_{N+1}\phi_{N+1}) e^{-V_1/\sigma} dx_N \\ &= \int_{\mathbb{T}^d} \left(F(x) - \sum_{i=1}^N \mathcal{L}_i \phi_i \right) e^{-V_1/\sigma} dx_N. \end{aligned} \tag{65}$$

At this point, the remainder terms will be of the form

$$r_{2N-2}^{(1)}, r_{2N-4}^{(2)}, \dots, r_{2N-2k}^{(k)}, \dots, r_2^{(1)},$$

such that $r_{2N-2i}^{(i)} \in \mathcal{H}_i^\perp$, is unspecified. Starting from $r_{2N-2}^{(1)}$ a necessary and sufficient condition for the remainder $r_{2N-2i}^{(i)}$ to exist is that the integral of the equation with respect to dx_{N-i} vanishes, i.e.

$$\begin{aligned} F(x)Z(x) &= \int_{(\mathbb{T}^d)^N} (\mathcal{A}_{2N-1}\phi_{2N-1} + \dots + \mathcal{A}_{N+1}\phi_{N+1}) e^{-V_1/\sigma} dx_N dx_{N-1} \dots dx_1 \\ &+ \int_{(\mathbb{T}^d)^N} (\mathcal{L}_N \phi_N + \dots + \mathcal{L}_1 \phi_1) e^{-V_1/\sigma} dx_N dx_{N-1} \dots dx_1, \end{aligned} \tag{66}$$

where

$$Z(x) = \int_{\mathbb{T}^d} \dots \int_{\mathbb{T}^d} e^{-V_1/\sigma} dx_N \dots dx_1.$$

As above, after simplification, (66) becomes

$$\nabla_{x_0} \cdot (\nabla_{x_N} \phi_N + \dots + \nabla_{x_0} \phi_0) = Z(x)F(x),$$

which can be written as

$$\frac{\sigma}{Z(x)} \nabla_{x_0} \cdot \left(\int_{(\mathbb{T}^d)^N} (I + \nabla_{x_N} \theta_N) \dots (I + \nabla_{x_1} \theta_1) e^{-V/\sigma} dx_N \dots dx_1 \nabla_{x_0} \phi_0 \right) = F(x),$$

or more compactly

$$F(x) = \frac{\sigma}{Z(x)} \nabla_{x_0} \cdot (\mathcal{K}_1(x) \nabla_{x_0} \phi_0(x)),$$

where the terms in the right hand side have been specified and are unique. Thus, the $O(1)$ equation (66) provides a unique expression for $F(x)$. Moreover, for each $i = 1, \dots, N - 1$, there exists a smooth unique solution $r_{2N-2i}^{(i)} \in \mathcal{H}_{i-1}$ and $\phi_{2N} \in \mathcal{H}_N$ by Proposition 5.

Note that we have not uniquely identified the functions ϕ_1, \dots, ϕ_{2N} , since after the above N steps there will be remainder terms which are still unspecified. However, conditions (47a)–(47c) will hold for any choice of remainder terms which are still unspecified. In particular, we can set all the remaining unspecified remainder terms to 0. Moreover, every Poisson equation we have solved in the above steps has been of the form:

$$\mathcal{S}_k u(x_0, \dots, x_k) = a(x_0, \dots, x_k) \cdot \nabla_{x_0} \phi_0(x_0) + A(x_0, \dots, x_k) : \nabla_{x_0}^2 \phi_0(x_0),$$

where \mathcal{S}_k is of the form (38), and a and A are uniformly bounded with bounded derivatives. In particular, from the remark following Proposition 5 the pointwise estimates (48) hold. \square

Remark 7 Note that we do not have an explicit formula for the test functions, for $i = 1, \dots, N$. However, by applying (63) recursively one can obtain an explicit expression for the gradient of ϕ_i in terms of the correctors θ_i :

$$\nabla_{x_i} \phi_i = \nabla_{x_i} \theta_i (I + \nabla_{x_{i-1}} \theta_{i-1}) \cdots \cdots (I + \nabla_{x_1} \theta_1) \nabla_{x_0} \phi_0.$$

Since these are the only terms required for the calculation of the homogenized diffusion tensor we thus obtain an explicit characterisation of the effective coefficients.

5.2 Tightness of Measures

In this section we establish the weak compactness of the family of measures corresponding to $\{X_t^\epsilon : 0 \leq t \leq T\}_{0 < \epsilon \leq 1}$ in $C([0, T]; \mathbb{R}^d)$ by establishing tightness. Following [43], we verify the following two conditions which are a slight modification of the sufficient conditions stated in [9, Theorem 8.3].

Lemma 3 *The collection $\{X_t^\epsilon : 0 \leq t \leq T\}_{0 < \epsilon \leq 1}$ is relatively compact in $C([0, T]; \mathbb{R}^d)$ if it satisfies:*

1. For all $\delta > 0$, there exists $M > 0$ such that

$$\mathbb{P} \left(\sup_{0 \leq t \leq T} |X_t^\epsilon| > M \right) \leq \delta, \quad 0 < \epsilon \leq 1.$$

2. For any $\delta > 0, M > 0$, there exists ϵ_0 and γ such that

$$\gamma^{-1} \sup_{0 < \epsilon < \epsilon_0} \sup_{0 \leq t_0 \leq T} \mathbb{P} \left(\sup_{t \in [t_0, t_0 + \gamma]} |X_t^\epsilon - X_{t_0}^\epsilon| \geq \delta; \sup_{0 \leq s \leq T} |X_s^\epsilon| \leq M \right) \leq \delta.$$

To verify condition 3 we follow the approach of [43] and consider a test function of the form $\phi_0(x) = \log(1 + |x|^2)$. The motivation for this choice is that while $\phi_0(x)$ is increasing, we have that

$$\sum_{l=1}^3 (1 + |x|)^l |\nabla_x^l \phi_0(x)|_F \leq C, \tag{67}$$

where $|\cdot|_F$ denotes the Frobenius norm. Let $\phi_1, \dots, \phi_{2N-1}$ be the first $2N - 1$ test functions constructed in Proposition 6. Consider the test function

$$\begin{aligned} \phi^\epsilon(x) &= \phi_0(x) + \epsilon \phi_1(x, x/\epsilon) + \dots + \epsilon^N \phi_N(x, x/\epsilon, \dots, x/\epsilon^N) \\ &\quad + \epsilon^{N+1} \phi_{N+1}(x, x/\epsilon, \dots, x/\epsilon^N) + \dots + \epsilon^{2N-1} \phi_{2N-1}(x, x/\epsilon, \dots, x/\epsilon^N). \end{aligned} \tag{68}$$

Applying Itô’s formula, we have that

$$\phi^\epsilon(X_t^\epsilon) = \phi^\epsilon(x) + \int_0^t G(X_s^\epsilon) ds + \sqrt{2\sigma} \sum_{i=0}^N \sum_{j=0}^{2N-1} \epsilon^{j-i} \int_0^t \nabla_{x_i} \phi_j dW_s,$$

where $G(x)$ is a smooth function consisting of terms of the form:

$$\epsilon^{k-(i+j)} e^{V/\sigma} \nabla_{x_i} \cdot \left(e^{-V/\sigma} \sigma \nabla_{x_j} \phi_k \right) (x, x/\epsilon, \dots, x/\epsilon^N), \tag{69}$$

where $k \geq i + j$, by construction of the test functions. Moreover, $\nabla_{x_i} \phi_j = 0$ for $j < i$. To obtain relative compactness we need to individually control the terms arising in the drift. More specifically, we must show that the terms

$$\mathbb{E} \sup_{0 \leq t \leq T} \int_0^t |e^{V/\sigma} \nabla_{x_i} \cdot (e^{-V/\sigma} \sigma \nabla_{x_j} \phi_k)(X_s^\epsilon, X_s^\epsilon/\epsilon, \dots, X_s^\epsilon/\epsilon^N) ds|, \tag{70}$$

$$\mathbb{E} \left| \sup_{0 \leq t \leq T} \int_0^t \nabla_{x_j} \phi_k(X_s^\epsilon, X_s^\epsilon/\epsilon, \dots, X_s^\epsilon/\epsilon^N) dW_s \right|^2, \tag{71}$$

and

$$\sup_{0 \leq t \leq T} |\phi_j(X_t^\epsilon)|. \tag{72}$$

are bounded uniformly with respect to $\epsilon \in (0, 1]$. Terms of the type (70) can be bounded above by:

$$\mathbb{E} \sup_{0 \leq t \leq T} \int_0^t \left| (\nabla_{x_i} V \cdot \nabla_{x_j} \phi_k)(X_s^\epsilon, \dots, X_s^\epsilon/\epsilon^N) \right| + \left| \sigma \nabla_{x_i} \cdot \nabla_{x_j} \phi_k(X_s^\epsilon, \dots, X_s^\epsilon/\epsilon^N) \right| ds.$$

If $i > 0$, then $\nabla_{x_i} V$ is uniformly bounded, and so the above expectation is bounded above by

$$\begin{aligned} & C \mathbb{E} \int_0^T \left| \nabla_{x_j} \phi_k(X_s^\epsilon, \dots, X_s^\epsilon/\epsilon^N) \right| + \left| \nabla_{x_i} \cdot \nabla_{x_j} \phi_k(X_s^\epsilon, \dots, X_s^\epsilon/\epsilon^N) \right| ds \\ & \leq C \mathbb{E} \int_0^T \sum_{m=1}^3 |\nabla_{x_0}^m \phi_0(X_s^\epsilon)|_F ds \leq KT, \end{aligned}$$

using (67), for some constant $K > 0$ independent of ϵ . For the case when $i = 0$, an additional term arises from the derivative $\nabla_{x_0} V_0$ and we obtain an upper bound of the form

$$\begin{aligned} & \mathbb{E} \int_0^T \sum_{m=1}^3 |\nabla_{x_0}^m \phi_0(X_t^\epsilon)|_F (1 + |\nabla_{x_0} V_0(X_t^\epsilon)|) dt \\ & \leq \mathbb{E} \int_0^T \sum_{m=1}^3 |\nabla_{x_0}^m \phi_0(X_t^\epsilon)|_F (1 + \|\nabla V_0\|_{L^\infty} |X_t^\epsilon|) dt, \end{aligned} \tag{73}$$

and which is bounded by Assumption 1 and (67). For (71), we have

$$\begin{aligned} \mathbb{E} \left| \sup_{0 \leq t \leq T} \int_0^t \nabla_{x_j} \phi_k(X_s^\epsilon, X_s^\epsilon/\epsilon, \dots, X_s^\epsilon/\epsilon^N) dW_s \right|^2 & \leq 4 \mathbb{E} \int_0^T |\nabla_{x_j} \phi_k(X_s^\epsilon, X_s^\epsilon/\epsilon, \dots, X_s^\epsilon/\epsilon^N)|^2 ds \\ & \leq C \mathbb{E} \int_0^T \sum_{m=1}^3 |\nabla_{x_0}^m \phi_0(X_s^\epsilon)|_F ds, \end{aligned}$$

which is again bounded. Terms of the type (72) follow in a similar manner. Condition 3 then follows by an application of Markov’s inequality.

To prove Condition 3, we set $\phi_0(x) = x$ and let $\phi_1, \dots, \phi_{2N-1}$ be the test functions which exist by Proposition 6. Applying Itô’s formula to the corresponding multiscale test function (68), so that for $t_0 \in [0, T]$ fixed,

$$X_t^\epsilon - X_{t_0}^\epsilon = \int_{t_0}^t G ds + \sqrt{2\sigma} \sum_{i=0}^N \sum_{j=0}^{2N-1} \epsilon^{j-i} \int_{t_0}^t \nabla_{x_i} \phi_j dW_s, \tag{74}$$

where G is of the form given in (69). Let $M > 0$, and let

$$\tau_M^\epsilon = \inf\{t \geq 0; |X_t^\epsilon| > M\}. \tag{75}$$

Following [43], it is sufficient to show that

$$\mathbb{E} \left[\sup_{t_0 \leq t \leq T} \int_{t_0 \wedge \tau_M^\epsilon}^{t \wedge \tau_M^\epsilon} \left| e^{V/\sigma} \nabla_{x_i} \cdot \left(e^{-V/\sigma} \nabla_j \phi_k \right) (X_s^\epsilon, X_s^\epsilon/\epsilon, \dots, X_s^\epsilon/\epsilon^N) ds \right|^{1+\nu} \right] < \infty, \tag{76}$$

and

$$\mathbb{E} \left(\sup_{t_0 \leq t \leq t_0 + \gamma} \left| \int_{t_0 \wedge \tau_M^\epsilon}^{t \wedge \tau_M^\epsilon} \nabla_{x_i} \phi_j (X_s^\epsilon, X_s^\epsilon/\epsilon, \dots, X_s^\epsilon/\epsilon^N) dW_s \right|^{2+2\nu} \right) < \infty, \tag{77}$$

for some fixed $\nu > 0$. For (76), when $i > 0$, the term $\nabla_{x_i} V$ is uniformly bounded. Moreover, since $\nabla \phi_0$ is bounded, so are the test functions $\phi_1, \dots, \phi_{2N+1}$. Therefore, by Jensen’s inequality one obtains a bound of the form

$$\begin{aligned} C\gamma^\nu \mathbb{E} \int_{t_0}^{t_0 + \gamma} \left| e^{V/\sigma} \nabla_{x_i} \cdot \left(e^{-V/\sigma} \nabla_j \phi_k \right) (X_s^\epsilon, X_s^\epsilon/\epsilon, \dots, X_s^\epsilon/\epsilon^N) \right|^{1+\nu} ds \\ \leq C\gamma^\nu \int_{t_0}^{t_0 + \gamma} |K|^{1+\nu} ds \leq K'\gamma^{1+\nu}. \end{aligned}$$

When $i = 0$, we must control terms involving $\nabla_{x_0} V_0$ of the form,

$$\mathbb{E} \left[\sup_{t_0 \leq t \leq t_0 + \gamma} \int_{t_0 \wedge \tau_M^\epsilon}^{t \wedge \tau_M^\epsilon} |\nabla V_0 \cdot \nabla_{x_j} \phi_k|^{1+\nu} ds \right],$$

where τ_M^ϵ is given by (75). However, applying Jensen’s inequality,

$$\begin{aligned} \mathbb{E} \left[\sup_{t_0 \leq t \leq t_0 + \gamma} \int_{t_0 \wedge \tau_M^\epsilon}^{t \wedge \tau_M^\epsilon} |\nabla V_0 \cdot \nabla_{x_j} \phi_k|^{1+\nu} ds \right] &\leq C\gamma^\nu \int_{t_0 \wedge \tau_M^\epsilon}^{(t_0 + \gamma) \wedge \tau_M^\epsilon} \mathbb{E} |\nabla V_0 \cdot \nabla_{x_j} \phi_k|^{1+\nu} ds \\ &\leq C\gamma^\nu \int_{t_0 \wedge \tau_M^\epsilon}^{(t_0 + \gamma) \wedge \tau_M^\epsilon} \mathbb{E} |\nabla V_0(X_s^\epsilon)|^{1+\nu} ds \\ &\leq C\gamma^\nu \|\nabla^2 V_0\|_\infty^{1+\nu} \int_{t_0 \wedge \tau_M^\epsilon}^{(t_0 + \gamma) \wedge \tau_M^\epsilon} \mathbb{E} |X_s^\epsilon|^{1+\nu} ds \\ &\leq CM\gamma^{1+\nu} \|\nabla^2 V_0\|_{L^\infty}^{1+\nu}, \end{aligned} \tag{78}$$

as required. Similarly, to establish (77) we follow a similar argument, first using the Burkholder–Gundy–Davis inequality to obtain:

$$\begin{aligned} \mathbb{E} \left(\sup_{t_0 \leq t \leq t_0 + \gamma} \int_{t_0}^t |\nabla_{x_i} \phi_j dW_s|^{2+2\nu} \right) &\leq \mathbb{E} \left(\int_{t_0}^{t_0 + \gamma} |\nabla_{x_i} \phi_j|^2 ds \right)^{1+\nu} \\ &\leq \gamma^\nu \int_{t_0}^{t_0 + \gamma} \mathbb{E} |\nabla_{x_i} \phi_j|^{2+2\nu} ds \\ &\leq C\gamma^{1+\nu}. \end{aligned}$$

We note that Assumption 1 (3) is only used to obtain the bounds (73) and (78). A straightforward application of Markov’s inequality then completes the proof of condition

3. It follows from Prokhorov’s theorem that the family $\{X_t^\epsilon; t \in [0, T]\}_{0 < \epsilon \leq 1}$ is relatively compact in the topology of weak convergence of stochastic processes taking paths in $C([0, T]; \mathbb{R}^d)$. In particular, there exists a process X^0 whose paths lie in $C([0, T]; \mathbb{R}^d)$ such that $\{X^{\epsilon_n}; t \in [0, T]\} \Rightarrow \{X^0; t \in [0, T]\}$ along a subsequence ϵ_n .

5.3 Identifying the Weak Limit

In this section we uniquely identify any limit point of the set $\{X_t^\epsilon; t \in [0, T]\}_{0 < \epsilon \leq 1}$. Given $\phi_0 \in C_c^\infty(\mathbb{R}^d)$ define ϕ^ϵ to be

$$\begin{aligned} \phi^\epsilon(x) &= \phi_0(x) + \epsilon \phi_1(x/\epsilon) + \dots + \epsilon^N \phi_N(x, x/\epsilon, \dots, x/\epsilon^N) \\ &+ \dots + \epsilon^{2N} \phi_{2N}(x, x/\epsilon, \dots, x/\epsilon^N), \end{aligned}$$

where ϕ_1, \dots, ϕ_N are the test functions obtained from Proposition 6. Since each test function is smooth, we can apply Itô’s formula to $\phi^\epsilon(X_t^\epsilon)$ to obtain

$$\mathbb{E} \left[\phi^\epsilon(X_t^\epsilon) - \int_s^t \mathcal{L}^\epsilon \phi^\epsilon(X_u) du \mid \mathcal{F}_s \right] = \phi^\epsilon(X_s^\epsilon). \tag{79}$$

We can now use (45) to decompose $\mathcal{L}\phi^\epsilon$ into an $O(1)$ term and remainder terms which vanish as $\epsilon \rightarrow 0$. Collecting together $O(\epsilon)$ terms we obtain

$$\mathbb{E} \left[\phi_0(X_t^\epsilon) - \int_s^t \frac{\sigma}{Z(X_u^\epsilon)} \nabla_{x_0} \cdot (Z(X_u^\epsilon) \mathcal{M}(X_u^\epsilon) \nabla \phi_0(X_u^\epsilon)) du + \epsilon R_\epsilon \mid \mathcal{F}_s \right] = \phi_0(X_s^\epsilon),$$

where R_ϵ is a remainder term which is bounded in $L^2(\mu^\epsilon)$ uniformly with respect to ϵ , and where the homogenized diffusion tensor $\mathcal{M}(x)$ is defined in Theorem 1. Taking $\epsilon \rightarrow 0$ we see that any limit point is a solution of the martingale problem

$$\mathbb{E} \left[\phi_0(X_t^0) - \int_s^t \frac{\sigma}{Z(X_u^0)} \nabla_{x_0} \cdot (Z(X_u^0) \mathcal{M}(X_u^0) \nabla \phi_0(X_u^0)) du \mid \mathcal{F}_s \right] = \phi_0(X_s^0).$$

This implies that X^0 is a solution to the martingale problem for \mathcal{L}^0 given by

$$\mathcal{L}_0 f(x) = \frac{\sigma}{Z(x)} \nabla \cdot (Z(x) \mathcal{M}(x) \nabla f(x)).$$

From Lemma 1, the matrix $\mathcal{M}(x)$ is smooth, strictly positive definite and has bounded derivatives. Moreover,

$$\begin{aligned} Z(x) &= \int_{\mathbb{T}^d} \dots \int_{\mathbb{T}^d} e^{-V(x, x_1, \dots, x_N)/\sigma} dx_1 \dots dx_N \\ &= e^{-V_0(x)/\sigma} \int_{\mathbb{T}^d} \dots \int_{\mathbb{T}^d} e^{-V_1(x, x_1, \dots, x_N)/\sigma} dx_1 \dots dx_N, \end{aligned}$$

where the term in the integral is uniformly bounded. It follows from Assumption 1, that for some $C > 0$,

$$|\mathcal{M}(x) \nabla \Psi(x)| \leq C(1 + |x|), \quad \forall x \in \mathbb{R}^d,$$

where $\Psi = -\log Z$. Therefore, the conditions of the Stroock-Varadhan theorem [51, Theorem 24.1] holds, and therefore the martingale problem for \mathcal{L}^0 possesses a unique solution. Thus X^0 is the unique (in the weak sense) limit point of the family $\{X^\epsilon\}_{0 < \epsilon \leq 1}$. Moreover, by [51, Theorem 20.1], the process $\{X_t^0; t \in [0, T]\}$ will be the unique solution of the SDE (18), completing the proof.

6 Further Discussion and Outlook

In this paper, we have shown the convergence of the multi-scale diffusion process (8) to the homogenized (effective) diffusion process (18), as well as the convergence of the corresponding equilibrium measures. We have employed the classical martingale approach based on a suitable construction of test functions and analysis of the related Poisson equations. A notable feature is that the effective (macroscopic) process is a multiplicative diffusion process where the diffusion tensor depends on the macroscopic variable, whereas the noise in the microscopic dynamics is additive. This is due to the full coupling between the macroscopic and the microscopic scales. As discussed in the introduction, both processes are reversible diffusion processes satisfying the detailed balance condition. Therefore, according to [1], the corresponding Fokker Planck equations at all scales are Wasserstein gradient flows for the corresponding free energy functionals [30]. Thus, the rigorous analysis presented in this work leads to the conclusion that the Wasserstein gradient flow structure is preserved under coarse-graining. This raises the interesting question whether coarse-graining and, in particular, homogenization can be studied within the framework of evolutionary Gamma convergence [4, 16, 35, 52]. Another interesting question is obtaining quantitative rates of convergence [17] and also understanding the effect of coarse-graining on the Poincaré and logarithmic Sobolev inequality constants, using the methodology of two-scale convergence [24, 41]. We will return to these questions in future work.

Acknowledgements The authors thank S. Kalliadas and M. Pradas for useful discussions. They also thank B. Zegarinski for useful discussions and for pointing out Ref. [25]. The authors are also very grateful to the anonymous referees whose comments have greatly improved the content of this work. GAP and ABD acknowledge financial support by the Engineering and Physical Sciences Research Council of the UK through Grants Nos. EP/J009636, EP/L020564, EP/L024926, EP/P031587/1 and EP/L025159. GAP was partially funded by JPMorgan Chase & Co under J.P. Morgan A.I. Research Awards in 2019 and 2021. ABD was supported by Wave 1 of The UKRI Strategic Priorities Fund under the EPSRC Grant EP/W006022/1, particularly the Ecosystems of Digital Twins theme within that grant and The Alan Turing Institute. MHD was supported by EPSRC Grants EP/W008041/1 and EP/V038516/1.

Data Availability Data sharing not applicable to this article as no datasets were generated or analysed during the current study.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

1. Adams, S., Dirr, N., Peletier, M., Zimmer, J.: Large deviations and gradient flows. *Philos. Trans. R. Soc. A Math. Phys. Eng. Sci.* **371**, 20120341 (2013)
2. Ané, C., et al.: *Sur les inégalités de Sobolev Logarithmiques*. Société mathématique de France, Paris (2000)
3. Ansari, A.: Mean first passage time solution of the Smoluchowski equation: application to relaxation dynamics in myoglobin. *J. Chem. Phys.* **112**, 2516–2522 (2000)
4. Arrich, S., Mielke, A., Peletier, M.A., Savaré, G., Veneroni, M.: Passing to the limit in a Wasserstein gradient flow: from diffusion to reaction. *Calc. Var. Part. Differ. Equ.* **44**, 419–454 (2012)

5. Bakry, D., Gentil, I., Ledoux, M.: *Analysis and Geometry of Markov Diffusion Operators*, vol. 348. Springer, New York (2013)
6. Banerjee, S., Biswas, R., Seki, K., Bagchi, B.: Diffusion in a rough potential revisited. Preprint at <http://arxiv.org/abs/1409.4581> (2014)
7. Ben Arous, G., Owhadi, H.: Multiscale homogenization with bounded ratios and anomalous slow diffusion. *Commun. Pure Appl. Math.* **56**, 80–113 (2003)
8. Bensoussan, A., Lions, J., Papanicolaou, G.: *Asymptotic Analysis for Periodic Structures*, vol. 5. North Holland, Amsterdam (1978)
9. Billingsley, P.: *Probability and Measure*. Wiley, Hoboken (2008)
10. Bryngelson, J.D., Wolynes, P.G.: Spin glasses and the statistical mechanics of protein folding. *Proc. Natl. Acad. Sci.* **84**, 7524–7528 (1987)
11. Bryngelson, J.D., Onuchic, J.N., Socci, N.D., Wolynes, P.G.: Funnels, pathways, and the energy landscape of protein folding: a synthesis, proteins: structure. *Funct. Bioinform.* **21**, 167–195 (1995)
12. Cioranescu, D., Donato, P.: *Introduction to Homogenization*. Oxford University Press, Oxford (2000)
13. Dean, D.S., Gupta, S., Oshanin, G., Rosso, A., Schehr, G.: Diffusion in periodic, correlated random forcing landscapes. *J. Phys. A Math. Theor.* **47**, 372001 (2014)
14. Delgadino, M.G., Gvalani, R.S., Pavliotis, G.A.: On the diffusive-mean field limit for weakly interacting diffusions exhibiting phase transitions. *Arch. Ration. Mech. Anal.* **241**, 91–148 (2021). <https://doi.org/10.1007/s00205-021-01648-1>
15. Duncan, A.B., Kalliadasis, S., Pavliotis, G.A., Pradas, M.: Noise-induced transitions in rugged energy landscapes. *Phys. Rev. E* **94**, 032107 (2016)
16. Duong, M.H., Lamacz, A., Peletier, M.A., Sharma, U.: Variational approach to coarse-graining of generalized gradient flows. *Calc. Var. Part. Differ. Equ.* **56**, 100 (2017)
17. Duong, M.H., Lamacz, A., Peletier, M.A., Schlichting, A., Sharma, U.: Quantification of coarse-graining error in langevin and overdamped langevin dynamics. *Nonlinearity* **31**, 4517–4566 (2018)
18. Evans, L.C.: *Partial Differential Equations*, Graduate Studies in Mathematics, vol. 19. American Mathematical Society, Providence (1998)
19. Evans, L.C., Tabrizian, P.R.: Asymptotics for scaled Kramers–Smoluchowski equations. *SIAM J. Math. Anal.* **48**, 2944–2961 (2016)
20. Gardiner, C.: *Stochastic Methods*, Springer Series in Synergetics, 4th edn. Springer, Berlin (2009)
21. Gilbarg, D., Trudinger, N.S.: *Elliptic Partial Differential Equations of Second Order*. Springer, Berlin (2015)
22. Gomes, S.N., Pavliotis, G.A.: Mean field limits for interacting diffusions in a two-scale potential. *J. Nonlinear Sci.* **28**, 905–941 (2018)
23. Gomes, S.N., Kalliadasis, S., Pavliotis, G.A., Yatsyshin, P.: Dynamics of the Desai–Zwanzig model in multiwell and random energy landscapes. *Phys. Rev. E* **99**, 032109 (2019)
24. Grunewald, N., Otto, F., Villani, C., Westdickenberg, M.G.: A two-scale approach to logarithmic Sobolev inequalities and the hydrodynamic limit. *Ann. de l’Inst. Henri Poincaré Probab. Stat.* **45**, 302–351 (2009)
25. Hebisch, W., Zegarliński, B.: Coercive inequalities on metric measure spaces. *J. Funct. Anal.* **258**, 814–851 (2010)
26. Holley, R., Stroock, D.: Logarithmic Sobolev inequalities and stochastic Ising models. *J. Stat. Phys.* **46**, 1159–1194 (1987)
27. Hütter, M., Öttinger, H.C.: Fluctuation–dissipation theorem, kinetic stochastic integral and efficient simulations. *J. Chem. Soc. Faraday Trans.* **94**, 1403–1405 (1998)
28. Hyeon, C.: Can energy landscape roughness of proteins and RNA be measured by using mechanical unfolding experiments? *Proc. Natl. Acad. Sci.* **100**, 10249–10253 (2003)
29. Jikov, V.V., Kozlov, S.M., Oleinik, O.A.: *Homogenization of Differential Operators and Integral Functionals*. Springer, New York (2012)
30. Jordan, R., Kinderlehrer, D., Otto, F.: The variational formulation of the Fokker–Planck equation. *SIAM J. Math. Anal.* **29**, 1–17 (1998)
31. Komorowski, T., Landim, C., Olla, S.: *Fluctuations in Markov Processes: Time Symmetry and Martingale Approximation*, vol. 345. Springer, New York (2012)
32. Landim, C., Seo, I.: Metastability of one-dimensional, non-reversible diffusions with periodic boundary conditions. *Ann. Inst. Henri Poincaré Probab. Stat.* **55**, 1850–1889 (2019)
33. Lifson, S., Jackson, J.L.: On the self-diffusion of ions in a polyelectrolyte solution. *J. Chem. Phys.* **36**, 2410–2414 (1962)
34. López-Alamilla, N.J., Jack, M.W., Challis, K.J.: Enhanced diffusion and the eigenvalue band structure of Brownian motion in tilted periodic potentials. *Phys. Rev. E* **102**, 042405 (2020)
35. Mielke, A.: *On Evolutionary Γ -Convergence for Gradient Systems*, pp. 187–249. Springer, Cham (2016)
36. Milton, G.W.: *The theory of composites*. *Mater. Technol.* **117**, 483–493 (1995)

37. Mondal, D., Ghosh, P.K., Ray, D.S.: Noise-induced transport in a rough ratchet potential. *J. Chem. Phys.* **130**, 074703 (2009)
38. Muckenhoupt, B.: Hardy's inequality with weights. *Stud. Math.* **44**, 31–38 (1972)
39. Müller, K.: Reaction paths on multidimensional energy hypersurfaces. *Angew. Chem. Int. Ed. Engl.* **19**, 1–13 (1980)
40. Onuchic, J.N., Luthey-Schulten, Z., Wolynes, P.G.: Theory of protein folding: the energy landscape perspective. *Annu. Rev. Phys. Chem.* **48**, 545–600 (1997)
41. Otto, F., Reznikoff, M.G.: A new criterion for the logarithmic Sobolev inequality and two applications. *J. Funct. Anal.* **243**, 121–157 (2007)
42. Papanicolaou, G.C., Stroock, D., Varadhan, S.R.S.: Martingale approach to some limit theorems. In: *Duke Turbulence Conference* (Duke Univ., Durham, NC, 1976), vol. 6 (1977)
43. Pardoux, È., Veretennikov, A.Y.: On the Poisson equation and diffusion approximation. I. *Ann. Probab.* **29**, 1061–1085 (2001)
44. Pardoux, È., Veretennikov, A.Y.: On Poisson equation and diffusion approximation. II. *Ann. Probab.* **31**, 1166–1192 (2003)
45. Pavliotis, G.A.: *Stochastic Processes and Applications*, Vol. 60 of *Texts in Applied Mathematics*. Springer, New York (2014)
46. Pavliotis, G.A., Stuart, A.M.: *Multiscale Methods: Averaging and Homogenization*. Springer, New York (2008)
47. Peletier, M.A., Schlottke, M.C.: Gamma-convergence of a gradient-flow structure to a non-gradient-flow structure. <https://doi.org/10.48550/ARXIV.2105.03401> (2021)
48. Peletier, M.A., Savaré, G., Veneroni, M.: From diffusion to reaction via γ -convergence. *SIAM J. Math. Anal.* **42**, 1805–1825 (2010)
49. Ren, W., Vanden-Eijnden, E.: Probing multi-scale energy landscapes using the string method. Preprint at <http://arXiv.org/0205528> (2002)
50. Rezakhanlou, F., Seo, I.: Scaling limit of small random perturbation of dynamical systems. <https://doi.org/10.48550/ARXIV.1812.02069> (2018)
51. Rogers, L.C.G., Williams, D.: *Diffusions, Markov Processes and Martingales: Volume 2, Itô Calculus*, vol. 2. Cambridge University Press, Cambridge (2000)
52. Sandier, E., Serfaty, S.: Gamma-convergence of gradient flows with applications to Ginzburg–Landau. *Commun. Pure Appl. Math.* **57**, 1627–1672 (2004)
53. Saven, J.G., Wang, J., Wolynes, P.G.: Kinetics of protein folding: the dynamics of globally connected rough energy landscapes with biases. *J. Chem. Phys.* **101**, 11037–11043 (1994)
54. Seo, I., Tabrizian, P.: Asymptotics for scaled Kramers–Smoluchowski equations in several dimensions with general potentials. *Calc. Var. Part. Differ. Equ.* **59**, 11 (2019)
55. Veretennikov, A.Y.: On Sobolev solutions of Poisson equations in \mathbb{R}^d with a parameter. *J. Math. Sci. N. Y.* **179**, 48–79 (2011)
56. Zwanzig, R.: Diffusion in a rough potential. *Proc. Natl. Acad. Sci.* **85**, 2029–2030 (1988)

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.