# UNIVERSITY BIRMINGHAM University of Birmingham Research at Birmingham

# Seasonal variation of serum potassium and related prescription pattern

Thayakaran, Rasiah; Hotham, Richard; Gokhale, Krishna; Adderley, Nicola; Chandan, Joht; Nirantharakumar, Krishnarajah

DOI: 10.1136/jcp-2023-208759

License: Creative Commons: Attribution-NonCommercial (CC BY-NC)

Document Version Publisher's PDF, also known as Version of record

#### Citation for published version (Harvard):

Thayakaran, R, Hotham, R, Gokhale, K, Adderley, N, Chandan, J & Nirantharakumar, K 2023, 'Seasonal variation of serum potassium and related prescription pattern: An ecological time series', *Journal of Clinical Pathology*. https://doi.org/10.1136/jcp-2023-208759

Link to publication on Research at Birmingham portal

#### **General rights**

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

•Users may freely distribute the URL that is used to identify this publication.

•Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.

•User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?) •Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

#### Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.



# Seasonal variation of serum potassium and related prescription pattern: an ecological time series

Rasiah Thayakaran, Richard Hotham, Krishna M Gokhale, Nicola J Adderley, Joht Singh Chandan 💿 , Krishnaraiah Nirantharakumar

Institute of Applied Health Research, University of Birmingham, Birmingham, UK

#### Correspondence to

Dr Joht Singh Chandan. University of Birmingham, Birmingham, UK; j.s.chandan. 1@bham.ac.uk

RT and RH are joint first authors. JSC and KN are joint senior authors.

Received 4 January 2023 Accepted 26 May 2023

# ABSTRACT

#### Aims

To assess if ambient temperature-related effects on serum potassium levels impact clinical decision-making. Methods

This study is an ecological time series consisiting of 1 218 453 adult patients with at least one ACE inhibitor (ACEI) prescription who participate in a large UK primary care dataset.

Descriptive statistics and a quasi-Poisson regression model using time series data at regular time intervals (monthly) were undertaken to examine the association between potassium measurements and ACEI/potassium supplement prescriptions.

**Results** It is noted that correlating with lower ambient temperature, serum potassium values follow a seasonal pattern; peaks in winter months and troughs in summer. During summer months, there are clear annual spikes in the number of potassium prescriptions suggesting a change in prescribing practice during periods of potentially spurious hyperkalaemia. The converse pattern is seen in the ACEI prescription proportion which spikes annually during the winter period with lower average ambient temperatures. Our time series modelling demonstrated that each one unit increase in potassium is associated with a 33% increased rate of ACEI prescriptions (risk ratio, RR 1.33; 95% CI 1.12 to 1.59) and 63% decreased rate of potassium supplements (RR 0.37; 95% CI 0.32 to 0.43).

**Conclusions** Our findings highlight the seasonal pattern in serum potassium and we observe a corresponding alteration in prescribing practice for potassium sensitive medications. These findings demonstrate the importance of educating clinicians on the presence of seasonal potassium variability in addition to standard measurement error, and its potential impact on their prescribing activity.

In the UK, clinicians working in primary and

secondary care have access to analytical labora-

tories where serum blood testing is commonly

conducted. In 2015/2016, an average of 4.5 labo-

ratory tests per patient were requested in primary

care, the majority of which were serum blood tests.

This represents a threefold increase in testing since

2000/2001, potentially in response to the emphasis

placed on blood monitoring in clinical performance

for clinicians to request to investigate and monitor

renal function. The standard reference range

Specifically, serum potassium is an important test

INTRODUCTION

measures.<sup>1</sup>

# Check for updates

© Author(s) (or their employer(s)) 2023. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

To cite: Thavakaran R. Hotham R, Gokhale KM, et al. J Clin Pathol Epub ahead of print: [please include Day Month Year]. doi:10.1136/ jcp-2023-208759

BMJ

#### Thayakaran R, et al. J Clin Pathol 2023;0:1-6. doi:10.1136/jcp-2023-208759

## WHAT IS ALREADY KNOWN ON THIS TOPIC

 $\Rightarrow$  Serum potassium is a useful test for clinicians to request to investigation renal function and disease. However, previous work on small samples has identified that spurious potassium results can occur (which can lead to pseudohypohypokalaemia or pseudohyperkalaemia) and occur more apparently during summer and winter months.

### WHAT THIS STUDY ADDS

 $\Rightarrow$  First, this study confirms on a large nationally representative sample the inverse association between ambient temperature and average serum potassium recordings. Second, for the first time, this study identifies an association between seasonal fluctuations in potassium values and prescribing activity for potassium supplements and ACE inhibitors.

#### HOW THIS STUDY MIGHT AFFECT RESEARCH. **PRACTICE OR POLICY**

 $\Rightarrow$  This study clearly identifies the need to educate clinicians as to the impact of ambient temperature on the accuracy of blood test results. Second, there are policy considerations regarding the potential need to adopt novel approaches such as centrifugation on site in general practices to avert ambient temperaturerelated clinical prescribing errors.

for serum potassium concentration in the UK is between 3.5 and 5.3 mEq/L (equivalent to mmol/L for potassium)<sup>2 3</sup> and abnormalities in potassium levels can be associated with significant morbidity and mortality, most notably cardiac anomalies and arrest in extreme cases.<sup>4</sup> There is no standard definition for hyperkalaemia (raised potassium in the blood) or hypokalaemia (low blood potassium), but typically clinical review would be triggered if levels were above 5.5 mEq/L or below 3.0 mEq/L.<sup>56</sup> When undertaking potassium testing, spurious results can occur, resulting in pseudohypokalaemia or pseudohyperkalaemia. This may be due to the transcellular movement of ions during phlebotomy or contamination of blood samples.<sup>7</sup> Phlebotomy techniques such as prolonged tourniquet application<sup>8</sup> or fist clenching<sup>9</sup><sup>10</sup> and can lead to significant increases in the serum concentrations of potassium in collected samples. Phlebotomist retraining and monitoring programmes appear to reduce the number of pseudohyperkalaemic samples, demonstrating



impact of erroneous results from poor practice and potential mitigation strategies.<sup>11 12</sup>

Of note, another reason for erroneous results could relate to the stark seasonal variation of potassium measurement values. It has been demonstrated in different global settings that background ambient temperature appears to affect potassium samples as they are transported to the centrifuge for testing.<sup>13</sup> During periods where temperature falls, such as winter, the measured mean daily serum potassium concentration rises.<sup>14</sup> Although the shift in mean potassium values appears modest (0.2–0.3 mEq/L) and is roughly in line with total allowable error,<sup>15</sup> it results in a doubling of the number of tests being classified as hyperkalaemic, and therefore, requiring urgent clinical review.<sup>14</sup> These erroneous readings caused by seasonal variation can lead to significant issues for primary care providers. Inaccurate results may lead to clinicians-making incorrect clinical decisions, or if an erroneous result is suspected, require rebleeding of the patient, all of which negatively affects patient care and practice resources.

The problem caused by these erroneous results is thought to be so significant that there have been movements to encourage centrifuges to be placed on site within general practice surgeries.<sup>16</sup> In the region covered by National Health Service (NHS) Grampian, where seasonal effects are more pronounced due to greater distance between practices and processing laboratories, this solution has been adopted.<sup>17</sup> This has reduced the effect of seasonal variation<sup>17</sup> and led to a significant reduction in samples displaying marked hyperkalaemia.<sup>18</sup> However, this approach has not been adopted universally across the UK, therefore, this problem may still persist. There has yet to be any literature exploring the relationship of the spurious results due to seasonal variation and the impact this may have on prescribing practice. Many medications, most notably ACE inhibitors (ACEIs), are known to impact renal potassium excretion, with regular blood sample monitoring undertaken and dose adjustment advised in patients who are experiencing hyperkalaemia.<sup>19 20</sup> ACEIs are the most commonly prescribed medication for the treatment of high blood pressure, with 9% of UK adults taking them.<sup>21</sup> AECIs are also used frequently for cardiovascular disease secondary prevention, heart failure and diabetic nephropathy.<sup>22</sup> Alternatively, for patients who are found to be hypokalaemic, even if mild, clinicians may treat with a course of potassium replacement agents.<sup>23</sup>

Since there is evidence to suggest seasonal inaccuracies in potassium values exist, it is important to identify what impact this has on prescribing practice. Given this particularly affects patients on ACEI we chose to investigate this in a population of patients on ACEI.

#### METHODS

#### Study design, data source and study population

To explore this phenomenon, we have undertaken an ecological time series analysis exploring the relationship between serum potassium measurements (in patients prescribed ACEIs) and in those prescribed potassium supplements) and Met Office recorded temperature levels. Clinical data were derived from a nationally representative UK primary care dataset (IQVIA Medical Research Data (IMRD-UK)) between the 1 January 2003 to 31 December 2016.<sup>24</sup>

The IMRD-UK database (previously named 'The Health Improvement Network') is an anonymised patient record database derived from UK general practices using Vision electronic medical records software. The IMRD-UK database contains records taken from over 800 general practices including over 15 million patients which are thought to be representative of the UK population in terms of demographic structure and prevalence of key comorbidities.<sup>24-26</sup> The database consists of clinically recorded information on patient characteristics, drug prescriptions, diagnoses, consultations and diagnostic test results including blood serum testing results. Data extraction, transformation and loading of IMRD-UK were facilitated by the automated clinical epidemiology Data extraction for epidemiological research (DExtER) platform.<sup>27</sup> Data relating to the monthly average temperature for the UK have been taken from 1 January 2003 to 31 December 2016 from the UK Meteorological Office.<sup>28</sup>

In order to reduce under-recording of events, general practices were included 12 months following their instalment of electronic practice records or from the practices' acceptable mortality recording date.<sup>29</sup> We only included adults (aged at least 18 years) on at least one ACEI or potassium supplement in our respective study population.

#### **Study variables**

To monitor seasonal trends, we aggregated for monthly data on all potassium concentration values (a variable that has been extensively studied in UK primary data)<sup>30-33</sup> in patients with at least one ACEi prescription or potassium supplement. Extreme values of potassium were excluded (<1 mEq/L, >10 mEq/L). Whist isolated potassium readings outside of this range have been reported in hospitalised cardiac arrest patients,<sup>34 35</sup> it is implausible that such samples would be taken in primary care and as such were assumed to be erroneous values. Primary care data recorded in the UK do not contain information on the specific dose change of a medication, but does include information on whether the medication was represcribed. Therefore, our primary outcome is a proportion describing the number of ACEI prescriptions given less than 15 days after a prior ACEI prescription divided by the total number of ACEI prescriptions in that calendar month. Fifteen days is a sufficient period for a clinician to receive the results of an investigation, for these results to be actioned and then for any changes in prescribing to be made. Where patients experience hyperkalaemia, the usual clinical decision in patients already taking ACEI would depend on the severity of hypokalaemia. The clinical response would be either a dose reduction, medication discontinuation or pause awaiting a repeat sample.<sup>30</sup>

Table 1	Potassium measurements and	l total num	ber of	f prescriptions f	for potassium supp	lement and ACEI
				1 1		

	England	Northern Ireland	Scotland	Wales	UK
Potassium measurements	7 357 129	565 818	1 406 1 39	1 351 739	10680825
ACEI prescriptions Repeated prescriptions within 15 days of a previous ACEI prescription	4 404 063	79 426	223 703	893 549	5600741
Potassium supplement prescriptions	64 438	3861	9134	5534	82 967
ACEI, ACE inhibitor.					

Alternatively, when patients on ACEI experience hypokalaemia, provided no other cause was identified they may be prescribed potassium supplements.<sup>23</sup> Therefore, our secondary analysis examined the aggregated frequency of potassium supplements given in each month.

## Analysis

The study analysis has been conducted using the R statistical package (R Core Team, 2013). Initially, a linear model was undertaken to describe the relationship between UK ambient temperature and average potassium values per month.

Following this, simple descriptive statistics have been used to present the characteristics of the data. The data are displayed graphically describing trends over time during the study period.

A quasi-Poisson regression model using time series data at regular time intervals (monthly) was then undertaken to examine the association between potassium measurements and the proportion of ACEI within 15 days of a previous ACEI. This model was used to estimate incidence rate ratios and their corresponding 95% CIs to describe this risk.

# RESULTS

# Trends in potassium measurement and ACEI prescriptions

During the study period, the eligible study population included 1218453 UK wide patients with at least one ACEI prescription, with a total of 5600741 and 82967 prescriptions for ACEI and potassium supplements, respectively, prescribed. A total of 10680825 potassium measurements were taken during this period for 1169400 included patients (1569 measurements were excluded due to extreme/implausible values). Table 1 describes the potassium measurements, supplements and ACEI/potassium supplement data per UK region; England, Scotland, Wales and Northern Ireland.

Figure 1 demonstrates that as in other recorded studies, correlating with lower ambient temperature, serum potassium values follow a seasonal pattern; peaks in winter months and troughs in summer.<sup>13 14</sup> Then, during summer months, there are clear annual spikes in the proportion of potassium prescriptions given suggesting a change in prescribing practice during periods of potentially spurious hyperkalaemia. The converse pattern is seen in the ACEI prescription proportion which annual spikes during the winter period with lower average ambient temperatures.



**Figure 1** Temperatures (degrees Celsius), potassium measurements (mEq/L), the proportion of Potassium supplements and ACE Inhibitor prescriptions within 15 days of a previous ACE Inhibitor prescriptions in that calendar month between January 1st 2003 and December 31st 2016.



Figure 2 Relationship between UK temperature and average potassium values.

To explore the relationship further between serum potassium and ambient temperature, we undertook a linear model (coefficient=-0.02, intercept=4.62). The relationship can be seen in figure 2 and R<sup>2</sup> was 0.8335 suggesting that potassium values explained 83.35% of the variation in the temperature in the model.

#### Time series models

The main aim of the time series regression model was to investigate whether some of the short-term variation in the outcome (proportion of ACEI/supplement prescriptions) is associated by changes in the exposures (aggregated potassium concentration).

The quasi-Poisson regression model for ACEI prescriptions under 15 days with offset as total prescriptions within potassium concentration as an independent variable shows that each 1 mEq/L increase in potassium concentration is associated with an ACEI risk ratio of 1.33 (95% CI 1.12 to 1.59, p=0.001). This translates into a 33% increase in the proportion of ACEI prescriptions given within 15 days of a previous ACEI when the serum potassium goes up by 1 mEq/L, which can be seen in figure 3.

The quasi-Poisson regression model for potassium supplements prescriptions under 15 days with offset as total prescriptions within potassium concentration as an independent associated variable, shows that each 1 mEq/L increase in potassium concentration is associated with an ACEI risk ratio of 0.37 (95% CI 0.32 to 0.43, p<0.001). This translates into a 63% decreased risk in the proportion of potassium supplement prescriptions given within 15 days of an ACEI prescription when the serum potassium goes up by 1 mEq/L, which can be seen in figure 4.

#### DISCUSSION

We see as in other recorded studies, correlating with lower ambient temperature, serum potassium values follow a seasonal pattern; peaks in winter months and troughs in summer. We also identify that, during summer months, there are annual spikes in the proportion of potassium prescriptions given suggesting a change in prescribing practice during these periods. The converse pattern is seen in the ACEI prescription proportion which annual spikes during the winter period with lower average ambient temperatures. These descriptive findings are confirmed when applied to time series modelling.

Our study is limited by the unavailability of free text in the medical notes, hence we could not make patient-level inferences as to exact reasons why their medications were amended. Second,



**Figure 3** The plot showing ACE inhibitor (ACEI) prescriptions under 15 days with predicted values over time from 2003 to 2016.



**Figure 4** The plot showing potassium supplement prescriptions under 15 days with predicted values over time from 2003 to 2016.

attempts to identify discontinuation of a prescription for a short period of time using anonymised dataset is challenging. Therefore, we will not be able to state general practitioners (GPs) are stopping or pausing patients on ACEi during the study period, but we are able to say that there has been some form of change to their prescription, indicating a prescribing decision has taken place. Third, our approved protocol and study dataset did not allow us to evaluate the concurrent effect of other blood parameters on potassium levels, such as renal function, platelet count and leucocyte count. Fourth, the effect of seasonal changes in blood pressure on prescribing activity was not part of our study protocol but could be influencing the prescribing patterns seen in ACEi and certainly warrants further investigation.<sup>37</sup>

Despite these limitations, our findings demonstrate the importance of educating clinicians on the extent of measurement error present in their serum sampling requests and subsequent prescribing decisions. Our findings also suggest further investigation is needed to explore the clinical effects of facilitating centrifugation or newer approaches such as multiple panel point of care testing at the site of venepuncture to minimise spurious potassium measurements.<sup>16 17</sup> It is clear that there is an associated clinical implication related to these erroneous results which is significant. Consideration should be given to the adoption of approaches in the region covered by NHS Grampian (where GP surgeries are provided centrifuges on site),<sup>17</sup> where significant reductions in erroneous samples have been achieved.<sup>18</sup> This would be aided by formal cost–benefit analyses of these approaches, which is so far lacking in the literature.

In conclusion, our study demonstrates there are observable shifts in prescribing activity temporally associated with ambient temperature-related changes in serum potassium levels. Clinicians should be educated as to the breadth of this issue and consider these findings when interpreting biochemistry results at different points throughout the year.

#### Handling editor Patrick J Twomey.

#### Twitter Joht Singh Chandan @johtchandan

**Contributors** RT, KN and JSC were responsible for study conception. RT, KN, KMG and JSC were responsible for data extraction and statistical analysis. RT, KN and JSC were responsible for initial drafting of the manuscript. All authors (RT, KN, KMG, NJA, RH and JSC) were responsible for ongoing drafts and final review of the manuscript. All authors have read and approved the final manuscript. JSC is the guarantor of the manuscript and work undertaken.

**Funding** The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

**Competing interests** JSC, KN and KMG are codirectors of DExtER operating division which is part of the University of Birmingham. DExtER operating division supports the extraction and preparing of healthcare data to support epidemiological analyses such as those seen in this article.

Patient consent for publication Not applicable.

**Ethics approval** Anonymised data were used throughout the study provided by the data provider to the University of Birmingham. Studies using IMRD-UK database have had initial ethical approval from the NHS South-East Multicentre Research Ethics Committee, subject to prior independent scientific review. The Scientific Review Committee (IQVIA) approved the study protocol (SRC Reference Number: SRC19THIN036).

Provenance and peer review Not commissioned; externally peer reviewed.

**Data availability statement** Data may be obtained from a third party and are not publicly available.

**Open access** This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: http://creativecommons.org/licenses/by-nc/4.0/.

#### ORCID iD

Joht Singh Chandan http://orcid.org/0000-0002-9561-5141

#### REFERENCES

- 1 O'Sullivan JW, Stevens S, Hobbs FDR, *et al*. Temporal trends in use of tests in UK primary care, 2000-15: retrospective analysis of 250 million tests. *BMJ* 2018;363:k4666.
- 2 Julian B, Kenneth RJ, Danielle F. Harmonisation of reference intervals. 2011.
- 3 Berg J. The approach to pathology harmony in the UK. *Clin Biochem Rev* 2012;33:89–93.
- 4 He FJ, MacGregor GA. Fortnightly review: beneficial effects of potassium. BMJ 2001;323:497–501.
- 5 National Institute for Health and Care Excellence. Diuretics. 2023. Available: https:// cks.nice.org.uk/topics/heart-failure-chronic/prescribing-information/diuretics/
- 6 National Institute of Health and Care Excellence. ACE-inhibitors. 2023. Available: https://cks.nice.org.uk/topics/heart-failure-chronic/prescribing-information/aceinhibitors/ [Accessed 8 Mar 2023].
- 7 Crook MA. Pseudohyperkalaemia or spurious Hyperkalaemia. Ann Clin Biochem 2013;50:180–1.
- 8 Saleem S, Mani V, Chadwick MA, et al. A prospective study of causes of Haemolysis during Venepuncture: Tourniquet time should be kept to a minimum. Ann Clin Biochem 2009;46:244–6.
- 9 Bailey IR, Thurlow VR. Is suboptimal Phlebotomy technique Impacting on potassium results for primary care. *Ann Clin Biochem* 2008;45(Pt 3):266–9.
- 10 Don BR, Sebastian A, Cheitlin M, et al. Pseudohyperkalemia caused by fist Clenching during Phlebotomy. N Engl J Med 1990;322:1290–2.
- 11 Greenblatt MB, Torre M, Means J, et al. Use of potassium concentrations as a quality-of-service metric for Phlebotomists Detects systematic Preanalytical biases and facilitates their correction. *Clin Chem* 2014;60:1453–5.
- 12 Lucas F, Mata DA, Greenblatt MB, et al. A potassium-based quality-of-service metric reduces Phlebotomy errors, resulting in improved patient safety and decreased cost. Am J Clin Pathol 2022;157:789–98.
- 13 Lim HS, Lee YK, Min WK. Effects of long distance transportation of specimens on test results. *Lab Med Online* 2011;1:72.
- 14 Sinclair D, Briston P, Young R, *et al.* Seasonal Pseudohyperkalaemia. *J Clin Pathol* 2003;56:385–8.
- 15 Desirable biological variation database specifications Westgard. Available: https:// www.westgard.com/biodatabase1.htm [Accessed 16 May 2023].
- 16 Masters PW, Ismail AA. Minimising Factitious Hyperkalaemia. samples should be centrifuged after collection in general practices. *BMJ* 1997;315:190. 10.1136/ bmj.315.7101.190a Available: http://www.ncbi.nlm.nih.gov/pubmed/9251570
- 17 Turner HE, Peake RWA, Allison JJ. Seasonal Pseudohyperkalaemia: no longer an issue Ann Clin Biochem 2012;49:94–6.
- 18 Turner HE, Peake RWA, Allison JJ. The impact of centrifugation in primary care on pseudohyperkalaemia: a retrospective evaluation. *Ann Clin Biochem* 2013;50:371–3. 10.1177/0004563212474566 Available: http://dx.doi.org/101177/ 0004563212474566
- 19 Ben Salem C, Badreddine A, Fathallah N, et al. Drug-induced Hyperkalemia. Drug Saf 2014;37:677–92.
- 20 Raebel MA. Hyperkalemia associated with use of angiotensin-converting enzyme inhibitors and angiotensin receptor blockers. *Cardiovascular Therapeutics* 2012;30:e156–66. 10.1111/j.1755-5922.2010.00258.x Available: http://doi.wiley.com/10.1111/cdr.2012.30.issue-3

# **Original research**

- 21 Rouette J, McDonald EG, Schuster T, *et al.* Original research: treatment and Prescribing trends of antihypertensive drugs in 2.7 million UK primary care patients over 31 years: a population-based cohort study. *BMJ Open* 2022;12:e057510e057510.
- 22 Drugs affecting the renin-angiotensin system | treatment summaries | BNF | NICE. Available: https://bnf.nice.org.uk/treatment-summaries/drugs-affecting-the-reninangiotensin-system/ [Accessed 16 May 2023].
- 23 Kardalas E, Paschou SA, Anagnostis P, et al. Hypokalemia: a clinical update. Endocr Connect 2018;7:R135–46.
- 24 Blak BT, Thompson M, Dattani H, et al. Generalisability of the health improvement network (THIN) database: demographics, chronic disease prevalence and mortality rates. Inform Prim Care 2011;19:251–5.
- 25 IQVIA. UK EMR IQVIA medical research data. Available: https://www.iqvia.com/ library/fact-sheets/uk-emr-iqvia-medical-research-data [Accessed 3 Jan 2022].
- 26 Candore G, Hedenmalm K, Slattery J, et al. Can we rely on results from IQVIA medical research data UK converted to the observational medical outcome partnership common data model *Clin Pharmacol Ther* 2020;107:915–25. 10.1002/cpt.1785 Available: https://onlinelibrary.wiley.com/toc/15326535/107/4
- 27 Gokhale KM, Chandan JS, Toulis K, et al. Data extraction for Epidemiological research (Dexter): a novel tool for automated clinical epidemiology studies. Eur J Epidemiol 2021;36:165–78.
- 28 Meteorological office. data. Available: https://www.metoffice.gov.uk/research/climate/ maps-and-data/index [Accessed 3 Jan 2022].

- 29 Maguire A, Blak BT, Thompson M. The importance of defining periods of complete mortality reporting for research using automated data from primary care. *Pharmacoepidemiol Drug Saf* 2009;18:76–83.
- 30 Furuland H, McEwan P, Evans M, et al. Serum potassium as a Predictor of adverse clinical outcomes in patients with chronic kidney disease: new risk equations using the UK clinical practice research Datalink. BMC Nephrol 2018;19:211.
- 31 Horne L, Ashfaq A, MacLachlan S, et al. Epidemiology and health outcomes associated with Hyperkalemia in a primary care setting in England. BMC Nephrol 2019;20:85.
- 32 Schmidt M, Mansfield KE, Bhaskaran K, et al. Serum creatinine elevation after reninangiotensin system blockade and long term Cardiorenal risks: cohort study. BMJ 2017;356:j791.
- 33 Michel A, Martín-Pérez M, Ruigómez A, et al. Incidence and risk factors for severe renal impairment after first diagnosis of heart failure: A cohort and nested case–control study in UK general practice. Int J Cardiol 2016;207:S0167-5273(16)30165-6:252–7.:.
- 34 Tran HA. Extreme Hyperkalemia. South Med J 2005;98:729–32.
- 35 A case of extreme hypokalaemia PubMed. Available: https://pubmed.ncbi.nlm.nih. gov/27905308/ [Accessed 16 May 2023].
- 36 ACE-inhibitors | Prescribing information | heart failure chronic | CKS | NICE. Available: https://cks.nice.org.uk/topics/heart-failure-chronic/prescribing-information/ ace-inhibitors/ [Accessed 16 May 2023].
- 37 Narita K, Hoshide S, Kario K. Seasonal variation in blood pressure: Current evidence and recommendations for hypertension management. *Hypertens Res* 2021;44:1363–72.