UNIVERSITY^{OF} BIRMINGHAM University of Birmingham Research at Birmingham

Bounding the row sum arithmetic mean by Perron roots of row-permuted matrices

Engel, Gernot; Sergeev, Sergei

DOI: 10.1016/j.laa.2023.05.014

License: Creative Commons: Attribution (CC BY)

Document Version Publisher's PDF, also known as Version of record

Citation for published version (Harvard):

Engel, G & Sergeev, S 2023, 'Bounding the row sum arithmetic mean by Perron roots of row-permuted matrices', *Linear Algebra and its Applications*, vol. 673, pp. 220-232. https://doi.org/10.1016/j.laa.2023.05.014

Link to publication on Research at Birmingham portal

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

•Users may freely distribute the URL that is used to identify this publication.

•Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.

•User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?) •Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.



Contents lists available at ScienceDirect

Linear Algebra and its Applications

journal homepage: www.elsevier.com/locate/laa

Bounding the row sum arithmetic mean by Perron roots of row-permuted matrices



LINEAR

ications

Gernot Michael Engel^a, Sergeĭ Sergeev^{b,*}

^a Transversal Networks Corp., 2753 Marshall Parkway, Madison, WI 53713, USA
 ^b University of Birmingham, School of Mathematics, Edgbaston B15 2TT, UK

ARTICLE INFO

Article history: Received 5 September 2022 Received in revised form 11 May 2023 Accepted 11 May 2023 Available online 16 May 2023 Submitted by R. Brualdi

MSC: 15A18

Keywords: Perron root Row sums Rearrangement inequality

АВЅТ КАСТ

 $\mathbb{R}^{n \times n}_+$ denotes the set of $n \times n$ non-negative matrices. For $A \in \mathbb{R}^{n \times n}_+$ let $\Omega(A)$ be the set of all matrices that can be formed by permuting the elements within each row of A. Formally:

$$\Omega(A) = \{ B \in \mathbb{R}^{n \times n}_+ : \forall i \exists a \text{ permutation} \}$$

 $\phi_i \text{ s.t. } b_{i,j} = a_{i,\phi_i(j)} \ \forall j \}.$

For $B \in \Omega(A)$ let $\rho(B)$ denote the spectral radius or largest non-negative eigenvalue of B. We show that the arithmetic mean of the row sums of A is bounded by the maximum and minimum spectral radius of the matrices in $\Omega(A)$. Formally, we show that

$$\min_{B\in\Omega(A)}\rho(B) \le \frac{1}{n}\sum_{i=1}^n\sum_{j=1}^n a_{i,j} \le \max_{B\in\Omega(A)}\rho(B).$$

For positive A we obtain necessary and sufficient conditions for these inequalities to become an equality. We also give criteria which an irreducible matrix C should satisfy so that $\rho(C) = \min_{B \in \Omega(A)} \rho(B)$ or $\rho(C) = \max_{B \in \Omega(A)} \rho(B)$. These

* Corresponding author.

https://doi.org/10.1016/j.laa.2023.05.014

E-mail addresses: engel@transversalnetworks.net (G.M. Engel), s.sergeev@bham.ac.uk (S. Sergeev).

^{0024-3795/}© 2023 The Authors. Published by Elsevier Inc. This is an open access article under the CC BY license (http://creativecommons.org/licenses/by/4.0/).

criteria are used to derive algorithms for finding such C when all the entries of A are positive.

© 2023 The Authors. Published by Elsevier Inc. This is an open access article under the CC BY license (http:// creativecommons.org/licenses/by/4.0/).

1. Introduction

In what follows, \mathbb{R}^n_+ denotes the set of non-negative vectors with length n and $\mathbb{R}^{n \times n}_+$ denotes the set of non-negative $n \times n$ matrices. For $x \in \mathbb{R}^n_+$ or, respectively, $A \in \mathbb{R}^{n \times n}_+$ we write x > 0 or, respectively, A > 0, if all entries of vector x or matrix A are positive. We will work with the following matrix set, which can be defined for any matrix A.

Definition 1.1. For $A \in \mathbb{R}^{n \times n}_+$, the matrix set $\Omega(A)$ consists of the *row-permuted* matrices, whose entries in each row are a permutation of entries in the corresponding row of A. Formally:

$$\Omega(A) = \{ B \in \mathbb{R}^{n \times n}_+ : \forall i \; \exists a \text{ permutation } \phi_i \text{ s.t. } b_{i,j} = a_{i,\phi_i(j)} \; \forall j \}.$$
(1)

We will use the following standard notation for the Perron roots of matrices.

Definition 1.2. The Perron root (i.e. the largest non negative eigenvalue, or spectral radius) of a matrix $B \in \Omega(A)$ will be denoted by $\rho(B)$.

For $A \in \mathbb{R}^{n \times n}_+$ the following row sum inequality

$$\min_{i=1}^{n} \sum_{j=1}^{n} a_{i,j} \le \rho(A) \le \max_{i=1}^{n} \sum_{j=1}^{n} a_{i,j}$$

was first observed by Frobenius. The geometric means of the row sums as bounds for $\rho(A)$ were explored by Al'pin [1] and Elsner and van Driessche [2], and further generalized by Engel et al. [5]. In this paper we are interested in establishing a different connection between Perron roots and row sums. Namely, we show that the arithmetic mean of the row sums satisfies

$$\min_{B\in\Omega(A)}\rho(B) \le \frac{1}{n}\sum_{i=1}^{n}\sum_{j=1}^{n}a_{i,j} \le \max_{B\in\Omega(A)}\rho(B).$$
(2)

For A > 0 we obtain necessary and sufficient conditions for any of these inequalities to turn into equalities. For $A \in \mathbb{R}^{n \times n}_+$ we also give necessary and sufficient criteria for an irreducible matrix $C \in \Omega(A)$ to have $\rho(C) = \min_{B \in \Omega(A)} \rho(B)$ or $\rho(C) = \max_{B \in \Omega(A)} \rho(B)$.

To obtain these results we make use, in particular, of the following well-known facts. These facts, which we are going to use throughout the paper, are closely related to the famous Collatz-Wielandt inequality and are summarized in the following proposition: **Proposition 1.3** (e.g., [3], Theorem 1.11). For $A \in \mathbb{R}^{n \times n}_+$ and constants $\alpha > 0, \beta > 0$ and nonzero vector $x \in \mathbb{R}^n_+$ we have:

(i) $\alpha x \leq Ax$ implies $\alpha \leq \rho(A)$,

(ii) $Ax \leq \beta x$ with x > 0 implies $\rho(A) \leq \beta$.

In addition, if A is irreducible then the following implications hold:

(iii) if $\alpha x \leq Ax$ and $\exists i$ such that $\alpha x_i < \sum_{j=1}^n a_{i,j} x_j$ then $\rho(A) > \alpha$, (iv) if $Ax \leq \beta x$ and $\exists i$ such that $\sum_{j=1}^n a_{i,j} x_j < \beta x_i$ then $\rho(A) < \beta$.

The next result, which we will use to derive the criteria for $\rho(C) = \max_{B \in \Omega(A)} \rho(B)$ and $\rho(C) = \min_{B \in \Omega(A)} \rho(B)$, is known as the *rearrangement inequality*.

Proposition 1.4 (e.g., [6], page 261). Let $x, y \in \mathbb{R}^n_+$ be such that $x_1 \leq x_2 \leq \ldots \leq x_n$ and $y_1 \leq y_2 \leq \ldots \leq y_n$, and let $\phi: \{1, \ldots, n\} \rightarrow \{1, \ldots, n\}$ be an arbitrary permutation. Then the following inequalities hold:

$$\sum_{i=1}^{n} x_i y_{n+1-i} \le \sum_{i=1}^{n} x_i y_{\phi(i)} \le \sum_{i=1}^{n} x_i y_i.$$

2. Preliminary lemmas

The following lemma, related to the rearrangement inequality, establishes that the maximum Perron root is achieved on a matrix with all positive entries for which the correlation between the order of the components of its Perron eigenvector and each of its row vectors is maximized. The minimum Perron root is achieved when the correlation between the order of the components of its Perron eigenvector and each of its row vectors is minimized.

Lemma 2.1. Let $A \in \mathbb{R}^{n \times n}_+$ be irreducible. Then the following implications hold:

(i) if $\rho(A) = \max_{B \in \Omega(A)} \rho(B)$ and x is a Perron eigenvector of A then for $1 \le i, j, k \le n$:

$$x_k < x_j \text{ implies } a_{i,k} \leq a_{i,j},$$

(ii) if $\rho(A) = \min_{B \in \Omega(A)} \rho(B)$ and x is a Perron eigenvector of A then for $1 \le i, j, k \le n$:

$$x_k < x_j \text{ implies } a_{i,k} \ge a_{i,j}.$$

Proof. (i): Assume that $A \in \mathbb{R}^{n \times n}_+$ is irreducible and $\rho(A) = \max_{B \in \Omega(A)} \rho(B)$. Then $\exists x > 0$ such that $Ax = \rho(A)x$. By contradiction, assume that there exist i, j, k such

that $x_j > x_k$ but $a_{i,j} < a_{i,k}$. Let B be the matrix formed by swapping the two entries $a_{i,j}$ and $a_{i,k}$ so that $b_{i,j} = a_{i,k}$ and $b_{i,k} = a_{i,j}$, with all other entries of B equal to the entries of A. Then B is in $\Omega(A)$. We have $\rho(A)x_i = \sum_{j=1}^n a_{i,j}x_j < \sum_{j=1}^n b_{i,j}x_j$ and $\sum_{j=1}^{n} a_{s,j} x_j = \sum_{j=1}^{n} b_{s,j} x_j$ for $s \neq i$. Thus by Proposition 1.3 part (iii), $\rho(B) > \rho(A)$. Since this contradicts that $\rho(A) = \max_{B \in \Omega(A)} \rho(B)$ it follows that for $1 \le i, j, k \le n x_k < n x$ x_i implies $a_{i,k} \leq a_{i,j}$ and (i) is established.

(ii): The proof is similar to the previous part, with the difference that here we assume that $\rho(A) = \min_{B \in \Omega(A)} \rho(B)$. Upon assuming by contradiction that there exist i, j, ksuch that $x_j > x_k$ but $a_{i,j} > a_{i,k}$ we define matrix B by swapping the entries $a_{i,j}$ and $a_{i,k}$ so that $b_{i,j} = a_{i,k}$ and $b_{i,k} = a_{i,j}$, with all other entries of B equal to the entries of A. Observing that $\rho(A)x_i = \sum_{j=1}^n a_{i,j}x_j > \sum_{j=1}^n b_{i,j}x_j$ and $\sum_{j=1}^n a_{s,j}x_j = \sum_{j=1}^n b_{s,j}x_j$ for $s \neq i$, we use Proposition 1.3 part (iv) to obtain $\rho(B) < \rho(A)$, a contradiction establishing part (ii). \Box

Proof of the next lemma follows the reasoning used in the proof of Tchebychef's inequality [6] page 43.

Lemma 2.2. Let $A \in \mathbb{R}^{n \times n}_+$ have a Perron eigenvector $x \in \mathbb{R}^n_+$ satisfying $\sum_{i=1}^n x_i = 1$. Then the following properties hold:

(i) if $\forall i, j, k \ x_k < x_j$ implies $a_{i,k} \leq a_{i,j}$ then

$$\forall i \ \frac{\sum_{j=1}^{n} a_{i,j}}{n} \le \sum_{j=1}^{n} a_{i,j} x_j = \rho(A) x_i,$$

(ii) if $\forall i, j, k \ x_k < x_j$ implies $a_{i,k} \ge a_{i,j}$ then

$$\forall i \ \frac{\sum_{j=1}^{n} a_{i,j}}{n} \ge \sum_{j=1}^{n} a_{i,j} x_j = \rho(A) x_i,$$

- (iii) if $\forall i, j, k \ x_k < x_j$ implies $a_{i,k} \leq a_{i,j}$ or $\forall i, j, k \ x_k < x_j$ implies $a_{i,k} \geq a_{i,j}$, then the following are equivalent:

 - (a) $\frac{\sum_{j=1}^{n} a_{i,j}}{n} = \sum_{j=1}^{n} a_{i,j} x_j = \rho(A) x_i$ for all i; (b) either $x_i = \frac{1}{n}$ for all i, or for each i there is c_i such that $c_i = a_{i,j}$ for all j.

Proof. (i): The property that $\forall 1 \leq i, j, k \leq n \ x_k < x_j$ implies $a_{i,k} \leq a_{i,j}$ is equivalent to $(a_{i,j} - a_{i,k})(x_j - x_k) \ge 0$. From this we obtain

224 G.M. Engel, S. Sergeev / Linear Algebra and its Applications 673 (2023) 220–232

$$\forall i \ 2nx_i\rho(A) = \sum_{j=1}^n \sum_{k=1}^n (a_{i,j}x_j + a_{i,k}x_k) \ge \sum_{j=1}^n \sum_{k=1}^n (a_{i,j}x_k + a_{i,k}x_j)$$
$$\ge 2\left(\sum_{j=1}^n a_{i,j}\right)\left(\sum_{j=1}^n x_j\right) = 2\sum_{j=1}^n a_{i,j}.$$

This implies

$$\forall i \ \frac{\sum_{i=1}^{n} a_{i,j}}{n} \le \sum_{i=1}^{n} a_{i,j} x_j = \rho(A) x_i$$

establishing part (i).

(ii): The proof of this part is similar to the proof of part (i). Here we first observe that the property that $\forall 1 \leq i, j, k \leq n \ x_k < x_j$ implies $a_{i,k} \geq a_{i,j}$ is equivalent to $(a_{i,j} - a_{i,k})(x_j - x_k) \leq 0$. Using this inequality in the same way as in the proof of part (i) the opposite inequality is used, we obtain

$$\forall i \ \frac{\sum_{i=1}^{n} a_{i,j}}{n} \ge \sum_{i=1}^{n} a_{i,j} x_j = \rho(A),$$

establishing part (ii).

(iii): To establish (3) (a) implies (b) assume $\frac{\sum_{i=1}^{n} a_{i,j}}{n} = \sum_{i=1}^{n} a_{i,j}x_j = \rho(A)x_i$ and that either $\forall 1 \leq i, j, k \leq n \ x_k < x_j$ implies $a_{i,k} \leq a_{i,j}$ or $\forall 1 \leq i, j, k \leq n \ x_k < x_j$ implies $a_{i,k} \geq a_{i,j}$.

In the first case for any i, j, k we have that $(a_{i,j} - a_{i,k})(x_j - x_k) \ge 0$ and in the second case we have that $(a_{i,j} - a_{i,k})(x_j - x_k) \le 0$. In the first case, if there exists i such that $(a_{i,j} - a_{i,k})(x_j - x_k) > 0$ for some j and k then $\frac{\sum_{i=1}^{n} a_{i,j}}{n} < \sum_{i=1}^{n} a_{i,j}x_j = \rho(A)x_i$. Similarly in the second case if there exists i such that $(a_{i,j} - a_{i,k})(x_j - x_k) < 0$, then $\frac{\sum_{i=1}^{n} a_{i,j}}{n} > \sum_{i=1}^{n} a_{i,j}x_j = \rho(A)x_i$. Since none of these strict inequalities holds, we have

$$\forall i, j, k \ (a_{i,j} - a_{i,k})(x_j - x_k) = 0.$$
(3)

For any i = 1, ..., n let t(i) and d(i) be defined (non-uniquely) by

$$a_{i,t(i)} = \min_{j} a_{i,j}, \quad a_{i,d(i)} = \max_{j} a_{i,j}$$
 (4)

and suppose that $x_i = x_k$ does not hold for all $i \neq k$. Our aim is to show that then the coefficients in every row of A are equal to each other. Since either $\forall i, j, k \ x_k < x_j$ implies $a_{i,k} \leq a_{i,j}$ or $\forall i, j, k \ x_k < x_j$ implies $a_{i,k} \geq a_{i,j}$, we can let t(i) and d(i) be defined in such a way that not only equalities (4) hold but also in the first case $x_{t(i)} = \min_j x_j$ and $x_{d(i)} = \max_j x_j$ and in the second case $x_{t(i)} = \max_j x_j$ and $x_{d(i)} = \min_j x_j$. In both cases (3) entails that $(a_{i,t(i)} - a_{i,d(i)})(x_{t(i)} - x_{d(i)}) = 0$ and hence $a_{i,t(i)} = a_{i,d(i)}$. By (4)

we obtain that all entries in the *i*th row of A are equal to each other, establishing the implication $(a) \Rightarrow (b)$.

To prove that (b) implies (a), first observe that obviously if $x_j = \frac{1}{n}$ then $\frac{\sum_{i=1}^n a_{i,j}}{n} = \sum_{i=1}^n a_{i,j} x_j = \rho(A) x_i$. If instead for each *i* there is c_i such that $a_{i,j} = c_i$ for all *j*, then the unique Perron eigenvector *x* with $\sum_{j=1}^n x_j = 1$ has coordinates $x_i = c_i / \sum_{j=1}^n c_j$ for all *i* and the Perron root is $\rho(A) = \sum_{j=1}^n c_j$. Indeed, we have

$$\sum_{j=1}^{n} a_{i,j} x_j = c_i \sum_{i=1}^{n} x_j = c_i = \sum_{j=1}^{n} c_j \cdot \frac{c_i}{\sum_{j=1}^{n} c_j} = \rho(A) x_i.$$

In this case $\frac{\sum_{i=1}^{n} a_{i,j}}{n} = c_i = \sum_{j=1}^{n} a_{i,j} x_j$, establishing (a). \Box

3. Main results

We begin this section by establishing the inequality between the arithmetic mean of the rows and the largest and smallest Perron roots of matrices in $\Omega(A)$.

Theorem 3.1. For any $A \in \mathbb{R}^{n \times n}_+$

$$\min_{B \in \Omega(A)} \rho(B) \le \frac{1}{n} \sum_{i=1}^{n} \sum_{j=1}^{n} a_{i,j} \le \max_{B \in \Omega(A)} \rho(B).$$
(5)

Proof. We first assume that A > 0 and establish $\frac{1}{n} \sum_{i=1}^{n} \sum_{j=1}^{n} a_{i,j} \leq \max_{B \in \Omega(A)} \rho(B)$ for such A. Select $C \in \Omega(A)$ such that $\rho(C) = \max_{B \in \Omega(A)} \rho(B)$. Let x > 0 be a Perron eigenvector of C such that $\sum_{i=1}^{n} x_i = 1$. By Lemma 2.1 part (i) we have that for $1 \leq i, j, k \leq n x_k < x_j$ implies $c_{i,k} \leq c_{i,j}$. Then by Lemma 2.2 part (i) we have $\forall i \ \frac{1}{nx_i} \sum_{j=1}^{n} c_{i,j} \leq \sum_{j=1}^{n} c_{i,j} \frac{x_j}{x_i} = \max_{B \in \Omega(A)} \rho(B)$ and since $\sum_{j=1}^{n} c_{i,j} = \sum_{j=1}^{n} a_{ij}$ for all i, we obtain

$$\frac{1}{n}\sum_{i=1}^{n}\sum_{j=1}^{n}a_{i,j} \le \sum_{i=1}^{n}\max_{B\in\Omega(A)}\rho(B)x_{i} = \max_{B\in\Omega(A)}\rho(B).$$

Still assuming A > 0, we can establish $\frac{1}{n} \sum_{i=1}^{n} \sum_{j=1}^{n} a_{i,j} \ge \min_{B \in \Omega(A)} \rho(B)$ in a similar way. For this we select $C \in \Omega(A)$ such that $\rho(C) = \min_{B \in \Omega(A)} \rho(B)$ and let x > 0 be a Perron eigenvector of C such that $\sum_{i=1}^{n} x_i = 1$. Combining Lemma 2.1 part (ii) with Lemma 2.2 part (ii) we obtain $\forall i \ \frac{1}{nx_i} \sum_{j=1}^{n} c_{i,j} \ge \sum_{j=1}^{n} c_{i,j} \frac{x_j}{x_i}$ and hence

$$\frac{1}{n}\sum_{i=1}^{n}\sum_{j=1}^{n}a_{i,j} \ge \sum_{i=1}^{n}\min_{B\in\Omega(A)}\rho(B)x_{i} = \min_{B\in\Omega(A)}\rho(B)x_{i}$$

Now for arbitrary $A \in \mathbb{R}^{n \times n}_+$ and $\epsilon > 0$ we define $A^{\epsilon} = (a_{i,j}^{\epsilon}) = (a_{i,j} + \epsilon)$. Then since $0 < A^{\epsilon}$ we have that $\min_{B \in \Omega(A^{\epsilon})} \rho(B) \leq \frac{1}{n} \sum_{i=1}^{n} \sum_{j=1}^{n} (a_{i,j} + \epsilon) \leq \max_{B \in \Omega(A^{\epsilon})} \rho(B)$.

225

Thus by continuity of the Perron root and letting ϵ go to zero we obtain the desired inequality for A. \Box

We now establish the conditions when any of the inequalities in Theorem 3.1 becomes an equality.

Theorem 3.2. For $0 < A \in \mathbb{R}^{n \times n}_+$ the following are equivalent:

- (i) $\frac{1}{n} \sum_{i=1}^{n} \sum_{j=1}^{n} a_{i,j} = \max_{B \in \Omega(A)} \rho(B).$
- (ii) Either the flat vector x = (x_i) where ∀i x_i = 1 is a Perron eigenvector of A or there exists a non singular diagonal matrix ∃ D ≥ 0 such that DA is a flat matrix (i.e. ∀i, j d_ia_{i,j} = 1).
- (iii) $\frac{1}{n} \sum_{i=1}^{n} \sum_{j=1}^{n} a_{i,j} = \min_{B \in \Omega(A)} \rho(B).$

Proof. We first establish (i) \Rightarrow (ii). By (i), $\frac{1}{n}\sum_{i=1}^{n}\sum_{j=1}^{n}a_{i,j} = \max_{B\in\Omega(A)}\rho(B)$. Since the set $\Omega(A)$ is finite, there exist $C \in \Omega(A)$ and $y \in \mathbb{R}^{n \times n}_+$ with $\sum_{i=1}^{n}y_i = 1$ such that $Cy = (\max_{B\in\Omega(A)}\rho(B))y$. Thus

$$\sum_{i=1}^{n} \sum_{j=1}^{n} c_{i,j} y_j = \left(\max_{B \in \Omega(A)} \rho(B) \right) \left(\sum_{i=1}^{n} y_i \right) = \max_{B \in \Omega(A)} \rho(B) = \sum_{i=1}^{n} \sum_{j=1}^{n} a_{i,j} \frac{1}{n}$$

By Lemma 2.1 part (i) $\forall i, j, k: 1 \leq i, j, k \leq n$ we have that $y_k < y_j$ implies $c_{i,k} \leq c_{i,j}$. Then by Lemma 2.2 part (i)

$$\forall i \ \frac{1}{n} \sum_{j=1}^{n} c_{i,j} \leq \sum_{j=1}^{n} c_{i,j} y_j = \max_{B \in \Omega(A)} \rho(B) y_i.$$

Since $\forall i \ \frac{1}{n} \sum_{j=1}^{n} a_{i,j} = \frac{1}{n} \sum_{j=1}^{n} c_{i,j}$ we can rewrite this as

$$\forall i \ \frac{1}{n} \sum_{j=1}^{n} a_{i,j} \le \max_{B \in \Omega(A)} \rho(B) y_i.$$

As by (i) we have

$$\frac{1}{n}\sum_{i=1}^{n}\sum_{j=1}^{n}a_{i,j} = \max_{B\in\Omega(A)}\rho(B) = \max_{B\in\Omega(A)}\rho(B)\sum_{i=1}^{n}y_i,$$

if there exists *i* such that $\frac{1}{n} \sum_{j=1}^{n} a_{i,j} < \max_{B \in \Omega(A)} \rho(B) y_i$ then there would have to exist *k* such that $\frac{1}{n} \sum_{j=1}^{n} a_{k,j} > \max_{B \in \Omega(A)} \rho(B) y_k$, which is a contradiction, hence

$$\forall i \max_{B \in \Omega(A)} \rho(B) y_i = \frac{1}{n} \sum_{j=1}^n a_{i,j} = \sum_{j=1}^n c_{i,j} y_j = \frac{1}{n} \sum_{j=1}^n c_{i,j}.$$

Applying Lemma 2.2 part (iii), we obtain that either $\forall i \ y_i = 1/n \text{ or } \forall i, j, k \ c_{i,j} = c_{i,k}$. If $\forall i \ y_i = 1/n$ then $\forall i \ \frac{1}{n} \sum_{j=1}^n a_{i,j} = \max_{B \in \Omega(A)} \rho(B) \frac{1}{n}$, from which it follows that flat vector $x = (x_i)$ where $\forall i \ x_i = 1$ is a Perron eigenvector of A. If $\forall i, j, k \ c_{i,j} = c_{i,k}$ then $\forall i, j, k \ a_{i,j} = c_{i,j} = c_{i,k} = a_{i,k}$. Let D be the diagonal matrix where $\forall i \ d_{i,i} = \frac{1}{a_{i,i}}$ and the rest of the entries of D are 0. Thus DA is the flat matrix such that $\forall i, j \ d_i a_{i,j} = 1$.

We now show (ii) \Rightarrow (i), (iii). Assume first that the flat vector $x = (x_i)$ where $\forall i \ x_i = 1$ is a Perron eigenvector of A. This is equivalent to all row sums of A being equal to each other. If this property holds for A then it also holds for all $B \in \Omega(A)$, so the flat vector is a Perron eigenvector of any such B with the same Perron root (equal to any of the row sums). Thus we have both (i) and (iii), i.e.,

$$\max_{B \in \Omega(A)} \rho(B) = \min_{B \in \Omega(A)} \rho(B) = \frac{1}{n} \sum_{i=1}^{n} \sum_{j=1}^{n} a_{i,j}.$$
 (6)

Now assume that there exists a non-singular diagonal matrix $D \ge 0$ such that DA is a flat matrix. In this case the entries in each row of A are equal to each other, implying that $\Omega(A) = \{A\}$. As the left hand side and the right hand side of (5) are equal to each other, we obtain (6).

Finally, the proof of (iii) \Rightarrow (ii) is similar to the proof of (i) \Rightarrow (ii) and will be described more briefly. By (ii), $\frac{1}{n} \sum_{i=1}^{n} \sum_{j=1}^{n} a_{i,j} = \min_{B \in \Omega(A)} \rho(B)$. Since the set $\Omega(A)$ is finite, there exist $C \in \Omega(A)$ and $y \in \mathbb{R}^{n \times n}_+$ with $\sum_{i=1}^{n} y_i = 1$ such that $Cy = (\min_{B \in \Omega(A)} \rho(B))y$. Thus

$$\sum_{i=1}^{n} \sum_{j=1}^{n} c_{i,j} y_j = \left(\min_{B \in \Omega(A)} \rho(B) \right) \left(\sum_{i=1}^{n} y_i \right) = \min_{B \in \Omega(A)} \rho(B) = \sum_{i=1}^{n} \sum_{j=1}^{n} a_{i,j} \frac{1}{n}.$$

Next, combining Lemma 2.1 part (ii) and Lemma 2.2 part (ii) and using that for each i the sum of the *i*th row of A equals the sum of the *i*th row of C, we obtain

$$\forall i \ \frac{1}{n} \sum_{j=1}^{n} a_{i,j} \ge \min_{B \in \Omega(A)} \rho(B) y_i.$$

Using condition (iii), however, we see that the strict inequality cannot hold for any i and therefore we have

$$\forall i \min_{B \in \Omega(A)} \rho(B) y_i = \frac{1}{n} \sum_{j=1}^n a_{i,j} = \sum_{j=1}^n c_{i,j} y_j = \frac{1}{n} \sum_{j=1}^n c_{i,j}.$$

Condition (ii) then follows by applying Lemma 2.2 part (iii) (see the end of the proof of $(i) \Rightarrow (ii)$ written above.) \Box

The following result applies the rearrangement inequality (Proposition 1.4) to yield a sufficient condition for establishing when $\rho(A) = \max_{B \in \Omega(A)} \rho(B)$ and $\rho(A) = \min_{B \in \Omega(A)} \rho(B)$.

Theorem 3.3. Let $A \in \mathbb{R}^{n \times n}_+$ and $0 \le x \in \mathbb{R}^n_+$ be a Perron eigenvector of A. Then

$$(\forall i, j, k \ 1 \le i, j, k \le n : x_k < x_j \implies a_{i,k} \le a_{i,j}) \Longrightarrow \rho(A) = \max_{B \in \Omega(A)} \rho(B)$$
(7)

$$(\forall i, j, k \ 1 \le i, j, k \le n : x_k < x_j \implies a_{i,k} \ge a_{i,j}) \Longrightarrow \rho(A) = \min_{B \in \Omega(A)} \rho(B)$$
(8)

Proof. Consider the condition on the left hand side of (7). Observe that we can assume without loss of generality that $a_{i,k} \leq a_{i,j} \Leftrightarrow a_{l,k} \leq a_{l,j}$ for any two rows *i* and *l* of *A*. Indeed, if $x_k < x_j$ then this is the case (by the condition), and if $x_k = x_j$ then the entries $a_{i,k}$ and $a_{i,j}$ or $a_{l,k}$ and $a_{l,j}$ can be swapped without changing Ax, so that the modified matrix belongs to $\Omega(A)$ and has the same Perron eigenvector *x* and the same Perron root $\rho(A)$. Then we can also assume without loss of generality that simultaneously $x_1 \leq x_2 \leq \ldots \leq x_n$ and $a_{i,1} \leq a_{i,2} \leq \ldots \leq a_{i,n}$ for all *i*. If we consider any matrix $B \in \Omega(A)$, then the rearrangement inequality implies that $Bx \leq Ax = \rho(A)x$ and hence $\rho(B) \leq \rho(A)$.

Similarly, to prove the sufficiency of the condition on the right hand side of (10), we can assume without loss of generality that $a_{i,k} \ge a_{i,j} \Leftrightarrow a_{l,k} \ge a_{l,j}$ for any two rows *i* and *l* of *A*. Indeed, if $x_k < x_j$ then this is the case (by the condition), and if $x_k = x_j$ then the corresponding non-aligning entries in any row can be swapped to obtain the alignment. Then we can also assume without loss of generality that simultaneously $x_1 \le x_2 \le \ldots \le x_n$ and $a_{i,1} \ge a_{i,2} \ge \ldots \ge a_{i,n}$ for all *i*. If we consider any matrix $B \in \Omega(A)$, then the rearrangement inequality implies that $Bx \ge Ax = \rho(A)x$ and hence $\rho(B) \ge \rho(A)$. \Box

The following result applies Lemma 2.1 to show that for irreducible matrices conditions (7) and (8) of Theorem 3.3 are necessary and sufficient for $\rho(A) = \max_{B \in \Omega(A)} \rho(B)$ or $\rho(A) = \min_{B \in \Omega(A)} \rho(B)$.

Theorem 3.4. Let $A \in \mathbb{R}^{n \times n}_+$ be irreducible and $0 < x \in \mathbb{R}^n_+$ be a Perron eigenvector of A. Then

$$\rho(A) = \max_{B \in \Omega(A)} \rho(B) \iff (\forall i, j, k \ 1 \le i, j, k \le n : x_k < x_j \implies a_{i,k} \le a_{i,j})$$
(9)

$$\rho(A) = \min_{B \in \Omega(A)} \rho(B) \iff (\forall i, j, k \ 1 \le i, j, k \le n : x_k < x_j \implies a_{i,k} \ge a_{i,j})$$
(10)

Proof. By Lemma 2.1, the conditions on the right hand sides of (9) and (10) are necessary. The fact that they are sufficient follows immediately from Theorem 3.3. \Box

4. Solving $\max_{B \in \Omega(A)} \rho(B)$ and $\min_{B \in \Omega(A)} \rho(B)$

Below we give two simple iterative procedures for solving $\max_{B \in \Omega(A)} \rho(B)$ and $\min_{B \in \Omega(A)} \rho(B)$. Note that the computation of the minimum and maximum spectral radius over sets more general than $\Omega(A)$ was investigated by Protasov [7] where similar iterative procedures were suggested.

Before presenting the iterative procedures we first establish the following lemmas.

Lemma 4.1. Let $A \in \mathbb{R}^{n \times n}_+$ and P be a permutation matrix. Then $\rho(PA) = \rho(AP)$.

Proof. It is easy to see that any eigenvalue of PA is an eigenvalue of AP and the other way around:

$$PAx = \alpha x \Rightarrow AP(P^{-1}x) = \alpha(P^{-1}x),$$
$$APy = \beta y \Rightarrow PA(Py) = \beta(Py). \quad \Box$$

Lemma 4.2. For $A \in \mathbb{R}^{n \times n}_+$ and all permutation matrices P

$$\max_{B \in \Omega(A)} \rho(B) = \max_{B \in \Omega(PA)} \rho(B)$$

and

$$\min_{B \in \Omega(A)} \rho(B) = \min_{B \in \Omega(PA)} \rho(B).$$

Proof. Take arbitrary $B \in \Omega(PA)$. Then B = PC, where $C \in \Omega(A)$, and by Lemma 4.1 $\rho(B) = \rho(CP)$, where $CP \in \Omega(A)$. This observation implies that

$$\max_{B \in \Omega(A)} \rho(B) \ge \max_{B \in \Omega(PA)} \rho(B), \quad \min_{B \in \Omega(A)} \rho(B) \le \min_{B \in \Omega(PA)} \rho(B).$$

The reverse inequalities follow from a similar argument where we start with $B = \Omega(A)$ and represent $B = P^{-1}C$ with $C \in \Omega(PA)$. \Box

Definition 4.3. A $n \times n$ matrix A is said to be fully indecomposable if PAQ is irreducible for all permutation matrices P and Q.

We now argue that Algorithm 1 (see below) is valid. Observe that if in step 6 vector x is not in ascending order and hence P is not the identity matrix, then $C(Px) \ge Cx = \rho(C)x$ with at least one strict inequality, since all rows of C as well as Px are aligned together in ascending order, but this is not true about all rows of C and vector x. Then we obtain $(PC)Px \ge \rho(C)Px$ with at least one strict inequality, and by Proposition 1.3 part (iii) $\rho(C) < \rho(PC)$. If P is the identity matrix then $\forall i, j, k \ 1 \le i, j, k \le n : x_k < x_j \implies c_{i,k} \le c_{i,j}$ and by Theorem 3.4 $\rho(C) = \max_{B \in \Omega(C)} \rho(B)$. By Lemma 4.2 it

Algorithm 1 Solving $\max_{B \in \Omega(A)} \rho(B)$.

Input: $A \in \mathbb{R}^{n \times n}_+$ with A fully indecomposable.

- 1: Define matrix $C_0 \in \Omega(A)$ by placing the entries in each row of A in ascending order. 2: Find a permutation matrix $Q \in \mathbb{R}^{n \times n}_+$ such that the Euclidean norms of the rows of QC are in ascending
- order. 3: $P \in \mathbb{R}^{n \times n}_+$ is the zero matrix, $C := QC_0$.
- 4: while P is not the identity matrix do
- 5: Find a Perron eigenvector $x \in \mathbb{R}^n_+$ of C
- 6:
- if the entries of x are not in ascending order then Find a permutation matrix $P \in \mathbb{R}^{n \times n}_+$ so that entries of Px are in ascending order. 7:
- 8: else
- 9: Set P to be the identity matrix.
- 10: end if

C := PC, Q := PQ.11:

12: end while

Output: C_0Q , $\rho(C_0Q) = \max_{B \in \Omega(A)} \rho(B)$

Algorithm 2 Solving $\min_{B \in \Omega(A)} \rho(B)$.

Input: $A \in \mathbb{R}^{n \times n}_+$ with A fully indecomposable.

- 1: Define matrix $C_0 \in \Omega(A)$ by placing the entries in each row of A in descending order.
- 2: Find a permutation matrix $Q \in \mathbb{R}^{n \times n}_+$ such that the Euclidean norms of the row sums of QC are in descending order.
- 3: $P \in \mathbb{R}^{n \times n}_+$ is the zero matrix, $C := QC_0$. 4: while P is not the identity matrix do
- Find a Perron eigenvector $x \in \mathbb{R}^n_+$ of C5:
- 6:
- if the entries of x are not in descending order then Find a permutation matrix $P \in \mathbb{R}_{+}^{n \times n}$ so that entries of Px are in descending order. 7:
- 8: else Set P to be the identity matrix. 9:
- 10: end if
- $C := PC, \, Q := PQ.$ 11.

12: end while

Output: C_0Q , $\rho(C_0Q) = \min_{B \in \Omega(A)} \rho(B)$

follows that $\rho(C) = \max_{B \in \Omega(A)} \rho(B)$. The algorithm terminates in a finite number of iterations since $\rho(C)$ is strictly increasing so matrices C do not repeat, and since the number of permutations is finite. Lemma 4.1 also implies that for the final matrix C we have $\rho(C) = \rho(C_0Q)$, implying that C_0Q solves the problem of maximizing $\rho(B)$ over $\Omega(A)$ (while belonging to $\Omega(A)$).

Algorithm 2 is valid for the reasons similar to those explained above for Algorithm 1. We now demonstrate the work of Algorithm 1 on the following small example.

Example 4.4. Consider matrix

$$A = \begin{pmatrix} 2 & 5 & 2 & 2 & 5 \\ 6 & 6 & 2 & 3 & 1 \\ 7 & 3 & 5 & 5 & 3 \\ 3 & 3 & 4 & 6 & 8 \\ 2 & 4 & 2 & 5 & 5 \end{pmatrix}$$

First we align all rows of this matrix in ascending order thus obtaining C_0 . The Euclidian norms of the row sums if C are 11, 12, 18, 27 and 31. Thus initially Q = I and C = $QC_0 = C_0$ with its Perron vector x:

231

$$C = QC_0 = \begin{pmatrix} 2 & 2 & 2 & 5 & 5\\ 1 & 2 & 3 & 6 & 6\\ 3 & 3 & 5 & 5 & 7\\ 3 & 3 & 4 & 6 & 8\\ 2 & 2 & 4 & 5 & 5 \end{pmatrix}, \quad x \approx \begin{pmatrix} 0.3561\\ 0.4098\\ 0.5091\\ 0.5301\\ 0.4063 \end{pmatrix}$$

The components of x are not ascending and we have:

$$P = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 \\ 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 \end{pmatrix}$$

The next while loop proceeds, since $P \neq I$. We compute the next matrix C and its Perron eigenvector x:

$$C := PQC_0 = \begin{pmatrix} 2 & 2 & 2 & 5 & 5 \\ 2 & 2 & 4 & 5 & 5 \\ 1 & 2 & 3 & 6 & 6 \\ 3 & 3 & 5 & 5 & 7 \\ 3 & 3 & 4 & 6 & 8 \end{pmatrix}, \quad x \approx \begin{pmatrix} 0.3595 \\ 0.3987 \\ 0.4116 \\ 0.5055 \\ 0.5355 \end{pmatrix}$$

Here, x is in the ascending order. The algorithm ends and returns

$$C_0 PQ = \begin{pmatrix} 2 & 2 & 5 & 5 & 2 \\ 1 & 3 & 6 & 6 & 2 \\ 3 & 5 & 5 & 7 & 3 \\ 3 & 4 & 6 & 8 & 3 \\ 2 & 4 & 5 & 5 & 2 \end{pmatrix}, \quad \rho(C_0 PQ) \approx 20.9863.$$

Remark 4.5. We conducted a number of numerical experiments, in which we increased the matrix dimension from 5 to 200. For each dimension we generated 50 random instances of A and counted the number of while loops that Algorithms 1 and 2 require before convergence. For the whole dimension range, the average number of while loops stayed with the maximum number of loops not exceeding 3. Finding a reasonable upper bound on the number of loops before convergence is an open problem. Note that Cvetković and Protasov [4] establish that a similar algorithm has local quadratic convergence (see [4], page 19).

Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Data availability

No data was used for the research described in the article.

References

- Yu.A. Al'pin, Bounds for the Perron root of a non negative matrix involving properties of its graph, Math. Notes (Moscow) 58 (1995) 1121–1123.
- [2] L. Elsner, P. van den Driessche, Bounds for the Perron root using max eigenvalues, Linear Algebra Appl. 428 (2005) 2000–2005.
- [3] A. Berman, R.J. Plemmons, Nonnegative Matrices in Mathematical Sciences, Academic Press, New York, 1979.
- [4] A. Cvetković, V.Yu. Protasov, The greedy strategy for optimizing the Perron eigenvalue, Math. Program. 193 (2022) 1–31.
- [5] G.M. Engel, H. Schneider, S. Sergeev, On sets of eigenvalues of matrices with prescribed row sums and prescribed graph, Linear Algebra Appl. 455 (2014) 187–209.
- [6] G.H. Hardy, J.E. Littlewood, G. Polya, Inequalities, 2nd edition, Cambridge University Press, 1953.
- [7] V.Yu. Protasov, Spectral simple method, Math. Program., Ser. A 156 (2016) 485-511.