

Novel Insights into the Dermal Bioaccessibility and Human Exposure to Brominated Flame Retardant Additives in Microplastics

Abafe, Ovokeroye; Harrad, Stuart; Abdallah, Mohamed

DOI:

[10.1021/acs.est.3c01894](https://doi.org/10.1021/acs.est.3c01894)

License:

Creative Commons: Attribution (CC BY)

Document Version

Publisher's PDF, also known as Version of record

Citation for published version (Harvard):

Abafe, O, Harrad, S & Abdallah, M 2023, 'Novel Insights into the Dermal Bioaccessibility and Human Exposure to Brominated Flame Retardant Additives in Microplastics', *Environmental Science and Technology*.
<https://doi.org/10.1021/acs.est.3c01894>

[Link to publication on Research at Birmingham portal](#)

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

Novel Insights into the Dermal Bioaccessibility and Human Exposure to Brominated Flame Retardant Additives in Microplastics

Ovokeroye A. Abafe,* Stuart Harrad, and Mohamed Abou-Elwafa Abdallah



Cite This: <https://doi.org/10.1021/acs.est.3c01894>



Read Online

ACCESS |



Metrics & More




Article Recommendations



Supporting Information

ABSTRACT: In this study, we optimized and applied an *in vitro* physiologically based extraction test to investigate the dermal bioaccessibility of polybrominated diphenyl ethers (PBDEs) and hexabromocyclododecane (HBCDD), incorporated as additives in different types of microplastics (MPs), and assess human dermal exposure to these chemicals. The dermal bioaccessibility of PBDEs in polyethylene (PE) MPs was significantly higher ($P < 0.05$) than in polypropylene (PP) MPs. Both $\log K_{ow}$ and water solubility influenced the dermal bioaccessibility of PBDEs. For HBCDDs in polystyrene MPs, the dermally bioaccessible fractions were 1.8, 2.0, and 1.6% of the applied dose for α -, β -, and γ -HBCDDs, respectively. MP particle size and the presence of cosmetic formulations (antiperspirant, foundation, moisturizer and sunscreen) influenced the bioaccessibility of PBDEs and HBCDDs in MP matrices at varying degrees of significance. Human exposure to Σ PBDEs and Σ HBCDDs via dermal contact with MPs ranged from 0.02 to 22.2 and 0.01 to 231 $\text{ng (kg bw)}^{-1} \text{d}^{-1}$ and from 0.02 to 6.27 and 0.2 to 65 $\text{ng (kg bw)}^{-1} \text{d}^{-1}$ for adults and toddlers, respectively. Dermal exposure to PBDEs and HBCDDs in MPs is substantial, highlighting for the first time the significance of the dermal pathway as a major route of human exposure to additive chemicals in microplastics.

KEYWORDS: microplastics, additive chemicals, polybrominated diphenyl ethers, hexabromocyclododecane, dermal bioaccessibility, cosmetics, particle size



1. INTRODUCTION

Microplastics (MPs) have been widely reported to be present in the marine and freshwater environment with concentrations up to 102000 particles m^{-3} in seawater and are also detected in sediment, fish, soil, dust, air, food, and drink.^{1–4} Such ubiquitous distribution of MPs in the environment and consumer products inevitably leads to human exposure to these particles, which has been confirmed by the recent detection of MPs in human blood,⁵ lungs,⁶ and stool.⁷ Meanwhile, animal studies have indicated MP exposure to elicit reproductive toxicity in oysters, reduced feeding in daphnia, and hepatotoxicity in zebrafish, as well as tissue accumulation and disturbance of lipid metabolism in mice.⁶ However, the toxicological impacts of human exposure to MPs are not well understood, which *inter alia* may be attributed to ethical constraints, strict biosecurity measures to handle human samples, and limited analysis techniques.

While particle exposure may lead to inflammatory lesions, bioaccumulation, and oxidative stress, there is also concern over the potential release of toxic chemical additives/adsorbed contaminants from MPs to human body fluids due to their small size and corresponding larger surface area.¹ A wide range of chemical additives such as flame retardants, plasticizers, pigments, and fillers are often incorporated in plastics during manufacture to impart specific properties. Most of these

additives, particularly in the flame retardants group, e.g. polybrominated diphenyl ethers (PBDEs) and hexabromocyclododecane (HBCDD), have been found to cause adverse health effects including endocrine disruption, reproductive toxicity, neurotoxicity, hepatotoxicity, and cancer.⁸ A recent study reported the leaching of toxic brominated flame retardants (BFRs) from acrylonitrile-butadiene-styrene copolymer (ABS) plastics to the *in vitro* simulated digestive fluids of birds, rendering them available for absorption, *i.e.* bioaccessible;² however, there is no available information for humans. Bioaccessibility is defined as the total amount of a chemical that is released from a solid matrix (e.g., dust, soil, MPs) into human or animal body fluids (e.g., saliva, gastric fluid, sweat) and thereby becoming available for absorption to the systemic circulation.⁹ In particular, there are no available data on the release and subsequent dermal uptake of toxic additive chemicals (e.g., flame retardants and plasticizers) to human skin surface film liquid (SSFL) upon contact of MPs with the

Received: March 14, 2023

Revised: June 30, 2023

Accepted: July 3, 2023

skin, which is the largest body organ.¹⁰ The human SSFL consists of a mixture of sweat and sebum at various proportions.¹¹ Sweat is an aqueous-based fluid secreted from sweat glands to regulate body temperature. It consists mainly of electrolytes, amino acids, organic acids, vitamins, and other nitrogenous substances. Sebum is a clear, oily liquid secreted from sebaceous glands to protect the skin from drying out. It mainly consists of triglycerides, squalene, wax esters, and fatty acids, with a small amount of cholesterol and cholesteryl esters.¹²

Dermal risk assessment has been traditionally based on the quantitative structure–activity relationship (QSAR)^{13–15} and pharmacokinetic (PK) modeling.¹⁶ These methods are limited by uncertainties related to (a) the fraction of chemicals available for absorption in PK modeling (i.e., bioaccessible fraction), (b) the partitioning between exposure doses in various solvent vehicles and the stratum corneum (the uppermost layer of the skin) especially for nonaqueous exposure vehicles, and (c) chemical diffusivity through the skin owing to differences in thickness of the stratum corneum among species.¹⁷ *In vitro* physiologically based extraction tests (PBETs) have emerged as realistic alternatives to the traditional methods for assessing the bioaccessibility of hazardous chemicals from solid matrices and have been incorporated in regulatory risk assessment frameworks (BS EN 1811, 2011) and applied successfully to study the bioaccessibility of polychlorinated biphenyls (PCBs)¹⁸ and BFRs in house dust.¹⁷

The current understanding is that human exposure to MPs occurs through a combination of ingestion, inhalation, and dermal contact due to the presence of MPs in indoor dust, consumer products, water, foodstuffs, and air.¹⁹ While preliminary assessments of human exposure to MPs via ingestion and inhalation exist,¹⁹ there are no data on dermal exposure to MPs or assessment of the risk arising from such exposure. Recent studies by our research group highlighted the significance of dermal contact with synthetic furniture fibers as a major contributor to human body burdens of PBDEs and HBCDD²⁰ and chlorinated organophosphate flame retardants.²¹ The adverse effects of flame retardants, combined with widespread dermal contact with MP particles (e.g., present in dust adhering to skin, atmospheric deposition of synthetic microfibers, and microbeads in personal care products), highlight the crucial need for further research into this area. Consequently, the current study assesses, for the first time, the dermal bioaccessibility of PBDEs and HBCDD incorporated in microplastics upon contact with synthetic human skin surface film liquid (i.e., sweat–sebum mixture). The factors influencing dermal bioaccessibility of these BFRs from MPs are evaluated, including: MP polymer type and particle size, BFR-specific physicochemical properties, and impact of topically applied cosmetics. Finally, the estimated bioaccessible fraction of the studied BFRs was applied to assess human dermal exposure to these toxic additive chemicals from various MP types, including polyethylene (PE), polypropylene (PP), and polystyrene (PS), using realistic but conservative exposure scenarios.

2. MATERIALS AND METHODS

2.1. Chemicals and Reagents. All solvents used for sample preparation and gas and liquid chromatographic analysis were of HPLC grade (Fisher Scientific, Loughborough, United Kingdom). Individual standards of BDEs

28, 47, 99, 100, 153, 154, 183, 209, 77, 128, ¹³C₁₂-BDE 100, and ¹³C₁₂-BDE 209, as well as single standards of α -, β -, and γ -HBCDD, *d*₁₈- γ -HBCDD, and ¹³C₁₂- α -, β -, and γ -HBCDD were purchased from Wellington Laboratories Inc., Ontario, Canada. Two standard reference materials, European Reference Material for polyethylene (ERM-EC-590) and polypropylene (ERM-EC-591) certified for polybrominated diphenyl ether (PBDE) concentrations, were purchased from the European Joint Research Centre Institute for Reference Materials and Measurements (Brussels, Belgium). An in-house laboratory reference material of extruded polystyrene with known concentrations of α -, β -, and γ -hexabromocyclododecane (HBCDD) was obtained from the National Institute for Environmental Studies (NIES, Tsukuba City, Ibaraki, Japan). Cosmetics including sunscreen, antiperspirant, moisturizing cream, and foundation were of popular commercial brands purchased from a local UK supermarket.

2.2. Microplastic Samples. Microplastics from ERM-EC-590 and ERM-EC-591 with an original pellet particle size of approximately 4 mm were used in the study with MPs of particle size <0.45 mm produced from the original pellets using a Fritsch Pulverisette 0 cryo-vibratory micro mill (Idar-Oberstein, Germany). Frozen (−80 °C) plastic pellets of approximately 4 mm were placed in the stainless steel grinding mortar (50 mL volume) together with a 25 mm diameter stainless steel ball and submerged in liquid nitrogen (−196 °C) to aid the pulverisation process. The sample was ground at a vibrational frequency of 30 Hz for 5 min and repeated 3 times, resulting in plastic particles that passed through a 0.45 mm mesh aluminum sieve.

Extruded polystyrene samples were cut into small pieces, grated with a fine particle diameter stainless steel kitchen grater, and filtered through a 0.45 mm mesh stainless steel sieve. Two particle sizes of 3.5–4 mm (original pellet size provided by the manufacturer) and <0.45 mm were used for polyethylene and polypropylene MP experiments, while extruded polystyrene of particle size <0.45 mm was used for experiments related to HBCDDs.

2.3. Preparation of Synthetic Sweat and Sebum Mixture. The preparation of a physiologically simulated synthetic sweat and sebum mixture (SSSM) was conducted according to a US patent (US20080311613A1) using more than 25 organic and inorganic components (see Table S1 in the Supporting Information).¹² The pH was adjusted to the physiological pH of the human skin surface film liquid (SSFL) (5.3 ± 0.1). Synthetic sweat and sebum mixtures were individually prepared and then mixed in different physiologically relevant ratios (Table S2) using drops of Tween-80 to mimic the naturally secreted SSFL and prevent phase separation.^{22,23}

2.4. Physiologically Based Extraction Test (PBET) Protocol. The PBET protocol in this study was adopted from Pawar et al.¹⁷ and Ertl and Butte.¹⁸ Briefly, approximately 60 mg of microplastic sample (PE, PP, and PS) and approximately 6 mg of cosmetic (when tested) were weighed into a predried and sterilized glass test tube. Since no definitive data on the ratio of MPs to SSSM are available, we adopted a ratio of 1:100 w/v MP to SSSM ratio (i.e., 60 mg of each MP to 6 mL SSSM) to mimic “wet skin conditions” as previously reported.¹⁷ The mixture was gently agitated on a magnetic stirrer hot plate maintained at the physiological temperature of the human skin (32 ± 3 °C). Following 1 h of agitation, the mixture was phase-separated by centrifugation at 3500 rpm for

10 min. The supernatant (SSSM) and the solute (MPs) were analyzed separately. All experiments were carried out in triplicate. Various physiologically relevant SSFL compositions (i.e., different sweat:sebum ratios) were tested (Table S2).

2.5. Sample Extraction and Cleanup. Sample extraction and cleanup were conducted according to a previously reported method²⁴ with slight modification. Briefly, each sample (PE and PP) was spiked with 100 ng of internal (surrogate) standard mixtures (BDE 128, ¹³C₁₂-BDE 100 and -BDE 209), while PS samples were spiked with 60 ng of a ¹³C₁₂- α -, β -, and γ -HBCDD internal standard mixture. This was followed by the addition of 3 mL of dichloromethane (DCM). The mixture was vortexed for 2 min followed by ultrasonication for 5 min and then centrifuged at 3500 rpm for 5 min. The organic phase was collected into a separate precleaned and sterilized glass test tube. The procedure was repeated twice. The collected extracts were evaporated to approximately 2 mL under a gentle stream of nitrogen set at 40 °C. 2 mL of hexane was added to precipitate any dissolved plastic and then reduced to approximately 1 mL and reconstituted in 2 mL of hexane to completely remove DCM followed by vortex mixing. Approximately 3 mL of concentrated sulfuric acid was added to samples and then vortexed for 1 min. The mixtures were left to stand for at least 5 h and then centrifuged at 3500 rpm for 5 min for phase separation. The organic layer was collected into a clean test tube. The sulfuric acid phase was further extracted twice with the addition of 2 mL of *n*-hexane, vortexed for 2 min, and centrifuged at 3500 rpm for 5 min. All the organic phases were combined and reduced to incipient dryness under a gentle stream of nitrogen at 40 °C. The extracts for PBDE analysis were reconstituted with 150 μ L of isooctane containing 250 pg μ L⁻¹ BDE-77 as a recovery determination (syringe) standard (RDS), while the extracts from the PS (i.e., for HBCDD analysis) were reconstituted in 150 μ L of methanol containing 250 pg μ L⁻¹ d₁₈- γ -HBCDD. Full details of the extraction protocol for each type of sample are presented in the Supporting Information.

2.6. Instrumental Determination of PBDEs and HBCDDs. The operating conditions of the GC-MS method

(for PBDEs analysis) and LC-MS/MS method (for HBCDDs analysis) used in the current study have been reported previously.^{24,17} Details of all instrumental parameters are presented in Table S3 in the Supporting Information.

2.7. Quality Assurance/Quality Control. A procedural blank containing SSSM without MPs was analyzed with every batch of 5 samples. None of the cosmetics were found to contain BFR concentrations above the LOQ of the target compounds; hence, results were not blank corrected. Method limits of detection (LODs) and limits of quantification (LOQs) were estimated based on S/N = 3:1 and 10:1, respectively. The LODs and LOQs ranged from 0.54 to 3.13 μ g kg⁻¹ and 1.87 to 9.50 μ g kg⁻¹ for PBDEs, while they ranged from 0.64 to 1.72 μ g kg⁻¹ and 1.94 to 5.22 μ g kg⁻¹ for HBCDDs, respectively.

The recovery of internal surrogate standards ranged from 115 to 117%, 61 to 97% and 40 to 127% for ERM-EC-590, ERM-EC-591, and ERM-EC-590 + cosmetics, respectively, while they ranged from 89 to 99% and 48 to 129% for PS and PS + cosmetics, respectively. To test the effectiveness of the method on the three MPs matrices following the bioaccessibility experiment, a mass balance was carried out as the sum of the concentrations of PBDEs and HBCDDs in the SSSM and in the residual MPs relative to the original concentrations determined in the MP matrices prior to bioaccessibility experiments. The mass balance recovery of PBDEs ranged from 80 to 152% and 56 to 103%, respectively, in PE and PP MPs at the physiologically relevant sweat:sebum (i.e., 1:1 sweat: sebum) mixture (Figure S1 in the Supporting Information). Whereas the mass balance recovery of HBCDDs ranged from 67 to 94% in polystyrene (Figure S2 in the Supporting Information), indicating the effectiveness of the analytical method for the determination of the target analytes in the respective matrices. Further quality control measures detailing the validation and reliability of the analytical method in this study are provided in Tables S4–S7 in the Supporting Information).

2.8. Bioaccessibility and Human Dermal Exposure to Additive BFRs in MPs. The bioaccessible fraction of each target chemical from each MP type is calculated from eq 1

$$f_{\text{bioaccessible}} = \frac{\text{concentration of analyte in SSSM}}{\text{concentration of analyte in SSSM} + \text{concentration of analyte in the solute (i.e. MP residue)}} \times 100 \quad (1)$$

where the concentration of analyte in SSSM is the amount of chemical released into the synthetic sweat:sebum mixture following the bioaccessibility experiment and the concentration of analytes in the solute is the amount of chemical remaining in the MP particles after the bioaccessibility experiment.

In assessing human dermal exposure to BFRs present as MP additives in the indoor environment, we employed 0.12 and 0.002 weight fractions of MPs in indoor dust (i.e., 0.12 g MPs/g dust for high-exposure and 0.02 g MPs/g dust for low-exposure scenario), as reported previously.²⁵ Other factors used for the assessment of the dermal exposure dose (DED) of BFRs in MPs were obtained from the USEPA exposure factor handbook²⁶ as presented in Table S8 in the Supporting Information.

The human dermal exposure dose (DED) to the target BFR was estimated using eq 2²⁶

$$\text{DED} = \frac{C \times \text{BSA} \times \text{MPAS} \times F_A \times \text{IEF} \times \text{MPF}}{\text{BW}} \quad (2)$$

where DED = daily exposure dose (ng kg⁻¹ bw day⁻¹), C = concentration of BFRs in MPs (ng/g), BSA = body surface area exposed (cm²), MPAS = MPs adhered to skin (mg/cm²), F_A = bioaccessible fraction (unitless), IEF = indoor exposure fraction (hours spent over a day in an indoor environment) (unitless), MPF = fraction of MPs in indoor dust (unitless), and BW = body weight (kg).

2.9. Statistical Analysis. The distribution of the data in this study was tested with the Shapiro–Wilk test and was found to be normally distributed. Descriptive statistics (e.g., mean, standard deviation etc.) were generated using Microsoft Office Excel 2016. Parametric tests (e.g., analysis of variance, Pearson's correlation, etc.) were performed with XLSTAT Version 2021.3.1. *p* < 0.05 was deemed significant.




Figure 1. Impact of sweat:sebum composition on the dermal bioaccessibility of PBDEs in (a) PE MP pellets and (b) PP MP pellets.

3. RESULTS AND DISCUSSION

3.1. Dermal Bioaccessibility of Polybrominated Diphenyl Ethers in Polyethylene and Polypropylene Microplastics.

The skin, being the largest organ of the human body, is well-known to protect against pollutants, bacteria, ultraviolet radiation, etc.; however, it is consistently exposed to a wide range of xenobiotics both intentionally through the application of cosmetics or drugs and unintentionally through exposure to indoor and outdoor particle-bound environmental pollutants.²⁷ While the penetration of a xenobiotic compound through the human skin is established to follow a passive diffusion process governed mainly by compound-specific physicochemical properties,²⁸ for chemicals bound to solid matrices (e.g., particulate matter, MPs), the chemical's release from the matrix into the human body fluid (i.e., SSFL) can be more important.^{18,29} In other words, a chemical within a solid matrix (e.g., MPs) must become *bioaccessible* first, in order to be available for absorption depending on its ability to penetrate the skin barrier.⁹

The results of dermal bioaccessibility of PBDEs in polyethylene and polypropylene MPs are presented in Figure 1a,b. The bioaccessibility of PBDEs in PE and PP MPs increased with an increasing sebum ratio for all PBDE congeners. At the most physiologically relevant sweat: sebum composition (1:1), $f_{\text{bioaccessible}}$ ranged from 1.7% for BDE 209 to 5.5% for BDE 153 and ~12% for BDE 183 to ~39% for BDE 209 in PP and PE microplastic pellets, respectively. These results prove, for the first time, the release of PBDEs from MPs into the SSFL, which is the first step in dermal uptake of these contaminants. BDE 209 (~39%) was the most bioaccessible PBDE in PE MP pellets, with $f_{\text{bioaccessible}}$ values of BDE-47, -99, -100, -153, and -154 ranging between 24 and 28%. The high bioaccessibility of BDE-209 may have been due to the higher concentration of this congener compared to the other BDE congeners in the PE and PP polymer matrices, which was at least 1 order of magnitude higher than the concentrations of most of the penta- and hexaBDE congeners, i.e. dose dependent. Also, BDE-209 being the most lipophilic of the PBDE congeners could have accumulated in the fat-rich (50%) synthetic skin surface film liquid. This is supported by the strong linear relationship between the bioaccessibility of PBDEs and fat contents in different food types reported by Yu et al.,³⁰ as well as the observations of Hornero-Mendez and Minguez-Mosquera,³¹ in which the addition of olive oil to carrot prior to a bioaccessibility experiment significantly

increased the % bioaccessible fraction of carotenoid ($\log K_{\text{ow}}$ 17.62).

The physicochemical parameters of PBDEs and HBCDDs are presented in Tables S9 and S10. The $f_{\text{bioaccessible}}$ values for individual PBDE congeners in PE MP pellets showed moderate correlation ($p < 0.05$) with $\log K_{\text{ow}}$ ($r = 0.45$), $\log K_{\text{oa}}$ ($r = 0.36$) and $\log K_{\text{oc}}$ ($r = 0.15$). There were no significant associations between the bioaccessibility of the studied PBDE congeners and their water solubility or vapor pressure. Similarly, no association ($p > 0.05$) was observed between $f_{\text{bioaccessible}}$ of PBDEs in PP MPs and their physicochemical properties ($\log K_{\text{oa}}$, $\log K_{\text{ow}}$, $\log K_{\text{oc}}$, water solubility, and vapor pressure). This indicates that while some physicochemical properties of PBDEs influenced their leaching from PE MPs, such influences were less influential drivers of the bioaccessibility of PBDEs in PP MP pellets.

The bioaccessibility of PBDEs in PE and PP microplastics differed significantly ($p = 0.0002$), with PE presenting higher $f_{\text{bioaccessible}}$ values of PBDEs compared to PP MPs. This observation could be related to the lower relative density (0.92 g/cm³) of PP MPs, compared to PE MPs (0.97 g/cm³), which limits their sinking/rising behavior in liquid media, resulting in less interaction between the PP MPs and the SSFL. The decreased diffusivity of penetrants in polypropylene plastics has been previously noted.³² The diffusion coefficients of several hydrophobic organic compounds have been shown to be consistently lower in PP than in PE,³³ a phenomenon ascribed to the higher degree of branched chain carbons in PP. This is further exacerbated by the rigidity and the strong oil resistance properties of polypropylene plastics,³⁴ which could repel the lipid components of human sweat:sebum mixtures.

Interestingly, the $f_{\text{bioaccessible}}$ values of PBDEs in MPs in the present study were generally lower than the $f_{\text{bioaccessible}}$ values reported for PBDEs in dust using an *in vitro* human gastrointestinal (GIT) PBET,³⁵ with the exception of BDE 209, which was slightly more bioaccessible in our study. Overall, while the $f_{\text{bioaccessible}}$ values of tri- to octaBDEs were generally higher in previous GIT PBET studies in indoor dust, the bioaccessibility of BDE 209 in MPs via the dermal pathway in the present study exceeded those estimated via the oral route using a colon-extended GIT model,³⁶ dialysis membrane with the Tenax method³⁷ and a fasting GIT model,³⁸ highlighting the significance of dermal uptake of this lipophilic group of chemicals.^{36–38}

3.1.1. Impact of Particle Size on the Bioaccessibility of PBDEs in PE and PP Microplastics. To determine the impact




Figure 2. Impact of the particle size on the bioaccessibility of PBDEs in (a) PE and (b) PP MPs.

of particle size on the bioaccessibility of PBDEs in MPs, the original PE and PP MP pellets (3.5–4 mm) were crushed and passed through a 0.45 mm sieve. The bioaccessibility of tri- to octaBDE in the 0.45 mm MP particles thus generated was approximately double that in the original pellets. The $f_{\text{bioaccessible}}$ values of PBDEs ranged from 12 to 39% and 37 to 82%, respectively, for the <4 and <0.45 mm polyethylene MP particles (Figure 2a). By comparison they ranged from 1.6 to 5.5% and 17 to 54% for the <4 and <0.45 mm polypropylene MP particles (Figure 2b), respectively. The $f_{\text{bioaccessible}}$ values of PBDEs in the <4 and 0.45 mm fractions differed significantly ($p < 0.05$) in both the PE and PP MP particles. These differences could be due to the greater exterior surface area of the smaller particles coming into contact with the sweat:sebum mixture, leading to increased leaching of PBDEs into the skin surface film liquid. Our results highlight the impact of MP particle size and polymer type on the leaching behavior of PBDEs into human skin surface film liquid. This is consistent with the findings of Guo et al.,² in which the leaching of PBDEs from plastics into the digestive fluids of birds increased with decreased particle size.

3.1.2. Impact of Cosmetics on the Dermal Bioaccessibility of PBDEs from MPs. Cosmetics and their ingredients potentially remain on the skin for a long period of time and could alter the composition and physicochemical properties of the skin surface film liquid, which could potentially influence the bioaccessibility of chemicals from solid matrices in contact with skin.¹⁷ To investigate the influence of commonly applied cosmetics, we determined the $f_{\text{bioaccessible}}$ values of PBDEs in PE and PP MPs into a 1:1 sweat:sebum mixture in the presence of antiperspirant, moisturizer, foundation, and sunscreen. The $f_{\text{bioaccessible}}$ values of PBDEs in the presence of each of these cosmetics were compared with the $f_{\text{bioaccessible}}$ value of the control group (i.e., same conditions but without any cosmetics). As shown in Figure 3, the bioaccessibility of BDE-209 increased with the introduction of antiperspirant and foundation but not with the moisturizer and sunscreen, which did not impact BDE-209 bioaccessibility. Though the cause of this observation is not fully understood, one plausible explanation could be that the ingredients in the foundations and antiperspirants impacted the surface tension of the sweat:sebum mixture, hence enhancing the solubility of the highly lipophilic BDE-209 in the 1:1 sweat:sebum mixture. This is supported by the strong positive correlation of the $f_{\text{bioaccessible}}$ values of PBDEs following the application of antiperspirant and foundation with $\log K_{\text{ow}}$ ($r = 0.86$; $r = 0.91$) and $\log K_{\text{oc}}$ ($r = 0.68$; $r = 0.77$). This is in line with previous studies in which the active ingredients in sunscreen




Figure 3. Impact of cosmetics on the bioaccessibility of PBDEs in MPs.

formulations were reported to enhance the dermal penetration of moderately lipophilic compounds, e.g. 2,4-dichlorophenoxyacetic acid.^{39–41}

Conversely, the application of deodorant, foundation, moisturizer, and sunscreen inhibited the release of tri- to octaBDE from MPs into the sweat:sebum (1:1) mixture (Figure 3). Previous studies have shown that the presence of certain cosmetics decreased the bioaccessibility of polychlorinated biphenyls.¹⁸ However, the causes of these observations require further investigation.

3.2. Bioaccessibility of HBCDDs from Polystyrene MPs. The $f_{\text{bioaccessible}}$ values of HBCDDs increased with an increase in sebum concentration. At lower sebum concentrations (i.e., at 100% sweat, 1% sebum, and 10% sebum), the $f_{\text{bioaccessible}}$ value of HBCDDs in PS MPs was highest for the α -isomer followed by the β - and γ -isomers. However, the reverse was the case upon increasing the sebum composition from 20 to 50%, in which the γ -isomer was the most bioaccessible. At 100% sebum, the bioaccessibility of the β -isomer was highest, reaching 6% of the applied dose. Under the most physiologically relevant SSFL composition (1:1 sweat:sebum), the $f_{\text{bioaccessible}}$ values were 1.6, 1.8, and 2.0%, respectively, for the α -, β -, and γ -HBCDD isomers (Figure 4). The bioaccessibility of HBCDDs showed a strong positive correlation with $\log K_{\text{ow}}$ ($r = 0.999$) and a fairly strong negative correlation with water solubility ($r = -0.766$), suggesting the influence of sebum on the release of HBCDDs from polystyrene MPs upon contact with the skin surface. In house dust, higher bioaccessibility of HBCDDs into a 1:1 sweat: sebum mixture (ranging from 41 to 50%) was reported previously.¹⁷ These values are an order of magnitude higher than the $f_{\text{bioaccessible}}$ values for PS MPs in the current study. One




Figure 4. Influence of SSM composition on the bioaccessibility of HBCDD in polystyrene MPs.

plausible explanation for this observation would be the differences in matrices and less facile leaching of HBCDDs incorporated into PS microplastics than for HBCDDs adsorbed onto the surface of dust particles.

The impact of cosmetics on the bioaccessibility of HBCDDs from PS MPs varied for the different isomers. The application of moisturizer increased the $f_{\text{bioaccessible}}$ values of α - and γ -HBCDD from 1.86 and 2.35% to 2.72 and 6.96%, respectively (Figure 5). Similarly, foundation cream increased the




Figure 5. Influence of cosmetics on the bioaccessibility of HBCDDs in polystyrene MPs.

bioaccessibility of γ -HBCDD from 2.35 to 7.23% but decreased the $f_{\text{bioaccessible}}$ values of α - and β -HBCDD from 1.86 and 1.91% to 1.16 and 0.25%, respectively. Both the sunscreen and antiperspirant decreased the $f_{\text{bioaccessible}}$ values of all three HBCDD isomers. Although the application of cosmetics influenced the bioaccessibility of HBCDDs differently, only sunscreen significantly ($p = 0.002$) decreased the $f_{\text{bioaccessible}}$ values of all 3 HBCDDs studied. Antiperspirant ($p = 0.06$), foundation ($p = 0.72$), and moisturizer ($p = 0.49$) did not significantly influence the bioaccessibility of HBCDDs. Similar decreases in the $f_{\text{bioaccessible}}$ values of HBCDDs in house dust has been previously noted¹⁷—a phenomenon ascribed to the retention of lipophilic chemicals by lipids present in the cosmetics.

The $f_{\text{bioaccessible}}$ values of HBCDDs following the application of cosmetics e.g. antiperspirant ($r = 0.862$), foundation ($r = 0.905$), and moisturizer ($r = 0.972$) showed strong correlation with isomer-specific $\log K_{\text{ow}}$ (Table S10).

3.3. Dermal Exposure to BFRs via Contact with MPs in Indoor Dust. The daily dermal exposure to PBDEs and HBCDDs (DED in units of $\text{ng (kg bw)}^{-1} \text{d}^{-1}$) via contact with microplastics in indoor dust for adults and toddlers is presented in Table 1. PentaBDE congeners (BDE-47 and -99) and BDE-209 were the main contributors to the total exposure to PBDEs arising from dermal contact with PE and PP MPs in indoor dust. The daily exposure dose for adults ranged from 0.05 to $22.34 \text{ ng (kg bw)}^{-1} \text{d}^{-1}$ and 0.001 to $0.37 \text{ ng (kg bw)}^{-1} \text{d}^{-1}$ for the high and low MP exposure scenarios, respectively (i.e., 0.12 g MPs/g dust for high-exposure and 0.02 g MPs/g dust for low-exposure scenario). While for toddlers, they ranged from 0.5 to $230 \text{ ng (kg bw)}^{-1} \text{d}^{-1}$ and 0.01 to $3.85 \text{ ng (kg bw)}^{-1} \text{d}^{-1}$, respectively, for high and low MP exposure scenarios. While the DED of PBDEs associated with polyethylene MPs exceeded their corresponding values in polypropylene MPs for the penta- and octaBDE congeners, polypropylene MPs delivered a higher exposure dose for BDE-209 in the indoor environment, albeit not statistically significant.

The DEDs of PBDEs in the current study were lower than the US EPA reference doses (RfD) of 2, 3, and $7 \mu\text{g (kg bw)}^{-1} \text{d}^{-1}$ for penta-, octa-, and deca-BDEs, respectively.⁴² While our exposure assessment for PBDEs in MPs was carried out with certified reference materials containing relatively high concen-

Table 1. Human Exposure to PBDEs ($\text{ng kg}^{-1} \text{bw d}^{-1}$) via Dermal Contact with MPs at Home

BFR	type of polymer											
	polyethylene				polypropylene				polystyrene			
	high intake (i.e., high MP intake)		low intake (i.e., low MPF)		high intake (i.e., high MP intake)		low intake (i.e., low MPF)		high Intake (i.e., high MP intake)		low intake (i.e., low MPF)	
	adult	toddler	adult	toddler	adult	toddler	adult	toddler	adult	toddler	adult	toddler
BDE-28	0.2	2	0.004	0.01	0.1	0.5	0.001	0.01				
BDE-47	13	131	0.2	2	9	89	0.14	1.5				
BDE-99	14	143	0.2	2	12	123	0.20	2				
BDE-100	3	31	0.1	0.5	2.3	23	0.04	0.4				
BDE-153	1	13	0.02	0.2	0.6	6	0.01	0.1				
BDE-154	0.7	7	0.01	0.1	0.3	3	0.01	0.1				
BDE-183	2	20	1.9	0.3	1	12	0.02	0.2				
BDE-209	20	211	0.34	3.5	22	231	0.4	4				
α -HBCDD									2.8	52	0.1	0.5
β -HBCDD									1.1	12	0.02	0.2
γ -HBCDD									6.3	65	0.1	1.1

trations of PBDEs, higher concentrations of BDE-209 ($260\text{--}2600\ \mu\text{g g}^{-1}$) have been previously reported in indoor dust from UK and US homes that were identified to contain Br-rich polymer fragments.⁴³ Another study also reported BFR concentrations ranging from 47000 to $69000\ \mu\text{g g}^{-1}$ for BDE 209 in dust containing fibers and particles abraded from BFR-treated polymeric materials.⁴⁴

For HBCDDs, DED values ranged from 1.13 to $6.27\ \text{ng (kg bw)}^{-1}\ \text{d}^{-1}$ and 0.02 to $0.10\ \text{ng (kg bw)}^{-1}\ \text{d}^{-1}$ for dermally exposed adults, while for toddlers, they ranged from 11.80 to $65.10\ \text{ng (kg bw)}^{-1}\ \text{d}^{-1}$ and 0.20 to $1.08\ \text{ng (kg bw)}^{-1}\ \text{d}^{-1}$ for the high- and low-exposure scenarios, respectively.

The results of the present study indicate that exposure to HBCDD via dermal uptake from MPs exceeds the lifetime average daily dose via dermal exposure through direct skin contact with flame-retarded curtains containing HBCDD concentrations an order of magnitude higher than those in the MPs in this study.⁴⁵ Similarly, results from previous studies on HBCDD exposure via frequent hand-to-mouth contact were significantly lower than their DEDs arising from MP exposure.⁴⁶ The present study highlights the significance of microplastics as a substantial source of human exposure to HBCDDs via the dermal pathway.

3.4. Study Limitations. This is the first study to experimentally assess human exposure to BFRs as chemical additives in MPs via any human exposure pathway; hence, it is not possible to compare the magnitude of dermal exposure from this pathway with other exposure pathways (e.g., inhalation and ingestion of MPs with BFR additives). While the DED values reported for these chemicals in this study highlight the significance of the dermal exposure pathway, the dermal exposure assessment in our study was conservative. We assumed that humans are only exposed to MPs via indoor dust containing as little as 0.2% (low-contact scenario) and 12% (high-contact scenario) of MPs⁴⁷ with a conservative fraction of 1% of these values adhered to a fixed body surface area and using a fixed bioaccessible fraction of additive BFRs obtained from a $1\ \text{h}$ experimental contact time. Also, the concentrations of BFRs in the certified reference materials are low compared with their concentrations in secondary MPs arising from flame-retarded plastics. These conservative scenarios are very unlikely, as the human skin is exposed daily to different kinds of MPs from various sources, including fabrics,⁴⁸ personal care products,⁴⁹ furniture,⁵⁰ outdoor dust, and atmospheric deposition.¹⁹

While dermal contact with MPs occurs via a variety of sources, it is important to state that the dermal exposure estimate in this study was based solely on the bioaccessible fraction of these chemicals, which ranged from 1.56 to 2% for HBCDDs and 17 to 82% for PBDEs. These amounts of chemicals may not cross the stratum corneum to the epidermis and dermis into systemic circulation to exert toxic effects (i.e., bioavailable). In reality, the amount of these chemicals that would be available for systemic circulation would likely be considerably lower owing to the hydrophilic nature of the epidermis and dermis as previously reported for the lipophilic chemical benzo[a]pyrene, which penetrated easily into the lipidic layers of the stratum corneum but for which diffusion through the hydrophilic epidermis and dermis was low.⁵¹ This is consistent with previous results from our research group that established an increasing dermal resistance to the penetration of the more lipophilic γ -HBCDD compared to α - and β -isomers²⁸ because as $\log K_{ow}$ increases, diffusive transport

through the aqueous barrier becomes more restricted, thereby decreasing absorption.

To the best of our knowledge, the current study provides the first insights into the dermal bioaccessibility of additive BFR chemicals from different types of MPs and assesses the subsequent human dermal exposure to these chemicals via skin contact with MPs in house dust. We established various factors influencing the bioaccessibility of BFRs—a major class of additive chemicals used in various plastic polymers. These include the physicochemical properties of the studied BFRs, the polymer type, and MP particle size, as well as the use/application of various cosmetic formulations. We found that humans can be substantially exposed to target BFRs via dermal contact with flame-retarded MPs. We recommend that future studies should focus on the dermal bioavailability of these chemicals (i.e., their transfer across the skin barrier into the systemic circulation) for a more accurate quantification of the risk arising from dermal exposure to MPs. Moreover, it is important to establish the magnitude of human exposure to these toxic chemicals in MPs via other pathways e.g. ingestion and inhalation, in order to accurately examine the cumulative exposure dose and risk arising from such exposure.

■ ASSOCIATED CONTENT

SI Supporting Information

The Supporting Information is available free of charge at <https://pubs.acs.org/doi/10.1021/acs.est.3c01894>.

Detailed description of the composition of the skin surface film liquids, detailed description of the extraction method and the optimized instrumental method parameters for gas chromatography-mass spectrometry and liquid chromatography with tandem mass spectrometry analyses, and detailed validation data for the experimental protocol and quality control (PDF)

■ AUTHOR INFORMATION

Corresponding Author

Okokeroye A. Abafe — School of Geography, Earth and Environmental Sciences, University of Birmingham, Birmingham B15 2TT, United Kingdom; orcid.org/0000-0002-5672-6463; Email: o.abafe@bham.ac.uk

Authors

Stuart Harrad — School of Geography, Earth and Environmental Sciences, University of Birmingham, Birmingham B15 2TT, United Kingdom; orcid.org/0000-0003-4650-0564

Mohamed Abou-Elwafa Abdallah — School of Geography, Earth and Environmental Sciences, University of Birmingham, Birmingham B15 2TT, United Kingdom

Complete contact information is available at:

<https://pubs.acs.org/10.1021/acs.est.3c01894>

Notes

The authors declare no competing financial interest.

■ ACKNOWLEDGMENTS

This project has received funding from the European Union's Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie Individual Fellowship Grant Agreement Number 101026229.

REFERENCES

- (1) Prata, J. C.; da Costa, J. P.; Lopes, I.; Duarte, A. C.; Rocha-Santos, T. Environmental Exposure to Microplastics: An Overview on Possible Human Health Effects. *Sci. Total Environ.* **2020**, *702*, 134455.
- (2) Guo, H.; Zheng, X.; Luo, X.; Mai, B. Leaching of Brominated Flame Retardants (BFRs) from BFRs-Incorporated Plastics in Digestive Fluids and the Influence of Bird Diets. *J. Hazard Mater.* **2020**, *393*, 122397.
- (3) Zhang, Q.; Xu, E. G.; Li, J.; Chen, Q.; Ma, L.; Zeng, E. Y.; Shi, H. A Review of Microplastics in Table Salt, Drinking Water, and Air: Direct Human Exposure. *Environ. Sci. Technol.* **2020**, *54* (7), 3740–3751.
- (4) Peng, J.; Wang, J.; Cai, L. Current Understanding of Microplastics in the Environment: Occurrence, Fate, Risks, and What We Should Do. *Integr. Environ. Assess. Manag.* **2017**, *13* (3), 476–482.
- (5) Leslie, H. A.; van Velzen, M. J. M.; Brandsma, S. H.; Vethaak, A. D.; Garcia-Vallejo, J. J.; Lamoree, M. H. Discovery and Quantification of Plastic Particle Pollution in Human Blood. *Environ. Int.* **2022**, *163*, 107199.
- (6) Jenner, L. C.; Rotchell, J. M.; Bennett, R. T.; Cowen, M.; Tentzeris, V.; Sadofsky, L. R. Detection of Microplastics in Human Lung Tissue Using MFTIR Spectroscopy. *Sci. Total Environ.* **2022**, *831*, 154907.
- (7) Schwabl, P.; Köppel, S.; Königshofer, P.; Bucsecs, T.; Trauner, M.; Reiberger, T.; Liebmann, B. Detection of Various Microplastics in Human Stool. *Ann. Int. Med.* **2019**, *171* (7), 453–457.
- (8) Hahladakis, J. N.; Velis, C. A.; Weber, R.; Iacovidou, E.; Purnell, P. An Overview of Chemical Additives Present in Plastics: Migration, Release, Fate and Environmental Impact during Their Use, Disposal and Recycling. *J. Hazard Mater.* **2018**, *344*, 179–199.
- (9) Ragnarsdóttir, O.; Abdallah, M. A. E.; Harrad, S. Dermal Uptake: An Important Pathway of Human Exposure to Perfluoroalkyl Substances? *Environ. Pollut.* **2022**, *307*, 119478.
- (10) Zimmermann, K. A. Skin: The Human Body's Largest Organ. *Live Science*. **2018**; October 22, 2018.
- (11) Nicolaides, N. Skin Lipids: Their Biochemical Uniqueness. *Science* (1979) **1974**, *186* (4158), 19.
- (12) Stefaniak, B. A.; Harvey, J. C. Artificial Skin Surface Film Liquids. US Patent 20080311613A1, December 18, 2008.
- (13) Balls, M.; Bogni, A.; Bremer, S.; Casati, S.; Coecke, S.; Eskes, C.; Prieto, P.; Sabbioni, E.; Worth, A.; Zuang, V.; Barratt, M.; Blaauboer, B.; Botham, P.; Combes, R.; Doehmer, J.; Fentem, J.; Liebsch, M.; Spielmann, H. Alternative (Non-Animal) Methods for Chemicals Testing: Current Status and Future Prospects. *ATLA Alternatives to Laboratory Animals* **2002**, *30*, 1.
- (14) Chen, L.; Han, L.; Lian, G. Recent Advances in Predicting Skin Permeability of Hydrophilic Solutes. *Adv. Drug Delivery Rev.* **2013**, *65*, 295.
- (15) Fitzpatrick, D.; Corish, J.; Hayes, B. Modelling Skin Permeability in Risk Assessment - The Future. *Chemosphere* **2004**, *55* (10), 1309.
- (16) Anissimov, Y. G.; Jepps, O. G.; Dancik, Y.; Roberts, M. S. Mathematical and Pharmacokinetic Modelling of Epidermal and Dermal Transport Processes. *Adv. Drug Delivery Rev.* **2013**, *65*, 169.
- (17) Pawar, G.; Abdallah, M. A. E.; de Sáa, E. V.; Harrad, S. Dermal Bioaccessibility of Flame Retardants from Indoor Dust and the Influence of Topically Applied Cosmetics. *J. Expo Sci. Environ. Epidemiol.* **2017**, *27* (1), 100–105.
- (18) Ertl, H.; Butte, W. Bioaccessibility of Pesticides and Polychlorinated Biphenyls from House Dust: In-Vitro Methods and Human Exposure Assessment. *J. Expo Sci. Environ. Epidemiol.* **2012**, *22* (6), 574–583.
- (19) Ageel, H. K.; Harrad, S.; Abdallah, M. A. E. Occurrence, Human Exposure, and Risk of Microplastics in the Indoor Environment. *Environ. Sci. Process Impacts* **2022**, *24* (1), 17–31.
- (20) Abdallah, M. A. E.; Harrad, S. Dermal Contact with Furniture Fabrics Is a Significant Pathway of Human Exposure to Brominated Flame Retardants. *Environ. Int.* **2018**, *118*, 26.
- (21) Abou-Elwafa Abdallah, M.; Harrad, S. Dermal Uptake of Chlorinated Organophosphate Flame Retardants via Contact with Furniture Fabrics; Implications for Human Exposure. *Environ. Res.* **2022**, *209*, 112847.
- (22) Stefaniak, A. B.; Harvey, C. J. Dissolution of Materials in Artificial Skin Surface Film Liquids. *Toxicology in Vitro*. **2006**, *20*, 1265.
- (23) Stefaniak, A. B.; Duling, M. G.; Geer, L.; Virji, M. A. Dissolution of the Metal Sensitizers Ni, Be, Cr in Artificial Sweat to Improve Estimates of Dermal Bioaccessibility. *Environmental Sciences: Processes and Impacts* **2014**, *16* (2), 341.
- (24) Abdallah, M. A. E.; Drage, D. S.; Sharkey, M.; Berresheim, H.; Harrad, S. A Rapid Method for the Determination of Brominated Flame Retardant Concentrations in Plastics and Textiles Entering the Waste Stream. *J. Sep. Sci.* **2017**, *40* (19), 3873–3881.
- (25) Liu, C.; Li, J.; Zhang, Y.; Wang, L.; Deng, J.; Gao, Y.; Yu, L.; Zhang, J.; Sun, H. Widespread Distribution of PET and PC Microplastics in Dust in Urban China and Their Estimated Human Exposure. *Environ. Int.* **2019**, *128*, 116–124.
- (26) US EPA. USEPA Exposure Factors Handbook Chapter 7: Dermal Exposure Studies. *Exposure Factors Handbook* 2011; EPA/600/R (September).
- (27) Das, C.; Olmsted, P. D. The Physics of Stratum Corneum Lipid Membranes. *Philos. Trans A Math Phys. Eng. Sci.* **2016**, *374* (2072), 20150126.
- (28) Abdallah, M. A. E.; Pawar, G.; Harrad, S. Evaluation of 3D-Human Skin Equivalents for Assessment of Human Dermal Absorption of Some Brominated Flame Retardants. *Environ. Int.* **2015**, *84*, 64–70.
- (29) Qiao, G. L.; Brooks, J. D.; Riviere, J. E. Pentachlorophenol Dermal Absorption and Disposition from Soil in Swine: Effects of Occlusion and Skin Microorganism Inhibition. *Toxicol. Appl. Pharmacol.* **1997**, *147* (2), 234–246.
- (30) Yu, Y.-X.; Li, J.-L.; Zhang, X.-Y.; Yu, Z.-Q.; Van de Wiele, T.; Han, S.-Y.; Wu, M.-H.; Sheng, G.-Y.; Fu, J.-M. Assessment of the Bioaccessibility of Polybrominated Diphenyl Ethers in Foods and the Correlations of the Bioaccessibility with Nutrient Contents. *J. Agric. Food Chem.* **2010**, *58* (1), 301–308.
- (31) Hornero-Méndez, D.; Mínguez-Mosquera, M. I. Bioaccessibility of Carotenes from Carrots: Effect of Cooking and Addition of Oil. *Innovative Food Science and Emerging Technologies* **2007**, *8* (3), 407.
- (32) Berens, A. R.; Hopfenberg, H. B. Diffusion of Organic Vapors at Low Concentrations in Glassy PVC, Polystyrene, and PMMA. *J. Membr. Sci.* **1982**, *10* (2–3), 283–303.
- (33) Lee, H.; Byun, D. E.; Kim, J. M.; Kwon, J. H. Desorption Modeling of Hydrophobic Organic Chemicals from Plastic Sheets Using Experimentally Determined Diffusion Coefficients in Plastics. *Mar. Pollut. Bull.* **2018**, *126*, 312.
- (34) Sastri, V. R. Commodity Thermoplastics: Polyvinyl Chloride, Polyolefins, Cycloolefins and Polystyrene. *Plastics in Medical Devices* **2022**, 113.
- (35) Fang, M.; Stapleton, H. M. Evaluating the Bioaccessibility of Flame Retardants in House Dust Using an In Vitro Tenax Bead-Assisted Sorptive Physiologically Based Method. *Environ. Sci. Technol.* **2014**, *48* (22), 13323–13330.
- (36) Abou-Elwafa Abdallah, M.; Tilston, E.; Harrad, S.; Collins, C. In Vitro Assessment of the Bioaccessibility of Brominated Flame Retardants in Indoor Dust Using a Colon Extended Model of the Human Gastrointestinal Tract. *Journal of Environmental Monitoring* **2012**, *14* (12), 3276.
- (37) Kademoglou, K.; Williams, A. C.; Collins, C. D. Bioaccessibility of PBDEs Present in Indoor Dust: A Novel Dialysis Membrane Method with a Tenax TA® Absorption Sink. *Sci. Total Environ.* **2018**, *621*, 1.
- (38) Yu, Y. X.; Pang, Y. P.; Li, C.; Li, J. L.; Zhang, X. Y.; Yu, Z. Q.; Feng, J. L.; Wu, M. H.; Sheng, G. Y.; Fu, J. M. Concentrations and Seasonal Variations of Polybrominated Diphenyl Ethers (PBDEs) in in- and out-House Dust and Human Daily Intake via Dust Ingestion

Corrected with Bioaccessibility of PBDEs. *Environ. Int.* **2012**, *42* (1), 124.

(39) Brand, R. M.; McMahon, L.; Jendrzejewski, J. L.; Charron, A. R. Transdermal Absorption of the Herbicide 2,4-Dichlorophenoxyacetic Acid Is Enhanced by Both Ethanol Consumption and Sunscreen Application. *Food Chem. Toxicol.* **2007**, *45* (1), 93–97.

(40) Pont, A. R.; Charron, A. R.; Brand, R. M. Active Ingredients in Sunscreens Act as Topical Penetration Enhancers for the Herbicide 2,4-Dichlorophenoxyacetic Acid. *Toxicol. Appl. Pharmacol.* **2004**, *195* (3), 348–354.

(41) Brand, R. M.; Charron, A. R.; Sandler, V. L.; Jendrzejewski, J. L. Moisturizing Lotions Can Increase Transdermal Absorption of the Herbicide 2,4-Dichlorophenoxyacetic Acid Across Hairless Mouse Skin. *Cutan Ocul Toxicol.* **2007**, *26* (1), 15–23.

(42) Wemken, N.; Drage, D. S.; Abdallah, M. A. E.; Harrad, S.; Coggins, M. A. Concentrations of Brominated Flame Retardants in Indoor Air and Dust from Ireland Reveal Elevated Exposure to Decabromodiphenyl Ethane. *Environ. Sci. Technol.* **2019**, *53* (16), 9826.

(43) Webster, T. F.; Harrad, S.; Millette, J. R.; Holbrook, R. D.; Davis, J. M.; Stapleton, H. M.; Allen, J. G.; McClean, M. D.; Ibarra, C.; Abdallah, M. A.-E.; Covaci, A. Identifying Transfer Mechanisms and Sources of Decabromodiphenyl Ether (BDE 209) in Indoor Environments Using Environmental Forensic Microscopy. *Environ. Sci. Technol.* **2009**, *43* (9), 3067–3072.

(44) Rauert, C.; Harrad, S.; Suzuki, G.; Takigami, H.; Uchida, N.; Takata, K. Test Chamber and Forensic Microscopy Investigation of the Transfer of Brominated Flame Retardants into Indoor Dust via Abrasion of Source Materials. *Sci. Total Environ.* **2014**, *493*, 639.

(45) Schrenk, D.; Bignami, M.; Bodin, L.; Chipman, J. K.; del Mazo, J.; Grasl-Kraupp, B.; Hogstrand, C.; Hoogenboom, L.; Leblanc, J. C.; Nebbia, C. S.; Nielsen, E.; Ntzani, E.; Petersen, A.; Sand, S.; Schwerdtle, T.; Wallace, H.; Benford, D.; Fürst, P.; Rose, M.; Ioannidou, S.; Nikolić, M.; Bordajandi, L. R.; Vleminckx, C. Update of the Risk Assessment of Hexabromocyclododecanes (HBCDDs) in Food. *EFSA Journal* **2021**, *19* (3), e06421.

(46) Roosens, L.; Abdallah, M. A.-E.; Harrad, S.; Neels, H.; Covaci, A. Exposure to Hexabromocyclododecanes (HBCDs) via Dust Ingestion, but Not Diet, Correlates with Concentrations in Human Serum: Preliminary Results. *Environ. Health Perspect.* **2009**, *117* (11), 1707–1712.

(47) Zhang, J.; Wang, L.; Kannan, K. Microplastics in House Dust from 12 Countries and Associated Human Exposure. *Environ. Int.* **2020**, *134*, 105314.

(48) Periyasamy, A. P.; Tehrani-Bagha, A. A Review on Microplastic Emission from Textile Materials and Its Reduction Techniques. *Polym. Degrad. Stab.* **2022**, *199*, 109901.

(49) Bashir, S. M.; Kimiko, S.; Mak, C.-W.; Fang, J. K.-H.; Gonçalves, D. Personal Care and Cosmetic Products as a Potential Source of Environmental Contamination by Microplastics in a Densely Populated Asian City. *Front Mar Sci.* **2021**, *8*, 1.

(50) Salthammer, T. Microplastics and Their Additives in the Indoor Environment. *Angew. Chem., Int. Ed.* **2022**, *61* (32), e202205713.

(51) Jacques, C.; Jamin, E. L.; Perdu, E.; Duplan, H.; Mavon, A.; Zalko, D.; Debrauwer, L. Characterisation of B(a) P Metabolites Formed in an Ex Vivo Pig Skin Model Using Three Complementary Analytical Methods. *Anal Bioanal Chem.* **2010**, *396* (5), 1691–1701.