

Extensive field evidence for the release of HONO from the photolysis of nitrate aerosols

Andersen, Simone T.; Carpenter, Lucy J.; Reed, Chris; Lee, James D.; Chance, Rosie; Sherwen, Tomás; Vaughan, Adam R.; Stewart, Jordan; Edwards, Pete M.; Bloss, William J.; Sommariva, Roberto; Crilley, Leigh R.; Nott, Graeme J.; Neves, Luis; Read, Katie; Heard, Dwayne E.; Seakins, Paul W.; Whalley, Lisa K.; Boustead, Graham A.; Fleming, Lauren T.

DOI:
[10.1126/sciadv.add6266](https://doi.org/10.1126/sciadv.add6266)

License:
Creative Commons: Attribution (CC BY)

Document Version
Publisher's PDF, also known as Version of record

Citation for published version (Harvard):
Andersen, ST, Carpenter, LJ, Reed, C, Lee, JD, Chance, R, Sherwen, T, Vaughan, AR, Stewart, J, Edwards, PM, Bloss, WJ, Sommariva, R, Crilley, LR, Nott, GJ, Neves, L, Read, K, Heard, DE, Seakins, PW, Whalley, LK, Boustead, GA, Fleming, LT, Stone, D & Fomba, KW 2023, 'Extensive field evidence for the release of HONO from the photolysis of nitrate aerosols', *Science Advances*, vol. 9, no. 3, eadd6266.
<https://doi.org/10.1126/sciadv.add6266>

[Link to publication on Research at Birmingham portal](#)

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.



ATMOSPHERIC SCIENCE

Extensive field evidence for the release of HONO from the photolysis of nitrate aerosols

Simone T. Andersen^{1+*}, Lucy J. Carpenter^{1*}, Chris Reed², James D. Lee^{1,3}, Rosie Chance¹, Tomás Sherwen^{1,3}, Adam R. Vaughan¹, Jordan Stewart¹, Pete M. Edwards¹, William J. Bloss⁴, Roberto Sommariva⁴, Leigh R. Crilley^{4‡}, Graeme J. Nott², Luis Neves⁵, Katie Read^{1,3}, Dwayne E. Heard⁶, Paul W. Seakins⁶, Lisa K. Whalley^{2,6}, Graham A. Boustead⁶, Lauren T. Fleming⁶, Daniel Stone⁶, Kanneh Wadinga Fomba⁷


Copyright © 2023 The Authors, some rights reserved; exclusive licensee American Association for the Advancement of Science. No claim to original U.S. Government Works. Distributed under a Creative Commons Attribution License 4.0 (CC BY).

Particulate nitrate (pNO_3^-) has long been considered a permanent sink for NO_x (NO and NO_2), removing a gaseous pollutant that is central to air quality and that influences the global self-cleansing capacity of the atmosphere. Evidence is emerging that photolysis of pNO_3^- can recycle HONO and NO_x back to the gas phase with potentially important implications for tropospheric ozone and OH budgets; however, there are substantial discrepancies in “renoxification” photolysis rate constants. Using aircraft and ground-based HONO observations in the remote Atlantic troposphere, we show evidence for renoxification occurring on mixed marine aerosols with an efficiency that increases with relative humidity and decreases with the concentration of pNO_3^- , thus largely reconciling the very large discrepancies in renoxification photolysis rate constants found across multiple laboratory and field studies. Active release of HONO from aerosol has important implications for atmospheric oxidants such as OH and O_3 in both polluted and clean environments.

INTRODUCTION

Nitrous acid (HONO) has a pivotal role in tropospheric chemistry as an important source of the hydroxyl radical (OH) (1–7). It has also been proposed as a substantial source of NO_x (NO and NO_2) to the remote marine environment (8–10). NO_x regulates the abundance of atmospheric oxidants and is essential for the formation of secondary atmospheric aerosols, and OH controls the self-cleansing capacity of the atmosphere via degradation of pollutants and greenhouse gases such as methane (1).

HONO is produced through the gas-phase reaction of NO and OH radicals (R1) and lost through photolysis (R2), reaction with OH radicals (R3), and dry deposition (R4)



In the remote oceanic troposphere, NO_x levels are too low to supply any notable levels of HONO from R1 and primary HONO emission sources such as vehicle exhaust, wild fires, and soils (2, 3, 11) are absent. Recent measurements also suggest the absence of an ocean surface source (8). Reactions on aerosol surfaces have historically been suggested to make only moderate contributions to daytime HONO formation (5), although there is evidence for NO_2 -to-HONO conversion on aerosols in polluted to semi-polluted regions (4, 6, 7). However, recent field observations in the marine atmosphere have indicated that photolysis of particulate nitrate (pNO_3^-) associated with sea salt aerosol can be an important source of HONO and NO_x (9, 10, 12), consistent with laboratory studies demonstrating that photolysis of pNO_3^- is significantly enhanced compared to photolysis of gaseous HNO_3 , with HONO and NO_x as the major products (13–20)



This “renoxification” process is important because it offers a rapid route for recycling of NO_x from inorganic nitrate, which has historically been thought to be slow because of the small photolysis frequency of gas phase HNO_3 . If renoxification supplies a substantial amount of NO_x to remote oceanic regions, where sources have been considered to be limited primarily to ship emissions and to transport and decomposition of peroxyacetyl nitrate, it could have a global-scale impact on production of tropospheric oxidants such as O_3 and OH and, hence, on methane removal (21).

The photolysis rate constant of renoxification, $j^{\text{pNO}_3^-}$, is typically expressed as a ratio to the gas-phase HNO_3 photolysis frequency, giving an enhancement factor $f = \frac{j^{\text{pNO}_3^-}}{j^{\text{HNO}_3}}$. There is a very high uncertainty in f , with laboratory and field studies reporting values spanning three orders of magnitude (Table 1). Field observations of

¹Wolfson Atmospheric Chemistry Laboratories, Department of Chemistry, University of York, York, UK. ²FAAM Airborne Laboratory, Cranfield, UK. ³National Centre for Atmospheric Science, University of York, York, UK. ⁴School of Geography, Earth and Environmental Sciences, University of Birmingham, Birmingham, UK. ⁵Instituto Nacional de Meteorologia e Geofísica, São Vicente (INMG), Mindelo, Cabo Verde. ⁶School of Chemistry, University of Leeds, Leeds, UK. ⁷Atmospheric Chemistry Department, Leibniz Institute for Tropospheric Research (TROPOS), Leipzig, Germany. [†]Present address: Max Planck Institute for Chemistry, Mainz, Germany. [‡]Present address: Department of Chemistry, York University, Toronto, Ontario, Canada.

*Corresponding author. Email: lucy.carpenter@york.ac.uk (L.J.C.); simone.andersen@york.ac.uk (S.T.A.).

HONO in the remote oceanic atmosphere, which offer a robust method to diagnose the presence of any missing sources but are so far limited to only a few days of measurements, have been reconciled with known sources and sinks using f of between ~25 and 450 (9, 12, 21, 22). This range is within reported values from laboratory studies on various surfaces (14, 18–20, 23) and aerosol filter samples (17) of between ~10 and 1700. However, recent experiments using suspended nitrate particles (23) and calculations derived from observed ratios of NO_x/HNO_3 in the polluted boundary layer (24) have derived a much smaller f of 1 to 30. Thus, there is as yet no consensus on whether renoxification offers a limited or a highly significant role in the NO_x and OH budgets of remote environments or field evidence for HONO production from renoxification occurring on ambient aerosol other than sea salt aerosol.

RESULTS

The ARNA (Atmospheric Reactive Nitrogen over the remote Atlantic) field campaigns took place over the tropical Atlantic Ocean in August 2019 and February 2020 using the FAAM BAe-146-301 atmospheric research aircraft and in August 2019 at the Cape Verde Atmospheric Observatory (CVAO; Fig. 1A). Twelve flights (four in summer and eight in winter; Fig. 1B) were conducted with in situ measurements including NO , NO_2 , HONO, O_3 , and aerosol surface area. pNO_3^- was determined from aerosol filters sampled over each straight-and-level run (SLR). Photolysis rates and OH radical concentrations were modeled using the global three-dimensional atmospheric chemistry model GEOS-Chem. At the CVAO, NO_x , pNO_3^- , O_3 , and photolysis rates are measured routinely, and these were supplemented by HONO measurements during the ARNA campaigns (10).

Figure 1 (C to G) shows the aircraft vertical profiles of NO , NO_2 , HONO, and pNO_3^- and total aerosol surface area. For NO , NO_2 , and pNO_3^- , the vertical profiles show clear enhancements between 1500 and 2500 m. The air sampled in this layer predominately originates from over Africa (figs. S1 and S2) and shows tracers of biomass burning and dust in the aerosol composition (figs. S3 and S4). The mean mixing ratios of HONO (± 1 SD) were 18.2 ± 5.9 parts per trillion by volume (pptv) in the marine boundary layer (MBL) and 14.2 ± 6.4 pptv above the MBL (0.5 to 3.0 km). Figure 2A shows that HONO levels measured at the CVAO were about a factor of 3 lower (4.7 ± 1.8 pptv at solar noon) than the MBL aircraft measurements but similar to previous measurements made at the CVAO (~3.5 pptv at solar noon) (10).

The lifetime of HONO during the ARNA and CVAO campaigns was approximately 12 min; thus, the steady-state HONO concentrations can be estimated from balancing the known in situ production and loss mechanisms described in R1 to R4 (see the Supplementary Materials). For the entire dataset, measured HONO levels were substantially larger than these calculated levels, which are negligible in this very low NO_x environment (Figs. 1F and 2A), demonstrating the presence of an additional source. Figure 2B shows that the missing HONO source required to balance the measured HONO concentrations was strongly correlated with solar irradiance (plotted as $j\text{HNO}_3$) in the CVAO data (r^2 of linear fit = 0.79), consistent with a photochemical mechanism such as reaction R5. For the CVAO campaign, HONO was measured over full diurnal cycles (i.e., a large range of solar irradiance values) in homogeneous air experiencing relatively invariant pNO_3^- concentrations (mean and SD of 19.0 ± 5.6 nmol m^{-3}). The aircraft sorties experienced a much larger range of pNO_3^- concentrations (33.5 ± 19.8 nmol m^{-3}), aerosol compositions, and relative humidities (RHs), and

Table 1. Overview of previous studies investigating the production of HONO from surface-adsorbed HNO_3 /nitrate. Concentrations of pNO_3^- given in other units than mol m^{-3} were converted using temperature at 298 K and pressure at 1 atm. $j\text{pNO}_3^-$ was determined from the production of HONO + NO_2 , and $j\text{pNO}_3^- \rightarrow \text{HONO}$ is determined from the production of HONO. Enhancement factors were estimated using $j\text{HNO}_3 = 7 \times 10^{-7} \text{ s}^{-1}$ if not reported in the study.

Study type	Surface	$[\text{pNO}_3^-]$ ($10^{-9} \text{ mol m}^{-3}$)	$j\text{pNO}_3^-$ (10^{-5} s^{-1})	$j\text{pNO}_3^- \rightarrow \text{HONO}$ (10^{-5} s^{-1})	Enhancement factor (f)	Airmass origin	Reference
Field	Aerosols	0.04–2.0		20	150–450	Marine	(9)
Field/ laboratory	Aerosols	0.7–39.5	0.62–50.0		8–700	Urban, rural, remote	(17)
Field/model	Aerosols	40–265*			1–30		(24)
Global model	Aerosols	0.4–40			25–100	Marine	(21)
Laboratory	Aerosols†	1600–9700			<10		(23)
Field/ laboratory	Aerosols	5–15			18–54‡	Marine	(12)
	Surface	Density of HNO_3 on surface ($10^{-7} \text{ mol m}^{-2}$)	$j\text{pNO}_3^-$ (10^{-5} s^{-1})	$j\text{pNO}_3^- \rightarrow \text{HONO}$ (10^{-5} s^{-1})	Enhancement factor (f)	Airmass origin	Reference
Laboratory	Urban Grime		120		1700	Urban	(14)
Laboratory	Aluminum	4–251	5.7–15.3		8–220		(20)
Laboratory	Oak	3–174	1.6–37.0		20–530		(20)
Laboratory	Maple	22–380	0.9–4.9		10–70		(20)

* pNO_3^- here is the sum of pNO_3^- and HNO_3 (5th to 95th percentiles).

†Pure nitrate salts added to Teflon chambers.

‡Twenty-fifth to 75th percentiles.




Fig. 1. Flight tracks and vertical profiles of $p\text{NO}_3^-$, NO, NO_2 , HONO, and aerosol surface area during ARNA-1 (August 2019) and ARNA-2 (February 2020). (A) Map over the region, where the red box is the area shown in (B). (B) Flight tracks from ARNA-1 in red colors and ARNA-2 in blue colors, alongside the location of the CVAO. The vertical profiles of (C) $p\text{NO}_3^-$, (D) NO, (E) NO_2 , (F) HONO, and (G) aerosol surface area are colored by their respective flight tracks as shown in (B). Each data point is an average of an SLR. The gray vertical line in (F) shows the range of calculated HONO gas-phase source-only concentrations for each SLR during the flights. The error bars represent the 1σ uncertainties described in Supplementary Text.

here, the relationship between the missing HONO source and $j\text{HNO}_3$ was weak. Potential reasons for the lower missing HONO source in the ground-based CVAO measurements are discussed below.

The missing HONO source is plotted against $j\text{HNO}_3 \times [p\text{NO}_3^-]$ in Fig. 3A. If the missing source was entirely due to renoxification ($P_{\text{HONO}_{\text{het}}}$), then it should be equal to the product of $j\text{pNO}_3^-$ and $p\text{NO}_3^-$ and the slope equal to f because $f = \frac{j\text{pNO}_3^-}{j\text{HNO}_3}$. Figure 3A shows, however, that there was no simple linear relationship between these parameters, unlike a previous field study of renoxification occurring on sea salt aerosol in the marine atmosphere (9). These results indicate that more factors were influencing $j\text{pNO}_3^-$ than simply the intensity of solar radiation.

The ARNA flights encompassed a range of RH, from low humidities ($<30\%$ RH), where sea salt particles are expected to be dry and solid, to those where they become deliquesced (>45 to 55% RH) and therefore contain some water on the surface, to high humidities ($>75\%$ RH), where sea salt aerosols will be present as suspended droplets (25). We find significant differences in f across these RH ranges, with average values of 156 ± 91 (1σ), 96 ± 49 , and 39 ± 29 for $>75\%$ RH, 53 to 75% RH, and $<30\%$ RH, respectively ($P < 0.0001$ when comparing the samples where RH $>75\%$ and RH $<30\%$). Figure 3B illustrates the general tendency for f to increase with RH; a simple linear fit has an r^2 of 0.44. Solid-phase nitrate aerosol exhibits a much lower quantum yield of photolysis compared to aqueous phase (26) because the products cannot diffuse out of the particle but instead will recombine. A marked increase in f on particulate sodium nitrate between dry conditions ($\sim 3\%$

RH), with negligible renoxification, and higher RH representing aqueous aerosol have been found in the laboratory (23). Our observations show that such effects are identifiable in the atmosphere.

We also find that f reduces with $[p\text{NO}_3^-]$, as shown in Fig. 3C, which shows both our aircraft and ground-based (CVAO) data. In air masses dominated by sea salt aerosol with generally low $[p\text{NO}_3^-]$, we find an average f calculated from the aircraft data of 157 (range, 54 to 297), similar to the airborne study of Ye *et al.* (9) who derived f_{obs} of 150 to 450 on sea salt aerosol. We observed similar enhancement factors for mixed sea salt/dust aerosol; however, significantly lower f was derived for biomass burning aerosol, which contained high $[p\text{NO}_3^-]$ (f of ~ 10 to 60). Laboratory measurements of HONO production from photolysis of aerosol filter samples collected in urban, suburban/rural, and remote areas (17) reported a notably similar relationship between the enhancement factor and $[p\text{NO}_3^-]$ as our study. The Ye *et al.* (17) fit to their laboratory data is shown in Fig. 3C as the gray dashed line. Their $j\text{pNO}_3^-_{\text{effective}}$ was converted to an enhancement factor using $j\text{HNO}_3 = 7 \times 10^{-7} \text{ s}^{-1}$, which corresponds to typical tropical summer conditions on the ground (solar elevation angle $\theta = 0^\circ$) as simulated in their light-exposure experiments, thus

$$f = \frac{6.1 \times 10^{-4} \times \ln(1 + 4.4 \times 10^{-1} \times [p\text{NO}_3^-]) - 3.5 \times 10^{-5}}{7 \times 10^{-7}} \quad (1)$$

where $[p\text{NO}_3^-]$ is the ambient concentration in units of $10^{-9} \text{ mol m}^{-3}$ (air). This empirical relationship was attributed by Ye *et al.* (17) to a surface catalysis mechanism for renoxification because it is consistent with that observed in the photolysis of surface-




Fig. 2. Evidence for a photochemical origin of the missing HONO source. (A) Average diurnal cycle of HONO mixing ratios measured at the CVAO in August 2019 (red), where the shaded area shows \pm the SE of the measurements, compared to calculated HONO photostationary state (PSS; see Supplementary Text) mixing ratios (gray). The deviation from PSS maximizes around local noon, indicating that the additional source term has a strong photolytic dependence. (B) Missing HONO source at the CVAO (hourly averages, red circles, r^2 of linear fit = 0.79) and for the aircraft data (hourly averages, black unfilled circles, r^2 of linear fit = 0.06) plotted against $j\text{HNO}_3$.

adsorbed HNO_3 /nitrate on various surfaces (20). Our observations span a much wider range of pNO_3^- concentrations than previous field studies (9, 10, 12), allowing the relationship of $[\text{pNO}_3^-]$ with the renoxification rate constant to be explored in the field. Given that the Ye *et al.* (17) experiments were carried out on aerosol collected primarily from urban or urban-influenced locations, it is not unexpected to find significant differences in the absolute values of f between and within these studies. However, Fig. 3C demonstrates a remarkable similarity in the general relationship of f with bulk nitrate.

Recent laboratory experiments and field observations of NO_x/HNO_3 ratios have derived enhancement factors of <30 and have suggested therefore that renoxification plays only a limited role in atmospheric chemistry (blue and gray boxes in Fig. 3D) (23, 24). However, these studies were carried out under very high pNO_3^- mass concentrations, where our observations would suggest only low enhancement factors that have only a small dependence on pNO_3^- across the concentration ranges explored.

We next further explore the relationship between f and $[\text{pNO}_3^-]$. A number of studies have indicated that renoxification is driven by photolysis of surface-bound rather than bulk nitrate. Nitrate or nitric acid adsorbed on surfaces can undergo a much more rapid photolysis compared to bulk aqueous nitrate or gas phase HNO_3 due to enhanced absorption cross sections arising from optimal alignment and orientation of HNO_3 molecules on surfaces (13, 15, 27) and high quantum yields (compared to the aqueous phase) due to reduced solvent cage effects (13). Therefore, if the nitric acid is located substantially on or near the aerosol surface, its photolysis rate could be enhanced by orders of magnitude over that of gas-phase nitric acid. While molecular dynamics simulations offer conflicting results as to whether nitrate anions prefer interfacial to bulk solvation, the presence of certain cations has been shown experimentally to lead to preferential distributions of nitrate ion at the interface (28–30). In addition, several experimental studies show that the products of nitrate photolysis are enhanced by the presence of halide ions, a phenomenon that has been attributed to the known surface affinity of halide ions pulling sodium cations closer, in turn, drawing NO_3^- to the interface (31–35). We illustrate

this surface-enhanced mechanism in Fig. 4. Potential synergisms between HNO_3 and organic films (36, 37) could further enhance the concentration and photochemistry of surface nitrate compared to bulk aerosol nitrate.

If renoxification is controlled by the availability of surface NO_3^- , then we would expect f to be dependent on the partitioning between the equilibrium surface and equilibrium bulk nitrate in liquid (deliquesced) aerosol. If nitrate behaves as a surfactant, as suggested by the theoretical and laboratory studies discussed above, then such partitioning can be described using a Langmuir adsorption isotherm

$$[\text{pNO}_3^-]_{\text{surface}} = \frac{Q^0 K_L [\text{pNO}_3^-]_{\text{bulk}}}{1 + K_L [\text{pNO}_3^-]_{\text{bulk}}} \quad (2)$$

where Q^0 is the maximum loading of adsorbate NO_3^- corresponding to complete monolayer coverage [reaching saturation under conditions of high solute ($[\text{pNO}_3^-]_{\text{bulk}}$) concentration] and K_L is the Langmuir equilibrium constant of NO_3^- .

Because we measured bulk pNO_3^- rather than surface-bound, we can define $P_{\text{HONO}_{\text{net}}}$ as $j\text{pNO}_3^-_{\text{effective}} \times [\text{pNO}_3^-]_{\text{bulk}}$ and the observationally derived enhancement factor, f_{obs} , as $\frac{j\text{pNO}_3^-_{\text{effective}}}{j\text{HNO}_3}$. Thus, it follows that

$$f_{\text{obs}} = \frac{j\text{pNO}_3^-_{\text{surface}}}{j\text{HNO}_3} \times \frac{[\text{pNO}_3^-]_{\text{surface}}}{[\text{pNO}_3^-]_{\text{bulk}}} \quad (3)$$

The dependence of f_{obs} on $[\text{pNO}_3^-]_{\text{bulk}}$ can then be estimated by combining Eqs. 2 and 3

$$f_{\text{obs}} = \frac{f \cdot [\text{pNO}_3^-]_{\text{surface}}}{[\text{pNO}_3^-]_{\text{bulk}}} = \frac{f \cdot Q^0 K_L}{1 + K_L [\text{pNO}_3^-]_{\text{bulk}}} \quad (4)$$

where $f = j\text{pNO}_3^-_{\text{surface}}/j\text{HNO}_3$.

Figure 3D shows how the Langmuir expression (Eq. 4) fits the enhancement factors derived from our field data under conditions where the aerosol was expected to be deliquesced (RH > 53%; colored circle and empty blue triangles), along with the Ye *et al.* (9) marine boundary layer data (orange box), the laboratory data of Ye *et al.* (17) (gray symbols) and of Shi *et al.* (23) (gray bar),




Fig. 3. Relationships of P_{HONOhet} and enhancement factors with various atmospheric parameters. (A) Calculated missing HONO source (P_{HONOhet}) versus the product of $[\text{pNO}_3^-]_{\text{bulk}}$ and the photolysis frequency of gaseous HNO_3 . Note the lack of a simple linear relationship. Data points are colored according to the dominant aerosol classification (Supplementary Materials). $[\text{pNO}_3^-]_{\text{bulk}}$ was converted from nmol m^{-3} to pptv using surface pressure and temperature. The $[\text{pNO}_3^-]_{\text{bulk}}$ used is a lower limit, meaning the derived f are upper limits, due to a potentially lower than 100% sampling efficiency of coarse mode aerosols ($>1 \mu\text{m}$) for aircraft measurements (56–58). (B) Derived enhancement factors as a function of RH. (C) Derived enhancement factors as a function of $[\text{pNO}_3^-]_{\text{bulk}}$. Also shown are the laboratory data of Ye *et al.* (17) using aerosol filters (gray triangles) and their fit to the data (gray dashed line) and the aircraft data of Ye *et al.* (9) from the remote North Atlantic marine boundary layer (orange rectangle). (D) Enhancement factors as a function of a wide range of $[\text{pNO}_3^-]_{\text{bulk}}$. Same as (C) but a wider comparison of f_{obs} with previously reported values and data restricted to conditions where the aerosol was expected to be deliquesced. The dotted line shows the Langmuir fit (Eq. 4) applied to all data.

and those derived from ambient measurements of the NO_x/HNO_3 ratio (blue bar) (24).

The best fit for Eq. 4 was found, using the mean f of 70 and $[\text{pNO}_3^-]_{\text{bulk}}$ concentrations in units of $10^{-9} \text{ mol m}^{-3}$ air (as determined from filter measurements), with $Q^0 = 29 \text{ nmol m}^{-3}$ and $K_L = 0.19 \text{ nmol}^{-1} \text{ m}^3$. Calculations using the aerosol thermodynamics module ISORROPIA II (38) using the ARNA aerosol composition, temperature, and humidity data (see the Supplementary Materials) show that for $\text{RH} > 53\%$, essentially all ($>99\%$) of the nitrate (gas phase HNO_3 and pNO_3^-) was in the particle phase. For equivalently humid conditions, the solute ($[\text{pNO}_3^-]_{\text{bulk}}$) concentrations are thus proportional to the $[\text{pNO}_3^-]$ concentrations in air, and we use the latter for ease of comparison across studies. Note that Eq. 4 only considers production of HONO and ignores any coproduction of NO_x . This is important to note when comparing our enhancement factors to laboratory studies, which have measured the production of all gaseous oxidized nitrogen products although is of limited consequence if the yield of HONO is >0.9 , as suggested from a budget analysis of field measurements (22).

Although there is significant variance in the data around the simple Langmuir fit of f , likely due to differences in humidity and aerosol composition as discussed above, the reasonable fit ($r^2 = 0.44$) demonstrates a potential explanation for the observed dependence of the renoxification enhancement factor on $[\text{pNO}_3^-]_{\text{bulk}}$ across multiple studies. Alternative mechanisms for these observed relationships include competing decomposition pathways of NO_2^- leading to greater conversion to non-HONO products at high nitrate concentrations (39) and a reduction in the quantum yield for NO_2^- formation at high nitrate concentrations (40). Noting that halide anions and sodium cations have been suggested to lead to enhancement of the surface concentration and photochemistry of surface nitrate compared to bulk aerosol nitrate, we also plot f against $[\text{pCl}^-]/[\text{pNO}_3^-]$ and $[\text{pNa}^+]/[\text{pNO}_3^-]$ in fig. S5. While the enhancement factor generally increases with these ratios, this could simply be a reflection of the relationship of f with RH (Fig. 3B) because RH was higher in air masses containing elevated sea salt. Ultimately, our field observations are not able to demonstrate what factors besides RH and $[\text{pNO}_3^-]$ control the renoxification efficiency but suggest that additional variables are affecting f to an important degree.

DISCUSSION

The observations shown here represent the most extensive field measurements of HONO in the marine atmosphere to date and confirm its widespread presence in the marine lower troposphere at mixing ratios in the range of ~ 5 to 20 pptv. The fact that HONO production occurs in association with a wide range of aerosol types shows that renoxification is not limited to sea salt aerosol although is strongly restricted under dry conditions (such as those in the free troposphere), where aerosols are not deliquesced.

Nevertheless, the relationships found between the enhancement factor f and aerosol composition, alongside the fact that the derived pNO_3^- photolysis rates are orders of magnitude higher than gas phase nitric acid photolysis rates, are consistent with a surface aerosol mechanism for renoxification. The observed reduction in the renoxification efficiency with increasing concentration of pNO_3^- helps reconcile the very large discrepancies in renoxification photolysis rate constants found across multiple laboratory and field studies. The observations emphasize the importance in laboratory renoxification studies of generating aerosol composition as representative as possible of the clean marine troposphere.

The relationships observed among f , humidity, and $[\text{pNO}_3^-]$ were undoubtedly affected by additional sources of variability. In sunlight, NO_3^- photolysis occurs predominantly via two channels: one producing the nitrite anion (NO_2^-) and $\text{O}(^3\text{P})$ and the other producing NO_2 and OH (39). Gaseous HONO production from the former pathway requires acidity and has been found to be strongly dependent on aerosol pH (16). Here, we note that the ground-based CVAO HONO measurements were associated with lower enhancements (shown in Fig. 3C) compared to the aircraft observations. One potential reason for this is that the surface measurements experience fresh rather than aged sea salt aerosol due to sampling within the surf zone. They would therefore likely be associated with higher aerosol pH, which disfavors HONO production. It has also been shown that the renoxification rate constant can be enhanced in the presence of organic matter, through various




Fig. 4. Potential surface-mediated mechanism for renoxification on nitrate aerosol. Nitrate photolysis can be enhanced by the presence of halide ions, which have a well-established affinity for the surface. This has been attributed to the enhanced partitioning of nitrate ions at the interface because of the existence of a double layer of interfacial halide ions and subsurface counterion (31–35).

mechanisms (16, 24, 36, 41–45). It is expected therefore to find a variation of f beyond that described by our simple relationships, although investigation of these additional factors is outside the scope of our study.

Because it has been previously determined that f of 25 to 50 occurring only on sea salt aerosol results in peak enhancements of 20 to 60% for OH, 10 to 30% for ozone, and up to a factor of 20 for NO_x concentrations in the tropical and subtropical marine boundary layer (21), it is clear that the rapid production of HONO observed in this study occurring on diverse aerosol types, showing a mean f of 70 across all samples, will have a substantial impact on our understanding of atmospheric oxidant cycling. We note additionally that nitrate aerosols have become increasingly more important in the atmosphere because of an increase in precursor ammonia emissions and a decline of ammonium sulfate aerosols (46). Thus, recycling of nitric acid to nitrogen oxides on nitrate aerosol could have important, increasing, and as yet unexplored implications for the trends and distributions of atmospheric oxidants.

MATERIALS AND METHODS

At the CVAO, HONO was measured using a long-path absorption photometer (LOPAP-03, QUMA GmbH) (47). The calibration and standard operating procedures are described by Kleffman and Wiesen (48). In 2019 (data shown in Fig 2), the instrument was deployed on top of a 7.5-m tower, and the detection limit was 1.1 pptv (2σ , 30 s). Because of the sampling time for the aerosol composition being 24 hours, the CVAO HONO data used in Fig. 3 were averaged over 1000 to 1500 local time to illustrate their relationship with pNO_3^- . Aerosol samples collected at the CVAO were analyzed for major ions using a standard ion chromatography technique as described by Fomba *et al.* (49). Filters were changed every 24 hours, and the composition was assumed to be uniform across the sampling period. NO_x has been measured continually at the CVAO since 2006 using a chemiluminescence instrument (Air Quality Design Inc.) (50, 51).

On the FAAM aircraft, NO_x ($\text{NO} + \text{NO}_2$) and HONO were measured using differential photolysis (52), where NO_2 and HONO are photolytically converted into NO, followed by NO chemiluminescence detection using a dual-channel instrument equipped with two custom-built photolytic converters. The HONO conversion efficiencies were calibrated against an ultraviolet-visible cavity enhanced absorption spectroscopy using HIRAC [The Highly Instrumented Reactor for Atmospheric Chemistry (53)] similarly to Reed *et al.* (52) (see the Supplementary Materials). The average HONO detection limit over the SLR was 4.2 pptv (2σ). Aerosol chemical composition was determined by off-line analysis of filter samples. Two identical inlets are mounted on the port side of the aircraft allowing collection of duplicate samples. During the ARNA campaigns, sampled aerosol were divided into two size fractions according to aerodynamic particle diameter (approximately corresponding to >1 and <1 μm). Filters were stored frozen (-20°C) until extraction. Anions (Cl^- , NO_3^- , NO_2^- , SO_4^{2-} , and $\text{C}_2\text{O}_4^{2-}$) and cations (Na^+ , K^+ , NH_4^+ , Ca^{2+} , and Mg^{2+}) were determined in the aqueous extracts using ion chromatography (Thermo Fisher Scientific, Dionex-1100; see the Supplementary Materials).

Photolysis rates and OH concentrations were extracted for all ground and airborne observations at the nearest point in space and time from the GEOS-Chem model (v12.9.0, DOI:10.5281/zenodo.3950327). The model was run at a nested horizontal resolution of $0.25^\circ \times 0.3125^\circ$ over the region (-32.0° to 15.0°E , 0.0° to 34.0°N), with boundary conditions provided by a separate global model run spun up for 1 year. The photolysis rates are calculated online in quadrature using Fast-JX code (54, 55). Comparison with available campaign observations is described in the Supplementary Materials.

Supplementary Materials

This PDF file includes:
Supplementary Text
Figs. S1 to S9

Table S1

References

[View/request a protocol for this paper from Bio-protocol.](#)

REFERENCES AND NOTES

- P. S. Monks, C. Granier, S. Fuzzi, A. Stohl, M. L. Williams, H. Akimoto, M. Amann, A. Baklanov, U. Baltensperger, I. Bey, N. Blake, R. S. Blake, K. Carslaw, O. R. Cooper, F. Dentener, D. Fowler, E. Fragkou, G. J. Frost, S. Generoso, P. Ginoux, V. Grewe, A. Guenther, H. C. Hansson, S. Henne, J. Hjorth, A. Hofzumahaus, H. Huntrieser, I. S. A. Isaksen, M. E. Jenkin, J. Kaiser, M. Kanakidou, Z. Klimont, M. Kulmala, P. Laj, M. G. Lawrence, J. D. Lee, C. Lioussse, M. Maione, G. McFiggans, A. Metzger, A. Mieville, N. Moussiopoulos, J. J. Orlando, C. D. O'Dowd, P. I. Palmer, D. D. Parrish, A. Petzold, U. Platt, U. Pöschl, A. S. H. Prévôt, C. E. Reeves, S. Reimann, Y. Rudich, K. Sellegri, R. Steinbrecher, D. Simpson, H. ten Brink, J. Theklo, G. R. van der Werf, R. Vautard, V. Vestreng, C. Vlachokostas, R. von Glasow, Atmospheric composition change—Global and regional air quality. *Atmos. Environ.* **43**, 5268–5350 (2009).
- N. Theys, R. Volkamer, J. F. Müller, K. J. Zarzana, N. Kille, L. Clarisse, I. De Smedt, C. Lerot, H. Finkenzeller, F. Hendrick, T. K. Koenig, C. F. Lee, C. Knote, H. Yu, M. Van Roozendael, Global nitrous acid emissions and levels of regional oxidants enhanced by wildfires. *Nat. Geosci.* **13**, 681–686 (2020).
- H. Su, Y. Cheng, R. Oswald, T. Behrendt, I. Trebs, F. X. Meixner, M. O. Andreae, P. Cheng, Y. Zhang, U. Pöschl, Soil nitrite as a source of atmospheric HONO and OH radicals. *Science* **333**, 1616–1618 (2011).
- X. Lu, Y. Wang, J. Li, L. Shen, J. C. H. Fung, Evidence of heterogeneous HONO formation from aerosols and the regional photochemical impact of this HONO source. *Environ. Res. Lett.* **13**, 114002 (2018).
- X. Li, F. Rohrer, A. Hofzumahaus, T. Brauers, R. Häseler, B. Bohn, S. Broch, H. Fuchs, S. Gomm, F. Holland, J. Jäger, J. Kaiser, F. N. Keutsch, I. Lohse, K. Lu, R. Tillmann, R. Wegener, G. M. Wolfe, T. F. Mentel, A. Kiendler-Scharr, A. Wahner, Missing gas-phase source of HONO inferred from Zeppelin measurements in the troposphere. *Science* **344**, 292–296 (2014).
- N. Zhang, X. Zhou, P. B. Shepson, H. Gao, M. Alaghmand, B. Stirn, Aircraft measurement of HONO vertical profiles over a forested region. *Geophys. Res. Lett.* **36**, (2009).
- Z. Liu, Y. Wang, F. Costabile, A. Amoroso, C. Zhao, L. G. Huey, R. Stickel, J. Liao, T. Zhu, Evidence of aerosols as a media for rapid daytime HONO production over China. *Environ. Sci. Technol.* **48**, 14386–14391 (2014).
- L. R. Crilley, L. J. Kramer, F. D. Pope, C. Reed, J. D. Lee, L. J. Carpenter, L. D. J. Hollis, S. M. Ball, W. J. Bloss, Is the ocean surface a source of nitrous acid (HONO) in the marine boundary layer? *Atmos. Chem. Phys.* **21**, 18213–18225 (2021).
- C. Ye, X. Zhou, D. Pu, J. Stutz, J. Festa, M. Spolario, C. Tsai, C. Cantrell, R. L. Mauldin, T. Campos, A. Weinheimer, R. S. Hornbrook, E. C. Apel, A. Guenther, L. Kaser, B. Yuan, T. Karl, J. Haggerty, S. Hall, K. Ullmann, J. N. Smith, J. Ortega, C. Knote, Rapid cycling of reactive nitrogen in the marine boundary layer. *Nature* **532**, 489–491 (2016).
- C. Reed, M. J. Evans, L. R. Crilley, W. J. Bloss, T. Sherwen, K. A. Read, J. D. Lee, L. J. Carpenter, Evidence for renoxification in the tropical marine boundary layer. *Atmos. Chem. Phys.* **17**, 4081–4092 (2017).
- J. N. Pitts, H. W. Biermann, A. M. Winer, E. C. Tuazon, Spectroscopic identification and measurement of gaseous nitrous acid in dilute auto exhaust. *Atmos. Environ.* **18**, 847–854 (1984).
- Y. Zhu, Y. Wang, X. Zhou, Y. F. Elshorbany, C. Ye, M. Hayden, A. J. Peters, An investigation into the chemistry of HONO in the marine boundary layer at Tudor Hill Marine Atmospheric Observatory in Bermuda. *Atmos. Chem. Phys.* **22**, 6327–6346 (2022).
- C. Zhu, B. Xiang, L. T. Chu, L. Zhu, 308 nm photolysis of nitric acid in the gas phase, on aluminum surfaces, and on ice films. *J. Phys. Chem. A* **114**, 2561–2568 (2010).
- A. M. Baergen, D. J. Donaldson, Photochemical renoxification of nitric acid on real urban grime. *Environ. Sci. Technol.* **47**, 815–820 (2013).
- J. Du, L. Zhu, Quantification of the absorption cross sections of surface-adsorbed nitric acid in the 335–365 nm region by Brewster angle cavity ring-down spectroscopy. *Chem. Phys. Lett.* **511**, 213–218 (2011).
- N. K. Scharko, A. E. Berke, J. D. Raff, Release of nitrous acid and nitrogen dioxide from nitrate photolysis in acidic aqueous solutions. *Environ. Sci. Technol.* **48**, 11991–12001 (2014).
- C. Ye, N. Zhang, H. Gao, X. Zhou, Photolysis of particulate nitrate as a source of HONO and NO_x. *Environ. Sci. Technol.* **51**, 6849–6856 (2017).
- C. Ye, N. Zhang, H. Gao, X. Zhou, Matrix effect on surface-catalyzed photolysis of nitric acid. *Sci. Rep.* **9**, 4351 (2019).
- X. Zhou, H. Gao, Y. He, G. Huang, S. B. Bertman, K. Civerolo, J. Schwab, Nitric acid photolysis on surfaces in low-NO_x environments: Significant atmospheric implications. *Geophys. Res. Lett.* **30**, (2003).
- C. Ye, H. Gao, N. Zhang, X. Zhou, Photolysis of nitric acid and nitrate on natural and artificial surfaces. *Environ. Sci. Technol.* **50**, 3530–3536 (2016).
- P. Kasibhatla, T. Sherwen, M. J. Evans, L. J. Carpenter, C. Reed, B. Alexander, Q. Chen, M. P. Sulprizio, J. D. Lee, K. A. Read, W. Bloss, L. R. Crilley, W. C. Keene, A. A. P. Szpenny, A. Hodzic, Global impact of nitrate photolysis in sea-salt aerosol on NO_x, OH, and O₃ in the marine boundary layer. *Atmos. Chem. Phys.* **18**, 11185–11203 (2018).
- C. Ye, D. E. Heard, L. K. Whalley, Evaluation of novel routes for NO_x formation in remote regions. *Environ. Sci. Technol.* **51**, 7442–7449 (2017).
- Q. Shi, Y. Tao, J. E. Krechmer, C. L. Heald, J. G. Murphy, J. H. Kroll, Q. Ye, Laboratory investigation of renoxification from the photolysis of inorganic particulate nitrate. *Environ. Sci. Technol.* **55**, 854–861 (2021).
- P. S. Romer, P. J. Wooldridge, J. D. Crouse, M. J. Kim, P. O. Wennberg, J. E. Dibb, E. Scheuer, D. R. Blake, S. Meinardi, A. L. Brosius, A. B. Thames, D. O. Miller, W. H. Brune, S. R. Hall, T. B. Ryerson, R. C. Cohen, Constraints on aerosol nitrate photolysis as a potential source of HONO and NO_x. *Environ. Sci. Technol.* **52**, 13738–13746 (2018).
- J. Zeng, G. Zhang, S. Long, K. Liu, L. Cao, L. Bao, Y. Li, Sea salt deliquescence and crystallization in atmosphere: An in situ investigation using x-ray phase contrast imaging. *Surf. Interface Anal.* **45**, 930–936 (2013).
- S. A. Asher, D. D. Tuschel, T. A. Vargson, L. Wang, S. J. Geib, Solid state and solution nitrate photochemistry: Photochemical evolution of the solid state lattice. *Chem. A Eur. J.* **115**, 4279–4287 (2011).
- C. Zhu, B. Xiang, L. Zhu, R. Cole, Determination of absorption cross sections of surface-adsorbed HNO₃ in the 290–330 nm region by Brewster angle cavity ring-down spectroscopy. *Chem. Phys. Lett.* **458**, 373–377 (2008).
- J. Cheng, C. D. Vecitis, M. R. Hoffmann, A. J. Colussi, Experimental anion affinities for the air/water interface. *J. Phys. Chem. B* **110**, 25598–25602 (2006).
- M. Xu, R. Spinney, H. C. Allen, Water structure at the air–aqueous interface of divalent cation and nitrate solutions. *J. Phys. Chem. B* **113**, 4102–4110 (2009).
- W. Hua, D. Verreault, H. C. Allen, Surface electric fields of aqueous solutions of NH₄NO₃, Mg(NO₃)₂, NaNO₃, and LiNO₃: Implications for atmospheric aerosol chemistry. *J. Phys. Chem. C* **118**, 24941–24949 (2014).
- L. M. Wingen, A. C. Moskun, S. N. Johnson, J. L. Thomas, M. Roeselová, D. J. Tobias, M. T. Kleinman, B. J. Finlayson-Pitts, Enhanced surface photochemistry in chloride–nitrate ion mixtures. *Phys. Chem. Chem. Phys.* **10**, 5668–5677 (2008).
- N. K. Richards, B. J. Finlayson-Pitts, Production of gas phase NO₂ and halogens from the photochemical oxidation of aqueous mixtures of sea salt and nitrate ions at room temperature. *Environ. Sci. Technol.* **46**, 10447–10454 (2012).
- N. K. Richards, L. M. Wingen, K. M. Callahan, N. Nishino, M. T. Kleinman, D. J. Tobias, B. J. Finlayson-Pitts, Nitrate ion photolysis in thin water films in the presence of bromide ions. *J. Phys. Chem. A* **115**, 5810–5821 (2011).
- R. Zhang, M. Gen, D. Huang, Y. Li, C. K. Chan, Enhanced sulfate production by nitrate photolysis in the presence of halide ions in atmospheric particles. *Environ. Sci. Technol.* **54**, 3831–3839 (2020).
- A. C. Hong, S. N. Wren, D. J. Donaldson, Enhanced surface partitioning of nitrate anion in aqueous bromide solutions. *J. Phys. Chem. Lett.* **4**, 2994–2998 (2013).
- S. R. Handley, D. Clifford, D. J. Donaldson, Photochemical loss of nitric acid on organic films: A possible recycling mechanism for NO_x. *Environ. Sci. Technol.* **41**, 3898–3903 (2007).
- N. Nishino, S. A. Hollingsworth, A. C. Stern, M. Roeselová, D. J. Tobias, B. J. Finlayson-Pitts, Interactions of gaseous HNO₃ and water with individual and mixed alkyl self-assembled monolayers at room temperature. *Phys. Chem. Chem. Phys.* **16**, 2358–2367 (2014).
- C. Fountoukis, A. Nenes, ISORROPIA II: A computationally efficient thermodynamic equilibrium model for K⁺, Ca²⁺, Mg²⁺, NH₄⁺, Na⁺, SO₄²⁻, NO₃⁻, Cl⁻, H₂O aerosols. *Atmos. Chem. Phys.* **7**, 4639–4659 (2007).
- M. Gen, Z. Liang, R. Zhang, B. R. Go Mabato, C. K. Chan, Particulate nitrate photolysis in the atmosphere. *Environ. Sci. Atmos.* **2**, 111–127 (2022).
- M. Roca, J. Zahardis, J. Bone, M. El-Maazawi, V. H. Grassian, 310 nm irradiation of atmospherically relevant concentrated aqueous nitrate solutions: Nitrite production and quantum yields. *J. Phys. Chem. A* **112**, 13275–13281 (2008).
- M. Ndour, P. Conchon, B. D'Anna, O. Ka, C. George, Photochemistry of mineral dust surface as a potential atmospheric renoxification process. *Geophys. Res. Lett.* **36**, (2009).
- D. I. Reeser, N.-O. A. Kwamena, D. J. Donaldson, Effect of organic coatings on gas-phase nitrogen dioxide production from aqueous nitrate photolysis. *J. Phys. Chem. C* **117**, 22260–22267 (2013).
- J. E. Dyson, G. A. Boustead, L. T. Fleming, M. Blitz, D. Stone, S. R. Arnold, L. K. Whalley, D. E. Heard, Production of HONO from NO₂ uptake on illuminated TiO₂ aerosol particles

- and following the illumination of mixed TiO₂ ammonium nitrate particles. *Atmos. Chem. Phys.* **21**, 5755–5775 (2021).
44. S. L. Mora Garcia, S. Pandit, J. G. Navea, V. H. Grassian, Nitrous acid (HONO) formation from the irradiation of aqueous nitrate solutions in the presence of marine chromophoric dissolved organic matter: Comparison to other Organic photosensitizers. *ACS Earth Space Chem.* **5**, 3056–3064 (2021).
 45. X. Wang, E. Z. Dalton, Z. C. Payne, S. Perrier, M. Riva, J. D. Raff, C. George, Superoxide and nitrous acid production from nitrate photolysis is enhanced by dissolved aliphatic organic matter. *Environ. Sci. Technol. Lett.* **8**, 53–58 (2021).
 46. S. E. Bauer, D. Koch, N. Unger, S. M. Metzger, D. T. Shindell, D. G. Streets, Nitrate aerosols today and in 2030: A global simulation including aerosols and tropospheric ozone. *Atmos. Chem. Phys.* **7**, 5043–5059 (2007).
 47. J. Heland, J. Kleffmann, R. Kurtenbach, P. Wiesen, A new instrument to measure gaseous nitrous acid (HONO) in the atmosphere. *Environ. Sci. Technol.* **35**, 3207–3212 (2001).
 48. J. Kleffmann, P. Wiesen, Technical note: Quantification of interferences of wet chemical HONO LOPAP measurements under simulated polar conditions. *Atmos. Chem. Phys.* **8**, 6813–6822 (2008).
 49. K. W. Fomba, K. Müller, D. van Pinxteren, L. Poulain, M. van Pinxteren, H. Herrmann, Long-term chemical characterization of tropical and marine aerosols at the Cape Verde Atmospheric Observatory (CVAO) from 2007 to 2011. *Atmos. Chem. Phys.* **14**, 8883–8904 (2014).
 50. S. T. Andersen, L. J. Carpenter, B. S. Nelson, L. Neves, K. A. Read, C. Reed, M. Ward, M. J. Rowlinson, J. D. Lee, Long-term NO_x measurements in the remote marine tropical troposphere. *Atmos. Meas. Tech.* **14**, 3071–3085 (2021).
 51. J. D. Lee, S. J. Moller, K. A. Read, A. C. Lewis, L. Mendes, L. J. Carpenter, Year-round measurements of nitrogen oxides and ozone in the tropical North Atlantic marine boundary layer. *J. Geophys. Res. Atmos.* **114**, (2009).
 52. C. Reed, C. A. Brumby, L. R. Crilley, L. J. Kramer, W. J. Bloss, P. W. Seakins, J. D. Lee, L. J. Carpenter, HONO measurement by differential photolysis. *Atmos. Meas. Tech.* **9**, 2483–2495 (2016).
 53. D. R. Glowacki, A. Goddard, K. Hemavibool, T. L. Malkin, R. Commane, F. Anderson, W. J. Bloss, D. E. Heard, T. Ingham, M. J. Pilling, P. W. Seakins, Design of and initial results from a highly instrumented reactor for atmospheric chemistry (HIRAC). *Atmos. Chem. Phys.* **7**, 5371–5390 (2007).
 54. J. Mao, D. J. Jacob, M. J. Evans, J. R. Olson, X. Ren, W. H. Brune, J. M. S. Clair, J. D. Crouse, K. M. Spencer, M. R. Beaver, P. O. Wennberg, M. J. Cubison, J. L. Jimenez, A. Fried, P. Weibring, J. G. Walega, S. R. Hall, A. J. Weinheimer, R. C. Cohen, G. Chen, J. H. Crawford, C. McNaughton, A. D. Clarke, L. Jaeglé, J. A. Fisher, R. M. Yantosca, P. Le Sager, C. Carouge, Chemistry of hydrogen oxide radicals (HO_x) in the Arctic troposphere in spring. *Atmos. Chem. Phys.* **10**, 5823–5838 (2010).
 55. H. Bian, M. J. Prather, Fast-J2: Accurate simulation of stratospheric photolysis in global chemical models. *J. Atmos. Chem.* **41**, 281–296 (2002).
 56. P. Formenti, W. Elbert, W. Maenhaut, J. Haywood, M. O. Andreae, Chemical composition of mineral dust aerosol during the Saharan Dust Experiment (SHADE) airborne campaign in the Cape Verde region, September 2000. *J. Geophys. Res. Atmos.* **108**, 8576 (2003).
 57. M. O. Andreae, W. Elbert, R. Gabriel, D. W. Johnson, S. Osborne, R. Wood, Soluble ion chemistry of the atmospheric aerosol and SO₂ concentrations over the eastern North Atlantic during ACE-2. *Tellus B* **52**, 1066–1087 (2000).
 58. A. Sanchez-Marroquin, D. H. P. Hedges, M. Hiscock, S. T. Parker, P. D. Rosenberg, J. Trembath, R. Walshaw, I. T. Burke, J. B. McQuaid, B. J. Murray, Characterisation of the filter inlet system on the FAAM BAE-146 research aircraft and its use for size-resolved aerosol composition measurements. *Atmos. Meas. Tech.* **12**, 5741–5763 (2019).
 59. L. J. Carpenter, Z. L. Fleming, K. A. Read, J. D. Lee, S. J. Moller, J. R. Hopkins, R. M. Purvis, A. C. Lewis, K. Müller, B. Heinold, H. Herrmann, K. W. Fomba, D. van Pinxteren, C. Müller, I. Tegen, A. Wiedensohler, T. Müller, N. Niedermeier, E. P. Achterberg, M. D. Patey, E. A. Kozlova, M. Heimann, D. E. Heard, J. M. C. Plane, A. Mahajan, H. Oetjen, T. Ingham, D. Stone, L. K. Whalley, M. J. Evans, M. J. Pilling, R. J. Leigh, P. S. Monks, A. Karunaharan, S. Vaughan, S. R. Arnold, J. Tschritter, D. Pöhler, U. Frieß, R. Holla, L. M. Mendes, H. Lopez, B. Faria, A. J. Manning, D. W. R. Wallace, Seasonal characteristics of tropical marine boundary layer air measured at the Cape Verde Atmospheric Observatory. *J. Atmos. Chem.* **67**, 87–140 (2010).
 60. K. A. Read, A. S. Mahajan, L. J. Carpenter, M. J. Evans, B. V. E. Faria, D. E. Heard, J. R. Hopkins, J. D. Lee, S. J. Moller, A. C. Lewis, L. Mendes, J. B. McQuaid, H. Oetjen, A. Saiz-Lopez, M. J. Pilling, J. M. C. Plane, Extensive halogen-mediated ozone destruction over the tropical Atlantic Ocean. *Nature* **453**, 1232–1235 (2008).
 61. M. J. A. Rijkenberg, C. F. Powell, M. Dall'Osto, M. C. Nielsdottir, M. D. Patey, P. G. Hill, A. R. Baker, T. D. Jickells, R. M. Harrison, E. P. Achterberg, Changes in iron speciation following a Saharan dust event in the tropical North Atlantic Ocean. *Mar. Chem.* **110**, 56–67 (2008).
 62. I. Chiappello, G. Bergametti, L. Gomes, B. Chatenet, F. Dulac, J. Pimenta, E. S. Soares, An additional low layer transport of Sahelian and Saharan dust over the north-eastern Tropical Atlantic. *Geophys. Res. Lett.* **22**, 3191–3194 (1995).
 63. C. A. Keller, K. E. Knowland, B. N. Duncan, J. Liu, D. C. Anderson, S. Das, R. A. Lucchesi, E. W. Lundgren, J. M. Nicely, E. Nielsen, L. E. Ott, E. Saunders, S. A. Strode, P. A. Wales, D. J. Jacob, S. Pawson, Description of the NASA GEOS Composition Forecast Modeling System GEOS-CF v1.0. *J. Adv. Model. Earth Syst.* **13**, e2020MS002413 (2021).
 64. I. B. Pollack, B. M. Lerner, T. B. Ryerson, Evaluation of ultraviolet light-emitting diodes for detection of atmospheric NO₂ by photolysis-chemiluminescence. *J. Atmos. Chem.* **65**, 111–125 (2010).
 65. C. Reed, M. J. Evans, P. D. Carlo, J. D. Lee, L. J. Carpenter, Interferences in photolytic NO₂ measurements: Explanation for an apparent missing oxidant? *Atmos. Chem. Phys.* **16**, 4707–4724 (2016).
 66. G. A. Boustead, "Measurement of nitrous acid production from aerosol surfaces using photo-fragmentation laser induced fluorescence," thesis, University of Leeds, (2019).
 67. D. Silvia, J. Skilling, *Data Analysis: A Bayesian Tutorial* (Oxford Univ. Press, ed. 2, 2006).
 68. R. Chance, T. D. Jickells, A. R. Baker, Atmospheric trace metal concentrations, solubility and deposition fluxes in remote marine air over the south-east Atlantic. *Mar. Chem.* **177**, 45–56 (2015).
 69. M. O. Andreae, Soot carbon and excess fine potassium: Long-range transport of combustion-derived aerosols. *Science* **220**, 1148–1151 (1983).
 70. A. R. Baker, T. D. Jickells, K. F. Biswas, K. Weston, M. French, Nutrients in atmospheric aerosol particles along the Atlantic Meridional Transect. *Deep-Sea Res. II Top. Stud. Oceanogr.* **53**, 1706–1719 (2006).
 71. W. Stumm, J. J. Morgan, in *Aquatic Chemistry* (John Wiley, 1996), pp. 1022.
 72. H. C. Price, K. J. Baustian, J. B. McQuaid, A. Blyth, K. N. Bower, T. Choularton, R. J. Cotton, Z. Cui, P. R. Field, M. Gallagher, R. Hawker, A. Merrington, A. Miltenberger, R. R. Neely III, S. T. Parker, P. D. Rosenberg, J. W. Taylor, J. Trembath, J. Vergara-Temprado, T. F. Whale, T. W. Wilson, G. Young, B. J. Murray, Atmospheric ice-nucleating particles in the dusty tropical Atlantic. *J. Geophys. Res. Atmos.* **123**, 2175–2193 (2018).
 73. P. D. Rosenberg, A. R. Dean, P. I. Williams, J. R. Dorsey, A. Minikin, M. A. Pickering, A. Petzold, Particle sizing calibration with refractive index correction for light scattering optical particle counters and impacts upon PCASP and CDP data collected during the Fennec campaign. *Atmos. Meas. Tech.* **5**, 1147–1163 (2012).
 74. OPAC (Optical Properties of Aerosols and Clouds) aerosol database, <https://geisa.aeris-data.fr/opac/> [accessed 4 June 2021].
 75. M. Hess, P. Koepke, I. Schult, Optical properties of aerosols and clouds: The software package OPAC. *Bull. Am. Meteorol. Soc.* **79**, 831–844 (1998).
 76. C. L. Ryder, F. Marengo, J. K. Brooke, V. Estelles, R. Cotton, P. Formenti, J. B. McQuaid, H. C. Price, D. Liu, P. Ausset, P. D. Rosenberg, J. W. Taylor, T. Choularton, K. Bower, H. Coe, M. Gallagher, J. Crosier, G. Lloyd, E. J. Highwood, B. J. Murray, Coarse-mode mineral dust size distributions, composition and optical properties from AER-D aircraft measurements over the tropical eastern Atlantic. *Atmos. Chem. Phys.* **18**, 17225–17257 (2018).
 77. B. Weinzierl, D. Sauer, M. Esselborn, A. Petzold, A. Veira, M. Rose, S. Mund, M. Wirth, A. Ansmann, M. Tesche, S. Gross, V. Freudenthaler, Microphysical and optical properties of dust and tropical biomass burning aerosol layers in the Cape Verde region—An overview of the airborne in situ and lidar measurements during SAMUM-2. *Tellus B* **63**, 589–618 (2011).
 78. C. L. Ryder, E. J. Highwood, A. Walser, P. Seibert, A. Philipp, B. Weinzierl, Coarse and giant particles are ubiquitous in Saharan dust export regions and are radiatively significant over the Sahara. *Atmos. Chem. Phys.* **19**, 15353–15376 (2019).
 79. K. Kandler, K. Lieke, N. Benker, C. Emmel, M. Küpper, D. Müller-Ebert, M. Ebert, D. Scheuvs, A. Schladitz, L. Schütz, S. Weinbruch, Electron microscopy of particles collected at Praia, Cape Verde, during the Saharan Mineral Dust Experiment: Particle chemistry, shape, mixing state and complex refractive index. *Tellus B* **63**, 475–496 (2011).
 80. T. Müller, A. Schladitz, K. Kandler, A. Wiedensohler, Spectral particle absorption coefficients, single scattering albedos and imaginary parts of refractive indices from ground based in situ measurements at Cape Verde Island during SAMUM-2. *Tellus B* **63**, 573–588 (2011).
 81. B. T. Johnson, S. R. Osborne, J. M. Haywood, M. A. J. Harrison, Aircraft measurements of biomass burning aerosol over West Africa during DABEX. *J. Geophys. Res. Atmos.* **113**, (2008).
 82. K. Lieke, K. Kandler, D. Scheuvs, C. Emmel, C. V. Glahn, A. Petzold, B. Weinzierl, A. Veira, M. Ebert, S. Weinbruch, L. Schütz, Particle chemical properties in the vertical column based on aircraft observations in the vicinity of Cape Verde Islands. *Tellus B* **63**, 497–511 (2011).
 83. S. Lance, C. A. Brock, D. Rogers, J. A. Gordon, Water droplet calibration of the Cloud Droplet Probe (CDP) and in-flight performance in liquid, ice and mixed-phase clouds during ARCPAC. *Atmos. Meas. Tech.* **3**, 1683–1706 (2010).

84. W. C. Hinds, *Aerosol Technology: Properties, Behavior, and Measurement of Airborne Particles* (Wiley, ed. 2, 1999).
85. K. R. Travis, C. L. Heald, H. M. Allen, E. C. Apel, S. R. Arnold, D. R. Blake, W. H. Brune, X. Chen, R. Commane, J. D. Crouse, B. C. Daube, G. S. Diskin, J. W. Elkins, M. J. Evans, S. R. Hall, E. J. Hints, R. S. Hornbrook, P. S. Kasibhatla, M. J. Kim, G. Luo, K. McKain, D. B. Millet, F. L. Moore, J. Peischl, T. B. Ryerson, T. Sherwen, A. B. Thames, K. Ullmann, X. Wang, P. O. Wennberg, G. M. Wolfe, F. Yu, Constraining remote oxidation capacity with ATOM observations. *Atmos. Chem. Phys.* **20**, 7753–7781 (2020).
86. R. R. Draxler, G. Hess, Description of the HYSPLIT4 modeling system (Air Resources Laboratory, 1997).
87. J. D. Lee, F. A. Squires, T. Sherwen, S. E. Wilde, S. J. Cliff, L. J. Carpenter, J. R. Hopkins, S. J. Bauguitte, C. Reed, P. Barker, G. Allen, T. J. Bannan, E. Matthews, A. Mehra, C. Percival, D. E. Heard, L. K. Whalley, G. V. Ronnie, S. Seldon, T. Ingham, C. A. Keller, K. E. Knowland, E. G. Nisbet, S. Andrews, Ozone production and precursor emission from wildfires in Africa. *Environ. Sci. Atmos.* **1**, 524–542 (2021).
88. D. L. Savoie, J. M. Prospero, E. S. Saltzman, Non-sea-salt sulfate and nitrate in trade wind aerosols at Barbados: Evidence for long-range transport. *J. Geophys. Res. Atmos.* **94**, 5069–5080 (1989).
89. B. Mason, *Principles of Geochemistry* (Wiley, ed. 3, 1966).
90. R. Schlitzer, E. Masferrer Dodas, M. Adjou, R. F. Anderson, F. Andre, D. M. Cockwell, C. Jeandel, W. Geibert, W. Landing, M. Lohan, M. T. Maldonado, A. Tagliabue, W. Abouchami, E. P. Achterberg, A. M. Agather, A. Aguilar-Islas, H. Amakawa, P. Andersson, A. Annett, C. Archer, K. Arrigo, L. Artigue, M. Auro, O. Baars, I. Baconnais, A. Baker, K. Bakker, W. Bam, H. W. Bange, K. Barbeau, C. Basak, M. Baskaran, N. R. Bates, D. Bauch, F. Baurand, S. M. Becker, P. van Beek, M. Behrens, M. Belhadi, M. Belton, B. Bergquist, E. Black, S. Blain, K. Bluhm, A. R. Bowie, K. L. Bowman, P. Boyd, M. Boye, E. A. Boyle, P. Branellec, L. Bridgestock, G. Brissebrat, B. K. A. T. J. Browning, K. W. Bruland, M. Brzezinski, K. N. Buck, N. J. Buck, C. Buck, K. Buesseler, A. J. Bull, R. Bundy, E. Butler, P. H. Cai, D. Cardinal, C. Carlson, N. Casacuberta Arola, K. Casciotti, M. Castrillejo, E. Chamizo Calvo, R. Chance, M. A. Charette, B. L. A. Charlier, Z. Chase, J. E. Chavez, M. Cheize, M. Chen, H. Cheng, F. Chever, V. Chinni, R. Chmiel, M. Christl, T. M. Church, I. Closset, A. Colman, M. Colombo, M. Conte, T. M. Conway, M. Corkill, D. Cossa, S. E. Cravatte, K. C. Crocket, P. Croot, J. Cullen, G. A. Cutter, H. J. de Baar, T. van de Flierdt, G. de Souza, J. S. De Vera, F. Dehairs, F. Deng, P. van der Merwe, K. Djaoudi, A. Dufour, G. Dulaquais, Y. Echegoyen Sanz, L. Edwards, C. Ehler, M. Ellwood, S. Fan, R. A. Fine, J. N. Fitzsimmons, M. Q. Fleisher, M. S. Forbes, R. Francois, M. Frank, J. Friedrich, F. Fripiat, H. Fröllje, J. Gagnon, S. J. G. Galer, M. Gallinari, T. Gamo, Y. Gao, J. Garcia Orellana, E. Garcia Solsona, M. Gault-Ringold, E. George, L. J. A. Gerringa, M. Gilbert, J. M. Godoy, M. Le Goff, S. L. Goldstein, A. Gourain, J. Granger, P. Grasse, M. Grenier, E. Grossteffan, C. Guieu, M. Gutjahr, R. Van Hale, C. R. Hammerschmidt, J. Happell, C. Hassler, E. C. Hathorne, M. Hatta, N. J. Hawco, C. T. Hayes, L. Heimburger, M. I. Heller, P. B. Henderson, G. Henderson, T. Henry, S. van Heuven, C. Holmden, T. M. Holmes, J. Hopkins, C. J. M. Hoppe, M. J. Hopwood, T. J. Horner, Y. Hsieh, K. Huang, M. P. Humphreys, S. L. Jackson, D. J. Janssen, W. J. Jenkins, L. T. Jensen, S. John, J. L. Jones, D. Kadko, R. Kayser, T. C. Kenna, J. Kenyon, R. Khondoker, D. Kieke, M. Kienast, T. Kim, L. Kipp, J. K. Klar, M. Klunder, Y. Kondo, K. Kreissig, S. Kretschmer, S. Krisch, K. Krupp, K. Kulinski, Y. Kumamoto, K. Kunde, P. Laan, F. Lacan, M. Lagarde, P. J. Lam, M. Lambelet, C. Lamborg, D. Lannuzel, P. Latour, G. Laukert, E. C. Laurenceau-Cornec, F. A. C. le Moigne, J. van Oijjen, E. Le Roy, E. van Weerlee, O. Lechtenfeld, J. Lee, N. Lehmann, N. Lemaître, P. Lherminier, X. Li, J. Li, B. T. Liguori Pires, S. H. Little, M. Lopez Lora, D. E. Lott, A. J. M. Lough, Y. Lu, C. Mahaffey, K. Maiti, F. Marin, C. Marsay, P. Masqué, E. Mawji, M. R. McIlvin, C. Measures, S. Mehic, S. Michael, R. Middag, A. Miere, A. Milne, H. Minami, J. Moffett, W. Moore, B. Moran, P. Morton, A. Mucci, P. Mukherjee, J. Nishioka, R. L. Nixon, T. Noble, M. G. Novak, S. H. Nunige, H. Obata, H. Ogawa, D. C. Ohnemus, K. Orians, S. Ossebaer, J. E. O'Sullivan, K. Pahnke, M. Paul, F. J. Pavia, H. Perez-Tribouillier, M. M. G. Perron, B. D. Peters, M. V. Petrova, V. Pham, P. Pinedo-Gonzalez, A. Piotrowski, Y. Plancherel, F. Planchon, H. Planquette, A. Plante, S. Pohle, D. Porcelli, C. Pradoux, B. C. Proemse, V. Puigcorbe, E. Pulido-Villena, P. Quay, P. Rafter, I. Rapp, L. Ratnarajah, S. Rauschenberg, C. Rees, M. Rehkamper, R. Rember, T. Remenyi, J. A. Resing, S. Retelletti Brogi, M. J. A. Rijkensberg, L. F. Robinson, M. Roca Marti, T. Roeske, S. Roig, J. Rolison, M. Rosenberg, A. R. S. Ross, M. Roy-Barman, A. Ruacho, M. M. Rutgers van der Loeff, M. A. Saito, P. Salau, V. Sanial, A. Santoro, G. Sarthou, C. Schallenberg, U. Schauer, H. Scher, C. Schlosser, N. Schuback, C. Schwanger, P. Scott, P. N. Sedwick, S. Selzer, K. Seyitmuhammedov, R. Shelley, R. Sherrill, K. Sherrin, A. M. Shiller, M. Sieber, D. Sigman, S. K. Singh, N. D. Singh, W. M. Smethie, A. J. R. Smith, Y. Sohrin, B. M. Sohst, J. E. Sonke, M. Souhait, S. Speich, R. Steinfeldt, T. Stichel, I. Stimac, C. Stirling, B. Summers, G. J. Swarr, J. H. Swift, V. Tailandier, M. Thomas, S. Tibben, R. Till, A. T. Townsend, P. Tréguer, J. Tremblay, T. Trull, R. Tuerena, B. Twining, O. Valk, D. Vance, C. Venchiarutti, L. H. Vieira, M. Villa Alfageme, S. M. Vivancos, A. H. L. Voelker, T. Wagener, E. Wallner Halewood, R. Wang, F. Wang, R. Wang, M. J. Warner, R. J. Watson, T. Weber, A. Wefing, R. Weisend, D. Weiss, L. M. Whitmore, K. Wong, E. M. S. Woodward, Y. Wu, J. Wu, K. Wuttig, N. J. Wyatt, Y. Xiang, R. C. Xie, S. Yu, P. Zhang, J. Zhang, L. Zheng, X. Zheng, C. M. Zurbrugg, The GEOTRACES Intermediate Data Product 2021 (GEOTRACES, 2021).
91. R. Atkinson, D. L. Baulch, R. A. Cox, J. N. Crowley, R. F. Hampson, R. G. Hynes, M. E. Jenkin, M. J. Rossi, J. Troe, Evaluated kinetic and photochemical data for atmospheric chemistry: Volume I—Gas phase reactions of O_x, HO_x, NO_x and SO_x species. *Atmos. Chem. Phys.* **4**, 1461–1738 (2004).
92. R. M. Harrison, A.-M. N. Kitto, Evidence for a surface source of atmospheric nitrous acid. *Atmos. Environ.* **28**, 1089–1094 (1994).
93. R. M. Harrison, J. D. Peak, G. M. Collins, Tropospheric cycle of nitrous acid. *J. Geophys. Res. Atmos.* **101**, 14429–14439 (1996).
94. I. Trebs, L. L. Lara, L. M. M. Zeri, L. V. Gatti, P. Artaxo, R. Dlugi, J. Slanina, M. O. Andreae, F. X. Meixner, Dry and wet deposition of inorganic nitrogen compounds to a tropical pasture site (Rondonia, Brazil). *Atmos. Chem. Phys.* **6**, 447–469 (2006).
95. J. Stutz, B. Alicke, A. Nefel, Nitrous acid formation in the urban atmosphere: Gradient measurements of NO₂ and HONO over grass in Milan, Italy. *J. Geophys. Res. Atmos.* **107**, LOP 5-1–LOP 5-15 (2002).
96. J. W. Deardorff, Convective velocity and temperature scales for the unstable planetary boundary layer and for Rayleigh convection. *J. Atmos. Sci.* **27**, 1211–1213 (1970).
97. J. Kleffmann, Daytime sources of nitrous acid (HONO) in the atmospheric boundary layer. *ChemPhysChem* **8**, 1137–1144 (2007).
98. V. Michoud, A. Colomb, A. Borbon, K. Miett, M. Beekmann, M. Camredon, B. Aumont, S. Perrier, P. Zapf, G. Siour, W. Ait-Helal, C. Afif, A. Kukui, M. Furger, J. C. Dupont, M. Haefelin, J. F. Doussin, Study of the unknown HONO daytime source at a European suburban site during the MEGAPOLI summer and winter field campaigns. *Atmos. Chem. Phys.* **14**, 2805–2822 (2014).
99. F. Spataro, A. Ianniello, Sources of atmospheric nitrous acid: State of the science, current research needs, and future prospects. *J. Air Waste Manage. Assoc.* **64**, 1232–1250 (2014).
100. M. Li, H. Su, G. Li, N. Ma, U. Pöschl, Y. Cheng, Relative importance of gas uptake on aerosol and ground surfaces characterized by equivalent uptake coefficients. *Atmos. Chem. Phys.* **19**, 10981–11011 (2019).

Acknowledgments: We would like to thank S. Swift (University of York) for assistance with the ion chromatography measurements, E. Matthews (University of Manchester) for help with running HYSPLIT back trajectories for the aircraft measurements, S. Cliff (University of York) for running the gas-phase biomass burning flag, L. Kramer (Ricardo AEA) for assistance with the Cape Verde HONO measurements and data analysis, and the remaining ARNA team. Airborne data were obtained using the BAe-146-301 Atmospheric Research Aircraft (ARA) flown by Airtask Ltd. and managed by the FAAM Airborne Laboratory, jointly operated by UKRI and the University of Leeds. We thank the staff from Airtask for the campaign organization, flight planning, and resolving any safety issues: B. Black, F. Brennan, S. James, M. Robinson, and D. Simpson. We thank the NASA Global Modeling and Assimilation Office (GMAO) for forecasting support during ARNA. **Funding:** This work was supported by National Environmental Research Council (NERC) grant NE/S000518/1 to L.J.C., W.J.B., and J.D.L. The Cape Verde Atmospheric Observatory is funded through the National Centre for Atmospheric Science (NCAS) to L.J.C. S.T.A.'s PhD was supported by the SPHERES Natural Environment Research Council (NERC) Doctoral Training Partnership (DTP) under grant NE/L002574/1. L.J.C. acknowledges funding from the European Research Council (ERC) under the European Union's Horizon 2020 programme (project O3-SML; grant agreement no. 833290). **Author contributions:** Conceptualization: L.J.C. Methodology: All authors. Investigation: All authors. Visualization: S.T.A. Funding acquisition: L.J.C., J.D.L., and W.J.B. Project administration: L.J.C., J.D.L., T.S., and W.J.B. Supervision: L.J.C., J.D.L., and W.J.B. Writing (original draft): L.J.C. and S.T.A. Writing (review and editing): All authors. **Competing interests:** The authors declare that they have no competing interests. **Data and materials availability:** CVAO NO_x—Norwegian Institute for Air Research (NILU) EBAS database (https://ebas-data.nilu.no/DataSets.aspx?nations=CV238CPV&projects=ACTRIS&InstrumentTypes=chemiluminescence_photolytic&fromDate=1970-01-01&toDate=2022-12-31). Aircraft measurements—CEDA Archive (<https://catalogue.ceda.ac.uk/uuid/1a7f76b65e774cf6abb09cdee642dd73>). GEOS-Chem code—The GEOS-Chem model (v12.9.0, DOI: 10.5281/zenodo.3950327). All data needed to evaluate the conclusions in the paper are present in the paper and/or the Supplementary Materials.

Submitted 23 June 2022
Accepted 19 December 2022
Published 18 January 2023
10.1126/sciadv.add6266

Extensive field evidence for the release of HONO from the photolysis of nitrate aerosols

Simone T. Andersen, Lucy J. Carpenter, Chris Reed, James D. Lee, Rosie Chance, Toms Sherwen, Adam R. Vaughan, Jordan Stewart, Pete M. Edwards, William J. Bloss, Roberto Sommariva, Leigh R. Crilley, Graeme J. Nott, Luis Neves, Katie Read, Dwayne E. Heard, Paul W. Seakins, Lisa K. Whalley, Graham A. Boustead, Lauren T. Fleming, Daniel Stone, and Khandeh Wadinga Fomba

Sci. Adv., **9** (3), eadd6266.
DOI: 10.1126/sciadv.add6266

View the article online

<https://www.science.org/doi/10.1126/sciadv.add6266>

Permissions

<https://www.science.org/help/reprints-and-permissions>

Use of this article is subject to the [Terms of service](#)

Science Advances (ISSN) is published by the American Association for the Advancement of Science. 1200 New York Avenue NW, Washington, DC 20005. The title *Science Advances* is a registered trademark of AAAS.

Copyright © 2023 The Authors, some rights reserved; exclusive licensee American Association for the Advancement of Science. No claim to original U.S. Government Works. Distributed under a Creative Commons Attribution License 4.0 (CC BY).