

Aging modulates homeostatic leukocyte trafficking to the peritoneal cavity in a sex-specific manner

Hopkin, Sophie J; Pezhman, Laleh; Begum, Jenefa; Kavanagh, Dean; McGettrick, Helen M; Iqbal, Asif J; Chimen, Myriam

DOI:

[10.1093/jleuko/qiad053](https://doi.org/10.1093/jleuko/qiad053)

License:

Creative Commons: Attribution (CC BY)

Document Version

Publisher's PDF, also known as Version of record

Citation for published version (Harvard):

Hopkin, SJ, Pezhman, L, Begum, J, Kavanagh, D, McGettrick, HM, Iqbal, AJ & Chimen, M 2023, 'Aging modulates homeostatic leukocyte trafficking to the peritoneal cavity in a sex-specific manner', *Journal of Leukocyte Biology*, pp. 1-14. <https://doi.org/10.1093/jleuko/qiad053>

[Link to publication on Research at Birmingham portal](#)

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.



When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

Aging modulates homeostatic leukocyte trafficking to the peritoneal cavity in a sex-specific manner

Sophie J. Hopkin,¹ Laleh Pezhman,² Jenefer Begum,¹ Dean Kavanagh,¹ Helen M. McGettrick,^{2,*†} Asif J. Iqbal,^{1,*†}  and Myriam Chimen^{2,*†} 

¹Institute of Cardiovascular Sciences, University of Birmingham, Birmingham, B15 2TT, United Kingdom

²Institute of Inflammation and Ageing, University of Birmingham, Birmingham, B15 2TT, United Kingdom

*Corresponding author: Institute of Inflammation and Ageing, University of Birmingham, Birmingham, B15 2TT, United Kingdom. Email: m.chimen@bham.ac.uk; Institute of Cardiovascular Sciences, University of Birmingham, Birmingham, B15 2TT, United Kingdom. Email: a.j.iqbal@bham.ac.uk; Institute of Inflammation and Ageing, University of Birmingham, Birmingham, B15 2TT, United Kingdom. Email: h.m.mcgettrick@bham.ac.uk

Abstract

Aging is associated with exacerbated systemic inflammation (inflammaging) and the progressive loss of immune system function (immunosenescence). Leukocyte migration is necessary for effective immunity; however, dysregulated trafficking of leukocytes into tissue contributes to inflammaging and the development of age-related inflammatory diseases. Aging modulates leukocyte trafficking under inflammatory conditions; however, whether aging modulates leukocyte trafficking under homeostatic conditions remains to be elucidated. Although immune responses are evidently sexually dimorphic, limited studies have investigated the effect of sex on age-related changes to leukocyte trafficking processes. Here, we investigated age-related and sex-specific changes to the leukocyte populations within the peritoneal cavity of young (3-mo), middle-aged (18-mo) and old (21-mo) male and female wild-type mice in the steady state. We found an age-related increase in the number of leukocytes within the peritoneal cavity of female mice, predominantly B cells, which may reflect increased trafficking through this tissue with age. This was accompanied by an increased inflammatory environment within the aged cavity, including increased levels of chemoattractants, including B cell chemoattractants CXCL13 and CCL21, soluble adhesion molecules, and proinflammatory cytokines, which was more pronounced in aged female mice. Intravital microscopy techniques revealed altered vascular structure and increased vascular permeability within the peritoneal membrane of aged female mice, which may support increased leukocyte trafficking to the cavity with age. Together, these data indicate that aging affects homeostatic leukocyte trafficking processes in a sex-specific fashion.

Keywords: aging, inflammation, leukocyte trafficking, sexual dimorphism, T cells

1 Introduction

Aging profoundly affects the mammalian immune system, with older mice and humans exhibiting increased circulating levels of proinflammatory mediators^{1–3} and changes to immune cell function^{4–8} when compared with their younger counterparts. Growing evidence suggests that the aging process significantly modulates leukocyte trafficking during inflammation in both humans and mice (reviewed by Hopkin et al.).⁹ For example, immunization of aged (>20 mo) C57Bl6 mice with ovalbumin resulted in reduced CD4⁺ T cell recruitment to the spleen compared with young (<3 mo) mice, which correlated with delayed induction of antigen-specific immunity.¹⁰ However, much less is known about the impact of age on leukocyte trafficking associated with immunosurveillance and tissue homeostasis.^{11–13} Given this, it is crucial for us to increase our understanding of the impact of age on leukocyte trafficking that leads to weakened immune surveillance in the elderly.

Although it is well established that immune responses are sexually dimorphic in both humans and mice,^{14–16} to date most in vivo studies have used either a single sex or a mixture of male and female participants to dissect the mechanisms underpinning immunological events.^{16,17} Therefore, the impact of biological sex on age-related changes to the immune system has been severely underresearched. There remains controversy over whether such

changes are more pronounced in males or females.^{18,19} While older men tend to present with an immune risk profile (i.e., indicative of increased mortality),²⁰ older women are more susceptible to developing autoimmune diseases.^{16,21,22} Indeed, during the 2020 SARS-CoV-2 pandemic, older men (63 to 76 yr old) had increased mortality rates following SARS-CoV-2 infection compared with older women.²³ However, healthy older women (50 to 60 yr old) have increased circulating levels of IL-6 and cholesterol when compared with older men (48 to 58 yr old), suggesting increased systemic inflammation and increased risk of cardiovascular events in otherwise healthy older women.²⁴ It remains unclear as to the extent hormones, sex chromosomes, or gender-dependent environmental cues influence the aging immune system,¹⁸ as well as the extent of their influence on age-related changes to leukocyte trafficking processes.^{9,13} To understand whether aging effects homeostatic leukocyte trafficking in a sex-dependent manner, we quantified leukocyte populations across multiple tissues in young (3 mo), middle-aged (18 mo) and old (21 mo) male and female mice using flow cytometry. The most striking age-related changes to tissue leukocyte composition occurred in the peritoneal cavity, and this age-dependent phenotype was more pronounced in female mice. Aging significantly increased the number of leukocytes within the peritoneal cavity, which was associated with increases in proinflammatory soluble mediators in the peritoneum indicative of age-related changes

[†] Authors contributed equally to this research.

Received: December 1, 2022. **Revised:** March 28, 2023. **Accepted:** April 28, 2023. **Corrected and Typeset:** June 13, 2023

© The Author(s) 2023. Published by Oxford University Press on behalf of Society for Leukocyte Biology.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted reuse, distribution, and reproduction in any medium, provided the original work is properly cited.

to homeostatic leukocyte trafficking. Additionally, we observed that the integrity of the peritoneal membrane was compromised with age and may assist leukocyte trafficking into the peritoneal cavity in the steady state. The sex-specific changes to homeostatic leukocyte trafficking that occur with age likely contribute to the dimorphic immune responses observed in older males and females.

2 Materials and methods

2.1 Murine tissue collection and processing

Experiments were conducted in accordance with UK Home Office regulations and appropriate ethics. Three-month-old (28.36 ± 2 g male; 22.04 ± 1 g female), 18-mo-old (41.26 ± 6 g male; 30.90 ± 3 g female) and 21-mo-old (32.73 ± 2 g male; 30.08 ± 1 g female) C57Bl/6J wild-type mice were purchased from Charles River Laboratories and were housed at the University of Birmingham animal unit with free access to food and water. Mice were maintained in a specific pathogen-free environment, determined by quarterly health screening of the unit using Federation of European Laboratory Animal Science Associations approved methods. Mice were fed the rodent 5LF2 diet (IPS LabDiet) and were housed in the animal facility for at least 2 mo prior to experimentation. Environmental conditions were 21 ± 2 °C, $55\% \pm 10\%$ relative humidity, and a 12-h light-dark cycle. Mice were culled by cardiac puncture with blood collected into EDTA-coated Eppendorfs. The peritoneal cavity was lavaged with ice cold 5 mM EDTA. Spleen, right hind limb, and inguinal lymph nodes (iLNs) were isolated and stored in phosphate-buffered saline (PBS). Peritoneal lavage fluid (PLF) was centrifuged at 400 *g* for 5 min: supernatant was stored at -80 °C and cells were resuspended in MACS buffer (0.1 mM EDTA, 0.6% bovine serum albumin in PBS; all from Sigma-Aldrich). The spleen and iLNs were crushed through a 40- μ m filter. Bone marrow was flushed out of the right hind limb using PBS. Spleen, iLN, bone marrow, and blood samples were all incubated in RBC lysis buffer (155 mM NH_4Cl , 12 mM NaHCO_3 , 0.1 mM EDTA; all from Sigma-Aldrich) for 10 min, followed by centrifugation at 400 *g* and then resuspension in MACS buffer.

2.2 Flow cytometric analysis

All samples were blocked with FcR blocker (Miltenyi Biotec) prior to staining with the following of antibodies for 20 min at 4 °C, after which samples were washed and fixed with 2% formaldehyde: anti-CD3 PE-Cy7 (clone 145-2c11), anti-CD4 eFluor450 (clone GK1.5), anti-CD8 PE-TexasRed (clone 5H10), anti-CD44 FITC (clone IM7), anti-CD25 AF700 (clone PC61.5), anti-KLRG1 APC-eFluor780 (clone 2F1), anti-CD62L PE (clone MEL-14), anti-CD19-APC (clone 1D3), anti-CD45 APC-CY7 (clone 104), F4/80 FITC (clone BM8), anti-CD11c PE-Cy7 (clone N418), anti-gp38 PE (clone 8.1.1) (all from Thermo Fisher Scientific); anti-CD45.2 BV605 (clone 104), anti-CD23 BV421 (clone B3B4), anti-CD93 BV650 (clone AA4.1), anti-CD43 PerCP-Cy5.5 (clone S7), anti-CD21/35 PE (clone 7G6), anti-Siglec F PE-CF594 (clone E50-2440), Ly6G APC (clone 1A8) (all from BD). Compensation controls were generated using cells isolated from the spleen. Immediately prior to analysis CountBright beads (Invitrogen) and Zombie Aqua (BioLegend) were added and samples were acquired using Fortessa-X20. Data were analyzed offline using FlowJo (V-10.2.6; TreeStar) using the flow cytometry gating strategy depicted in [Supplementary Fig. 1](#). Leukocyte populations were quantified using CountBright beads according to the manufacturer's instructions. Data are presented as the total number of each leukocyte population within a given tissue.

2.3 Intravital microscopy

Three-month-old and 21-mo-old male and female C57Bl6 mice were anesthetized by intraperitoneal injection of ketamine hydrochloride (100 mg/kg; Pfizer) and medetomidine hydrochloride (10 mg/kg; Pfizer). The trachea and right common carotid artery were cannulated, and the peritoneal membrane was exposed. Prior to imaging, 50 μ L 20 kDa dextran-TRITC and 50 μ L 70 kDa dextran-FITC (both at 50 mg/mL; both from Sigma-Aldrich) were administered via the carotid artery. The peritoneal membrane was imaged at 60 min post-dextran administration using an Olympus IX81 inverted microscope. After imaging, the mice were sacrificed via cervical dislocation and the PLF was collected as described previously. The concentration of fluorescent dextrans in the PLF was quantified using a BioTek Synergy HT plate reader and the data presented as μ g dextran per mL PLF. Images were analyzed offline using Angiogenesis Analyzer plug-in on Fiji Image J (version 2.1.0; Cambridge) to calculate the number of vessels and total vessel length, number of nodes (points of bifurcation), number of mesh structures (vessel circuits), and total mesh area (all expressed per mm^2).

2.4 Cytokine array

The secretome of the PLF was quantified using the Mouse XL Cytokine Array Kit according to the manufacturer's instructions (ARY028; R&D Systems). Membranes were exposed for 1 min and developed using the Compact X4 Automatic X-ray Film Processor (Xograph). The developed membrane was scanned ([Supplementary Fig. 2](#)), and the images were analyzed using ImageJ v2.1.0 (National Institutes of Health). Briefly, regions of interest were drawn around each analyte and the intensity value was extrapolated. As each analyte was present in duplicate on the membrane, intensity values were averaged and the background value was deducted ([Supplementary Table 1](#)). The intensity values of the 21-mo-old mice were then divided by the intensity values of 3-mo-old mice to determine the age-related fold-change in analytes for a specific sex group. Additionally, the intensity values of the female mice were then divided by the intensity values of the male mice to determine the sex-specific fold-change in analytes for a specific age group. Each data plot represents pooled samples from 3 mice per age group for each sex.

2.5 Enzyme-linked immunosorbent assay

The concentration of CCL21 (DY457) and CXCL13 (DY470) in the PLF and TNF α (DY410) and IL-6 (DY406) in the serum was quantified using enzyme-linked immunosorbent assay (ELISA) (all from Bio-Techne) according to the manufacturer's instructions. For the PLF, neat samples were added in duplicate to the 96-well plate. For the serum samples, singular neat samples were added to the 96-well plate due to limited serum. OD₄₅₀ values were determined using the BioTek Synergy HT plate reader. Background signal was deducted from the OD₄₅₀ values of the samples, standard curves were generated for each chemokine/cytokine, and the concentration of each chemokine/cytokine was then interpolated for the PLF and serum samples. Data are presented as pg/mL.

2.6 Statistical analysis

Data were analyzed using GraphPad v8.4.3 Prism (GraphPad Software) and presented as mean \pm SEM for *n* mice per group for *n* independent experiments. Normality was assessed using Shapiro-Wilk test. Univariate analysis was performed using unpaired *t* test. Multivariate analysis was performed using analysis of variance, with Bonferroni posttest. *P* < 0.05 was deemed statistically significant.

3 Results

3.1 Leukocyte trafficking to the naïve peritoneal cavity is dysregulated by aging in a sex-specific manner

In order to determine whether aging or biological sex affects homeostatic leukocyte trafficking processes, we initially investigated the leukocyte composition of the peritoneal cavity in young (3-mo), middle aged (18-mo) and old (21-mo) noninflamed mice for both sexes. Low numbers of CD45⁺ leukocytes were detected in the naïve peritoneal cavity of 3- and 18-mo-old mice, and these were comparable across both sexes (Fig. 1A). Aging (above 21 mo) increased the accumulation of CD45⁺ leukocytes within the cavity in both sexes, although this was only significant within the female group (Fig. 1A). Interestingly, significantly higher numbers of CD45⁺ leukocytes were seen in the peritoneum of aged female mice compared with aged male mice (Fig. 1A), indicating age-associated sexual dimorphism in leukocyte trafficking to the cavity under homeostatic conditions.

Further analysis of leukocyte subsets revealed similar age-related trends for peritoneal Ly6G⁺ neutrophils (Fig. 1B), CD11c⁺ dendritic cells (DCs) (Fig. 1C), CD3⁺ T cells (Fig. 1D), and CD3⁺ KLRG1⁺ senescent T cells (Fig. 1E)—in which numbers for these subsets were all significantly elevated in the 21-mo-old mice and were broadly comparable between the sexes when compared with 3-mo-old mice. Similar trends were also seen for CD3⁺ T cell subsets, including CD3⁺ CD8⁺ T cells, CD8⁺ CD62L⁺ CD44⁺ central memory T cells, and CD8⁺ CD62L[−] CD44⁺ effector memory T cells (Supplementary Fig. 3). Interestingly, we observed significantly more Ly6G⁺ neutrophils and CD11c⁺ DCs in male mice at 21 mo-old compared with their 18-mo-old counterparts (Fig. 1B and C). In contrast, CD3⁺ T cell (Fig. 1D) numbers, including CD3⁺ KLRG1⁺ senescent T cell (Fig. 1E) and CD8⁺ CD62L[−] CD44⁺ effector memory T cell (Supplementary Fig. 3) numbers, were higher in 21-mo-old female mice compared with 18-mo-olds. Other peritoneal leukocyte subsets displayed different age-related trends. Significantly more CD19⁺ B cells and CD19⁺ CD93[−] CD43[−] CD21[−] CD23[−] age-associated B cells (ABCs) were detected in the peritoneum of 21-mo-old female mice compared with 3- and 18-mo-old mice (Fig. 1F and G). Interestingly, we observed no age-associated change in these populations in male mice, indicating sexual dimorphism in the trafficking of B cells and ABCs in aged female mice. In contrast to the previous populations, peritoneal F4/80^{hi} macrophages significantly declined with age for both sexes, with significant reductions seen at both 18 and 21 mo (Fig. 1H). Altogether, these observations indicate that aging skews the peritoneal environment toward a proinflammatory state that supports the recruitment of neutrophils, antigen-presenting cells (DCs, B cells), and T cells, including senescent T cells, into the cavity, which may exacerbate the inflammatory environment even further. Moreover, the aging peritoneal environment appears to promote tissue resident macrophage egress from the cavity, which is indicative of the proinflammatory state of the tissue.

3.2 Aging modulates the composition of leukocytes in the blood and lymphoid organs in the steady state, indicating leukocyte mobilization

The age-associated changes in leukocyte trafficking into the peritoneal cavity may simply mirror the absolute numbers of these subpopulations circulating in the blood as a consequence of inflammation. Indeed, circulating levels of TNF α and IL-6 increased with age; however, this was only statistically significant for

21-mo-old female mice (Supplementary Fig. 4) and not for male mice. While age tended to increase circulating numbers of CD45⁺ leukocytes, this was only statistically significant in male mice (Fig. 2A). The numbers of CD19⁺ B cells and Ly6G⁺ neutrophils were also significantly increased in the blood of 21-mo-old male and female mice compared with 3- and 18-mo-old animals (Fig. 2B and Supplementary Fig. 5A) and broadly corresponded with increased numbers of these leukocytes within the aged peritoneal cavity. In contrast, increased circulating numbers of CD11c⁺ DCs were observed in 21-mo-old male mice but not in female mice when compared with 3- and 18-mo-old mice (Supplementary Fig. 5B), which corresponded with increased CD11c⁺ DC numbers in the peritoneal cavity of these mice. It is therefore possible that the age-related increase in these leukocyte populations within the peritoneal cavity may be due to the extravasation of these leukocytes from the blood.

Aging is known to reduce the pool of immune cells within lymphoid tissues, with reduced numbers of B cells in the bone marrow,²⁵ T cells in the thymus²⁶ and both cell types in the spleen and LNs²⁷ all reported. For example, the T cell population within the iLNs was reported to be reduced by approximately 40% in older (18 to 21 mo) mice compared with young (2 to 3 mo) mice.²⁸ As such, we investigated the effect of age and sex on the composition of leukocytes within the lymphoid tissues of the bone marrow, spleen, and iLNs. We observed no difference in CD45⁺ numbers in the bone marrow of mice with age or with sex (Fig. 2C). We observed a significant reduction in CD19⁺ B cell numbers and increase in CD11c⁺ DC numbers within the bone marrow of 21-mo-old female mice, but not in male mice, compared with 3-mo-old animals (Supplementary Fig. 6A and B). While the numbers of CD45⁺ cells remained unchanged with age in the spleens of female mice, we observed a significant age-associated reduction in the number of these cells in the spleens of male mice at 18 and 21 mo-old compared with 3-mo-old animals (Fig. 2D). Aging resulted in reduced CD3⁺ T cell numbers within the spleens of male and female mice (Supplementary Fig. 7A). In contrast, splenic CD19⁺ B cell, CD11c⁺ DC, and F4/80^{hi} macrophage numbers were all increased in 21-mo-old female mice compared with their young counterparts (Fig. 2E and Supplementary Fig. 7B and C). Surprisingly, we observed an almost complete loss of CD45⁺ cells, including CD3⁺ T cells, CD19⁺ B cells, and CD11c⁺ DCs, within the iLNs at 18- and 21 mo-old mice compared with 3-mo-old mice for both sexes (Fig. 2F and Supplementary Fig. 8A to C). Collectively, these data indicate that aging dysregulates leukocyte trafficking through secondary lymphoid tissues, except the bone marrow, and is sexually dimorphic for some tissues (e.g., spleen).

3.3 Local chemokine levels are increased in the aging peritoneum

Next, we examined whether the age-related and sex-specific differences in leukocyte recruitment to the peritoneal cavity under homeostatic conditions could be explained by variations in the levels of chemokines/chemoattractants, leukocyte survival/growth factors, and soluble adhesion molecules within the PLF, reminiscent of previous studies.^{15,29} Indeed, we observed a ≥ 3 -fold increase in the levels of chemokines in the peritoneal cavity of aged male and female mice compared with young mice, with these age-associated changes being more pronounced in female mice (Fig. 3A and Supplementary Fig. 9). Interestingly, some chemokines (CCL12, CCL19) were restricted to age-associated increases in their expression in a sex-specific manner. For example, a ≥ 3 -fold increase in the expression of CCL12,




Fig. 1. Baseline trafficking of leukocytes into the resting peritoneum is dysregulated by aging in a sex-specific manner. Total number of peritoneal (A) CD45⁺ leukocytes, (B) Ly6G⁺ neutrophils, (C) CD11c⁺ DCs, (D) CD3⁺ T cells, (E) CD3⁺ KLRG1⁺ senescent T cells, (F) CD19⁺ B cells, (G) CD19⁺ CD93⁻ CD43⁻ CD21⁻ CD23⁻ ABCs, and (H) F4/80^{hi} macrophages from resting 3-, 18-, and 21-mo-old female (♀) and male (♂) C57Bl6 mice were quantified using flow cytometry. Analysis of variance showed a significant effect of age on the number of each leukocyte subset ($P < 0.01$) and of sex on the number of leukocytes ($P < 0.001$), macrophages ($P < 0.01$), and B cells ($P < 0.05$) cells. Data are presented as mean \pm SEM using $n = 5$, $n = 8$, and $n = 6$ for 3-, 18-, and 21-mo-old mice, respectively, from $n = 2$ independent experiments per age group for each sex. Color-coded data points indicate independent experiments per age group. * $P < 0.05$, ** $P < 0.01$, and *** $P < 0.001$ by Bonferroni multiple comparison posttest.




Fig. 2. Leukocyte trafficking through blood and lymphoid organs are altered with age, displaying sexual dimorphism. (A, C, D, F) CD45⁺ leukocytes and (B, E) CD19⁺ B-cells within the (A, B) blood, (C) bone marrow, (D, E) spleen, and (F) iLNs of naive 3-, 18- and 21-mo-old female (♀) and male (♂) C57Bl6 mice were quantified using flow cytometry. Analysis of variance showed a significant effect of age on the number of leukocytes in the blood, spleen, and iLNs ($P < 0.05$), and of sex on leukocyte numbers in the blood ($P < 0.001$). Analysis of variance analysis also showed a significant effect of age and sex on the number of B cells in the blood and spleen ($P < 0.01$). Data are presented as mean \pm SEM using $n = 5$, $n = 8$, and $n = 6$ for 3-, 18-, and 21-mo-old mice, respectively, from $n = 2$ independent experiments per age group for each sex. Color-coded data points indicate independent experiments per age group. * $P < 0.05$, ** $P < 0.01$, and *** $P < 0.001$ by Bonferroni multiple comparison posttest.




Fig. 3. Aging skews the peritoneal secretome toward a proinflammatory microenvironment. An abundance of proinflammatory mediators in the peritoneal fluid of naïve 3- and 21-mo-old female (♀) and male (♂) C57Bl6 mice were analyzed using a cytokine array. Fluids from 3 mice per age group for each sex were pooled, in which $n = 1$. Heatmaps depict the fold change in (A) chemokines/chemoattractants, growth/survival factors, and soluble adhesion molecules, with (D) cytokines of aged mice relative to young mice for each sex. Analytes with <3 -fold change are represented by white squares. Quantification of (B) CCL21 and (C) CXCL13 in the peritoneal lavage fluid of 3-, 18-, and 21-mo-old female and male wild-type mice by ELISA. Analysis of variance showed a significant effect of age ($P < 0.001$) and sex ($P < 0.001$) on the concentration of CCL21 in the peritoneal fluid and of age ($P < 0.001$), but not sex, on the concentration of CXCL13. Data are presented as mean \pm SEM using $n = 4$ to 5 per age group, from $n = 1$ independent experiment. * $P < 0.05$, ** $P < 0.01$, and *** $P < 0.001$ by Bonferroni multiple comparison posttest.




Fig. 4. Aging was associated with increased permeability and modulated structure of the peritoneal membrane vasculature. (A) Relative abundance of angiogenic factors in the peritoneal fluid of naive 3- and 21-mo-old female (♀) and male (♂) C57Bl6 mice were analyzed using a cytokine array. Fluids from 3 mice per age group were pooled, in which n = 1. Heatmaps depict the fold change in cytokines and angiogenic factors of aged mice relative to young mice for each sex group. The vasculature of the peritoneal membrane from (B, D) young (3 mo) and (C, E) aged (21 mo) (B, C) female and (D, E) male mice were imaged using intravital microscopy. Images were quantified to assess (F) the leakage of fluorescent dextran into the extravascular space, expressed as mean number of gray pixels; (I) average number of vessels; (J) total length of vessels; (K) number of nodes; (L) number of meshes; and (M) total mesh area all expressed per mm². Quantification of the concentration of (G) 20 kDa and (H) 70 kDa fluorescent dextran within the peritoneal lavage fluid postimaging. Data are mean ± SEM for n = 5 independent experiments, using 5 mice per age group. *P < 0.05, **P < 0.01, ***P < 0.001 by Bonferroni multiple comparison test. Scale bar = 10 µm.

contributing to age-related changes in peritoneal leukocyte composition.

4 Discussion

Homeostatic leukocyte trafficking is key to maintain tissue homeostasis and immunosurveillance.¹³ Although growing evidence supports the notion that the aging process affects leukocyte trafficking patterns during inflammation (reviewed by Hopkin et al.),⁹ a limited number of studies have investigated the impact of chronological aging on homeostatic leukocyte trafficking, and even fewer have considered sexual dimorphism within these age-related changes.^{32–35} Here, we found that aging was associated with significantly increased numbers of leukocytes and proinflammatory mediators in the peritoneal cavity of mice in the steady state. Importantly, these age-related changes occurred in a sex-specific manner, with increased vascular changes observed in female mice.

Peritonitis is a well-established and widely used model to study leukocyte trafficking during acute inflammation, in which the effects of age^{36,37} and sex^{15,38,39} on leukocyte recruitment to the peritoneal cavity in this model have been investigated. But studies investigating age-related changes to peritoneal leukocyte populations in the steady state are often limited to male mice.^{7,40} In aged mice, we observed increased numbers of peritoneal leukocytes, including neutrophils and B cells, and reduced macrophage levels, which is in line with Mogilenko et al.,⁷ who reported a similar profile in the peritoneal cavity of aged (17 to 24 mo) male C57Bl6 mice compared with young (3 to 6 mo) using single-cell RNA sequencing. Moreover, Linehan et al.⁴⁰ also described reduced frequencies of peritoneal macrophages in aged (15 to 20 mo) male C57Bl6 mice compared with young mice (<3 mo). We extend and report for the first time the impact of sexual dimorphism in the context of aging at the steady state. Older female mice displayed an exacerbated increase in peritoneal leukocyte numbers, especially B cells, compared with aged male mice, suggesting that females may be more susceptible to age-related changes in leukocyte trafficking patterns into the peritoneal cavity.

Increased numbers of peritoneal leukocytes in aged mice fits the paradigm of inflammaging, whereby the development of chronic inflammation during the aging process alters the inflammation status of peripheral tissues.^{41–43} We found a significant age-associated reduction in the number of leukocytes residing in the spleen and iLNs and an increase in circulating leukocytes, suggesting mobilization of leukocytes from lymphoid tissues. The structure and function of LNs are influenced by the aging process, including the leukocyte composition of the tissue (reviewed by Turner and Mabbott).²⁷ Here, we report an age-related reduction in the number of B cells within the iLNs of 21-mo-old mice compared with 3-mo-old mice. In contrast, Turner and Mabbott²⁸ reported an increase in B cells in LNs from aged female C57Bl6 mice compared with young mice, but these analyses were obtained from cervical, axillary, brachial, and inguinal LNs pooled into a single-cell suspension, which confounds interpretation. Although no direct comparisons have been made for male mice, studies utilizing rats suggest that aging has a similar impact on architecture of LN in male animals, including distorted tissue structure and cellular composition and reduced numbers of germinal center reactions.^{44,45}

Leukocyte mobilization from lymphoid tissues into the circulation was described by Kay et al.,¹⁵ whereby increased numbers of circulating Gr1^{high} neutrophils, coupled with reduced levels in the spleen, correlated with increased recruitment of neutrophils into

the peritoneal cavity upon induction of peritonitis. While acute inflammation (e.g., peritonitis) is not comparable to age-associated chronic inflammation,⁴⁶ our data from steady-state aged animals mirrors inflammation induced leukocyte trafficking between lymphoid tissue and the peritoneal cavity, thus suggesting that aging influences the dynamics of trafficking between these compartments. Indeed, increased numbers of B cells in the peritoneal cavity of aged female mice could be the result of age-related changes to B cell trafficking patterns, indicated by reduced B cell numbers in lymphoid tissues and increased B cell numbers in the circulation. That said, other factors could also potentially explain these findings (e.g., aging may influence leukocyte proliferation, apoptosis, or retention rates within the peritoneal cavity). Although there is currently no specific literature on age-related changes to proliferation/apoptosis rates of peritoneal leukocytes, there is an abundance of evidence supporting age-related changes to leukocyte proliferation^{47–49} and apoptosis^{49–52} rates across multiple tissues in humans and mice. For example, naïve CD4⁺ T cells isolated from the spleens of older (18 to 20 mo) C57Bl6 mice exhibited increased rates of apoptosis in response to antigen stimulation *ex vivo* compared with those isolated from young (2 to 3 mo) mice.⁵³ Whether aging specifically alters peritoneal leukocyte proliferation and apoptosis rates remains to be answered.

Published datasets regarding the sex-specific differences in peritoneal leukocyte populations of noninflamed mice have primarily utilized young mice and are usually a byproduct of models of inflammation.^{15,38,39} In agreement with previous studies,^{15,38} we report increased leukocyte numbers, including increased F4/80^{hi} macrophages, in the peritoneal cavity of 3-mo-old female mice compared with 3-mo-old male mice. Interestingly, Scotland et al.³⁸ reported higher numbers of peritoneal leukocytes in young (<5 mo) female C57Bl6 mice compared with young male mice, and importantly, these changes were not due to sex differences in leukocyte apoptosis or proliferation rates, but rather were attributed to the effects of sex steroids. Indeed, ovariectomized female mice had comparable numbers of peritoneal leukocytes to young male mice, suggesting that female sex hormones play a direct role in the regulation of leukocyte trafficking to the peritoneal cavity. Both estrogen and progesterone reportedly suppress endothelial activation (e.g., estrogen downregulates VCAM-1 expression on TNF α -activated endothelial cells),^{54–58} while progesterone inhibits IL-6, IL-8, CXCL2, and CXCL1 production by lipopolysaccharide-stimulated human umbilical vein endothelial cells *in vitro*.⁵⁵ In the steady state, levels of soluble ICAM-1 and VCAM-1 within serum were reportedly higher in the early phases of the menstrual cycle, in which circulating levels of estrogen and progesterone are at their lowest, and correlated with reduced numbers of circulating monocytes.⁵⁹ Therefore, sex hormones likely play a role in regulating leukocyte trafficking in young mice but not in aged mice due to the age-related decline in circulating sex hormone levels (reviewed by Horstman et al).⁶⁰

Increased levels of chemokines and soluble adhesion molecules within the aging peritoneal cavity may have facilitated the increased trafficking of various leukocytes that we observed. The approach used was an unbiased analysis of pooled samples, which may be skewed by individual mice and therefore requires validating using alternative methodologies. Indeed, ELISA analysis confirmed increased levels of the B cell chemoattractants CCL21 and CXCL13 in the PLF of aged female mice compared with young, offering a potential mechanism of action for increased B cell recruitment to the aging cavity. The changes in leukocyte numbers observed here could also be caused by an age-related increase in the survival and growth of certain

administration, resources, and supervision. S.J.H., H.M.M., A.J.I., and M.C. contributed to writing—original draft. All authors contributed to the writing—review and editing.

Supplementary material

Supplementary materials are available at *Journal of Leukocyte Biology* online.

Funding

S.J.H., L.P., and J.B. were supported by PhD studentships funded by the Royal Society PhD Studentship (RGF/R1/180008), Rosetree Foundation (A2092), and the British Heart Foundation (FS/20/2/34799), respectively. M.C. was supported by a Royal Society Dorothy Hodgkin Fellowship (DH16044). This work was also supported by a Birmingham Fellowship to A.J.I. and Medical Research Council project grant MR/T028025/1.

Conflict of interest

A.J.I., H.M.M., and M.C. hold patents related to an immunomodulatory peptide. A.J.I. has received funding from F. Hoffmann-La Roche AG. All other authors have no conflicts of interest to declare.

References

- Fagiolo U, Cossarizza A, Scala E, Fanales-Belasio E, Ortolani C, Cozzi E, Monti D, Franceschi C, Paganelli R, et al. Increased cytokine production in mononuclear cells of healthy elderly people. *Eur J Immunol*. 1993;23(9):2375–2378. <https://doi.org/10.1002/eji.1830230950>
- Ershler WB, Keller ET. Age-associated increased interleukin-6 gene expression, late-life diseases, and frailty. *Annu Rev Med*. 2000;51(1):245–270. <https://doi.org/10.1146/annurev.med.51.1.245>
- Hu L, Mauro TM, Dang E, Man G, Zhang J, Lee D, Wang G, Feingold KR, Elias PM, Man M-Q, et al. Epidermal dysfunction leads to an age-associated increase in levels of Serum inflammatory cytokines. *J Invest Dermatol*. 2017;137(6):1277–1285. <https://doi.org/10.1016/j.jid.2017.01.007>
- Pereira BI, De Maeyer RPH, Covre LP, Nehar-Belaid D, Lanna A, Ward S, Marches R, Chambers ES, Gomes DCO, Riddell NE, et al. Sestrins induce natural killer function in senescent-like CD8(+) T cells. *Nat Immunol*. 2020;21(6):684–694. <https://doi.org/10.1038/s41590-020-0643-3>
- Nakajima T, Schulte S, Warrington KJ, Kopecky SL, Frye RL, Goronzy JJ, Weyand CM, et al. T-cell-mediated lysis of endothelial cells in acute coronary syndromes. *Circulation*. 2002;105(5):570–575. <https://doi.org/10.1161/hc0502.103348>
- Nicoletti C, Borghesi-Nicoletti C, Yang XH, Schulze DH, Cerny J. Repertoire diversity of antibody response to bacterial antigens in aged mice. II. Phosphorylcholine-antibody in young and aged mice differ in both VH/VL gene repertoire and in specificity. *J Immunol*. 1991;147(8):2750–2755. <https://doi.org/10.4049/jimmunol.147.8.2750>
- Mogilenko DA, Shpynov O, Andhey PS, Arthur L, Swain A, Esaulova E, Brioschi S, Shchukina I, Kerndl M, Bambouskova M, et al. Comprehensive profiling of an aging immune system reveals clonal GZMK(+) CD8(+) T cells as conserved hallmark of inflammation. *Immunity*. 2021;54(1):99–115.e12. <https://doi.org/10.1016/j.immuni.2020.11.005>
- Tabula Muris C. A single-cell transcriptomic atlas characterizes ageing tissues in the mouse. *Nature*. 2020;583(7817):590–595. <https://doi.org/10.1038/s41586-020-2496-1>
- Hopkin S, Lord JM, Chimen M. Dysregulation of leukocyte trafficking in ageing: causal factors and possible corrective therapies. *Pharmacol Res*. 2021;163:105323. <https://doi.org/10.1016/j.phrs.2020.105323>
- Lefebvre JS, Maue AC, Eaton SM, Lanthier PA, Tighe M, Haynes L. The aged microenvironment contributes to the age-related functional defects of CD4 T cells in mice. *Aging Cell*. 2012;11(5):732–740. <https://doi.org/10.1111/j.1474-9726.2012.00836.x>
- Wols HA, Johnson KM, Ippolito JA, Birjandi SZ, Su Y, Le PT, Witte PL. Migration of immature and mature B cells in the aged microenvironment. *Immunology*. 2010;129(2):278–290. <https://doi.org/10.1111/j.1365-2567.2009.03182.x>
- Ritzel RM, Craspe J, Patel AR, Verma R, Grenier JM, Chauhan A, Jellison ER, McCullough LD. Age-Associated resident memory CD8 T cells in the central nervous system are primed to potentiate inflammation after ischemic brain injury. *J Immunol*. 2016;196(8):3318–3330. <https://doi.org/10.4049/jimmunol.1502021>
- Reglero-Real N, Rolas L, Nourshargh S. Leukocyte trafficking: time to take time seriously. *Immunity*. 2019;50(2):273–275. <https://doi.org/10.1016/j.immuni.2019.01.013>
- Marquez EJ, Chung C-h, Marches R, Rossi RJ, Nehar-Belaid D, Eroglu A, Mellert DJ, Kuchel GA, Banchereau J, Ucar D, et al. Sexual-dimorphism in human immune system aging. *Nat Commun*. 2020;11(1):751. <https://doi.org/10.1038/s41467-020-14396-9>
- Kay E, Gomez-Garcia L, Woodfin A, Scotland RS, Whiteford JR. Sexual dimorphisms in leukocyte trafficking in a mouse peritonitis model. *J Leukoc Biol*. 2015;98(5):805–817. <https://doi.org/10.1189/jlb.3A1214-601RR>
- Dodd KC, Menon M. Sex bias in lymphocytes: implications for autoimmune diseases. *Front Immunol*. 2022;13:945762. <https://doi.org/10.3389/fimmu.2022.945762>
- Lee SK. Sex as an important biological variable in biomedical research. *BMB Rep*. 2018;51(4):167–173. <https://doi.org/10.5483/BMBRep.2018.51.4.034>
- Gubbels Bupp MR. Sex, the aging immune system, and chronic disease. *Cell Immunol*. 2015;294(2):102–110. <https://doi.org/10.1016/j.cellimm.2015.02.002>
- Rubtsov AV, Rubtsova K, Fischer A, Meehan RT, Gillis JZ, Kappler JW, Marrack P. Toll-like receptor 7 (TLR7)-driven accumulation of a novel CD11c(+) B-cell population is important for the development of autoimmunity. *Blood*. 2011;118(5):1305–1315. <https://doi.org/10.1182/blood-2011-01-331462>
- Strindhall J, Skog M, Ernerudh J, Bengner M, Löfgren S, Matussek A, Nilsson BO, Wikby A. The inverted CD4/CD8 ratio and associated parameters in 66-year-old individuals: the Swedish HEXA immune study. *Age (Dordr)*. 2013;35(3):985–991. <https://doi.org/10.1007/s11357-012-9400-3>
- Group DS. Age- and sex-specific prevalences of diabetes and impaired glucose regulation in 13 European cohorts. *Diabetes Care*. 2003;26(1):61–69. <https://doi.org/10.2337/diacare.26.1.61>
- Vitale C, Mendelsohn ME, Rosano GM. Gender differences in the cardiovascular effect of sex hormones. *Nat Rev Cardiol*. 2009;6(8):532–542. <https://doi.org/10.1038/nrcardio.2009.105>
- Zhou F, Yu T, Du R, Fan G, Liu Y, Liu Z, Xiang J, Wang Y, Song B, Gu X, et al. Clinical course and risk factors for mortality of adult inpatients with COVID-19 in Wuhan, China: a retrospective cohort study. *Lancet*. 2020;395(10229):1054–1062. [https://doi.org/10.1016/S0140-6736\(20\)30566-3](https://doi.org/10.1016/S0140-6736(20)30566-3)

54. Nilsson BO. Modulation of the inflammatory response by estrogens with focus on the endothelium and its interactions with leukocytes. *Inflamm Res*. 2007;56(7):269–273. <https://doi.org/10.1007/s00011-007-6198-z>
55. Goddard LM, Ton AN, Org T, Mikkola HKA, Iruela-Arispe ML. Selective suppression of endothelial cytokine production by progesterone receptor. *Vascul Pharmacol*. 2013;59(1-2):36–43. <https://doi.org/10.1016/j.vph.2013.06.001>
56. Caulin-Glaser T, Watson CA, Pardi R, Bender JR. Effects of 17beta-estradiol on cytokine-induced endothelial cell adhesion molecule expression. *J Clin Invest*. 1996;98(1):36–42. <https://doi.org/10.1172/JCI118774>
57. Mukherjee TK, Nathan L, Dinh H, Reddy ST, Chaudhuri G. 17-epiestriol, An estrogen metabolite, is more potent than estradiol in inhibiting vascular cell adhesion molecule 1 (VCAM-1) mRNA expression. *J Biol Chem*. 2003;278(14):11746–11752. <https://doi.org/10.1074/jbc.M207800200>
58. Simoncini T, Maffei S, Basta G, Barsacchi G, Genazzani AR, Liao JK, De Caterina R. Estrogens and glucocorticoids inhibit endothelial vascular cell adhesion molecule-1 expression by different transcriptional mechanisms. *Circ Res*. 2000;87(1):19–25. <https://doi.org/10.1161/01.RES.87.1.19>
59. Bonello N, Norman RJ. Soluble adhesion molecules in serum throughout the menstrual cycle. *Hum Reprod*. 2002;17(9):2272–2278. <https://doi.org/10.1093/humrep/17.9.2272>
60. Horstman AM, Dillon EL, Urban RJ, Sheffield-Moore M. The role of androgens and estrogens on healthy aging and longevity. *J Gerontol A Biol Sci Med Sci*. 2012;67(11):1140–1152. <https://doi.org/10.1093/gerona/gls068>
61. Amezcua Vesely MC, Schwartz M, Bermejo DA, Montes CL, Cautivo KM, Kalergis AM, Rawlings DJ, Acosta-Rodríguez EV, Gruppi A, et al. FcγRIIb and BAFF differentially regulate peritoneal B1 cell survival. *J Immunol*. 2012;188(10):4792–4800. <https://doi.org/10.4049/jimmunol.1102070>
62. Bruunsgaard H, Andersen-Ranberg K, Jeune B, Pedersen AN, Skinhoj P, Pedersen BK. A high plasma concentration of TNF-alpha is associated with dementia in centenarians. *J Gerontol A Biol Sci Med Sci*. 1999;54(7):M357–M364. <https://doi.org/10.1093/gerona/54.7.M357>
63. Wei J, Xu H, Davies JL, Hemmings GP. Increase of plasma IL-6 concentration with age in healthy subjects. *Life Sci*. 1992;51(25):1953–1956. [https://doi.org/10.1016/0024-3205\(92\)90112-3](https://doi.org/10.1016/0024-3205(92)90112-3)
64. Daynes RA, Araneo BA, Ershler WB, Maloney C, Li GZ, Ryu SY. Altered regulation of IL-6 production with normal aging. Possible linkage to the age-associated decline in dehydroepiandrosterone and its sulfated derivative. *J Immunol*. 1993;150(12):5219–5230. <https://doi.org/10.4049/jimmunol.150.12.5219>
65. Spaulding CC, Walford RL, Effros RB. Calorie restriction inhibits the age-related dysregulation of the cytokines TNF-alpha and IL-6 in C3B10RF1 mice. *Mech Ageing Dev*. 1997;93(1-3):87–94. [https://doi.org/10.1016/S0047-6374\(96\)01824-6](https://doi.org/10.1016/S0047-6374(96)01824-6)
66. Wang C, Qin L, Manes TD, Kirkiles-Smith NC, Tellides G, Pober JS. Rapamycin antagonizes TNF induction of VCAM-1 on endothelial cells by inhibiting mTORC2. *J Exp Med*. 2014;211(3):395–404. <https://doi.org/10.1084/jem.20131125>
67. Luu NT, Rainer GE, Nash GB. Differential ability of exogenous chemotactic agents to disrupt transendothelial migration of flowing neutrophils. *J Immunol*. 2000;164(11):5961–5969. <https://doi.org/10.4049/jimmunol.164.11.5961>
68. Chimen M, Yates CM, McGettrick HM, Ward LSC, Harrison MJ, Apta B, Dib LH, Imhof BA, Harrison P, Nash GB, et al. Monocyte subsets coregulate inflammatory responses by integrated signaling through TNF and IL-6 at the endothelial cell interface. *J Immunol*. 2017;198(7):2834–2843. <https://doi.org/10.4049/jimmunol.1601281>
69. Watson C, Whittaker S, Smith N, Vora AJ, Dumonde DC, Brown KA. IL-6 acts on endothelial cells to preferentially increase their adherence for lymphocytes. *Clin Exp Immunol*. 1996;105(1):112–119. <https://doi.org/10.1046/j.1365-2249.1996.d01-717.x>
70. Yousef H, Czupalla CJ, Lee D, Chen MB, Burke AN, Zera KA, Zandstra J, Berber E, Lehallier B, Mathur V, et al. Aged blood impairs hippocampal neural precursor activity and activates microglia via brain endothelial cell VCAM1. *Nat Med*. 2019;25(6):988–1000. <https://doi.org/10.1038/s41591-019-0440-4>
71. Morisaki N, Saito I, Tamura K, Tashiro J, Masuda M, Kanzaki T, Watanabe S, Masuda Y, Saito Y. New indices of ischemic heart disease and aging: studies on the serum levels of soluble intercellular adhesion molecule-1 (ICAM-1) and soluble vascular cell adhesion molecule-1 (VCAM-1) in patients with hypercholesterolemia and ischemic heart disease. *Atherosclerosis*. 1997;131(1):43–48. [https://doi.org/10.1016/S0021-9150\(97\)06083-8](https://doi.org/10.1016/S0021-9150(97)06083-8)
72. Richter V, Rassoul F, Purschwitz K, Hentschel B, Reuter W, Kuntze T. Circulating vascular cell adhesion molecules VCAM-1, ICAM-1, and E-selectin in dependence on aging. *Gerontology*. 2003;49(5):293–300. <https://doi.org/10.1159/000071710>
73. Miles EA, Thies F, Wallace FA, Powell JR, Hurst TL, Newsholme EA, Calder PC. Influence of age and dietary fish oil on plasma soluble adhesion molecule concentrations. *Clin Sci (Lond)*. 2001;100(1):91–100. <https://doi.org/10.1042/CS20000198>
74. Miles EA, Rees D, Banerjee T, Cazzola R, Lewis S, Wood R, Oates R, Tallant A, Cestaro B, Yaqoob P, et al. Age-related increases in circulating inflammatory markers in men are independent of BMI, blood pressure and blood lipid concentrations. *Atherosclerosis*. 2008;196(1):298–305. <https://doi.org/10.1016/j.atherosclerosis.2006.11.002>
75. Eschen O, Christensen J, Decaterina R, Schmidt E. Soluble adhesion molecules in healthy subjects: a dose-response study using n-3 fatty acids. *Nutr Metab Cardiovasc Dis*. 2004;14(4):180–185. [https://doi.org/10.1016/S0939-4753\(04\)80002-4](https://doi.org/10.1016/S0939-4753(04)80002-4)
76. Demerath E, Towne B, Blangero J, Siervogel RM. The relationship of soluble ICAM-1, VCAM-1, P-selectin and E-selectin to cardiovascular disease risk factors in healthy men and women. *Ann Hum Biol*. 2001;28(6):664–678. <https://doi.org/10.1080/03014460110048530>
77. Otero K, Turnbull IR, Poliani PL, Vermi W, Cerutti E, Aoshi T, Tassi I, Takai T, Stanley SL, Miller M, et al. Macrophage colony-stimulating factor induces the proliferation and survival of macrophages via a pathway involving DAP12 and beta-catenin. *Nat Immunol*. 2009;10(7):734–743. <https://doi.org/10.1038/ni.1744>
78. Tran TM, Temkin V, Shi B, Pagliari L, Daniel S, Ferran C, Pope RM. TNFalpha-induced macrophage death via caspase-dependent and independent pathways. *Apoptosis*. 2009;14(3):320–332. <https://doi.org/10.1007/s10495-009-0311-4>
79. Deshpande A, Elliott J, Kari N, Jiang B, Michel P, Toosizadeh N, Fahadan PT, Kidwell C, Wintermark M, Laksari K, et al. Novel imaging markers for altered cerebrovascular morphology in aging, stroke, and Alzheimer's disease. *J Neuroimaging*. 2022;32(5):956–967. <https://doi.org/10.1111/jon.13023>
80. Edelberg JM, Reed MJ. Aging and angiogenesis. *Front Biosci*. 2003;8(6):s1199–s1209. <https://doi.org/10.2741/1166>
81. Swift ME, Kleinman HK, DiPietro LA. Impaired wound repair and delayed angiogenesis in aged mice. *Lab Invest*. 1999;79(12):1479–1487.
82. Edelberg JM, Lee SH, Kaur M, Tang L, Feirt NM, McCabe S, Bramwell O, Wong SC, Hong MK, et al. Platelet-derived growth factor-AB limits the extent of myocardial infarction in a rat

- model: feasibility of restoring impaired angiogenic capacity in the aging heart. *Circulation*. 2002;105(5):608–613. <https://doi.org/10.1161/hc0502.103672>
83. Chen YM, Zhang JS, Duan XL. Changes of microvascular architecture, ultrastructure and permeability of rat jejunal villi at different ages. *World J Gastroenterol*. 2003;9(4):795–799. <https://doi.org/10.3748/wjg.v9.i4.795>
84. Huynh J, Nishimura N, Rana K, Peloquin JM, Califano JP, Montague CR, King MR, Schaffer CB, Reinhart-King CA, et al. Age-related intimal stiffening enhances endothelial permeability and leukocyte transmigration. *Sci Transl Med*. 2011;3(112):112–122. <https://doi.org/10.1126/scitranslmed.3002761>
85. Karaman S, Buschle D, Luciani P, Leroux J-C, Detmar M, Proulx ST. Decline of lymphatic vessel density and function in murine skin during aging. *Angiogenesis*. 2015;18(4):489–498. <https://doi.org/10.1007/s10456-015-9479-0>
86. Owen-Woods C, Joulia R, Barkaway A, Rolas L, Ma B, Nottebaum AF, Arkill KP, Stein M, Girbl T, Golding M, et al. Local microvascular leakage promotes trafficking of activated neutrophils to remote organs. *J Clin Invest*. 2020;130(5):2301–2318. <https://doi.org/10.1172/JCI133661>
87. Hao Y, O'Neill P, Naradikian MS, Scholz JL, Cancro MP. A B-cell subset uniquely responsive to innate stimuli accumulates in aged mice. *Blood*. 2011;118(5):1294–1304. <https://doi.org/10.1182/blood-2011-01-330530>
88. Ratliff M, Alter S, Frasca D, Blomberg BB, Riley RL. In senescence, age-associated B cells secrete TNFalpha and inhibit survival of B-cell precursors. *Aging Cell*. 2013;12(2):303–311. <https://doi.org/10.1111/ace1.12055>
89. Mouat IC, Goldberg E, Horwitz MS. Age-associated B cells in autoimmune diseases. *Cell Mol Life Sci*. 2022;79(8):402. <https://doi.org/10.1007/s00018-022-04433-9>