UNIVERSITY^{OF} BIRMINGHAM University of Birmingham Research at Birmingham

Infant death from accidental suffocation and strangulation in bed in England and Wales

Garstang, Joanna Jane; Menka, Marivjena

DOI: 10.1136/bmjpo-2023-002419

License: Creative Commons: Attribution-NonCommercial (CC BY-NC)

Document Version Publisher's PDF, also known as Version of record

Citation for published version (Harvard):

Garstang, JJ & Menka, M 2024, 'Infant death from accidental suffocation and strangulation in bed in England and Wales: rare or unrecognised events?', *BMJ Paediatrics Open*, vol. 8, no. 1, e002419. https://doi.org/10.1136/bmjpo-2023-002419

Link to publication on Research at Birmingham portal

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

•Users may freely distribute the URL that is used to identify this publication.

•Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.

•User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?) •Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

BMJ Paediatrics Open

Infant death from accidental suffocation and strangulation in bed in England and Wales: rare or unrecognised events?

Joanna Jane Garstang 💿 ,^{1,2} Marivjena Menka 💿 ¹

ABSTRACT

To cite: Garstang JJ, Menka M. Infant death from accidental suffocation and strangulation in bed in England and Wales: rare or unrecognised events? *BMJ Paediatrics Open* 2024;**8**:e002419. doi:10.1136/ bmjpo-2023-002419

Received 26 November 2023 Accepted 24 January 2024

Check for updates

© Author(s) (or their employer(s)) 2024. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

¹College of Medical and Dental Sciences, University of Birmingham, Birmingham, UK ²Children and Family Services, Birmingham Community Healthcare NHS Trust, Aston, UK

Correspondence to

Dr Joanna Jane Garstang; j. garstang@bham.ac.uk

Background Mandatory joint police and healthcare investigations of sudden unexpected death in infancy (SUDI) have been in place since 2008 in England. These include death scene examination with cause of death determined at multiprofessional case conference. Detailed evidence on sleep arrangements is available for most cases potentially leading to more being identified as due to accidental suffocation. SUDI remaining unexplained following investigation are classified as SIDS (sudden infant death syndrome) or unspecified deaths. Our objective was to determine whether detailed SUDI investigation has led to an increase in deaths classified as accidental suffocation or strangulation in bed (ASSB)? Methods We obtained official mortality data for England and Wales for infants dying aged 0-364 days for International Statistical Classification of Diseases and Related Health Problems, 10th revision codes R95 (SIDS), R96, R98, R99 (unspecified causes of mortality) and W75 (ASSB) for the years 2000-2019.

We calculated the mortality rate for ASSB, SIDS and unspecified causes based on total live births each year. Results Unexplained SUDI decreased from 353 in 2000 to 175 in 2019, with the mortality rate falling from 0.58 to 0.29 per 1000 live births. The total postneonatal mortality rate fell during this time from 1.9 to 0.9 per 1000 live births suggesting this is a genuine fall. SIDS accounted for 70% of unexplained SUDI in 2000 falling to 49% in 2020 with a corresponding increase in R99 unspecified deaths. Few deaths were recorded as ASSB (W75), ranging between 4 in 2010 and 24 in 2001. The rate for ASSB ranged from 0.6 to 4.0 per 100000 live births. **Conclusions** There is a shift away from SIDS (R95) towards unspecified causes of death (R96, R98, R99). Improved investigation of deaths has not led to increased numbers of death identified as due to ASSB. There needs to be clear guidelines on accurate classification of deaths from ASSB to facilitate learning from deaths and inform prevention efforts.

INTRODUCTION

Sudden unexpected death in infancy (SUDI) is the sudden and unexpected death of an infant which would not have been reasonably expected to occur 24 hours previously and where no pre-existing medical cause of death is apparent.¹ There are around 350 such deaths in England each year, of which

WHAT IS ALREADY KNOWN ON THIS TOPIC

- \Rightarrow Sudden infant death syndrome has declined significantly since the 1990s and continues to fall.
- \Rightarrow Since 2008, in England, all sudden infant deaths have detailed multiagency investigation.
- ⇒ Many sudden infant deaths occur in hazardous sleep circumstances involving cosleeping with parents who have consumed alcohol or drugs.

WHAT THIS STUDY ADDS

- ⇒ Detailed multiagency investigation of sudden infant deaths has not led to an increased recognition of deaths from accidental suffocation or strangulation in bed (ASSB).
- ⇒ It is likely that deaths from ASSB are being missed as rates in England and Wales are much lower than other countries with detailed sudden infant death investigation.

HOW THIS STUDY MIGHT AFFECT RESEARCH, PRACTICE OR POLICY

- \Rightarrow There needs to be clear guidance for classifying cause of death for sudden infant deaths.
- ⇒ Recognition of ASSB deaths could help inform efforts to prevent sudden infant deaths, by empowering professionals to discuss risks and mechanisms for death with parents.

approximately half have a cause subsequently identified, for example, congenital malformation or infection, and half remain unexplained.² Unexplained deaths may be classified as sudden infant death syndrome (SIDS) although the term 'unascertained' is also used frequently when pathologists have been unable to establish a cause of death. SIDS is defined as the sudden unexpected death of an infant with the onset of the lethal episode occurring during sleep, which remains unexplained after thorough investigation. This includes a complete postmortem examination, review of circumstances of death and clinical history.³

The risk factors for SIDS are well known, many relate to the sleep environment as well as exposure to tobacco smoke antenatally and postnatally.⁴ The highest risks relate to tobacco-exposed infants sharing sleep surfaces with adults who have consumed alcohol or drugs. A metaanalysis determined an adjusted OR of 151 (95% CI 50 to 448) for death of infants cosleeping in bed with parents who were tobacco smokers and had consumed alcohol or drugs in the previous 24 hours.⁵ The comparison was infants room-sharing with non-smoking, non-drug or alcohol using parents but these control infants have a very low risk of SUDI potentially exaggerating the OR. In contrast, a pooled analysis of two case-control studies determined an OR of 18.3 (95% CI 7.7 to 43.5) for infants cosleeping on a sofa with an adult who had consumed more than two units of alcohol compared with infants sleeping alone.⁶ Some of these deaths may be due to accidental suffocation. However, this is difficult to determine as postmortem examination for both SIDS and accidental suffocation have similar non-specific findings,⁷ and there are no diagnostic findings.⁸ The diagnosis of accidental suffocation relies on a careful history and death scene examination. There is no international consensus on the diagnosis of accidental suffocation in infants.

As SIDS rates have declined significantly since the 1990s, the proportion in highly hazardous sleep environments has increased.⁹ UK pathologists prefer to classify deaths with sleep environment risk factors as unascertained rather than SIDS¹⁰ and now nearly half of UK unexplained infant deaths are classified as 'unascertained'.¹¹ There is considerable international variation in the classification of cause of death following SUDI.¹² Reanalysis of cases originally determined as SIDS in New Zealand and New South Wales identified that up to 20% may have been due to accidental suffocation.^{13 14}

In the UK, the determination of the cause of death following SUDI rests with coroners, death certificates are based on the coronial process. Since 2008, in England, this should be informed by a detailed multiagency investigation, including a joint home visit by police and healthcare professionals with death scene analysis. The process concludes with a multiprofessional case conference documenting all relevant factors and an opinion as to the cause of death.¹ A similar process exists in Wales, although there are no joint home visits and police visit scenes of death alone.¹⁵ The National Child Mortality Database (NCMD) has collated information from reviews of all child deaths in England since April 2019. Joint home visits took place for 65% of SUDI between 2019 and 2021, 48% of unexplained deaths occurred in hazardous cosleeping environments involving parental drugs, alcohol, smoking, preterm or low birthweight infants.² Previous research has identified that coroners frequently rely on the conclusion of the postmortem examination alone and not the wider multiagency investigation,¹⁶ and child death overview panels may disagree with pathologists on the cause of death.¹⁷ The aim of this study is to identify if improvements in the investigation of SUDI and identification of hazardous sleep environments has resulted in any changes in the classification of causes of death.

The research question was: has detailed SUDI investigation led to an increase in infant deaths classified as accidental suffocation or strangulation in bed (ASSB)?

METHODS

We searched official mortality statistics for data on deaths of infants aged from 0 to 364 days in the years 2000–2019. These data are based on death certificates registered in England and Wales with cause of death as determined by coroners. Underlying causes of death were coded according to International Statistical Classification of Diseases and Related Health Problems, 10th revision (ICD-10).¹⁸ We included deaths certified due to the following ICD-10 codes: R95 SIDS, R96 other sudden death, cause unknown, R98 unattended death, R99 other ill-defined and unspecified causes of mortality and W75 accidental suffocation and strangulation in bed (ASSB). We have referred to these combined codes R95–98 and W75 as sleep-related SUDI.

We excluded codes W76–84 other accidental threats to breathing, as these may not have occurred in a sleep environment and W78 inhalation of gastric contents may represent an agonal process.

Data for the years 2013–2019 were obtained directly from the Office for National Statistics website using the 'Query Data' feature.¹⁹ We obtained data for deaths for 2000–2012 directly from Office for National Statistics which supplied this from their archive.

We obtained live birth numbers, neonatal and the postneonatal mortality rate for all years directly from the Office for National Statistics website.

We calculated the total sleep-related SUDI, the proportions linked to each ICD-10 code and the mortality rate for ASSB for each year.

We did not involve bereaved families directly in this project although the findings have been discussed with them.

RESULTS

The total number of sleep-related SUDI fell from 365 in 2000, to its lowest of 185 in 2019. Unexplained infant deaths (R95–99) decreased from 353 in 2000 to 175 in 2019. There were few deaths recorded in any year due to ASSB, the least was 4 in 2010 and the most was 24 in 2001.

The numbers of deaths for each year by ICD-10 code and infant mortality rates are shown in table 1. There were no deaths recorded under R98 for the entire period, so this code has been omitted. The fall in numbers of unexplained infant deaths is unlikely to be due to these deaths being classified under other ICD-10 codes as there has been an overall decline in total neonatal and postneonatal deaths over the same period. The decline in the number of sleep-related SUDI is illustrated in figure 1.

The proportion of sleep-related SUDI classified as R95 SIDS fell slightly from 68% in 2000 to 46% in 2019 with a corresponding increase in R99. SIDS accounted for 70%

<table-container>IndexIndexIndexIndexIndexIndexIndexIndexIndexIndexIndexIndexIndex11<</table-container>	Table 1 Number of sleep-related SUDI for each ICD-10 code and infant mortality rates 2000–2019									
Year R89% R89% R95%-96% R75% R75% <		Number of deaths for ICD-10 codes % of total sleep-related SUDI						Neonatal		
<table-container><table-container><table-container><table-container><table-container><table-container><table-container><table-container><table-container><table-container><table-container><table-container><table-container><table-container><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-container></table-container></table-container></table-container></table-container></table-container></table-container></table-container></table-container></table-container></table-container></table-container></table-container></table-container>	Year	R95%	R96%	R99%	R95%-99%	W75%	Total sleep-related SUDI (R95–99 and W75)	mortality rate/1000 live births	Postneonatal mortality rate/1000 live births	
<table-container><table-container><table-container><table-container><table-container><table-container><table-container><table-container><table-container><table-container><table-container><table-container><table-container><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-container></table-container></table-container></table-container></table-container></table-container></table-container></table-container></table-container></table-container></table-container></table-container></table-container>	2000	247	0	106	353	12	365	3.9	1.8	
<table-container> 245 9</table-container>		67.7%	0.0%	29.0%	96.7%	3.3%				
<table-container><table-container><table-container><table-container><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-container></table-container></table-container></table-container>	2001	245	0	102	347	24	371	3.6	1.9	
<table-container> 193 9 111 94 9 24 34 11 2604 640 0.0% 643% 620% <</table-container>		66.0%	0.0%	27.5%	93.5%	6.5%				
<table-container>initial stateinitial stateinitial stateinitial stateinitial state2001500.001.45030.0021.00<</table-container>	2002	193	0	111	304	20	324	3.6	1.7	
<table-container><table-container><table-container><table-container><table-container><table-container><table-container><table-container><table-container><table-container><table-container><table-container><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-container></table-container></table-container></table-container></table-container></table-container></table-container></table-container></table-container></table-container></table-container></table-container>		59.6%	0.0%	34.3%	93.8%	6.2%				
<table-container><table-container><table-container><table-container><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-container></table-container></table-container></table-container>	2003	185	0	145	330	21	351	3.7	1.8	
<table-container>21201931124353531.661800.0%3.7.%9.6.%3.6.5.7.%5.7.%5.7.%5.7.%5.7.%200613.4%0.3%3.7.%9.5.%4.5.%5.7.%5.7.%5.7.%5.7.%200714.4%0.3%3.7.%9.5.%4.7.%5.7.%5.7.%5.7.%5.7.%200715.4%0.3%3.7.%9.5.%4.7.%5.7.%5.7.%5.7.%5.7.%201610.40.4%2.941.43.7.%5.7.%5.7.%5.7.%5.7.%202610.50.4%9.6.%0.4%3.7.%5.7.%5.7.%5.7.%5.7.%203710.5%0.4%0.6%0.6%0.6%0.6%6.7.%5.7.%5.7.%5.7.%203810.6%0.6%0.6%0.6%0.6%0.6%6.7.%5.7.%5.7.%5.7.%203910.6%0.6%0.6%0.6%0.6%0.6%5.7.%5.7.%5.7.%5.7.%203010.6%0.6%0.6%0.6%0.6%0.6%5.7.%5.7.%5.7.%5.7.%203010.6%0.6%0.6%0.6%0.6%0.6%5.7.%5.7.%5.7.%5.7.%203010.6%0.6%0.6%0.6%0.6%0.6%5.7.%5.7.%5.7.%5.7.%5.7.%203010.6%0.6%0.6%0.6%0.6%0.6%<</table-container>		52.7%	0.0%	41.3%	94.0%	6.0%				
<table-container><table-container><table-container><table-container><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-container></table-container></table-container></table-container>	2004	212	0	119	331	12	343	3.5	1.6	
<table-container>2004 61% 61% 61% 61% 61%14144338163643.51.72006 60% 60%0.0%34.7% 94.7%55.3%4.7%1.53.51.52017 60.7%0.0%30.7%96.1%3.9%1.7%1.61.62017 60.7%0.0%30.2%96.1%3.9%1.7%3.9%1.62017 70%190.0%30.2%96.1%3.9%3.9%3.9%3.9%3.9%2018 70%190.0%36.9%90.9%4.0%3.9%<!--</td--><td>61.8%</td><td>0.0%</td><td>34.7%</td><td>96.5%</td><td>3.5%</td></table-container>		61.8%	0.0%	34.7%	96.5%	3.5%				
<table-container><table-container><table-container><table-container><table-container><table-container><table-container><table-container><table-container><table-container><table-container><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-container></table-container></table-container></table-container></table-container></table-container></table-container></table-container></table-container></table-container></table-container>	2005	224	1	114	338	16	354	3.5	1.7	
<table-container>1940111305153203.51.560.6%0.0%34.7%95.3%4.7%77%77%78%78%78%2050.0%34.7%2991271%71%78%</table-container>		63.1%	0.3%	32.1%	95.5%	4.5%				
<table-container>index 60%0%94.7%95.3%4.7%207 65.9%0.6%0.0%30.2%96.1%3.9%3.1%3.3%1.665.9%0.0%30.2%96.1%3.9%</table-container>	2006	194	0	111	305	15	320	3.5	1.5	
<table-container>2007205094299122113.31.665.9%0.0%30.2%96.1%3.9%319319319319319200819401183087313131931931920101941133087313131319313120101630.3%35.9%97.8%2.2%2131313131201016311127542931313120111720.4%39.8%96.9%1.4%2831313120121631.4%39.8%96.9%1.4%2931313120141720.4%39.8%96.9%1.4%2931313120151631.4%39.8%96.9%1.4%2931313120161740.4%39.8%96.9%1.4%2931313120171640.4%39.8%96.9%1.8%2931313120181751.8%1.8%1.8%1.8%1.2%1.2%1.2%20191840.4%39.8%96.9%4.7%252.8%1.2%1.2%20191840.6%36.9%96.9%4.7%252.8%1.2%1.2%20141840.6%36.</table-container>		60.6%	0.0%	34.7%	95.3%	4.7%				
<table-container>index 6.9%0.0%30.2%96.1%3.9%2008 52%19001830813 60%213.31.552%0.0%36.8%96.0%4.0%2.0%1.6%1.5%3.2%1.5%2010 61.6%141133087 7.8%3.2%3.2%1.5%3.2%1.5%2010 61.6%13112754 8.6%2.2%3.2%1.4%3.2%3.2%1.4%2011 61.6%0.4%38.9%96.0%1.4%2.2%3.2%3.2%1.4%2012 61.6%0.4%38.4%96.0%1.4%2.3%3.2%3.2%3.2%2014 61.6%0.6%36.3%95.1%4.9%2.7%3.2%3.2%3.2%2015 61.6%0.6%30.1%97.3%2.7%2.5%2.6%1.4%2014 70.1%130.1%92.3%1.8%3.6%1.8%3.6%<!--</td--><td rowspan="2">2007</td><td>205</td><td>0</td><td>94</td><td>299</td><td>12</td><td rowspan="2">311</td><td rowspan="2">3.3</td><td rowspan="2">1.6</td></table-container>	2007	205	0	94	299	12	311	3.3	1.6	
<table-container>1901900118308133213.31.552.9%0.0%36.8%96.0%4.0%4.0%4.0%200919411133087.313.21.561.6%0.3%35.9%97.8%2.2%4291.41.470116.6%0.3%39.8%98.6%1.4%71.41.470117097.5%4291.41.4701170.9%36.9%1.4%283.11.3701170.9%36.9%95.9%4.9%291.4701170.9%36.9%95.9%7292.91.3701170.9%36.9%97.9%2.9%1.31.37011740.9%97.3%2.7%2.91.21.370110.9%30.1%97.3%2.7%2.91.21.27011740.9%95.3%1.6%2.9%1.41.470110.9%36.3%95.3%1.6%1.41.21.27011840.9%36.9%1.6%1.6%1.41.47011840.9%95.3%4.7%1.41.47011849.6%9.6%4.7%1.41.47011849.6%1.6%1.6%1.41.47011849.6%1.6%1.6%1.4</table-container>		65.9%	0.0%	30.2%	96.1%	3.9%				
<table-container><table-container><table-container><table-container><table-container><table-container><table-container><table-container><table-container><table-container><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-container></table-container></table-container></table-container></table-container></table-container></table-container></table-container></table-container></table-container>	2008	190	0	118	308	13	321	3.3	1.5	
200919411330871532.91.561.6%0.3%35.9%97.8%2.2%2.2%2.2%3.2%3.6%3.2%20101631112754273.1%1.4%54.4%0.4%38.8%98.6%1.4%273.1%1.4%20101720.4%97.9%1.4%283.1%1.4%20111740.4%34.3%95.1%4.9%3.1%3.1%3.1%20121740.4%30.1%95.1%4.9%2.7%3.1%3.1%20131700.1%10.4%27453.1%1.2%1.2%20141300.1%10.4%2.7%2.8%1.2%1.2%2015130.1%0.0%3.3%9.5%3.7%2.6%1.2%20141240.0%3.8%9.5%4.9%3.1%3.1%20151240.0%3.8%9.6%4.5%3.1%3.1%20161240.0%3.8%3.6%3.6%3.1%3.1%20171360.0%3.2%9.6%3.6%3.6%3.6%20181360.0%3.6%3.6%3.6%3.6%3.6%20141360.0%3.6%3.6%3.6%3.6%3.6%20151430.0%3.6%3.6%3.6%3.6%3.6%20161360.0%3.6%<		59.2%	0.0%	36.8%	96.0%	4.0%				
<table-container>indexindexindexindexindexindexindexindex2010163111275427314isdim0.4%38.8%98.6%1.4%293142011172097269142831320111740.6%34.3%95.1%4.9%2831320121740.6%78252729291320131740.6%31.4%97.3%2.7%29291320141700.6%104274531220151360.6%1042745361220141360.6%37.3%98.2%1.8%291220151360.6%12.9%96.9%1.6%1220161360.6%36.4%96.9%4.7%1220161360.6%36.4%96.9%1.6%122017110.6%36.4%96.9%261.6%2018110.6%36.9%1626122018110.6%36.9%1626262019110.6%36.9%16262620191310.6%36.9%36.9%262620191310.6%36.9%36.9%262620191310.6%36.9%</table-container>	2009	194	1	113	308	7	315	3.2	1.5	
1631111275427931454.%00.4%039.8%096.6%01.4%021212114212		61.6%	0.3%	35.9%	97.8%	2.2%				
<table-container><table-container><table-container><table-container><table-container><table-container><table-container><table-row><table-container><table-row><table-row><table-row><table-container><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-container></table-row></table-row></table-row></table-container></table-row></table-container></table-container></table-container></table-container></table-container></table-container></table-container>	2010	163	1	111	275	4	279	3	1.4	
1720972691428331.360.8%0.0%34.3%95.1%4.9%4.9%1.0%1.0%1.0%20121740782527291.367.2%0.0%30.1%97.3%2.7%2.9%1.9%1.9%201317001042745292.81.260.9%0.0%37.3%98.2%1.8%2.7%2.81.2201413301092421222.61.220151240.0%38.4%95.3%4.7%2.61.220151240.0%38.4%95.8%4.2%2.61.120161180.0%38.4%95.8%4.2%2.61.120161180.0%52.2%98.4%1.6%2.61.120171110.994.9992.42.81.120181110.0%43.9%95.6%4.4%2.42.81.120181110.0%43.9%96.6%2.42.42.81.12019860.0%36.9%96.6%2.42.81.1		58.4%	0.4%	39.8%	98.6%	1.4%				
Index 608%0.0%34.3%95.1%4.9%20121740782527292.91.367.2%0.0%30.1%97.3%2.7%2.7%2.81.220131700.0%37.3%98.2%1.8%272.81.220141.3%0.0%37.3%98.2%1.8%242.9%2.61.220151.3%0.0%42.9%95.3%4.7%2.61.21.220161.40.0%36.4%95.3%4.7%2.61.21.220171.80.0%36.4%95.8%4.2%2.61.11.220181.1%0.0%52.5%96.4%1.6%2.61.11.120171.10.0%94.0%95.6%4.4%2.42.41.120181.110.0%94.0%95.6%4.4%2.42.42.61.120181.110.0%94.0%95.6%95.6%2.62.61.12019860.0%43.9%95.6%95.6%2.62.61.12019860.0%89.0%95.6%95.6%2.6%2.6%1.1	2011	172	0	97	269	14	283	3	1.3	
17407825272592.91.367.2%0.0%30.1%97.3%2.7%2.7%2.7%2.9%1.32013170010427452.9%2.8%1.2%60.9%0.0%37.3%98.2%1.8%2.7%2.6%1.2%20141330109242122.4%2.6%1.2%201512.4%0.0%42.9%95.3%4.7%2.6%1.2%201512.4%0.0%83.4%95.8%4.2%2.7%1.12016180.0%38.4%95.8%4.2%2.6%1.6%20171110.9%9.8%9.6%2.6%2.8%1.120181110.0%43.9%95.6%4.4%2.4%2.8%1.120181110.0%43.9%9.6%9.6%2.6%2.8%1.12019860.0%89.9%175101852.8%1.3%		60.8%	0.0%	34.3%	95.1%	4.9%				
Index 67.2%0.0%30.1%97.3%2.7%2013 60.9%1700.0%1042745292.81.260.9%0.0%37.3%98.2%1.8%7261.22014 52.4%0.0%42.9%95.3%4.7%261.21.22015 57.4%0.0%38.4%95.8%4.2%212.71.12016 61.4%180.0%38.4%95.8%4.2%2.61.22017 51.9%180.0%52.9%98.4%1.6%2.81.12018 61.9%1110.0%43.9%95.6%4.4%2.61.12018 61.9%1110.0%43.9%95.6%9.6%2.62.81.12019 7019660.0%89.0%1.5%9.6%2.62.81.12019 7019660.0%89.0%9.6%5.6%5.6%2.81.1	2012	174	0	78	252	7	259	2.9	1.3	
2013 0.0%1700.0%10427452792.81.260.9%0.0%37.3%98.2%1.8%71.8%1.8%2014133010924212242.61.252.4%0.0%42.9%95.3%4.7%2.61.2201512408320792162.71.157.4%0.0%38.4%95.8%4.2%2.61.1201611801332514252.81.1201711109499242.42.420181110.0%43.9%95.6%4.4%2.42.81.120181110.0%43.9%95.6%9.6%2.42.81.22019860.0%89175101852.81		67.2%	0.0%	30.1%	97.3%	2.7%				
indexindexindexindexindex2014133010924212242420141330.0%42.9%95.3%4.7%1.220151240.0%83.0%20792162.71.1201613095.3%4.2%2162.71.120171180.0%52.9%96.4%1.6%252.81.120171110.0%52.2%98.4%1.6%2042.81.120171110.0%43.9%9.6%4.4%2.81.120181110.0%9.6%9.6%2.81.12019660.0%9.6%9.6%2.81.12019860.0%89.0%1.75101852.81.2	2013	170	0	104	274	5	279	2.8	1.2	
2014 2014 52.4% 109242122542.61.252.4\%0.0%42.9%95.3%4.7%2015 57.4% 12408320792162.757.4%0.0%38.4%95.8%4.2%2016 4.3% 18013325142017 4.3% 111094992018 51.9% 111094992018 51.9% 111094992018 51.9% 1110.0%43.9%95.6%95.6%2018 51.9% 10.0%43.9%95.6%95.6%		60.9%	0.0%	37.3%	98.2%	1.8%				
idealidealidealidealidealidealideal20151240832079262.71.157.4%0.0%38.4%95.8%4.2%2.71.1201611801332514252.81.12017111094992042.81.120171110.0%43.9%95.6%4.4%2.11.120181110.0%43.9%95.6%9.6%2.81.12019860.0%43.9%95.6%9.6%2.81.1	2014	133	0	109	242	12	254	2.6	1.2	
$ \begin{array}{ c c c c c } \hline 2015 & 124 & 0 & 83 & 207 & 9 & 216 & 2.7 & 1.1 \\ \hline 57.4\% & 0.0\% & 38.4\% & 95.8\% & 4.2\% & 2.7 & 1.1 \\ \hline 57.4\% & 0.0\% & 133 & 251 & 4 & 255 & 2.8 & 1.1 \\ \hline 46.3\% & 0.0\% & 52.2\% & 98.4\% & 1.6\% & 2 & 2 & 2.8 & 1.1 \\ \hline 46.3\% & 0.0\% & 52.2\% & 98.4\% & 1.6\% & 2 & 2 & 1.1 \\ \hline 2017 & 111 & 0 & 94 & 9 & 9 & 9 & 24 & 2.8 & 1.1 \\ \hline 51.9\% & 0.0\% & 43.9\% & 95.6\% & 4.4\% & 2 & 2 & 1 & 1 \\ \hline 2018 & 111 & 0 & 94 & 9 & 9 & 9 & 24 & 2.8 & 1 & 1 \\ \hline 51.9\% & 0.0\% & 43.9\% & 95.6\% & 95.6\% & 2.8 & 1 & 1 \\ \hline 2019 & 86 & 0 & 89 & 175 & 10 & 185 & 2.8 & 1 \\ \hline \end{array}$		52.4%	0.0%	42.9%	95.3%	4.7%				
$ \begin{array}{ c c c c c c } \hline $5.4\% & $0.0\% & $38.4\% & $95.8\% & $4.2\% & $1.2\% & 1.1 \\ \hline $2016 & 1.8 & $0.0\% & $5.2\% & $98.4\% & $1.6\% & $1.6\% & $1.6\% & $1.6\% & $1.6\% & $1.6\% & $1.6\% & $1.6\% & $1.6\% & $1.6\% & 1.1 \\ \hline $1.11 & $0.0\% & $94.0\% & $9.6\% & $1.6\% & $1.6\% & $1.6\% & 1.1 \\ \hline $1.9\% & $0.0\% & $43.9\% & $9.6\% & $4.4\% & $1.6\% & 1.1 \\ \hline $1.9\% & $0.0\% & $43.9\% & $9.6\% & $4.4\% & $1.6\% & 1	2015	124	0	83	207	9	216	2.7	1.1	
$ \begin{array}{c c c c c c c c c c c c c c c c c c c $		57.4%	0.0%	38.4%	95.8%	4.2%				
46.3% 0.0% 52.2% 98.4% 1.6% 2017 111 0 94 9 9 204 2.8 1.1 51.9% 0.0% 43.9% 95.6% 4.4% - <td< td=""><td rowspan="2">2016</td><td>118</td><td>0</td><td>133</td><td>251</td><td>4</td><td rowspan="2">255</td><td rowspan="2">2.8</td><td rowspan="2">1.1</td></td<>	2016	118	0	133	251	4	255	2.8	1.1	
$ \begin{array}{c ccccccccccccccccccccccccccccccccccc$		46.3%	0.0%	52.2%	98.4%	1.6%				
51.9% 0.0% 43.9% 95.6% 4.4% 2018 111 0 94 9 9 204 2.8 1 51.9% 0.0% 43.9% 95.6% 95.6% 2.8 1 2019 86 0 89 175 10 185 2.8 1	2017	111	0	94	9	9	204	2.8	1.1	
2018111094992042.8151.9%0.0%43.9%95.6%95.6%95.6%2.81201986089175101852.81		51.9%	0.0%	43.9%	95.6%	4.4%				
51.9% 0.0% 43.9% 95.6% 2019 86 0 89 175 10 185 2.8 1	2018	111	0	94	9	9	204	2.8	1	
2019 86 0 89 175 10 185 2.8 1		51.9%	0.0%	43.9%	95.6%	95.6%				
	2019	86	0	89	175	10	185	2.8	1	
46.5% 0.0% 48.1% 94.6% 5.4%		46.5%	0.0%	48.1%	94.6%	5.4%				

R95=SIDS.

R96=other sudden death, cause unknown.

R99=other ill-defined and unspecified causes of mortality. W75=ASSB.

ASSB, accidental suffocation and strangulation in bed; ICD-10, International Statistical Classification of Diseases and Related Health Problems, 10th revision; SIDS, sudden infant death syndrome; SUDI, sudden unexpected death in infancy.

of unexplained infant deaths in 2000 falling to 49% in 2020. The proportion of sleep-related SUDI classified as W75 ASSB fluctuated between 1.4% and 6.0% over time. This is shown in figure 2.

The total unexplained infant mortality rate for R95–99 combined fell from 0.58 per 1000 live births in 2000 to 0.29 in 2019. There was no increase in W75 deaths in this time, with W75 deaths ranging from 0.6



Figure 1 Decline in sleep-related SUDI and infant mortality rates 2000–2019. SUDI, sudden unexpected death in infancy.

per 100 000 live births in 2016 to 4.0 in 2001. This is shown in figure 3.

DISCUSSION

Overall, sleep-related SUDI has fallen by over 40% from 365 to 185 deaths per year from 2000 to 2019, this is not just a diagnostic shift as evidenced by the accompanying decline in postneonatal mortality rate. The decrease in sleep-related SUDI is likely due to the ongoing safe-sleep campaigns and reduction in parental smoking.²⁰ The fall is largely in cases classified as SIDS, there were 247 SIDS cases in 2000 and 86 in 2019, an overall reduction of over 65%. The number of sleep-related SUDI classified as 'unascertained' has fluctuated over time ranging between 78 and 145 deaths per year. There has been no increase in the number of deaths classified as accidental suffocation and strangulation in bed (ASSB) despite statutory multiagency SUDI investigation.

This study included all sleep-related SUDI in England and Wales over a 20-year period, a total of 5793 cases. Postmortem examination has been mandatory for SUDI throughout this period, with all postmortems conducted by specialist paediatric pathologists required by law



Figure 2 Proportion of sleep-related SUDI by ICD-10 codes 2000–2019. ICD-10, International Statistical Classification of Diseases and Related Health Problems, 10th revision; SUDI, sudden unexpected death in infancy.



Figure 3 Unexplained infant mortality rate (R95–99), W75 rate and postneonatal mortality rate 2000–2019.

from 2008 although many had paediatric examinations before then.²¹ Similarly, multiagency investigation has improved considerably since 2008, with 65% of SUDI now having joint home visits.² Given the detailed investigation of SUDI, we can be reasonably assured that this study only included deaths that remained unexplained or due to ASSB. Deaths from medical or criminal causes are unlikely to have been misreported as SIDS, unascertained or ASSB; however, this remains a possibility given the limits of current knowledge and expertise particularly as SIDS is a diagnosis based on excluding other causes for death. The inclusion of sleep-related SUDI from birth will have included unexplained infant deaths that may occur shortly after birth.

Our findings are broadly similar to those reported across the European Union, where rates of ASSB range from 0 to 5.5 per 100 000 live births for the period 2005–2015, with the UK reported as 3.0 per 100000.²² However, this combined all deaths from ICD-10 codes W75-84, many of which will not have occurred in a sleep environment such as deaths of infants from choking on food. The low rates of W75-84 could reflect that much of the European Union does not have robust SUDI investigation, in keeping with best international practice.²³ France requires all SUDI cases to be referred to specialist regional hospitals for investigation, although provision of death scene investigation is variable.²⁴ In Norway, providing there are no criminal concerns, parents are offered voluntary death scene examination by forensic investigators.²⁵ In the Netherlands, a statutory process for detailed review of child deaths is yet to be established.²⁶ It is likely in Europe that cases of ASSB may not be identified as this diagnosis relies on circumstantial rather than pathological findings.

Although England has robust data collection on child deaths in the NCMD, this has not yet reliably reported infant deaths from ASSB. A recent NCMD analysis of SUDI and childhood did not record any deaths from ASSB.² The NCMD trauma report recorded 13 infant deaths from suffocation and strangulation but this included non-sleep-related incidents such as strangulation by blind cords.²⁷ Data from Welsh Child Death

Reviews do not show ASSB deaths separately from other causes. $^{\ensuremath{^{28}}}$

The low rates of ASSB found in this study are in sharp contrast to New Zealand and the USA, where there has been considerable investment in SUDI investigation including death scene analysis. Analysis of death certification data in the USA showed ASSB rates increased from 3.4 per 100 000 live births in 1993 to 23.0 in 2015, while overall sleep-related SUDI decreased from 154.6 to 92.4 per 100 000 live births.²⁹ This could well be an overestimation of ASSB due to potential conflation of risk factors for SIDS, such as hazardous cosleeping environments, as cause of death. In comparison, an analysis of data from US Child Death Review teams, using detailed case information determined that 899/4929 (18%) of sleep-related SUDI were due to ASSB (17.7 per 100000 live births).³⁰ This may still be an overestimate due to potential overinterpretation of risk factors such as softbedding as causal for ASSB. SIDS infants are commonly found face down in bedding despite being fully able to turn their head to the side, this is hypothesised to represent a failure of autoresuscitation in response to hypoxia or hypercarbia.³¹ An infant dying face-down in soft bedding may well be SIDS and not ASSB. Accurate data for ASSB may come from the New Zealand SUDI case-control study where all cases had detailed death scene analysis conducted by health trained investigators, paediatric postmortem examination and multiprofessional case conferences. They reported 20/137 (14.5%) SUDI cases were attributed to ASSB.³² Given the much higher rates of ASSB recognised in countries with detailed death scene analysis, it is probable that in England and Wales, we are not identifying such deaths and labelling them as unexplained instead.

Correctly recognising deaths from ASSB, while avoiding overdiagnosis is important. The San Diego SIDS definition³ is open to (mis)interpretation as category II SIDS includes deaths with 'mechanical asphyxia or suffocation ... not determined with certainty', particularly as it is not possible to prove asphyxia or suffocation at postmortem examination.⁷ The triple risk model hypothesises that SIDS occurs when an intrinsically vulnerable infant is exposed to an external stressor during a critical period of development.³³ Infants dying from ASSB do not need to be intrinsically vulnerable, the hazardous situation alone leads to death; these infants are, therefore, not the same as SIDS infants, so research combining both ASSB and SIDS deaths could be misleading. Inconsistent classification of infant deaths is likely to hamper research efforts and interpretation of risk factors³⁴ and prevent meaningful comparison between countries.³⁵ One aim of the original SIDS diagnosis was to assist research by enabling similar unexplained infant deaths to be studied. International consensus on standardising the diagnosis of ASSB could also be beneficial, by avoiding overdiagnosis but empowering accurate recognition based

on death scene examination and detailed caregiver accounts. Death scene examination should include photography as this enables the scene findings to be accurately recorded.³⁶ Although the use of dolls for re-enactment is often recommended,³⁶ these should be used with some caution as there is considerable potential for distress.³⁷ Greater diagnostic accuracy would enable further research efforts into both SIDS and ASSB.

Rates of unexpected infant deaths fell significantly from the 1990s with the safe sleep campaigns but have now plateaued in many countries,^{22 29} and deaths now occur increasingly in vulnerable socially deprived infants.²⁰ An in-depth review of English SUDI cases with serious child safeguarding concerns identified hazardous cosleeping in 38/40 with several involving parental drug or alcohol misuse and mental health problems.³⁸ The review concluded that highrisk families, need tailored information and support to implement safe sleep practices. It is also important to discuss the potential risks of accidental suffocation in some sleep environments. It can be easier for parents to understand risks of accidental suffocation, with clearly describable mechanisms such as airway obstruction or chest compression, than the complex interplay of intrinsic vulnerability and external stressors in SIDS. However, it is difficult for professionals to have these discussions with families if we cannot acknowledge the frequency with which deaths from ASSB take place. We need to empower professionals to have meaningful, supportive conversations with families, and accurate knowledge and understanding of risks, causes and mechanisms for infant deaths is the basis of these discussions. We also should recognise that despite robust investigation some infant deaths remain unexplained even in unsafe sleeping environments, and overdiagnosing ASSB is not helpful and could potentially increase parents distress given that self-blame is a common feature of parental grief.³⁹

We need to remove barriers to identifying deaths from ASSB, as under-recognition hinders research and prevention efforts. The standard of SUDI investigation in England has improved over the last few years.² The multiagency guidance for investigating SUDI in England¹ is about to be updated, as part of this the conclusions of multiagency SUDI investigations should be better integrated with coroners inquests to enable accurate death certification. We need to develop practical standards for diagnosing ASSB, based on death scene examination and carers accounts. This would probably lead to a diagnostic shift away from unexplained infant deaths, so it would be important to include ASSB in future monitoring of SUDI statistics. Clear, national guidance on classification of SUDI will promote effective research, inform families and help prevent future deaths.

Open access

Twitter Joanna Jane Garstang @jogarstang

Contributors JJG conceived the idea and drafted the final paper. MM collated and analysed data and contributed to writing the paper. Both authors agreed the final draft and submission for publication. JJG is guarantor.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests All authors have completed the ICMJE uniform disclosure form at http://www.icmje.org/disclosure-of-interest/ and declare: no support from any organisation for the submitted work; JGG has received research grants from NIHR for research into improving parental involvement in child death review (NIHR 203045), she is paid by the National Child Mortality Database as a specialist clinical advisor. JGG has been paid for developing and delivering educational presentations relating to Child Death for the Primary Care Conference, Stirling Events. JGG acts as expert witness for legal cases related to sudden child death, JGG has received funding from the Lullaby Trust to attend scientific meetings overseas; JGG is Chair of the Association of Child Death Review Professionals (UK), medical advisor to SUDC-UK (Sudden Unexplained Death in Childhood) and a member of the Lullaby Trust scientific committee no other relationships or activities that could appear to have influenced the submitted work. MM has no conflicts of interest to declare.

Patient and public involvement Patients and/or the public were not involved in the design, or conduct, or reporting, or dissemination plans of this research.

Ethics approval Ethical approval was not required for this project as it only used publicly available data from the Office of National Statistics (England and Wales).

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement Data are available on reasonable request. Original data are publicly available from Office for National Statistics or from the corresponding author on request.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: http://creativecommons.org/licenses/by-nc/4.0/.

ORCID iDs

Joanna Jane Garstang http://orcid.org/0000-0001-9268-0581 Marivjena Menka http://orcid.org/0000-0002-1966-2333

REFERENCES

- Royal College of Pathologists, Royal College of Paediatrics and Child Health. Sudden unexpected death in infancy and childhood. multi-agency guidelines for care and investigation. London; 2016.
- 2 National Child Mortality Database. Sudden and unexpected deaths in infancy and childhood, national child mortality database programme. Thematic Report. University of Bristol; 2022.
- 3 Krous HF, Beckwith JB, Byard RW, et al. Sudden infant death syndrome and unclassified sudden infant deaths: a definitional and diagnostic approach. *Pediatrics* 2004;114:234–8.
- 4 Horne RSC, Hauck FR, Moon RY. Sudden infant death syndrome and advice for safe sleeping. *BMJ* 2015;350:h1989.
- 5 Carpenter R, McGarvey C, Mitchell EA, et al. Bed sharing when parents do not smoke: is there a risk of SIDS? An individual level analysis of five major case-control studies. *BMJ Open* 2013;3:e002299.
- 6 Blair PS, Sidebotham P, Pease A, et al. Bed-sharing in the absence of hazardous circumstances: is there a risk of sudden infant death syndrome? An analysis from two case-control studies conducted in the UK. PLoS One 2014;9:e107799.
- 7 Mitchell E, Krous HF, Byard RW. Pathological findings in overlaying. *J Clin Forensic Med* 2002;9:133–5.
- 8 Becroft DMO, Thompson JMD, Mitchell EA. Nasal and intrapulmonary haemorrhage in sudden infant death syndrome. Arch Dis Child 2001;85:116–20.
- 9 Blair PS, Sidebotham P, Evason-Coombe C, et al. Hazardous cosleeping environments and risk factors amenable to change: casecontrol study of SIDS in south west England. *BMJ* 2009;339:b3666.
- 10 Gould SJ, Weber MA, Sebire NJ. Variation and uncertainties in the classification of sudden unexpected infant deaths among paediatric

pathologists in the UK: findings of a National Delphi Study. *J Clin Pathol* 2010;63:796–9.

- 11 Office for National Statistics. Main tables: unexplained deaths in infancy; 2022. England and Wales
- 12 Taylor BJ, Garstang J, Engelberts A, et al. International comparison of sudden unexpected death in infancy rates using a newly proposed set of cause-of-death codes. Arch Dis Child 2015;100:1018–23.
- 13 Hayman RM, McDonald G, Baker NJ de C, et al. Infant suffocation in place of sleep: New Zealand national data 2002-2009. Arch Dis Child 2015;100:610–4.
- 14 Shipstone RA, Young J, Thompson JMD, et al. An evaluation of pathologists' application of the diagnostic criteria from the San Diego definition of SIDS and unclassified sudden infant death. Int J Legal Med 2020;134:1015–21.
- 15 Public Health Wales. Procedural response to unexpected deaths in childhood. 2010.
- 16 Garstang J, Ellis C, Griffiths F, et al. Unintentional asphyxia, SIDS, and medically explained deaths: a descriptive study of outcomes of child death review (CDR) investigations following sudden unexpected death in infancy. *Forensic Sci Med Pathol* 2016;12:407–15.
- 17 Garstang J, Cohen M, Mitchell EA, et al. Classification of sleeprelated sudden unexpected death in infancy: A national survey. Acta Paediatr 2021;110:869–74.
- 18 World Health Organization. International statistical classification of diseases and related health problems 10th. 2019 Available: https:// icd.who.int/browse10/2019/en
- 19 Office for National Statistics. Nomis offical census and labour market Statistics. 2023. Available: https://www.nomisweb.co.uk/
- 20 Pease A, Turner N, Ingram J, *et al.* Changes in background characteristics and risk factors among SIDS infants in England: cohort comparisons from 1993 to 2020. *BMJ Open* 2023;13:e076751e076751.
- 21 Royal college of pathologists. Sudden unexpected death in infancy; a multi-agency protocol for care and investigation. Royal college of pathologists, royal college of paediatrics and child health, 2004.
- 22 de Visme S, Chalumeau M, Levieux K, et al. National variations in recent trends of sudden unexpected infant death rate in western Europe. J Pediatr 2020;226:179–85.
- 23 Garstang J, Ellis C, Sidebotham P. An evidence-based guide to the investigation of sudden unexpected death in infancy. *Forensic Sci Med Pathol* 2015;11:345–57.
- 24 Levieux K, Patural H, Harrewijn I, *et al.* The French prospective multisite registry on sudden unexpected infant death (OMIN): rationale and study protocol. *BMJ Open* 2018;8:e020883.
- 25 Boylestadt L. Trends in sudden death in infants and small children in Norway. ISPID annual scientific meeting Soria Moria, Oslo,
- 26 Gijzen S, L'Hoir MP, Boere-Boonekamp MM, et al. Stakeholders' opinions on the implementation of Child Death Review in the Netherlands. *BMC Res Notes* 2016;9:228.
- 27 National Child Mortality Database. Deaths of children and young people due to traumatic incidents. University of Bristol, 2022.
- 28 Public Health Wales. Patterns and trends of child deaths in Wales, 2011-2020. Child Death Review Programme April 2022.
- 29 Erck Lambert AB, Parks SE, Shapiro-Mendoza CK. National and State Trends in Sudden Unexpected Infant Death: 1990-2015. *Pediatrics* 2018;141:e20173519.
- 30 Parks SE, Erck Lambert AB, Hauck FR, et al. Explaining sudden unexpected infant deaths, 2011-2017. *Pediatrics* 2021;147:e2020035873.
- Garcia AJ III, Koschnitzky JE, Ramirez J-M. The physiological determinants of Sudden Infant Death Syndrome. *Respiratory Physiology & Neurobiology* 2013;189:288–300.
 Mitchell EA, Thompson JMD, Zuccollo J, *et al*. The combination
- 32 Mitchell EA, Thompson JMD, Zuccollo J, et al. The combination of bed sharing and maternal smoking leads to a greatly increased risk of sudden unexpected death in infancy: the New Zealand SUDI Nationwide Case Control Study. *The New Zealand Medical Journal* 2017;130:52–64.
- 33 Filiano JJ, Kinney HC. A perspective on neuropathologic findings in victims of the sudden infant death syndrome: the triple-risk model. *Biol Neonate* 1994;65:194–7.
- 34 Byard RW, Shipstone RA, Young J. Continuing major inconsistencies in the classification of unexpected infant deaths. *J Forensic Leg Med* 2019;64:20–2.
- 35 Goldstein RD, Blair PS, Sens MA, et al. Inconsistent classification of unexplained sudden deaths in infants and children hinders surveillance, prevention and research: recommendations from The 3rd International Congress on Sudden Infant and Child Death. Forensic Sci Med Pathol 2019;15:622–8.

6

- 36 Bundock EA, Corey TS, Andrew TA, et al. Unexplained Pediatric Deaths: Investigation, Certification, and Family. Needs: Academic Forensic Pathology International, 2019.
- 37 Kristensen P. How do newly bereaved parents experience a doll reenactment shortly after the sudden, unexpected death of their child? A mixed-methods study. ISPID international conference on Stillbirth, SIDS and SUDI Florence,
- 38 Child safeguarding practice review panel. Out of routine: A review of sudden unexpected death in infancy (SUDI) in families where the children are considered at risk of Signifcant harm; 2020Jul.
- 3 Gold KJ, Sen A, Leon I. Whose fault is it anyway? guilt, blame, and death attribution by mothers after stillbirth or infant death. *Illness, Crisis & Loss* 2018;26:40–57.