# **UNIVERSITY** OF BIRMINGHAM University of Birmingham Research at Birmingham

### Specialist healthcare services for concussion/mild traumatic brain injury in England

Karvandi, Elika; Helmy, Adel; Kolias, Angelos G; Belli, Antonio; Ganau, Mario; Gomes, Clint; Grey, Michael; Griffiths, Michael; Griffiths, Timothy; Griffiths, Philippa; Holliman, Damian; Jenkins, Peter; Jones, Ben; Lawrence, Tim; McLoughlin, Terence; McMahon, Catherine; Messahel, Shrouk; Newton, Joanne; Noad, Rupert; Raymont, Vanessa

DOI: 10.1136/bmjopen-2023-077022

License: Creative Commons: Attribution (CC BY)

Document Version Publisher's PDF, also known as Version of record

#### Citation for published version (Harvard):

Karvandi, E, Helmy, A, Kolias, AG, Belli, A, Ganau, M, Gomes, C, Grey, M, Griffiths, M, Griffiths, T, Griffiths, P, Holliman, D, Jenkins, P, Jones, B, Lawrence, T, McLoughlin, T, McMahon, C, Messahel, S, Newton, J, Noad, R, Raymont, V, Sharma, K, Sylvester, R, Tadmor, D, Whitfield, P, Wilson, M, Woodberry, E, Parker, M & Hutchinson, PJ 2023, 'Specialist healthcare services for concussion/mild traumatic brain injury in England: a consensus statement using modified Delphi methodology', *BMJ open*, vol. 13, no. 12, e077022. https://doi.org/10.1136/bmjopen-2023-077022

Link to publication on Research at Birmingham portal

#### **General rights**

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

•Users may freely distribute the URL that is used to identify this publication.

•Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.

•User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?) Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

#### Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

## **BMJ Open** Specialist healthcare services for concussion/mild traumatic brain injury in England: a consensus statement using modified Delphi methodology

Elika Karvandi <sup>(i)</sup>, <sup>1</sup> Adel Helmy, <sup>1</sup> Angelos G Kolias <sup>(i)</sup>, <sup>1</sup> Antonio Belli, <sup>2</sup> Mario Ganau, <sup>3</sup> Clint Gomes, <sup>4,5</sup> Michael Grey, <sup>6</sup> Michael Griffiths, <sup>7,8</sup> Timothy Griffiths, <sup>9,10</sup> Philippa Griffiths, <sup>11</sup> Damian Holliman, <sup>12</sup> Peter Jenkins, <sup>13,14</sup> Ben Jones <sup>(i)</sup>, <sup>15,16</sup> Tim Lawrence, <sup>3</sup> Terence McLoughlin, <sup>4</sup> Catherine McMahon, <sup>17</sup> Shrouk Messahel <sup>(i)</sup>, <sup>18</sup> Joanne Newton, <sup>19</sup> Rupert Noad, <sup>20</sup> Vanessa Raymont <sup>(i)</sup>, <sup>21</sup> Kanchan Sharma, <sup>22</sup> Richard Sylvester, <sup>23,24</sup> Daniel Tadmor, <sup>25,26</sup> Peter Whitfield, <sup>20</sup> Mark Wilson, <sup>14,27</sup> Emma Woodberry, <sup>28</sup> Michael Parker, <sup>29</sup> Peter John Hutchinson <sup>(i)</sup>

#### ABSTRACT

**To cite:** Karvandi E, Helmy A, Kolias AG, *et al.* Specialist healthcare services for concussion/mild traumatic brain injury in England: a consensus statement using modified Delphi methodology. *BMJ Open* 2023;**13**:e077022. doi:10.1136/ bmjopen-2023-077022

Prepublication history and additional supplemental material for this paper are available online. To view these files, please visit the journal online (http://dx.doi.org/10.1136/ bmjopen-2023-077022).

Received 23 June 2023 Accepted 14 November 2023

#### Check for updates

© Author(s) (or their employer(s)) 2023. Re-use permitted under CC BY. Published by BMJ.

For numbered affiliations see end of article.

Correspondence to Elika Karvandi; eek34@cam.ac.uk **Objective** To establish a consensus on the structure and process of healthcare services for patients with concussion in England to facilitate better healthcare quality and patient outcome.

**Design** This consensus study followed the modified Delphi methodology with five phases: participant identification, item development, two rounds of voting and a meeting to finalise the consensus statements. The predefined threshold for agreement was set at  $\geq$ 70%. **Setting** Specialist outpatient services.

**Participants** Members of the UK Head Injury Network were invited to participate. The network consists of clinical specialists in head injury practising in emergency medicine, neurology, neuropsychology, neurosurgery, paediatric medicine, rehabilitation medicine and sports and exercise medicine in England.

**Primary outcome measure** A consensus statement on the structure and process of specialist outpatient care for patients with concussion in England.

**Results** 55 items were voted on in the first round. 29 items were removed following the first voting round and 3 items were removed following the second voting round. Items were modified where appropriate. A final 18 statements reached consensus covering 3 main topics in specialist healthcare services for concussion; care pathway to structured follow-up, prognosis and measures of recovery, and provision of outpatient clinics. **Conclusions** This work presents statements on how the healthcare services for patients with concussion in

England could be redesigned to meet their health needs. Future work will seek to implement these into the clinical pathway.

#### INTRODUCTION

In England and Wales alone, more than a million people suffer a head injury every year.<sup>1</sup> These injuries occur as a result of direct or indirect impacts to the head and

#### STRENGTHS AND LIMITATIONS OF THIS STUDY

- ⇒ This consensus statement outlines structural and process-based factors needed to move towards standardised best-practice care for concussions in England to improve healthcare quality and patient outcomes.
- $\Rightarrow$  To our knowledge, no studies have reported on the healthcare system for concussion in England.
- ⇒ We engaged with various stakeholders involved in concussion care to ensure different perspectives are represented.
- ⇒ It cannot be excluded that some disagreement on statements was in part due to differences in definitions and terminology use despite probable broad agreement on the statement in principle.
- ⇒ The statements reached agreement in this panel, but we cannot exclude the possibility or to what degree the results are influenced by the experience and expertise of these individuals both for included and excluded statements.

are commonly encountered in sports, both at professional and grassroots levels. Injuries to the head, particularly in the context of sport, are often labelled 'concussions'. While there is a recently published definition available,<sup>2</sup> there is no widespread consensus for what constitutes a concussion, and it is often used to describe both the symptoms suffered following a head injury as well as the pathophysiological mechanisms causing these symptoms.<sup>3 4</sup> Despite some controversies in the field, concussion is widely recognised by the public and those playing sport, and for this reason, we have used the term for the purpose of this paper. A concussion is diagnosed on clinical grounds with evidence of immediate and transient neurological dysfunction that may or may not present with a wide range of signs and symptoms, such as loss of consciousness, post-traumatic amnesia, headache and vomiting.<sup>5</sup> However, most concussions do not present with obvious neurological signs, such as loss of consciousness, making recognition and diagnosis challenging.<sup>6–9</sup>

The post-concussion sequelae are assumed to recover spontaneously within a few days to a couple of weeks. Yet a proportion of individuals experience persistent symptoms for months to years after injury that requires timely and appropriate medical assessment and treatment.<sup>10</sup> Despite this known fact, guidance on clinical practice in the outpatient setting after the initial medial review is limited and ambiguous leaving the concussion population vulnerable to variations and suboptimal care.<sup>11</sup> This results in delayed intervention, resource waste, care inequalities and poorer quality of care.<sup>12–14</sup> There is no standardised, structured follow-up for those who experience persistent symptoms and current clinical services largely fail to meet the healthcare needs of these patients.<sup>15</sup>

Nonetheless, the past two decades have seen an exponential growth in concussion research which has improved our understanding of the injury and strategies to enhance recovery and quality of care.<sup>16</sup> Additionally, there is increasingly more awareness around the potential lasting effects of concussions, especially in sports, where there are growing concerns and calls for action.<sup>17</sup> The government and sports' national governing bodies are attempting to improve concussion management in all levels of sports by standardising protocols, allocating more funds for concussion research and incorporating new playing rules to reduce concussion risk and improve management.<sup>18</sup> Consequently, the demand for care is evolving and to meet patient needs changes to clinical services are required to provide evidence-based best-practice care while being cost-effective, delivering better healthcare value and quality and improving health outcome for patients with concussion.<sup>7 19</sup> This necessitates fundamental changes to the healthcare system in concussion care to facilitate the delivery of best-practice outpatient care and improve quality of care.<sup>20</sup> This consensus statement addresses the healthcare structure and processes in outpatient concussion care required to facilitate the delivery of best-practice care in specialist outpatient clinics to meet healthcare needs and improve patient outcome in the concussion population.

#### **METHODS**

The guidance on Conducting and REporting DElphi Studies was followed for reporting this consensus study.<sup>21</sup>

The work was undertaken as part of wider work by the UK Head Injury Network. There was no direct involvement with patients or the public. The study did not meet the criteria for requiring ethical approval.

#### Study design

A modified Delphi approach was used for this study. We followed a five-stage process: (1) panel identification, (2) literature search for initial statement development, (3, 4) two voting rounds followed by item removal and modifications and (5) consensus meeting and live voting to finalise the statements. The meeting was chaired by a working group member with previous experience in the Delphi consensus method.

#### **Panel selection**

The UK Head Injury Network is a panel of clinical and academic experts with extensive experience in head injury care in England, and setup to improve access to and quality of concussion outpatient services-for sportsrelated and non-sports-related concussions-and provide a network to foster research collaborations. The network consists of a multidisciplinary panel from academia and clinical medicine with specialists in emergency medicine, neurology, neuropsychology, neurosurgery, neuroscience, paediatric medicine, rehabilitation medicine and sports and exercise medicine. Members are invited to join the UK Head Injury Network if they meet the following criteria based on peer-identification (all members in the network were identified or recommended by existing members): (1) healthcare professionals with extensive first-hand clinical experience in concussions care in any care setting or specialty; and/or (2) academic expertise in concussions. The group work with Love of the Game, a national sports concussion charity, to achieve these goals.

All members of the UK Head Injury Network at the time of the consensus process were invited to vote and attend the final consensus meeting. The participating members were geographically diverse including representatives from a range of healthcare settings across England. In addition to the various specialties involved, this ensures a more representative and informed decision-making for the different stages and settings of care across England.

#### **Evidence review**

The need for a consensus statement had been discussed in previous meetings and highlighted in work completed by the group (unpublished observations). As the field of concussion has several contentious issues, we took a wide approach to topics covered in the initial statement development.

As the literature on implementation science in concussion is scarce, the literature search provided the evidence for clinical tasks that could be implemented. The panel used this literature combined with their first-hand local experience in various clinical settings to provide guidance on the structures and processes required to feasibly deliver these clinical tasks in England. Thus, statement development was guided by a literature search, topics of discussion within the network and results from previous work. Medline and Embase were searched for relevant literature in October 2022, primarily published national and international clinical guidelines and consensus statements (see online supplemental material 1 for search strategy). The search was limited to literature published in English.

A subgroup of network members was responsible for this stage of the process (EK, PJH, AH). The initial statements were circulated to all network members for comments on the scope, structure and wording prior to starting the official rounds of voting.

#### **Consensus process**

Two rounds of online surveys were administered between October and December 2022 followed by a virtual meeting in February 2023. The surveys were distributed for voting using the Qualtrics XM platform (Qualtrics, Provo, Utah).

All voting was anonymous to ensure uninhibited voting. Participants were asked to vote on each statement and had the binary options agree–disagree. The cut-off threshold for agreement was set at  $\geq$ 70% for the two rounds of voting.<sup>22–24</sup> A statement was only included if agreement was above the predefined threshold for consensus. Free text comments were incorporated to allow members to suggest modifications or refinements of the statements. Statements that did not meet the threshold were excluded after each round of voting and statements were modified for further considerations if suggestions were made by the participants.

All participants were invited to a virtual meeting (Zoom Video Communications, San Jose, USA). The meeting was scheduled for 3 hours to provide sufficient time to discuss each statement in detail. The purpose of the meeting was to finalise the statements. Statements reaching agreement and disagreement were discussed where alternatives were considered for the latter kind. The threshold for agreement was raised to  $\geq 85\%$  for this meeting to ensure a robust degree of consensus. The final consensus statements were discussed and, where necessary, modifications were made to ensure clarity and accuracy before a final anonymous vote took place during the meeting.

#### Patient and public involvement

No patient involvement.

#### RESULTS

#### **Evidence review**

While several papers were identified on therapeutic interventions, educational interventions and biomarkers for diagnosis and prognosis in concussion, as expected the literature on healthcare processes and structures in concussion is scarce. Thus, the final statements were primarily guided by previous discussions, interpretation of how research evidence on best-practice could be formulated into the structure and process of the healthcare system, and research work by the network and patient focus group sessions (unpublished observation).

Key papers identified during the literature review included clinical practical guidelines on concussion management<sup>25–28</sup> and systematic reviews on therapeutic interventions,<sup>29–34</sup> prognosis,<sup>35–44</sup> recovery<sup>45–46</sup> and biomarkers.<sup>47–48</sup> The clinical practice guidelines included a synthesis of practical guidelines primarily focused on four published papers: Center for Disease Control and Prevention Guideline on the Diagnosis and Management of Mild Traumatic Brain Injury Among Children (2019),<sup>49</sup> Consensus Statement on Concussion in Sport—the fifth International Conference on Concussion in Sport (2016), Ontario Neurotrauma Foundation Guideline for Concussion/Mild Traumatic Brain Injury & Persistent Symptoms (third edition; 2018), and Department of Veterans Affairs/Department of Defense Clinical Practice Guideline for the Management of Concussion-Mild Traumatic Brain Injury (2016).<sup>26</sup>

While it was agreed that covering a wide range of topics was appropriate during the initial statement development, the statements in this consensus report were narrowed down to specifically focus on the delivery of healthcare provisions through processes and structures in concussion care. This was deemed to be the most appropriate and relevant due to (1) some disagreement on clinical practice and management issues in concussion among the panel, (2) determining that these statements were beyond the purpose of this group and some of this guidance is featured in the most recent international Consensus Statement on Concussion in Sport (sixth edition),<sup>50</sup> and (3) recognition that the need for the healthcare service to be improved at a system level prior to changes in day-to-day clinical practice. These points became clear after the first round of voting and the majority of these statement were removed after this stage. The decision to remove the remaining statements on day-to-day practice was made during the final meeting.

#### **Consensus process**

Both rounds of voting had 21 votes (75% of network members at the time; see table 1 for demographic information on participating members). Following the initial statement development, the first round of voting had 55 items (figure 1, online supplemental material 2.1). Nine statements were eliminated after the first round of voting as they were below the threshold of agreement and six statements were amended based on comments from voting members. Of the 55 statements included in the first round of voting, 22 statements were included that did not follow the standard consensus methodology (online supplemental material 2.2). A different process was used for these statements and contained single-select or multiselect multiple choice options rather than the binary agree-disagree format used for the consensus statements. These statements are an important part of understanding the structure and processes for service delivery. Based on the responses from the first round of voting, three of these statements were amended and added as consensus statements with the binary agree-disagree response options in the following round of voting. As a result, the second round of voting included 26 statements that were

Table 1     Demographics for participating panel members			
Characteristics N (%)			
Sex, female		6 (21)	
Specialty			
Emergency medicine		2 (7)	
Neurology		3 (11)	
Neuropsychiatry		1 (4)	
Neuropsychology		3 (11)	
Neuros	surgery	12 (43)	
Paedia	tric medicine	1 (4)	
Rehab	ilitation medicine	1 (4)	
Neuros	science	2 (7)	
Sports	and exercise medicine	2 (7)	
Other		1 (4)	
Region			
East N	lidlands	1 (4)	
East of	f England	5 (18)	
Greate	r London	3 (11)	
North	East England	4 (14)	
North	West England	5 (18)	
South	East England	4 (14)	
South	West England	3 (11)	
West N	lidlands	1 (4)	
Yorksh	ire and the Humber	2 (7)	

put to a vote (online supplemental material 3). Three items did not reach the threshold of agreement following the second round of voting and were removed, leaving 23 statements to be discussed during the meeting. No statements were amended at this stage.

The virtual consensus meeting resulted in four statements being eliminated, two pairs of statements were combined and one statement was added. Of the eliminated statements, three were deemed to be beyond the scope of this consensus statement. The statement added during the meeting was suggested by one member and voted on twice as the level of agreement was just below the threshold following the first vote. The statement was modified and included following the second vote as the level of agreement was above the threshold.

This resulted in a final 18 statements (table 2), all above the raised threshold of 85% agreement. Fourteen statements achieved unanimous agreement. The final statements were organised into three themes: (1) care pathway to structured follow-up, (2) prognosis and measures of recovery, and (3) provision of outpatient clinics.

#### DISCUSSION

Healthcare services for patients with concussion need to be transformed to advance quality of care and patient outcome. Current healthcare services fail to meet the needs of a significant number of patients with concussion which results in suboptimal recovery and failure of the healthcare system to fulfil its purpose of improving quality of life through enhancing health. There is an urgent



Figure 1 Consensus process.

Agreement %

Table 2 Final consensus statements

1	Care pathway to structured follow-up	
	Patients with concussion with persistent symptoms should be able to access specialist clinics.	100
	There should be dedicated concussion clinics, or where not possible, clinics with dedicated time for patients with concussion.	100
	Specialist head injury clinics should comprise a multidisciplinary team that should include, as a minimum, a clinician, neuropsychologist and vestibular physiotherapist.	100
	There should be a direct patient pathway for patients with persistent symptoms to specialist outpatient services from the ED, GP and other sources.	100
2 Prognosis and measures of recovery		
	There should be a system for patients to be triaged to a specialist in head injury clinics based on persistent symptoms.	100
	There may be a role for digital tools (eg, platforms, apps) to be used for screening for persistent symptoms following concussion.	100
	Symptom scales can aid the assessment of symptom burden.	100
	Screening for persistent symptoms can be aided by symptom scales.	100
	A new patient-reported tool for the assessment of symptoms and outcome should be developed and validated.	90
	Prolonged symptoms may develop even in the absence of specific acute signs, such as loss of consciousness, post-traumatic amnesia and vomiting.	100
	There are no prognostic tools currently available with sufficient clinical accuracy to predict the risk of developing persistent symptoms.	100
	There is currently insufficient evidence to support the clinical application of objective biomarkers (ie, fluid, imaging) for identifying and screening for persistent symptoms after a concussion.	100
	Symptomatic patients with abnormal cranial/brain imaging should be considered for follow-up in a specialist clinic.	89
	There is not yet sufficient evidence to support the routine use of advanced imaging techniques (eg, DTI, fMRI, MEG) in clinical practice.	90
Provision of outpatient clinics		
	All patients should be directed to online information on concussion to help with expectations and recommendations.	100
	All clinics should have access to standardised protocols for assessing pituitary dysfunction.	100
	Comorbidities, socioeconomic status and social support should be factored into clinical decision-making on management.	90
	The underlying cause(s) of persistent symptoms should be established and treated on a case-by-case basis.	100
		L

DTI, diffusion tensor imaging; ED, emergency department; fMRI, functional MRI; GP, general practice; MEG, magnetoencephalography.

need to improve the delivery of and access to concussion care in order to reduce healthcare inequalities and errors in management. Guidelines exist, but healthcare services need to restructure so to facilitate the delivery of these guidelines.

The recommendations in this consensus statement primarily focus on the structure and process of specialist healthcare services in concussion care with the purpose of improving healthcare quality and patient outcome.<sup>20</sup> These are the components in healthcare services that facilitate the implementation and delivery of best-practice care to achieve optimal health for the target patient population.<sup>51</sup> Ultimately, patient outcome is determined by the structure and process of care which describe the care context and the sum of actions in the delivery of care, respectively.<sup>52</sup> This consensus statement prioritises a patient-centric approach with a focus on timely, integrated, effective, efficient and evidence-based care and urge standardisation of these practices nationally.

Given the recent developments in the field, translating basic and clinical evidence into the healthcare system requires work on feasibility, implementation and service redesign.<sup>53 54</sup> Without a systematic approach, the promise and desired goal of improving patient outcomes can be challenging to fulfil. Simultaneously, there are still gaps in evidence. Consensus statements serve as an important tool in bridging areas where evidence is underdeveloped.<sup>55</sup> This consensus statement provides guidance outlines how to redesign the structures and processes in care delivery for patients with concussion in England.

#### **Open access**

This is through expert opinion based on the current level of evidence and first-hand local experience in the various clinical settings which patients with concussion encounter and academic expertise. The guidance needs to be updated as the evidence changes, but this ensures patient outcome to improve without delay.

The statements included in the first voting round that did not follow the consensus format will help guide system redesign in future work through providing further detailed guidance on fundamental factors in concussion healthcare delivery. This includes the structure of clinical teams, timelines for care tasks and other processes in the clinical pathway from injury to specialist outpatient services.

While the consensus statements are aimed at all concussions, sports-related and non-sports-related concussions, the implementation of these services into the National Health Services in England requires reallocation and reprioritisation of resources, as well as a restructuring of the system. Applying these changes for the subgroup of concussions occurring in sports-at both professional and grassroot levels-could be more practical and achievable at this stage, especially accounting for the high number of all-cause concussions. For example, direct patient pathways from local sports clubs could feasibly be implemented with quick-access clinics given the manageable number of sports-related concussions. A short-term goal of the UK Head Injury Network is the development of regional sports concussion clinics across the country to alleviate the burden of concussions in all levels of sports. These clinics aim to provide evidencebased care. However, it is intended that these services will ultimately be expanded and offered to all-cause patients with concussion. Thus, while the implementation of these statements will primarily change clinical services for sports-related concussions in the near future, the statements are directed at national health services offered to all patients in the longer term.

#### **Clinical implications**

#### Care pathway to structured follow-up

Patients with persistent symptoms require further medical attention following the initial medical review—which is typically performed in emergency departments, general practice surgeries or by sport physicians—to recover to their preinjury level of health and function. Yet, in a Collaborative European NeuroTrauma Effectiveness Research in Traumatic Brain Injury (CENTER-TBI) study, 90% of 71 neurotrauma centres involved in the Europe-based study do not routinely schedule follow-up appointments for patients with head injury discharged from the emergency department.<sup>56</sup>

In the UK, there is no care pathway to ensure access to structured follow-up which is likely a fundamental underlying cause for the unmet needs in concussion care. The absence of such a service may be attributed to the heterogeneity of concussions and the vast differences in patient needs. However, a flexible and multidisciplinary approach is needed to personalise care and meet patient needs across the heterogenous recovery trajectories in the concussion population. Evidence suggests that providing multidisciplinary interventions can improve return to work and quality of life in this patient population.<sup>57</sup> Appropriate resources-time, equipment and staffing-need to be allocated to these services to ensure there is capacity to meet the diversity and scale of need. As a minimum, we suggest the inclusion of, though not limited to, a clinician (ie, specialist nurse, neurologist, neurosurgeon, sport and exercise physician), neuropsychologist and vestibular physiotherapist. This is due to the incidence of neurological, cognitive, psychological, cervicogenic and vestibular symptoms in post-concussive patients.<sup>58-60</sup> Moreover, due to the knowledge and skillset of physiotherapists, it may be beneficial to develop a new subspecialty in 'concussion physiotherapy'.<sup>28 61</sup>

Providing timely care is fundamental to meet patients' health needs. Delaying intervention can prolong recovery further and result in poorer outcome. One study found patients who sought care more than 30 days after a concussion had worse outcome at 3months compared with patients who sought care earlier.<sup>62</sup> Concussion services should be redesigned with care pathways that offer access to structured follow-up for all patients with concussion as necessary to ensure these patients receive timely and appropriate care. These statements echo the recommendations in the recent Lancet Commission on traumatic brain injury.<sup>63</sup>

#### Prognosis and measures of recovery

Objective measures of disease and prognostic factors provide a means for directing care.<sup>64</sup> One of the greatest challenges in concussion management is the lack of objective markers and the absence of reliable prognostic factors for recovery. Thus, early identification of patients at risk of prolonged recovery is not possible and therefore assessing the need for follow-up for individuals at an early stage is challenging. Consequently, no predictive tool can be used reliably in the clinical setting to assess the risk of prolonged recovery and the need for follow-up in the outpatient setting.<sup>40</sup> Even patients without marked acute signs and symptoms are at risk of prolonged recovery and should not be dismissed inadvertently. Of note, the panel recommends that patients with abnormal scans during the acute phase should always be offered follow-up as there is significant observable pathology during the acute phase to warrant further medical assessment.<sup>65</sup> There is evidence to suggest that some gross structural abnormalities found on acute CT imaging may be associated with worse outcomes.<sup>66</sup> However, we suggest that anything beyond standard structural imaging modalities-to rule out other pathologies-should not be used in the clinical setting. While advanced imaging methods are showing promise in research, they are yet to demonstrate reliable clinical utility and we do not recommend using these tools in clinical practice.<sup>38 47 48</sup>

In the absence of validated objective measures, ongoing self-reporting is currently the most reliable primary indicator of health status and method for identification of patients with persistent symptoms.<sup>67</sup> Of note, non-disclosure of concussion symptoms is a concern that needs to be addressed and is particularly prevalent in professional and community sports.<sup>68</sup> <sup>69</sup> To filter out people who have recovered and identify individuals who need further medical review, we recommend a screening system. This would need to be cost-effective and scalable due to the size of the concussion population. We suggest digital tools as a platform for the screening system as they offer a remote, convenient, efficient and scalable solution.<sup>70</sup> Once set up, costs remain comparatively low while also overcoming traditional barriers related to geographical, financial and resource limitations.<sup>71 72</sup> Such a system could also be efficiently and effectively used to triage patients to the most appropriate clinician in the multidisciplinary clinic based on patient-reported symptoms and disabilities.

Standardising evaluation of recovery in the absence of objective markers is challenging. However, symptom scales may aid assessment of symptom burden and provide a means for screening patients, particularly as history taking and patient-reported symptoms are currently the only methods for measuring recovery.<sup>67</sup> To achieve this long-term, we recommend developing and validating a patient-reported outcome measure for the concussion population to help assess recovery and determine care needs. This may include assessment of symptom burden, functional status, return to work or activity and guality of life. However, the specifics need to be determined in an appropriate study. While this study focuses on clinically actionable statements, we believe this is a research priority that is important to include as it is required to achieve many of the clinical statements recommended. A validated patient-reported outcome measure will facilitate the clinical pathway for patients with concussion to specialist outpatient services and will help move the field forward.

In the era of patient-centric and value-based healthcare, patient-reported outcome measures are increasingly important to capture the patient's perspective of their health. Even when objective measures are identified in the future, these tools should be used to capture a holistic patient-centric assessment of health status and recovery.

#### Provision of outpatient clinics

Providing early information and awareness is essential to managing expectations and can enhance recovery. A Transforming Research and Clinical Knowledge in Traumatic Brain Injury (TRACK-TBI) study found that only 42% of patients reported receiving head injury information at discharge from the emergency department.<sup>73</sup> Providing proper discharge information can be challenging in the emergency department due to the busy environment in these clinical settings and the cognitive state of the patients initially after injury which may prevent them from comprehending and retaining information.<sup>74</sup> Thus, we suggest directing patients to a single online platform for evidence-based information. The online modality allows anyone to access the information at any time, provides the same information to all regardless of geographical region and is easier to keep up to date with the evidence, which is particularly important in a fast-evolving field.

For patients with persistent symptoms, determining the most appropriate therapy can be challenging, especially in the absence of sufficiently strong evidence on therapeutic efficacy and timing of interventions. Given the vast heterogeneity in cases, no single treatment plan can be used for all patients with concussion and clinicians must evaluate the most appropriate intervention(s) on a case-by-case basis. While further research is required in this domain some therapeutic interventions have shown promise. Recommendations for therapeutic approaches are beyond the scope of this consensus statement and we recommend referring to other literature on this topic.<sup>29–34</sup> Biopsychosocial factors should be considered when assessing clinical and social care needs as these have been shown consistently to influence patient recovery.<sup>41</sup>

Lastly, we suggest that concussion clinics should have access to a standardised protocol for assessing pituitary function. Pituitary dysfunction can result in a range of different symptoms, many of which overlap with concussion symptoms, including fatigue, anxiety and mood disturbances. The literature on the incidence of pituitary dysfunction after concussions and screening protocols is still an area of active research. We suggest referring to a guidance paper by the British Neurotrauma Group on screening and management of pituitary dysfunction after traumatic brain injury.<sup>75</sup>

#### Recent developments in sports concussion guidelines

The sixth edition of the consensus statement on concussion in sport was published this year.<sup>50</sup> This statement also advises that individuals with persistent symptoms after a concussion (suggested as symptom persisting for longer than 4 weeks) should undergo a multimodal evaluation, including the use of standardised, validated symptom scales for assessment and to guide referrals. The statement recommends referral to specialist clinicians ideally part of a multidisciplinary network of practitioners, where possible, for targeted treatment for persistent symptoms including cervicogenic symptoms, migraine and headache, cognitive and psychological difficulties, balance disturbances, vestibular signs and oculomotor manifestations. Regarding the multidisciplinary network, the statement proposes the inclusion of sports medicine physicians, athletic trainers/therapists, physiotherapists, occupational therapists, sports chiropractors, neurologists, neurosurgeons, neuropsychologists, ophthalmologists, optometrists, physiatrists, psychologists and psychiatrists. Furthermore, biopsychosocial and comorbidities should also be considered in the context of persistent symptoms after concussions. Additionally, similar guidance has been provided regarding the use of advanced neuroimaging and other objective biomarkers in clinical practice.

#### Limitations

The literature on concussion, though rapidly evolving, is contentious and there are still numerous areas of disagreement among the panel. As discussed above, a wide approach was taken for the initial statement development which was later narrowed to focus on the healthcare structure and processes, in part. It cannot be excluded that some disagreement on statements was in part due to differences in definitions and terminology use despite probable broad agreement on the statement in principle. However, establishing agreement on terminology is a different consensus process on its own and was beyond the scope of this study.

We engaged with different stakeholders including academics, clinicians from different specialties and charities, and drew from previous patient-focus group sessions (unpublished observations). To ensure more robust statements, we chose a high threshold for consensus. While these statements reached agreement in this panel, we cannot exclude the possibility or to what degree the results are influenced by the experience and expertise of these individuals both for the included and excluded statements. The number of participants was in line with recommendations for Delphi studies<sup>76</sup>; however, more representatives from each specialty involved could have benefitted the consensus process.<sup>77</sup> The study was also limited in female representation. However, due to the high level of agreement and consistency with the literature, we believe the statements can be integrated reliably in the national healthcare system.

General practitioners are typically the first point of contact for patients with persistent problems seeking medical attention after a concussion and are a key player in the patient pathway. While the perspective of general practitioners is crucial for the wider concussion healthcare services, this study focused on the structures and processes in specialist outpatient services. However, the involvement of general practitioners is crucial for implementation and redesigning the concussion pathway.

#### **CONCLUSION**

This consensus statement outlines structural and processbased factors needed to move towards standardised best-practice care in concussion. This includes a final 18 statements on the care pathway for structured follow-up, prognosis and measures of recovery and provision of outpatient clinics. The statements are aimed at both the sports-related and all-cause concussion populations; however, they should be implemented for sports-related concussions as the first step before expanding to all patients with concussion given the greater feasibility of this approach. Further research and stakeholder engagement is needed to determine best-practice clinical concussion management in outpatient clinics and the recommendations in this statement provide the structures and processes whereby best-practice patient-centred care can be delivered in a timely, integrated, efficient and effective manner.

#### Author affiliations

<sup>1</sup>Department of Neurosurgery, University of Cambridge, Cambridge, UK <sup>2</sup>Inflammation and Ageing, University of Birmingham, Birmingham, UK <sup>3</sup>Nuffield Department of Clinical Neurosciences, University of Oxford, Oxford, UK <sup>4</sup>Royal Liverpool University Hospital, Liverpool, UK <sup>5</sup>UK Sports Institute, Liverpool, UK

<sup>6</sup>School of Sport, Exercise and Health Sciences, Loughborough University, Loughborough, UK

<sup>7</sup>Department of Clinical Infection, Microbiology and Immunology, Institute of Infection, Veterinary and Ecological Sciences, University of Liverpool, Liverpool, UK <sup>8</sup>Department of Paediatric Neurology, Alder-Hey Children's NHS Trust, Liverpool, UK <sup>9</sup>Department of Cognitive Neurology, Newcastle University, Newcastle Upon Tyne, UK

<sup>10</sup>Institute of Neurology, University College London, London, UK

 <sup>11</sup>Sunderland & South Tyneside Community Acquired Brain Injury Service, Northumberland Tyne and Wear NHS Foundation Trust, Newcastle upon Tyne, UK
<sup>12</sup>Department of Neurosurgery, Royal Victoria Infirmary, Newcastle upon Tyne, UK
<sup>13</sup>Wessex Neuroscience Centre, Southampton General Hospital, Southampton, UK
<sup>14</sup>Imperial College London, London, UK

<sup>15</sup>Carnegie Applied Rugby Research (CARR) Centre, Leeds Beckett University— Headingley Campus, Leeds, UK

<sup>16</sup>England Performance Unit, Rugby Football League Ltd, Leeds, UK

<sup>17</sup>Manchester Centre for Clinical Neurosciences (MCCN), Salford Royal Infirmary, Northern Care Alliance, Liverpool, UK

<sup>18</sup>Alder Hey Children's Hospital NHS Foundation Trust, Liverpool, UK

<sup>19</sup>Newcastle Upon Tyne Hospitals NHS Foundation Trust, Newcastle Upon Tyne, UK

- <sup>20</sup>University Hospitals Plymouth NHS Trust, Plymouth, UK
- <sup>21</sup>Department of Psychiatry, University of Oxford, Oxford, UK
- <sup>22</sup>Department of Neurology, North Bristol NHS Trust, Westbury on Trym, UK
- <sup>23</sup>National Hospital for Neurology and Neurosurgery, London, UK
- <sup>24</sup>Institute of Exercise and Health, University College London, London, UK
- <sup>25</sup>Carnegie School of Sport, Leeds Beckett University, Leeds, UK

<sup>26</sup>Medical, Leeds Rhinos Rugby League Club, Leeds, UK

<sup>27</sup>Department of Neurosurgery, Imperial College Healthcare NHS Trust, London, UK
<sup>28</sup>Department of Neuropsychology, University of Cambridge, Cambridge, UK
<sup>29</sup>Love of the Game, London, UK

Twitter Michael Grey @drmichaeljgrey, Ben Jones @23benjones and Shrouk Messahel @shroukmessahel

Acknowledgements We would like to thank Love of the Game for their support with this work. Peter Hutchinson would also like to thank the NIHR Senior Investigator Award, Cambridge Biomedical Research Centre, Medtech Brain Injury Co-operative, Global Health Research Group on Acquired Brain and Spine Injury and Royal College of Surgeons of England. AH is supported by the Cambridge Biomedical Research Campus, the Royal College of Surgeons of England, and is a Theme Lead in the NIHR Brain Injury MedTech Collaborative.

**Contributors** All authors have made substantial contributions to the manuscript in the following ways: EK, AH, AK, MP and PJAH led the delivery and planning of this consensus statement. AB, CG, MGre, MGri, TG, PG, AH, DH, PJAH, PJ, BJ, AK, EK, TL, TM, CH, SM, JN, RN, VR, KS, RS, DT, PW, MW, EW were responsible for the conception and design of the study. AK moderated the consensus meeting. AB, CG, MGre, MGri, TG, PG, AH, DH, PJAH, PJ, BJ, AK, EK, TL, TM, CH, SM, JN, RN, VR, KS, RS, DT, PW, MW, EW participated in the acquisition, analysis and interpretation of data. EK drafted the manuscript and CG, MGa, TG, AH, PJAH, PJ, BJ, AK, EK, CH, RN, DT, PW, and EW revised it thoroughly. All authors gave final approval of the submitted manuscript version. EK, AH, and PJAH are the guarantors for the overall study.

**Funding** The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

**Competing interests** AB serves as an expert advisor on concussion for England's Rugby Football Union, Football Association and Premier League. He also consults and has stocks in Marker Diagnostics LTD and licenses in Salivary Biomarker

of Concussion. AH served on the NICE Guideline Committee Trauma Guidelines for 2023. BJ consults for Premiership Rugby, London, and Rugby Football League, Manchester. DT consults for the Leeds Rhinos Rugby League Club and Hull City Tigers Football Club. MG serves on the advisory board for the Players Football Association. RN has undertaken work as a medicolegal expert in sports concussions. RS has served as an independent expert in legal cases in which head injury has occurred, some of these were mild head injuries. He also has received funding from the Rugby Football Association and Football Association for the Advanced Brainhealth project. TG has advised on legal cases where head injury has occurred, some of which were mild head injuries.

Patient and public involvement Patients and/or the public were not involved in the design, or conduct, or reporting, or dissemination plans of this research.

Patient consent for publication Not applicable.

**Ethics approval** The study was exempt from requiring ethical approval in accordance with the University of Cambridge Policy on the Ethics of Research Involving Human Participants and Personal Data.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement Data are available upon reasonable request.

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

**Open access** This is an open access article distributed in accordance with the Creative Commons Attribution 4.0 Unported (CC BY 4.0) license, which permits others to copy, redistribute, remix, transform and build upon this work for any purpose, provided the original work is properly cited, a link to the licence is given, and indication of whether changes were made. See: https://creativecommons.org/ licenses/by/4.0/.

#### **ORCID iDs**

Elika Karvandi http://orcid.org/0000-0003-1287-8200 Angelos G Kolias http://orcid.org/0000-0003-3992-0587 Ben Jones http://orcid.org/0000-0002-4274-6236 Shrouk Messahel http://orcid.org/0000-0003-0645-3070 Vanessa Raymont http://orcid.org/0000-0001-8238-4279 Peter John Hutchinson http://orcid.org/0000-0002-2796-1835

#### REFERENCES

- National Institute for Health and Care Excellence. Head injury: assessment and early management. (NICE guideline [NG232]).
  London: NICE, 2023. Available: https://www.nice.org.uk/guidance/ ng232
- 2 Silverberg ND, Iverson GL, et al, ACRM Brain Injury Special Interest Group Mild TBI Task Force members. The American Congress of rehabilitation medicine diagnostic criteria for mild traumatic brain injury. Arch Phys Med Rehabil 2023;104:1343–55.
- 3 Sharp DJ, Jenkins PO. Concussion is confusing us all. *Pract Neurol* 2015;15:172–86. 10.1136/practneurol-2015-001087 Available: https://www.scopus.com/inward/record.uri?eid=2-s2. 0-84929330344&doi=10.1136%2fpractneurol-2015-001087& partnerID=40&md5=c2ae641c33158dcbe06dc0b311be793f
- 4 Ganau L, Ligarotti GKI, Prisco L, et al. Letter: A Scoping review of registered clinical studies on mild traumatic brain injury and concussion (2000-2019). Neurosurgery 2021;89:E178–9.
- 5 Levin HS, Diaz-Arrastia RR. Diagnosis, prognosis, and clinical management of mild traumatic brain injury. *Lancet Neurol* 2015;14:506–17.
- 6 Marshall SW, Guskiewicz KM, Shankar V, et al. Epidemiology of sports-related concussion in seven US high school and collegiate sports. Inj Epidemiol 2015;2:13.
- 7 Langer L, Levy C, Bayley M. Increasing incidence of concussion: true epidemic or better recognition J Head Trauma Rehabil 2020;35:E60–6.
- 8 McCrea M, Guskiewicz KM, Marshall SW, *et al.* Acute effects and recovery time following concussion in collegiate football players. *JAMA* 2003;290:2556–63.

- 9 McCrory P, Meeuwisse WH, Echemendia RJ, et al. What is the lowest threshold to make a diagnosis of concussion Br J Sports Med 2013;47:268–71.
- 10 Cancelliere C, Verville L, Stubbs JL, et al. Post-concussion symptoms and disability in adults with mild traumatic brain injury: A systematic review and meta-analysis. J Neurotrauma 2023;40:1045–59.
- 11 Stern RA, Seichepine D, Tschoe C, et al. Concussion care practices and utilization of evidence-based guidelines in the evaluation and management of concussion: A survey of New England emergency departments. J Neurotrauma 2017;34:861–8.
- 12 Ross J, Santhirapala R, MacEwen C, et al. Helping patients choose wisely. BMJ 2018;361:k2585.
- 13 Maughan D, Ansell J. Protecting resources, promoting value: a doctors guide to cutting waste in clinical care. Academy of Medical Royal Colleges, 2014.
- 14 Williams E, Buck D, Babalola G. What are health inequalities? London: The King's Fund, 2020. Available: https://www.kingsfund. org.uk/publications/what-are-health-inequalities
- 15 Dasic D, Morgan L, Panezai A, et al. A Scoping review on the challenges, improvement programs, and relevant output Metrics for Neurotrauma services in major trauma centers. Surg Neurol Int 2022;13:171.
- 16 National Institute for Health and Excellence. Practical steps to improving the quality of care and services using NICE guidance, 2018Available: https://www.nice.org.uk/Media/Default/About/whatwe-do/Into-practice/Practical-steps-improving-quality-of-care-andservices-using-NICE-guidance.pdf
- 17 Wiebe DJ, Comstock RD, Nance ML. Concussion research: a public health priority. *Inj Prev* 2011;17:69–70.
- 18 Department for Culture M and SUG. UK concussion guidelines for non-elite (grassroots) sport. 2023. Available: http://sramedia.s3. amazonaws.com/media/documents/9ced1e1a-5d3b-4871-9209bff4b2575b46.pdf
- 19 Veliz P, McCabe SE, Eckner JT, *et al.* Trends in the prevalence of concussion reported by US adolescents, 2016-2020. *JAMA* 2021;325:1789–91.
- 20 Hannawa AF, Wu AW, Kolyada A, et al. The aspects of Healthcare quality that are important to health professionals and patients: A qualitative study. Patient Educ Couns 2022;105:1561–70.
- 21 Jünger S, Payne SA, Brine J, et al. Guidance on conducting and reporting delphi studies (credes) in palliative care: recommendations based on a methodological systematic review. *Palliat Med* 2017;31:684–706.
- 22 Diamond IR, Grant RC, Feldman BM, et al. Defining consensus: a systematic review recommends Methodologic criteria for reporting of Delphi studies. J Clin Epidemiol 2014;67:401–9.
- 23 Slade SC, Dionne CE, Underwood M, et al. Standardised method for reporting exercise programmes: protocol for a modified Delphi study. BMJ Open 2014;4:e006682.
- 24 Vogel C, Zwolinsky S, Griffiths C, *et al.* A Delphi study to build consensus on the definition and use of big data in obesity research. *Int J Obes* 2019;43:2573–86.
- 25 Giza CC, Kutcher JS, Ashwal S, et al. Summary of evidence-based guideline update: evaluation and management of concussion in sports: report of the guideline development subcommittee of the American Academy of neurology. *Neurology* 2013;80:2250–7.
- 26 Silverberg ND, laccarino MA, Panenka WJ, et al. Management of concussion and mild traumatic brain injury: A synthesis of practice guidelines. Arch Phys Med Rehabil 2020;101:382–93.
- 27 Marshall S, Bayley M, McCullagh S, et al. Updated clinical practice guidelines for concussion/mild traumatic brain injury and persistent symptoms. *Brain Inj* 2015;29:688–700.
- 28 Quatman-Yates CC, Hunter-Giordano A, Shimamura KK, et al. Physical therapy evaluation and treatment after concussion/ mild traumatic brain injury. J Orthop Sports Phys Ther 2020;50:CPG1–73.
- 29 Nygren-de Boussard C, Holm LW, Cancelliere C, et al. Nonsurgical interventions after mild traumatic brain injury: A systematic review. results of the International collaboration on mild traumatic brain injury prognosis. Arch Phys Med Rehabil 2014;95:S257–64.
- 30 Schneider KJ, Leddy JJ, Guskiewicz KM, et al. Rest and treatment/ rehabilitation following sport-related concussion: a systematic review. Br J Sports Med 2017;51:930–4.
- 31 Murray DA, Meldrum D, Lennon O. Can vestibular rehabilitation exercises help patients with concussion? A systematic review of efficacy, prescription and progression patterns. *Br J Sports Med* 2017;51:442–51.
- 32 Lal A, Kolakowsky-Hayner SA, Ghajar J, *et al.* The effect of physical exercise after a concussion: A systematic review and meta-analysis. *Am J Sports Med* 2018;46:743–52.

#### **Open access**

- 33 Reid SA, Farbenblum J, McLeod S. Do physical interventions improve outcomes following concussion: a systematic review and meta-analysis *Br J Sports Med* 2022;56:292–8.
- 34 Heslot C, Azouvi P, Perdrieau V, et al. A systematic review of treatments of post-concussion symptoms. J Clin Med 2022;11:6224.
- 35 Rausa VC, Anderson V, Babl FE, et al. Predicting concussion recovery in children and adolescents in the emergency Department. Curr Neurol Neurosci Rep 2018;18:78.
- 36 Mercier E, Tardif P-A, Cameron PA, et al. Prognostic value of neuronspecific Enolase (NSE) for prediction of post-concussion symptoms following a mild traumatic brain injury: a systematic review. Brain Inj 2018;32:29–40.
- 37 Zetterberg H, Morris HR, Hardy J, et al. Update on fluid biomarkers for concussion. Concussion 2016;1:CNC12.
- 38 McCrea M, Meier T, Huber D, et al. Role of advanced neuroimaging, fluid biomarkers and genetic testing in the assessment of sportrelated concussion: a systematic review. Br J Sports Med 2017;51:919–29.
- 39 O'Connell B, Kelly ÁM, Mockler D, et al. Use of blood biomarkers in the assessment of sports-related concussion – A systematic review in the context of their biological significance. Clin J Sport Med 2018;28:561–71.
- 40 Iverson GL, Gardner AJ, Terry DP, et al. Predictors of clinical recovery from concussion: a systematic review. Br J Sports Med 2017;51:941–8.
- 41 van Ierssel J, Pennock KF, Sampson M, et al. Which Psychosocial factors are associated with return to sport following concussion? A systematic review. J Sport Health Sci 2022;11:438–49.
- 42 Karantali E, Kazis D, McKenna J, *et al.* Neurofilament light chain in patients with a concussion or head impacts: a systematic review and meta-analysis. *Eur J Trauma Emerg Surg* 2022;48:1555–67.
- 43 Silverberg ND, Gardner AJ, Brubacher JR, et al. Systematic review of multivariable Prognostic models for mild traumatic brain injury. J Neurotrauma 2015;32:517–26.
- 44 Cancelliere C, Hincapié CA, Keightley M, et al. Systematic review of prognosis and return to play after sport concussion: results of the International collaboration on mild traumatic brain injury prognosis. Arch Phys Med Rehabil 2014;95:S210–29.
- 45 Kamins J, Bigler E, Covassin T, et al. What is the physiological time to recovery after concussion? A systematic review. Br J Sports Med 2017;51:935–40.
- 46 Manley G, Gardner AJ, Schneider KJ, et al. A systematic review of potential long-term effects of sport-related concussion. Br J Sports Med 2017;51:969–77.
- 47 Lindsey HM, Hodges CB, Greer KM, et al. Diffusion-weighted imaging in mild traumatic brain injury: A systematic review of the literature. *Neuropsychol Rev* 2023;33:42–121.
- 48 Khong E, Odenwald N, Hashim E, et al. Diffusion Tensor imaging findings in post-concussion syndrome patients after mild traumatic brain injury: A systematic review. Front Neurol 2016;7:156.
- 49 Lumba-Brown A, Yeates KO, Sarmiento K, et al. Centers for disease control and prevention guideline on the diagnosis and management of mild traumatic brain injury among children. JAMA Pediatr 2018;172:e182853.
- 50 Patricios JS, Schneider KJ, Dvorak J, *et al.* Consensus statement on concussion in sport: the 6th international conference on concussion in sport-Amsterdam 2022. *Br J Sports Med* 2023;57:695–711.
- 51 Rademakers J, Delnoij D, de Boer D. Structure, process or outcome: which contributes most to patients' overall assessment of Healthcare quality *BMJ Qual Saf* 2011;20:326–31.
- 52 NHS England and NHS Improvement. A model for measuring quality care. London: NHS England, 2021. Available: https://www.england. nhs.uk/wp-content/uploads/2022/02/qsir-measuring-quality-care.pdf
- 53 Grol R, Grimshaw J. From best evidence to best practice: effective implementation of change in patients' care. *Lancet* 2003;362:1225–30.
- 54 Bauer MS, Damschroder L, Hagedorn H, et al. An introduction to implementation science for the non-specialist. BMC Psychol 2015;3:32.
- 55 La Brooy C, Pratt B, Kelaher M. What is the role of consensus statements in a risk society *Journal of Risk Research* 2020;23:664–77.
- 56 Foks KA, Cnossen MC, Dippel DWJ, et al. Management of mild traumatic brain injury at the emergency Department and hospital admission in Europe: A survey of 71 Neurotrauma

centers participating in the CENTER-TBI study. *J Neurotrauma* 2017;34:2529–35.

- 57 Jensen KL, Wachner LG, Storm van den Brekel J, et al. Multidisciplinary interventions increase weekly working hours and quality of life in persons with post-concussion syndrome. *Neurol Res* 2023;45:926–35.
- 58 Hiploylee C, Dufort PA, Davis HS, et al. Longitudinal study of Postconcussion syndrome: not everyone recovers. J Neurotrauma 2017;34:1511–23. 10.1089/neu.2016.4677 Available: http://www. liebertonline.com/neu
- 59 Polinder S, Cnossen MC, Real RGL, et al. A multidimensional approach to post-concussion symptoms in mild traumatic brain injury. Front Neurol 2018;9:1113. 10.3389/fneur.2018.01113 Available: https://www.scopus.com/inward/record.uri?eid=2-s2. 0-85063738495&doi=10.3389%2ffneur.2018.01113&partnerID=40& md5=7da2daff1fa8712a1dffea114975fb89
- 60 Faulkner JW, Theadom A, Snell DL, et al. Network analysis applied to post-concussion symptoms in two mild traumatic brain injury samples. *Front Neurol* 2023;14:1226367.
- 61 Brown L, Camarinos J. The role of physical therapy in concussion rehabilitation. *Semin Pediatr Neurol* 2019;30:68–78.
- 62 Bunt SC, LoBue C, Hynan LS, *et al*. Early vs. delayed evaluation and persisting concussion symptoms during recovery in adults. *Clin Neuropsychol* 2023;37:1410–27.
- 63 Maas AIR, Menon DK, Manley GT, et al. Traumatic brain injury: progress and challenges in prevention, clinical care, and research. *Lancet Neurol* 2022;21:1004–60.
- 64 Déry J, De Guise É, Bussières È-L, et al. Prognostic factors for persistent symptoms in adults with mild traumatic brain injury: protocol for an overview of systematic reviews. Syst Rev 2021;10:254.
- 65 Voormolen DC, Haagsma JA, Polinder S, *et al.* Post-concussion symptoms in complicated vs. uncomplicated mild traumatic brain injury patients at three and six months post-injury: results from the CENTER-TBI study. *J Clin Med* 2019;8:1921:11.:.
- 66 Yuh EL, Jain S, Sun X, et al. Pathological computed tomography features associated with adverse outcomes after mild traumatic brain injury: A TRACK-TBI study with external validation in CENTER-TBI. JAMA Neurol 2021;78:1137–48.
- 67 Yeates KO, Räisänen AM, Premij Z, et al. What tests and measures accurately diagnose persisting post-Concussive symptoms in children, adolescents and adults following sport-related concussion? A systematic review. Br J Sports Med 2023;57:780–8.
- 68 Longworth T, McDonald A, Cunningham C, et al. Do Rugby League players under-report concussion symptoms? A cross-sectional study of elite teams based in Australia. BMJ Open Sport Exerc Med 2021;7:e000860.
- 69 Tadmor D, Till K, Phillips G, et al. I won't let you down; why 20% of men's and women's super League players Underreported suspected concussions. J Sci Med Sport 2023:S1440-2440(23)00416-4.
- 70 NIHR Evidence. What is digital health technology and what can it do for me? London, 2022. Available: https://evidence.nihr.ac.uk/ collection/what-is-digital-health-technology/
- 71 WHO Team: Digital Health and Innovation GRCS. Recommendations on Digital interventions for health system strengthening. 2019. Available: https://iris.who.int/bitstream/handle/10665/311941/ 9789241550505-eng.pdf?sequence=31
- 72 Powell D, Stuart S, Godfrey A. Sports related concussion: an emerging era in Digital sports technology. *NPJ Digit Med* 2021;4:164. 10.1038/s41746-021-00538-w Available: https://www.nature.com/ npjdigitalmed/
- 73 Seabury SA, Gaudette É, Goldman DP, et al. Assessment of followup care after emergency Department presentation for mild traumatic brain injury and concussion. JAMA Netw Open 2018;1:e180210.
- 74 Bergman K, Louis S. Discharge instructions for concussion: are we meeting the patient needs J Trauma Nurs 2016;23:327–33.
- 75 Tan CL, Alavi SA, Baldeweg SE, et al. The screening and management of pituitary dysfunction following traumatic brain injury in adults: British Neurotrauma group guidance. J Neurol Neurosurg Psychiatry 2017;88:971–81.
- 76 Birko S, Dove ES, Özdemir V, et al. Evaluation of nine consensus indices in Delphi foresight research and their dependency on Delphi survey characteristics: A simulation study and debate on Delphi design and interpretation. *PLoS ONE* 2015;10:e0135162.
- 77 de Villiers MR, de Villiers PJT, Kent AP. The Delphi technique in health sciences education research. *Med Teach* 2005;27:639–43.