

## Categorical Proof-theoretic Semantics

Pym, David; Ritter, Eike; Robinson, Edmund P.

DOI:

[10.1007/s11225-024-10101-9](https://doi.org/10.1007/s11225-024-10101-9)

License:

Creative Commons: Attribution (CC BY)

*Document Version*

Publisher's PDF, also known as Version of record

*Citation for published version (Harvard):*

Pym, D, Ritter, E & Robinson, EP 2024, 'Categorical Proof-theoretic Semantics', *Studia Logica*.  
<https://doi.org/10.1007/s11225-024-10101-9>

[Link to publication on Research at Birmingham portal](#)

### General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.




When citing, please reference the published version.

### Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact [UBIRA@lists.bham.ac.uk](mailto:UBIRA@lists.bham.ac.uk) providing details and we will remove access to the work immediately and investigate.



DAVID PYM   
EIKE RITTER   
EDMUND ROBINSON 

# Categorical Proof-theoretic Semantics

**Abstract.** In proof-theoretic semantics, model-theoretic validity is replaced by proof-theoretic validity. Validity of formulae is defined inductively from a base giving the validity of atoms using inductive clauses derived from proof-theoretic rules. A key aim is to show completeness of the proof rules without any requirement for formal models. Establishing this for propositional intuitionistic logic raises some technical and conceptual issues. We relate Sandqvist's (complete) base-extension semantics of intuitionistic propositional logic to categorical proof theory in presheaves, reconstructing categorically the soundness and completeness arguments, thereby demonstrating the naturality of Sandqvist's constructions. This naturality includes Sandqvist's treatment of disjunction that is based on its second-order or elimination-rule presentation. These constructions embody not just validity, but certain forms of objects of justifications. This analysis is taken a step further by showing that from the perspective of validity, Sandqvist's semantics can also be viewed as the natural disjunction in a category of sheaves.

*Keywords:* Proof-theoretic semantics, Base-extension semantics, Categorical logic, Kripke Semantics, Presheaves, Sheaves.

## 1. Introduction

In model-theoretic semantics, logical languages are interpreted in mathematical structures that carry appropriate axiomatizations and support a notion of model-theoretic consequence. Specifically, a propositional sentence  $\phi$  is a model-theoretic consequence of a set  $\Gamma$  of propositional sentences iff every model  $\mathcal{M}$  of  $\Gamma$  is also a model of  $\phi$ ; that is,

$$\Gamma \models \phi \text{ iff for all } \mathcal{M}, \text{ if, for all } \psi \in \Gamma, \text{ if } \mathcal{M} \models \psi, \text{ then } \mathcal{M} \models \phi$$

As Schroeder-Heister [34] explains, in this definition of consequence we have a transmission of truth from premisses to conclusion in which transmission is determined by classical implication in the meta-theory. This definition

---

David Pym, Eike Ritter and Edmund Robinson contributed equally to this work.

Presented by **Heinrich Wansing**; *Received* March 6, 2023

of consequence supports inductive characterizations of the meanings of the logical constants. For example,

$$\mathcal{M} \models \phi \wedge \psi \text{ iff } \mathcal{M} \models \phi \text{ and } \mathcal{M} \models \psi$$

Proof-theoretic consequence is usually defined as derivability in a formal system: a propositional sentence  $\phi$  is a proof-theoretic consequence of a set of propositional sentences  $\Gamma$  in a formal system  $S$  — that is,  $\Gamma \vdash_S \phi$  — if it can be derived from the elements of  $\Gamma$  using the axioms and inference rules of  $S$ . This definition of consequence also supports inductive characterizations of the meanings of the logical constants. For example,

$$\text{if } \Gamma \vdash_S \phi \text{ and } \Gamma \vdash_S \psi, \text{ then } \Gamma \vdash_S \phi \wedge \psi$$

Note that the invertibility of these characterizations is a delicate matter (see, for example, [22]).

From the perspective of model-theoretic semantics, and its primary notion of truth, the correctness of inferences in a formal system  $S$  is given by a soundness assertion,

$$\text{if } \Gamma \vdash_S \phi, \text{ then } \Gamma \models \phi$$

If completeness — the converse — also holds, then model-theoretic consequence and proof-theoretic consequence coincide. Note that completeness does not imply the invertibility of the proof-theoretic characterizations.

Proof-theoretic semantics provides an alternative account of the meaning of the logical constants to that which is provided by model-theoretic semantics. Roughly speaking, there are two approaches to proof-theoretic semantics. First, there is what we call Dummett–Prawitz proof-theoretic semantics [34] and, second, there is what we call base-extension semantics [23, 24, 32]. The former has temporal priority, but the latter may be seen as being the more general perspective [12].

The Dummett–Prawitz view arises from a philosophical reading by Dummett [4] of the normalization results by Prawitz [27]. It has subsequently been developed substantially both mathematically and philosophically with Schroeder-Heister [33] giving a general account that clearly separates the semantic and computational considerations. It has largely been developed for IL. In this context, the rules of the natural deduction system NJ for IL are taken to be *a priori* valid. An argument is a tree of formulas whose leaves are called its assumptions, some of which may be labelled as discharged. An argument is *indirect* (i.e., not direct) if it contains *detours* in which case it may be made *direct* by reduction *à la* Prawitz [27]; for example, the reduction of an indirect argument via  $\supset$  may be reduced as follows:

$$\begin{array}{ccc}
 & [\phi] & \\
 & \mathcal{D}_2 & \mathcal{D}_1 \\
 \mathcal{D}_1 & \frac{\psi}{\phi \supset \psi} & \phi \\
 \phi & \frac{\phi \supset \psi}{\psi} & \mathcal{D}_2 \\
 \hline
 \psi & \rightsquigarrow & \psi
 \end{array}$$

Arguments without assumptions and without detours are said to be *canonical proofs*, they are inherently valid. The validity of an arbitrary argument is determined by whether or not it represents according to some fixed operations (e.g., through reduction) one of these canonical proofs. In this way, proof-theoretic validity in the Dummett–Prawitz tradition is a semantics of proofs. This view is not our concern here.

Base-extension semantics is a characterization of consequence given by an inductively defined judgment whose base case is given by proof rather than by truth. Crucially, while in model-theoretic semantics the base case of the judgment is given by truth (i.e., in the model  $\mathcal{M}$  with interpretation  $I$ , the judgment  $w \Vdash p$  obtains iff  $w \in I(p)$ ), in proof-theoretic semantics it is instead given by provability in an atomic system; that is, by the following clause in which  $\mathcal{B}$  is an arbitrary atomic system:

$$\Vdash_{\mathcal{B}} p \quad \text{iff} \quad p \text{ is provable in } \mathcal{B}$$

Note that base-extension semantics relies on a (proof-irrelevant) provability judgement not on the judgement that a proof-object establishes a consequence.

Of course, one can give base-extension semantics that simply mimic model-theoretic semantics. Goldfarb’s completeness proof works by taking a base-extension semantics that encodes the possible worlds structure of a Kripke countermodel — see Goldfarb [16]. This approach allows completeness to be obtained while still using the usual frame semantics for disjunction in IPL as given by Kripke [19],

$$w \Vdash \phi \vee \psi \quad \text{iff} \quad w \Vdash \phi \text{ or } w \Vdash \psi,$$

where  $w$  denotes an arbitrary world. Similarly, recent work by Stafford and Nascimento [36] obtains completeness by constructing a Kripke model from sets of base rules.

By contrast, in the complete base-extension semantics for IPL given by Sandqvist [32], disjunction has the following clause in which  $\mathbb{A}$  is the set of atomic propositions:

$$\Vdash_{\mathcal{B}} \phi \vee \psi \quad \text{iff} \quad \text{for any } \mathcal{C} \supseteq \mathcal{B} \text{ and any } p \in \mathbb{A}, \\ \phi \Vdash_{\mathcal{C}} p \text{ and } \psi \Vdash_{\mathcal{C}} p \text{ implies } \emptyset \Vdash_{\mathcal{C}} p$$

Completeness is obtained by using a base that mimics propositions and their proofs, rather than the elements of a Kripke model.

While, in the spirit of Dummett–Prawitz proof-theoretic semantics, the form of the semantics for disjunction given above is naturally seen as a representation of the  $\vee$ -elimination rule of natural deduction [11, 27], it is also naturally seen as the well-known definition of  $\vee$  in second-order propositional logic; see, for example, [6, 14, 31, 32, 39]). This view is also discussed in Section 4.

In both cases, the definition restricts the conclusion of the hypothetical assumptions and the conclusion of the definition itself to be atomic. That is,

$$\frac{\begin{array}{cc} [\phi] & [\psi] \\ \vdots & \vdots \\ \phi \vee \psi & p \quad p \end{array}}{p}$$

and, writing  $\bigwedge$  for the second-order propositional universal quantifier,

$$\phi \vee \psi = \bigwedge p. (\phi \supset p) \supset (\psi \supset p) \supset p$$

Sandqvist’s semantics and our category-theoretic analysis of it can both be understood in both these ways. (?)

This way of interpreting disjunction has been studied in the context of the atomic fragment  $F_{at}$  [5, 7–9, 26] of System F [13, 15]. In [7, 9] it is shown that this interpretation is faithful; that is, for any formula  $\phi$  of IPL,  $\phi$  is provable iff its  $F_{at}$ -interpretation is provable in  $F_{at}$ . This result might, from a perspective that lies outwith our present scope, be seen as a proof-theoretic counterpart to Sandqvist’s completeness result. The link between  $F_{at}$  and our account is explained in Section 4. Furthermore, this form of the semantics for disjunction is closely related to Beth’s semantics see, for example, the discussion of Kripke–Beth–Joyal semantics and the relevant historical context in [20].

Gheorghiu and Pym [12] have shown that base-extension semantics may be regarded as the declarative counterpart to the operational paradigm of proof-theoretic validity. In particular, it is shown in [12] that Dummett–Prawitz validity can be recovered from base-extension semantics for intuitionistic propositional logic.

Category theory is a general theory of mathematical structures and their relations that was introduced by Eilenberg and Mac Lane in the middle of the 20th century in their foundational work on algebraic topology. It provides

a unifying language for studying mathematical structures that supports the use of one theory to explain another. Categorical logic is a highly developed area that studies logic from the point of view of categorical structures.

The relationships between model-theoretic semantics and categorical logic and between proof theory and categorical logic are rich and highly developed, especially in the world of intuitionistic and modal logics. See, for example, [2, 17, 21, 35]. The relationship with the proof theory of classical logic is less well developed, but see, for example, [1, 10, 30].

In this paper, we begin an exploration of the relationship between proof-theoretic semantics and categorical logic. There are two main motivations for this. First, in general, to bring proof-theoretic semantics into the framework of categorical logic. Second, more specifically, to explore certain technical aspects of the formulation of the base-extension semantics of intuitionistic propositional logic [16, 23, 24, 32, 36]:

- The formal naturality of the semantics
- The choices of semantics for disjunction.

Here, formal naturality refers to the existence of natural transformations between functors (as defined in [17, 20, 21], for example): if  $F$  and  $G$  are functors between categories  $C$  and  $D$ , then a natural transformation  $\eta$  between  $F$  and  $G$  is family of morphisms that satisfies the following:

1.  $\eta$  must associate to every object  $x$  in  $C$  an arrow  $\eta_x : F(x) \rightarrow G(x)$
2. For every  $f : x \rightarrow y$  in  $C$ ,  $\eta_y \circ F(f) = G(f) \circ \eta_x$ , where  $\circ$  denotes composition of morphisms.

Informally, the notion of a natural transformation captures that a given map between functors can be done consistently over an entire category. In the situation above, we refer to the structure being ‘natural in  $x$ ’.

Our primary focus will be Sandqvist’s base-extension semantics for intuitionistic propositional logic [32]. We give a concise introduction to this work in Section 2.

We now summarize the structure of the remainder of this paper. In Section 2, we summarize Sandqvist’s base-extension semantics for intuitionistic propositional logic, including the soundness and completeness of consequences derivable in NJ (recalled in Figure 1), denoted  $\Gamma \vdash \phi$ , for base-extension validity, denoted  $\Gamma \Vdash \phi$ . We proceed, in Section 3, to give a categorical formulation of Sandqvist’s constructions. This uses well-established (essentially type-theoretic) ideas from categorical logic (e.g., [17, 20, 21, 35]), working in a presheaf category. We establish the correctness (soundness and

completeness) of our algebraic constructions relative to Sandqvist’s semantics. Next, in Section 4, we discuss the relationships between base-extension semantics, Kripke models of intuitionistic logic, and presheaf models of type theory. In Section 5, we reconstruct categorically the logical metatheory of the base-extension semantics for intuitionistic propositional logic, and establish the familiar soundness and completeness theorems with respect to NJ: that is, that  $\Gamma \vdash \phi$  iff  $\Gamma \models \phi$ , where  $\models$  denotes the consequence relation derived from our category-theoretic model. The paper concludes with a reflection, in Section 6, on the semantics of disjunction, explaining how the choice of its interpretation — essentially between the Kripke-style interpretation and the Sandqvist-style interpretation, which adopts the second-order formulation that corresponds to the proof-theoretic interpretation offered by the elimination rule — affects completeness.

A preliminary version of this work has been presented at the 11th Scandinavian Logic Society Symposium, 2022 [28].

## 2. Base-extension Semantics for Intuitionistic Propositional Logic

Sandqvist [32] gives a base-extension proof-theoretic semantics for IPL for which natural deduction is sound and complete. We refer the reader to [32] for the detailed motivation and technical development. Here we give a very brief summary.

In the sequel, Romans  $p$ ,  $P$ , etc., respectively denote atoms and sets of atoms; Greeks  $\phi$ ,  $\Gamma$ , etc., respectively denote formulae and sets of formulae.

A base  $\mathcal{B}$  is a set of atomic rules (for  $\vdash_{\mathcal{B}}$ ), as in Definition 1, which also defines the application of base rules, and satisfaction in a base ( $\models_{\mathcal{B}}$ ).

DEFINITION 1. (Base) Base rules  $\mathcal{R}$ , application of base rules, and satisfaction of formulae in a (possibly finite) countable base  $\mathcal{B}$  of rules  $\mathcal{R}$  are defined as follows: an atomic rule  $\mathcal{R}$  is given as a second-order implication (denoted  $\Rightarrow$ ) involving (sets of) atoms:  $((P_1 \Rightarrow q_1), \dots, (P_n \Rightarrow q_n)) \Rightarrow r$  (upper case  $P_i$  denote sets of atoms, while lower case  $q_j$  denote individual atoms). This can be thought of as denoting the proof fragment with discharged hypotheses:

$$\frac{\begin{array}{c} [P_1] \quad [P_n] \\ q_1 \quad \dots \quad q_n \end{array} \mathcal{R}}{r}$$

---


$$\begin{array}{c}
 \frac{\Gamma \vdash \phi \quad \Gamma \vdash \psi}{\Gamma \vdash \phi \wedge \psi} \wedge I \qquad \frac{\Gamma \vdash \phi \wedge \psi}{\Gamma \vdash \phi} \wedge E \quad \frac{\Gamma \vdash \phi \wedge \psi}{\Gamma \vdash \psi} \wedge E \\
 \frac{\Gamma \vdash \phi}{\Gamma \vdash \phi \vee \psi} \vee I \quad \frac{\Gamma \vdash \psi}{\Gamma \vdash \phi \vee \psi} \vee I \quad \frac{\Gamma \vdash \phi \vee \psi \quad \Gamma, \phi \vdash \chi \quad \Gamma, \psi \vdash \chi}{\Gamma \vdash \chi} \vee E \\
 \frac{\Gamma, \phi \vdash \psi}{\Gamma \vdash \phi \supset \psi} \supset I \qquad \frac{\Gamma \vdash \phi \quad \Gamma \vdash \phi \supset \psi}{\Gamma \vdash \psi} \supset E
 \end{array}$$


---

 Figure 1. The calculus NJ (in sequential form and eliding  $\top$ ) [27]

We can now define consequence for atoms with respect to a base:

- (Ref)  $P, p \vdash_{\mathcal{B}} p$   
 (App $_{\mathcal{R}}$ ) if  $((P_1 \Rightarrow q_1), \dots, (P_n \Rightarrow q_n)) \Rightarrow r$  is a rule in the base and,  
 for all  $i \in [1, n]$ ,  $P, P_i \vdash_{\mathcal{B}} q_i$ , then  $P \vdash_{\mathcal{B}} r$

Here, the combinators (Ref and App $_{\mathcal{R}}$ ) specify how to construct proofs in a base using rules  $\mathcal{R}$  expressed using the second-order implication  $\Rightarrow$  as discussed above (see also Sandqvist [32]).

**DEFINITION 2.** (Validity in a Base) Validity in a base is defined inductively as follows:

- (At) For atomic  $p$ ,  $\Vdash_{\mathcal{B}} p$  iff  $\vdash_{\mathcal{B}} p$   
 ( $\supset$ )  $\Vdash_{\mathcal{B}} \phi \supset \psi$  iff  $\phi \Vdash_{\mathcal{B}} \psi$   
 ( $\wedge$ )  $\Vdash_{\mathcal{B}} \phi \wedge \psi$  iff  $\Vdash_{\mathcal{B}} \phi$  and  $\Vdash_{\mathcal{B}} \psi$   
 ( $\vee$ )  $\Vdash_{\mathcal{B}} \phi \vee \psi$  iff, for every atomic  $p$  and every  $\mathcal{C} \supseteq \mathcal{B}$ , if  $\phi \Vdash_{\mathcal{C}} p$  and  $\psi \Vdash_{\mathcal{C}} p$ , then  $\Vdash_{\mathcal{C}} p$   
 ( $\perp$ )  $\Vdash_{\mathcal{B}} \perp$  iff, for all atomic  $p$ ,  $\Vdash_{\mathcal{B}} p$   
 and  
 (Inf) for  $\Theta \neq \emptyset$ ,  $\Theta \Vdash_{\mathcal{B}} \phi$  iff, for every  $\mathcal{C} \supseteq \mathcal{B}$ , if  $\Vdash_{\mathcal{C}} \theta$  for every  $\theta \in \Theta$ , then  $\Vdash_{\mathcal{B}} \phi$

**DEFINITION 3.** (Validity) Define (cf. [32])  $\Gamma \Vdash \phi$  as: for all  $\mathcal{B}$ , if  $\Vdash_{\mathcal{B}} \psi$  for all  $\psi \in \Gamma$ , then  $\Vdash_{\mathcal{B}} \phi$ .

Write  $\Gamma \vdash \phi$  to denote that  $\phi$  is provable from  $\Gamma$  in the natural deduction systems NJ (cf. [32]). For reference, NJ is summarized in Figure 1.

**THEOREM 4.** (Soundness) If  $\Gamma \vdash \phi$ , then  $\Gamma \Vdash \phi$ .



**THEOREM 5.** (Completeness) If  $\Gamma \Vdash \phi$ , then  $\Gamma \vdash \phi$ .

Sandqvist's completeness theorem [32] makes essential use of *flattening*.

**DEFINITION 6.** (The operations  $-^b$  and  $-^{\natural}$ ) Let  $\Delta$  contain all elements of  $\Gamma \cup \{\phi\}$  and their subformulae. With every non-atomic  $\delta \in \Delta$ , associate a distinct atomic  $\delta^b \notin \Delta$  and, for every atomic  $q \in \Delta$ , take  $q^b = q$ . We write  $\Delta^b$  for the flattening of  $\Delta$ . We also require the inverse operation. For any atom  $p$ , define  $p^{\natural}$  to be  $\phi$  if  $\phi^b = p$  and  $p^{\natural} = p$ , otherwise.

Sandqvist [32] defines a special base  $\mathcal{N}$  depending on  $\Delta^b$  as follows:

**DEFINITION 7.** (The base  $\mathcal{N}_{\Delta^b}$ )  $\mathcal{N}_{\Delta^b}$  is defined as the base containing exactly the following rules, corresponding to NJ (Figure 1):

$$\begin{aligned}
\supset\text{I: } & (\phi^b \Rightarrow \psi^b) \Rightarrow (\phi \supset \psi)^b \\
\supset\text{E: } & (\phi \supset \psi)^b, (\Rightarrow \phi^b) \Rightarrow \psi^b \\
\wedge\text{I: } & (\Rightarrow \phi^b), (\Rightarrow \psi^b) \Rightarrow (\phi \wedge \psi)^b \\
\wedge\text{E: } & (\Rightarrow (\phi \wedge \psi)^b) \Rightarrow \phi^b \\
\wedge\text{E: } & (\Rightarrow (\phi \wedge \psi)^b) \Rightarrow \psi^b \\
\vee\text{I: } & (\Rightarrow \phi^b) \Rightarrow (\phi \vee \psi)^b \\
\vee\text{I: } & (\Rightarrow \psi^b) \Rightarrow (\phi \vee \psi)^b \\
\vee\text{E: } & (\Rightarrow (\phi \vee \psi)^b), (\phi^b \Rightarrow p), (\psi^b \Rightarrow p) \Rightarrow p \\
\perp\text{: } & (\Rightarrow \perp^b) \Rightarrow p
\end{aligned}$$

Before embarking on our category-theoretic analysis, we remark that in Appendix A we discuss the relationship between the base  $\mathcal{N}_-$  and how it is that validity is represented by consequence relations.

### 3. A Categorical Interpretation

There is a well-established way of interpreting natural deduction proofs in intuitionistic logic (NJ) in categories that have structure corresponding to the logical connectives [17, 20, 35]. Specifically, we interpret proofs in NJ in bicartesian closed categories, in which products are used to interpret conjunction, exponentials (function spaces) are used to interpret implication, and coproducts are used to interpret disjunction. Thus we obtain an interpretation of the following form: a morphism

$$[[\Gamma]] \xrightarrow{[[\Phi]]} [[\phi]]$$

interprets a proof  $\Phi$  of the consequence  $\Gamma \vdash \phi$ . Here,  $\Phi$  is a proof in NJ of the consequence, represented using the terms of a language such as the typed  $\lambda$ -calculus with products and sums.

In this set-up, we work with judgements of the form

$$x_1 : \phi_1, \dots, x_i : \phi_i, \dots, x_m : \phi_m \vdash \Phi(x_1, \dots, x_m) : \phi$$

which are read, in the sense of the propositions-as-types interpretation, as follows: if the  $x_i$ s are witnesses for proofs of the  $\phi_i$ s, then  $\Phi(x_1, \dots, x_m)$  denotes a proof of  $\phi$  constructed using the rules of NJ.

Now, if  $\Phi_i$  is a specific proof of  $\phi_i$ , then it can be substituted for  $x_i$  throughout this judgement to give

$$x_1 : \phi_1, \dots, x_m : \phi_m \vdash \Phi(x_1, \dots, x_m)[\Phi_i/x_i] : \phi$$

where the assumption  $x_i : \phi_i$  has been removed and the occurrence of  $x_i$  in  $\Phi$  has been replaced by  $\Phi_i$ .

In the setting of proof-theoretic semantics, we are concerned in the first instance with derivations that are restricted to the rules of a base. We introduce terms for derivations — which can be seen as a restricted class of the terms described above — in a base as follows:

$$\Phi := x \mid \Phi_{\mathcal{R}}(\Phi_1, \dots, \Phi_m)$$

where, as in Definition 1,  $\mathcal{R}$  is a rule

$$\frac{\begin{array}{c} [P_1] \quad [P_n] \\ q_1 \quad \dots \quad q_n \\ r \end{array}}{\mathcal{R}}$$

of a base  $\mathcal{B}$ ,  $x$  is a witness for a derivation, and the  $\Phi_i$ s denote, inductively, derivations of atoms  $p$ , as given in Definition 1.

If  $P = p_1, \dots, p_m$  and  $X = x_1, \dots, x_m$ , we write  $(X : P)$  for  $x_1 : p_1, \dots, x_m : p_m$ . Using this notation, well-formed derivations in a base are given inductively by

$$\frac{}{(X : P), x : p \vdash_{\mathcal{B}} x : p} \text{(Ref)}$$

$$\frac{(X : P), (X_i : P_i) \vdash_{\mathcal{B}} \Phi_i : q_i \quad i = 1, \dots, n}{(X : P) \vdash_{\mathcal{B}} \Phi_{\mathcal{R}}(\Phi_1, \dots, \Phi_n) : r} \text{(App}_{\mathcal{R}})$$

Note that in the rule  $\text{App}_{\mathcal{R}}$  the variables  $X_i$  are bound in the right-hand side of the conclusion.

Writing  $\Psi[\Phi/x]$  for the term obtained by substituting  $\Phi$  for  $x$  in  $\Psi$ , we have two key substitution lemmas. First, substitution preserves consequence:

LEMMA 8. If  $(X : P) \vdash_{\mathcal{B}} \Phi : q$  and  $(X : P), y : q \vdash_{\mathcal{B}} \Psi : r$ , then  $(X : P) \vdash_{\mathcal{B}} \Psi[\Phi/y] : r$ .

PROOF. *By induction over the derivation of  $\Psi$ .*

$\Psi = y$  Suppose  $(X : P), y : q \vdash_{\mathcal{B}} y : q$ . By definition,  $\Psi[\Phi/y] = \Phi$ . Hence, by assumption,  $(X : P) \vdash_{\mathcal{B}} \Psi[\Phi/y] : r$ .

$\Psi = \Phi_{\mathcal{R}} (\Phi_1, \dots, \Phi_n)$  By the induction hypothesis,  $(X : P), (X_i : P_i) \vdash_{\mathcal{B}} \Phi_i[\Phi/y] : q_i$ . Hence also  $(X : P) \vdash_{\mathcal{B}} \Phi_{\mathcal{R}}(\Phi_1[\Phi/y], \dots, \Phi_m[\Phi/y]) : r$ . ■

Second, substitution is associative:

LEMMA 9. For all derivations  $(X : P) \vdash_{\mathcal{B}} \Phi_1 : q_1$ ,  $(X : P), y_1 : q_1 \vdash_{\mathcal{B}} \Phi_2 : q_2$ , and  $(X : P), y_2 : q_2 \vdash_{\mathcal{B}} \Psi : r$ , where, without loss of generality,  $y_1 \neq y_2$ , we have

$$\Psi[\Phi_2[\Phi_1/y_1]/y_2] = (\Psi[\Phi_2/y_2])[\Phi_1/y_1]$$

PROOF. *By induction over the structure of  $\Psi$ .* ■

Functor categories of the form  $\mathbf{Set}^{\mathcal{W}^{op}}$ , where the  $(-)^{op}$  operation takes a category and reverses its morphisms, can be used to interpret proofs in NJ. The use of the  $(-)^{op}$  operator is in line with the convention in the category theory community. The reason is that  $\mathcal{W}$  embeds in  $\mathbf{Set}^{\mathcal{W}^{op}}$ , and indeed  $\mathbf{Set}^{\mathcal{W}^{op}}$  can be characterized as a completion of  $\mathcal{W}$ . For more details see Section 4.

For suitable choices of  $\mathcal{W}$ , these functor categories can also be used to interpret derivations in a base.

The basic idea is that the interpretation of an atomic proposition  $p$  in  $\mathbf{Set}^{\mathcal{W}^{op}}$  is the functor whose value at world  $(\mathcal{B}, (X : P))$  is the set of derivations of  $p$  in  $\mathcal{B}$  from hypotheses  $(X : P)$ . The action on morphisms of  $\mathcal{W}$  is given by substitution. We use the following definition:

DEFINITION 10. (Bases and Contexts) Define a category  $\mathcal{W}$  as follows:

- Objects of  $\mathcal{W}$  are pairs  $(\mathcal{B}, (X : P))$ , where  $\mathcal{B}$  is a base and  $(X : P)$  is a context;
- A morphism from  $(\mathcal{B}, (X : P))$  to  $(\mathcal{C}, (Y : Q))$  is given by an inclusion of the base  $\mathcal{C}$  into  $\mathcal{B}$  and a set of derivations  $X : P \vdash_{\mathcal{B}} \Phi_i : q_i$ , where  $Q = \{q_1, \dots, q_m\}$ . We write such a morphism as  $(\Phi_1, \dots, \Phi_m)$ ;
- The identity morphism on  $(X : P)$  is  $(x_1, \dots, x_n)$ ;

- The composition of a morphism  $(\Phi_1, \dots, \Phi_m)$  from  $(X:P)$  to  $(Y:Q)$  and a morphism  $(\Psi_1, \dots, \Psi_k)$  from  $(Y:Q)$  to  $(Z:R)$  is

$$(\Psi_1[\Phi_1/y_1, \dots, \Phi_m/y_m], \dots, \Psi_k[\Phi_1/y_1, \dots, \Phi_m/y_m])$$

Lemma 9 implies that composition is associative.

Now we extend the interpretation of atomic propositions  $p$  to the interpretation of formulae using categorical products for conjunction and exponentials in functor categories for implication. We use products, exponentials, and a quantification over atoms to represent the form of disjunction employed by Sandqvist, which corresponds to its second-order definition (or, alternatively, the elimination rule for  $\vee$  in NJ). See, for example, [3].

Key to understanding our categorical formulation of the exponentials in functor categories is the Yoneda lemma (see [21]): let  $\mathcal{C}$  be a (locally small) category,  $\mathbf{Set}$  be the category of sets, and  $F \in [\mathcal{C}^{op}, \mathbf{Set}]$  (the category of presheaves over  $\mathcal{C}$ ); then, for each object  $C$  of  $\mathcal{C}$ , with  $h^C = \text{hom}(-, C) : \mathcal{C}^{op} \rightarrow \mathbf{Set}$ , the natural transformations  $h^C \rightarrow F$  are in bijection with the elements of  $F(C)$  and this bijection is natural in  $C$  and  $F$ .

We will use the Yoneda lemma to calculate the values of certain functors between presheaf categories. These functors are defined as right adjoints. Given a functor  $G : \mathcal{C} \rightarrow \mathcal{D}$ , a *right adjoint* to  $G$  is a functor  $H : \mathcal{D} \rightarrow \mathcal{C}$ , whose defining characteristic is that  $\mathcal{D}(Gc, d)$  is naturally isomorphic to  $\mathcal{C}(c, Hd)$ . This notion has a dual. Given  $H : \mathcal{D} \rightarrow \mathcal{C}$ , a *left adjoint* is a functor  $G : \mathcal{C} \rightarrow \mathcal{D}$ , whose defining characteristic is that  $\mathcal{D}(Gc, d)$  is naturally isomorphic to  $\mathcal{C}(c, Hd)$ . The Yoneda lemma tells us that if  $F$  is defined as a right adjoint, then its values are certain natural transformations between functors defined from the corresponding left adjoint.

LEMMA 11. Suppose  $L : [\mathcal{C}^{op}, \mathbf{Set}] \rightarrow [\mathcal{D}^{op}, \mathbf{Set}]$  has a right adjoint denoted  $R : [\mathcal{D}^{op}, \mathbf{Set}] \rightarrow [\mathcal{C}^{op}, \mathbf{Set}]$ . Let  $G : \mathcal{D}^{op} \rightarrow \mathbf{Set}$ . Then for any  $C \in \mathcal{C}$ ,  $RG(C)$  can be taken to be the set of natural transformations  $L(\text{hom}(-, C))$  to  $G$ .

PROOF. By the Yoneda lemma,  $RG(C) \cong [\mathcal{C}^{op}, \mathbf{Set}](h^C, RG)$  and, by the adjunction, this is isomorphic to  $[\mathcal{C}^{op}, \mathbf{Set}](L(h^C), G)$ , the set of natural transformations from  $L(\text{hom}(-, C))$  to  $G$ . ■

Shortly, we shall apply this to characterize exponentials, with  $L = (-) \times F$ . In this case, we get that the value of  $F \supset G$  at  $C$  is the set of natural transformations from  $\text{hom}(-, C) \times F$  to  $G$ .

Products in the functor category  $[\mathcal{W}^{op}, \mathbf{Set}]$  are given component-wise. For functors  $F$  and  $G$ , we write  $F \supset G$  for the exponential functor. This functor is defined as the right adjoint to  $(-) \times F$ , using the characterization given above:

$$(F \supset G)(\mathcal{B}, (X : P)) = \text{Nat}(h^{(\mathcal{B}, (X : P))} \times F, G)$$

where  $\text{Nat}(\phi, \psi)$  denotes the set of natural transformations between the set-valued functors  $\phi$  and  $\psi$ .

We can now describe formally how disjunction is treated in terms of products and exponentials. Let  $\mathcal{A}$  be the discrete category of atomic propositions (its set of objects is the set of atomic propositions and its only morphisms are identities). Define a functor  $\Delta_{\mathcal{A}} : (\mathcal{W}^{op} \rightarrow \mathbf{Set}) \rightarrow ((\mathcal{W}^{op} \times \mathcal{A}) \rightarrow \mathbf{Set})$  by mapping a functor  $H : \mathcal{W}^{op} \rightarrow \mathbf{Set}$  to the functor  $H' : \mathcal{W}^{op} \times \mathcal{A} \rightarrow \mathbf{Set}$  such that  $H'((\mathcal{B}, (X : P)), p) = H((\mathcal{B}, (X : P)))$ . Call the right adjoint to this functor  $\forall_{\mathcal{A}}$ . As a right adjoint, we can again characterize its values as natural transformations. Specifically, if  $K : (\mathcal{W}^{op} \times \mathcal{A}) \rightarrow \mathbf{Set}$ , then  $(\forall_{\mathcal{A}} K)(\mathcal{B}, (X : P))$  is the set of natural transformations from  $\Delta_{\mathcal{A}} h^{(\mathcal{B}, (X : P))}$  to  $K$ . Since  $\mathcal{A}$  is discrete, this amounts to giving a natural transformation from  $h^{(\mathcal{B}, (X : P))}$  to  $K(-, p)$  for each atomic proposition  $p$ .

Definition 12 defines the interpretation of formulae in presheaves. The key cases for our purposes are the base case, which captures the derivation of atoms in a base, the case for disjunction, as discussed at length elsewhere, and the case for falsity, which is interpreted as nullary disjunction.

**DEFINITION 12.** (Interpretation Functor) Define a functor  $\llbracket \phi \rrbracket : \mathcal{W}^{op} \rightarrow \mathbf{Set}$  by induction over the structure of  $\phi$  as follows:

- $\llbracket p \rrbracket(\mathcal{B}, (X : P))$  is the set of derivations  $(X : P) \vdash_{\mathcal{B}} \Phi : p$ . Any morphism  $(\Phi_1, \dots, \Phi_m)$  from  $(\mathcal{B}, (X : P))$  to  $(\mathcal{C}, (Y : Q))$  maps a derivation  $(Y : Q) \vdash_{\mathcal{C}} \Phi : p$ , which is also a derivation  $(Y : Q) \vdash_{\mathcal{B}} \Phi : p$ , to the derivation  $(X : P) \vdash_{\mathcal{B}} \Phi[\Phi_1/x_1, \dots, \Phi_m/x_m] : p$ .
- $\llbracket \phi \wedge \psi \rrbracket$  is the product of the functors  $\llbracket \phi \rrbracket$  and  $\llbracket \psi \rrbracket$
- $\llbracket \phi \supset \psi \rrbracket$  is defined as  $\llbracket \phi \rrbracket \supset \llbracket \psi \rrbracket$
- $\llbracket \phi \vee \psi \rrbracket$  is defined as follows: let  $F = \llbracket \phi \rrbracket$ ,  $G = \llbracket \psi \rrbracket$ , and  $K((\mathcal{B}, (X : P)), p) = (F \supset \llbracket p \rrbracket) \supset ((G \supset \llbracket p \rrbracket) \supset \llbracket p \rrbracket)(\mathcal{B}, (X : P))$ . This can be extended to a functor  $\mathcal{W}^{op} \times \mathcal{A} \rightarrow \mathbf{Set}$ . Then  $\llbracket \phi \vee \psi \rrbracket$  is defined as  $\forall_{\mathcal{A}} K$
- $\llbracket \perp \rrbracket$  is defined as follows: let  $K((\mathcal{B}, (X : P)), p) = \llbracket p \rrbracket(\mathcal{B}, (X : P))$ . This can be extended to a functor  $\mathcal{W}^{op} \times \mathcal{A} \rightarrow \mathbf{Set}$ . Then  $\llbracket \perp \rrbracket$  is defined as  $\forall_{\mathcal{A}} K$ .

Note that in this interpretation, unlike in the basic interpretation of NJ proofs in bicartesian closed categories, disjunction does not correspond to coproduct in  $[\mathcal{W}^{op}, \mathbf{Set}]$ . We discuss this point in the sequel.

In Section 5, we establish soundness and completeness for our categorical formulation of Sandqvist's semantics. The proof of soundness uses the

existence of a natural transformation corresponding to  $\Vdash: \Gamma \Vdash \phi$  iff there exists a natural transformation from  $\llbracket \Gamma \rrbracket$  to  $\llbracket \phi \rrbracket$ . The proof of completeness uses a special base, as in [32], which is naturally extended via  $\llbracket - \rrbracket$  to the full consequence relation.

DEFINITION 13. (Categorical Base-extension Validity) We write  $\Gamma \models \phi$  iff there is a natural transformation  $\eta : \llbracket \Gamma \rrbracket \rightarrow \llbracket \phi \rrbracket$ .

For each base  $\mathcal{B}$  we show soundness and completeness relative to this base by considering a suitable full subcategory. Soundness and completeness then arise by considering the empty base.

For any base  $\mathcal{B}$ , we write  $\mathcal{W}_{\mathcal{B}}$  for the full subcategory of  $\mathcal{W}$  where the objects are pairs  $(\mathcal{C}, (Y : Q))$  such that  $\mathcal{B} \subseteq \mathcal{C}$ . For any full subcategory  $\mathcal{W}'$  of  $\mathcal{W}$ , we write  $\llbracket - \rrbracket^{\mathcal{W}'}$  for the functor  $\llbracket - \rrbracket$  restricted to objects and morphisms in  $\mathcal{W}'$ .

In Lemma 14, we assume, without loss of generality, that all  $\phi^b$  in  $\mathcal{N}_{\Delta^b}$  and the  $\phi_1^b, \dots, \phi_n^b$  are distinct from the atoms in  $\mathcal{B}$

LEMMA 14. Let  $\Gamma = \phi_1, \dots, \phi_n$ . Let  $\mathcal{B}' = \mathcal{B} \cup \mathcal{N}_{\Delta^b} \cup \{\Rightarrow \phi_1^b, \dots, \Rightarrow \phi_n^b\}$ . Let  $\mathcal{W}' = \mathcal{W}_{\mathcal{B}'}$ . If  $P \vdash_{\mathcal{B}'} p$ , then there exists a natural transformation  $\eta : \llbracket \Gamma \rrbracket^{\mathcal{W}'} \times \llbracket P^{\natural} \rrbracket^{\mathcal{W}'} \rightarrow \llbracket p^{\natural} \rrbracket^{\mathcal{W}'}$ .

PROOF. *The proof proceeds by induction over the structure of the derivation of  $P \vdash_{\mathcal{B}'} p$  structured by cases on the last rule used in the derivation. For the case of an  $\vee$ -elimination rule, we use again an induction over the structure of the  $\phi$ . Because the semantics of  $\vee$  does not use a coproduct, we cannot use a universal property in this case and by using this induction consider in the proof only all objects which are in the range of the semantics  $\llbracket - \rrbracket$ .*

*Rules in  $\{\Rightarrow \phi_1^b, \dots, \Rightarrow \phi_n^b\}$ :*

- *Suppose the last rule is  $\Rightarrow \phi^b$ . Then  $\phi^{\natural} = \phi$  and the projection from  $\llbracket \Gamma \rrbracket^{\mathcal{W}'} \times \llbracket P^{\natural} \rrbracket^{\mathcal{W}'}$  to  $\llbracket \phi \rrbracket^{\mathcal{W}'}$  yields the claim.*

*Rules in  $\mathcal{B}$ :*

- *Suppose the derivation is  $P, p \vdash_{\mathcal{B}'} p$ . The projection from  $\llbracket P, p^{\natural} \rrbracket^{\mathcal{W}'}$  to  $\llbracket p^{\natural} \rrbracket^{\mathcal{W}'}$  has the desired properties.*
- *Suppose the last rule of the derivation is an application of the rule  $\mathcal{R} = ((P_1 \Rightarrow p_1), \dots, (P_n \Rightarrow p_n)) \Rightarrow p$ . By the induction hypothesis, there are natural transformations  $\eta_i$  from  $\llbracket \Gamma \rrbracket^{\mathcal{W}'} \times \llbracket P_i^{\natural}, P_i \rrbracket^{\mathcal{W}'}$  to  $\llbracket p_i \rrbracket^{\mathcal{W}'}$ . Hence there are also natural transformations  $\eta'_i$  from  $\llbracket \Gamma \rrbracket^{\mathcal{W}'} \times \llbracket P_i^{\natural} \rrbracket^{\mathcal{W}'}$  to  $\llbracket P_i \supset p_i \rrbracket^{\mathcal{W}'}$ . Now consider any object of  $\mathcal{W}'$ . This is a pair  $(\mathcal{C}, (Y : Q))$  such that  $\mathcal{B} \subseteq \mathcal{C}$ . Let  $f$  be an element of  $\llbracket \Gamma \rrbracket^{\mathcal{W}'}(\mathcal{C}, (Y : Q))$  and let  $g$  be an element*

of  $\llbracket P^{\natural} \rrbracket^{\mathcal{W}'}(\mathcal{C}, (Y : Q))$ . Hence  $\eta'_{i(\mathcal{C}, (Y : Q))}(f, g)$  is a natural transformation  $\kappa_i$  from  $\text{Hom}(-, (\mathcal{C}, (Y : Q))) \times \llbracket P_i \rrbracket^{\mathcal{W}'}$  to  $\llbracket p_i \rrbracket^{\mathcal{W}'}$ . Let  $\pi_i$  the projection from  $(Y : Q), (Z_i : P_i)$  to  $P_i$ . By definition,  $\kappa_i(\text{Id}, \pi_i)$  is a derivation  $\Phi_i : (Y : Q), (Z : P_i) \vdash_{\mathcal{C}} p_i$ . Hence  $\Phi_{\mathcal{R}}(\Phi_1, \dots, \Phi_n)$  is a derivation of  $Y : Q \vdash_{\mathcal{C}} p$ .

Rules in  $\mathcal{N}_{\Delta^b}$ :

- Suppose the last rule is  $\psi_1^b, \psi_2^b \Rightarrow (\psi_1 \wedge \psi_2)^b$ . By the induction hypothesis, there are two natural transformations  $\eta_1$  and  $\eta_2$ , where  $\eta_i$  is a natural transformation from  $\llbracket \Gamma \rrbracket^{\mathcal{W}'} \times \llbracket P^{\natural} \rrbracket^{\mathcal{W}'}$  to  $\llbracket \psi_i \rrbracket^{\mathcal{W}'}$ . Hence  $\langle \eta_1, \eta_2 \rangle$  is a natural transformation from  $\llbracket \Gamma \rrbracket^{\mathcal{W}'} \times \llbracket P^{\natural} \rrbracket^{\mathcal{W}'}$  to  $\llbracket \psi_1 \wedge \psi_2 \rrbracket^{\mathcal{W}'}$ .
- Suppose the last rule is  $(\psi_1 \wedge \psi_2)^b \Rightarrow \psi_1^b$ . By the induction hypothesis, there exists a natural transformation  $\eta$  from  $\llbracket \Gamma \rrbracket^{\mathcal{W}'} \times \llbracket P^{\natural} \rrbracket^{\mathcal{W}'}$  to  $\llbracket \psi_1 \wedge \psi_2 \rrbracket^{\mathcal{W}'}$ . Now compose this natural transformation with the projection from  $\llbracket \psi_1 \wedge \psi_2 \rrbracket^{\mathcal{W}'}$  to  $\llbracket \psi_1 \rrbracket^{\mathcal{W}'}$ .
- Suppose the last rule is  $(\psi_1^b \Rightarrow \psi_2^b) \Rightarrow (\psi_1 \supset \psi_2)^b$ . By assumption, there is a natural transformation from  $\llbracket \Gamma \rrbracket^{\mathcal{W}'} \times \llbracket P^{\natural} \rrbracket^{\mathcal{W}'}$  to  $\llbracket \psi_2 \rrbracket^{\mathcal{W}'}$ . Hence there is also a natural transformation from  $\llbracket \Gamma \rrbracket^{\mathcal{W}'} \times \llbracket P^{\natural} \rrbracket^{\mathcal{W}'}$  to  $\llbracket \psi_1 \supset \psi_2 \rrbracket^{\mathcal{W}'}$ .
- Suppose the last rule is  $\psi_1^b, (\psi_1 \supset \psi_2)^b \Rightarrow \psi_2^b$ . By assumption, we have natural transformations  $\eta_1$  and  $\eta_2$  from  $\llbracket \Gamma \rrbracket^{\mathcal{W}'} \times \llbracket P^{\natural} \rrbracket^{\mathcal{W}'}$  to  $\llbracket \psi_1 \rrbracket^{\mathcal{W}'}$  and  $\llbracket \psi_1 \supset \psi_2 \rrbracket^{\mathcal{W}'}$  respectively. By the definition of function spaces for functor categories, there exists also a natural transformation from  $\llbracket \Gamma \rrbracket^{\mathcal{W}'} \times \llbracket P^{\natural} \rrbracket^{\mathcal{W}'}$  to  $\llbracket \psi_2 \rrbracket^{\mathcal{W}'}$ .
- Suppose the last rule is  $\psi_1^b \Rightarrow (\psi_1 \vee \psi_2)^b$ . By definition of the interpretation of disjunction there is a natural transformation  $\kappa$  from  $\llbracket \psi_1 \rrbracket^{\mathcal{W}'}$  to  $\llbracket \psi_1 \vee \psi_2 \rrbracket^{\mathcal{W}'}$ . By the induction hypothesis, there is a natural transformation  $\eta$  from  $\llbracket \Gamma \rrbracket^{\mathcal{W}'} \times \llbracket P^{\natural} \rrbracket^{\mathcal{W}'}$  to  $\llbracket \psi_1 \rrbracket^{\mathcal{W}'}$ .  $\kappa \circ \eta$  is therefore a natural transformation from  $\llbracket \Gamma \rrbracket^{\mathcal{W}'} \times \llbracket P^{\natural} \rrbracket^{\mathcal{W}'}$  to  $\llbracket \psi_1 \vee \psi_2 \rrbracket^{\mathcal{W}'}$ .
- Suppose the last rule is  $(\psi_1 \vee \psi_2)^b, (\psi_1^b \Rightarrow p), (\psi_2^b \Rightarrow p) \Rightarrow p$ . By the induction hypothesis, there are natural transformations  $\eta_{\psi_1 \vee \psi_2}$  from  $\llbracket \Gamma \rrbracket^{\mathcal{W}'} \times \llbracket P^{\natural} \rrbracket^{\mathcal{W}'}$  to  $\llbracket \psi_1 \vee \psi_2 \rrbracket^{\mathcal{W}'}$ ,  $\eta_{\psi_1}$  from  $\llbracket \Gamma \rrbracket^{\mathcal{W}'} \times \llbracket P^{\natural} \rrbracket^{\mathcal{W}'}$  to  $\llbracket p \rrbracket^{\mathcal{W}'}$  and  $\eta_{\psi_2}$  from  $\llbracket \Gamma \rrbracket^{\mathcal{W}'} \times \llbracket P^{\natural} \rrbracket^{\mathcal{W}'}$  to  $\llbracket p \rrbracket^{\mathcal{W}'}$ . Now we use an induction over the structure of  $p^{\natural}$ .
  - $p^{\natural} = p$ : By the definition of the interpretation of disjunction there is also a natural transformation from  $\llbracket \psi_1 \vee \psi_2 \rrbracket^{\mathcal{W}'} \times \llbracket \psi_1 \supset p \rrbracket^{\mathcal{W}'}$  to

- $\llbracket \psi_2 \supset p \rrbracket^{\mathcal{W}'}$  to  $\llbracket p \rrbracket^{\mathcal{W}'}$ . Hence there is also a natural transformation from  $\llbracket \Gamma \rrbracket^{\mathcal{W}'} \times \llbracket P^{\natural} \rrbracket^{\mathcal{W}'}$  to  $\llbracket p \rrbracket^{\mathcal{W}'}$ .
- $p^{\natural} = \phi_1 \wedge \phi_2$ : We also have  $P, \psi_1^{\flat} \vdash_{\mathcal{B}'} \phi_i^{\flat}$  and  $P, \psi_2^{\flat} \vdash_{\mathcal{B}'} \phi_i^{\flat}$  for  $i = 1, 2$ . By the induction hypothesis, there are natural transformations  $\eta_i$  from  $\llbracket \Gamma \rrbracket^{\mathcal{W}'} \times \llbracket P^{\natural} \rrbracket^{\mathcal{W}'}$  to  $\llbracket \phi_i \rrbracket^{\mathcal{W}'}$ . The natural transformation  $\langle \eta_1, \eta_2 \rangle$  is a natural transformation from  $\llbracket \Gamma \rrbracket^{\mathcal{W}'} \times \llbracket P^{\natural} \rrbracket^{\mathcal{W}'}$  to  $\llbracket \phi_1 \wedge \phi_2 \rrbracket^{\mathcal{W}'}$ .
  - $p^{\natural} = \phi_1 \supset \phi_2$ : We also have  $P, \phi_1^{\flat}, \psi_i^{\flat} \vdash_{\mathcal{B}'} \phi_2^{\flat}$  and  $P, \phi_1^{\flat} \vdash_{\mathcal{B}'} (\psi_1 \vee \psi_2)^{\flat}$ . By the induction hypothesis, there is a natural transformation from  $\llbracket \Gamma \rrbracket^{\mathcal{W}'} \times \llbracket P^{\natural} \rrbracket^{\mathcal{W}'} \times \llbracket \phi_1 \rrbracket^{\mathcal{W}'}$  to  $\llbracket \phi_2 \rrbracket^{\mathcal{W}'}$ . Hence there is also a natural transformation from  $\llbracket \Gamma \rrbracket^{\mathcal{W}'} \times \llbracket P^{\natural} \rrbracket^{\mathcal{W}'}$  to  $\llbracket \phi_1 \supset \phi_2 \rrbracket^{\mathcal{W}'}$ .
  - $p^{\natural} = \phi_1 \vee \phi_2$ : For any atom  $q$  and any  $F_1: \llbracket \phi_1 \rrbracket^{\mathcal{W}'} \supset \llbracket q \rrbracket^{\mathcal{W}'}$ ,  $F_2: \llbracket \phi_2 \rrbracket^{\mathcal{W}'} \supset \llbracket q \rrbracket^{\mathcal{W}'}$ , we define a natural transformation  $\mu^{\psi_1}: \llbracket \Gamma \rrbracket^{\mathcal{W}'} \times \llbracket P^{\natural} \rrbracket^{\mathcal{W}'} \times \llbracket \psi_1 \rrbracket^{\mathcal{W}'} \rightarrow \llbracket q \rrbracket^{\mathcal{W}'}$  by  $\mu^{\psi_1}(\gamma, f, t) = \eta_{\psi_1}(\gamma, f, t)pF_1F_2$ . We similarly define a natural transformation  $\mu^{\psi_2}: \llbracket \Gamma \rrbracket^{\mathcal{W}'} \times \llbracket P^{\natural} \rrbracket^{\mathcal{W}'} \times \llbracket \psi_2 \rrbracket^{\mathcal{W}'} \rightarrow \llbracket q \rrbracket^{\mathcal{W}'}$  by  $\mu^{\psi_2}(\gamma, f, t) = \eta_{\psi_2}(\gamma, f, t)pF_1F_2$ . Now we define, using an informal  $\lambda$ -calculus notation a natural transformation  $\eta: \llbracket \Gamma \rrbracket^{\mathcal{W}'} \times \llbracket P^{\natural} \rrbracket^{\mathcal{W}'}$  to  $\llbracket \phi_1 \vee \phi_2 \rrbracket^{\mathcal{W}'}$  by

$$\eta(\gamma, f) = \lambda q. \lambda F_1: \llbracket \phi_1 \rrbracket^{\mathcal{W}'} \supset \llbracket q \rrbracket^{\mathcal{W}'}. \lambda F_2: \llbracket \phi_2 \rrbracket^{\mathcal{W}'} \supset \llbracket q \rrbracket^{\mathcal{W}'}. \\ \eta_{\psi_1 \vee \psi_2}(\gamma, f)p(\text{Cur}(\mu^{\phi})(\gamma, f))(\text{Cur}(\mu^{\psi})(\gamma, f))$$

■

LEMMA 15. (Algebraic Soundness) Suppose  $\Gamma \Vdash_{\mathcal{B}} \phi$ . Let  $\mathcal{W}'$  be the category  $\mathcal{W}_{\mathcal{B}}$ . Then there exists a natural transformation  $\eta_{\mathcal{B}}: \llbracket \Gamma \rrbracket^{\mathcal{W}'} \rightarrow \llbracket \phi \rrbracket^{\mathcal{W}'}$ .

PROOF. Let  $\Gamma = \phi_1, \dots, \phi_n$ . Let  $\mathcal{B}' = \mathcal{B} \cup \mathcal{N}_{\Delta^{\flat}} \cup \{\Rightarrow \phi_1^{\flat}, \dots, \Rightarrow \phi_n^{\flat}\}$ . Using the proof of Sandqvist's theorem 5.1, we have  $\Vdash_{\mathcal{B}'} \Gamma$ . Hence we have by assumption also  $\Vdash_{\mathcal{B}'} \phi$ , and, using the proof of Sandqvist's theorem 5.1 again, we have  $\Vdash_{\mathcal{B}'} \phi^{\flat}$ . Lemma 14 now yields the claim. ■

Now we turn to completeness, which is formulated as the converse of soundness.

LEMMA 16. (Algebraic Completeness) Consider any base  $\mathcal{B}$ . Let  $\mathcal{W}'$  be the category  $\mathcal{W}_{\mathcal{B}}$ . If there exists a natural transformation  $\eta_{\mathcal{B}}: \llbracket \Gamma \rrbracket^{\mathcal{W}'} \rightarrow \llbracket \phi \rrbracket^{\mathcal{W}'}$ , then  $\Gamma \Vdash_{\mathcal{B}} \phi$ .

PROOF. The proof proceeds by induction over the structure of  $\Gamma$  and  $\phi$ .

$\Gamma = P$ ,  $\phi = p$  Let  $id$  be the identity on  $P$ . By definition,  $\eta_{\mathcal{B}, (\mathcal{B}, (X:P))}(id)$  is a derivation — i.e., a proof according to Definition 2 —  $P \Vdash_{\mathcal{B}} p$ .



$\phi = \phi_1 \wedge \phi_2$  By definition, there exists natural transformations  $\eta_i: [\Gamma]^{W'} \rightarrow [\phi_i]^{W'}$ , and hence, by the induction hypothesis, we have  $\Gamma \Vdash_{\mathcal{B}} \phi_i$  and therefore  $\Gamma \Vdash_{\mathcal{B}} \phi_1 \wedge \phi_2$ .

$\Gamma = \Delta, \phi_1 \wedge \phi_2$  By the induction hypothesis, there exists a derivation  $\Delta, \phi_1, \phi_2 \Vdash_{\mathcal{B}} \phi$ , which is the required derivation, as  $[\Delta, \phi_1 \wedge \phi_2] = [\Delta, \phi_1, \phi_2]$ .

$\phi = \phi_1 \supset \phi_2$  By the definition of  $\supset$  as a right adjoint, there exists a natural transformation  $\eta': [\Gamma, \phi_1]^{W'} \rightarrow [\phi]^{W'}$ . By the induction hypothesis, there exists a derivation  $\Gamma, \phi_1 \Vdash_{\mathcal{B}} \phi_2$ , and therefore also a derivation  $\Gamma \Vdash_{\mathcal{B}} \phi_1 \supset \phi_2$ .

$\Gamma = \Delta, \phi_1 \supset \phi_2$  Suppose  $\Vdash_{\mathcal{C}} \Gamma$ . By algebraic soundness, for all  $\mathcal{D}$  such that  $\mathcal{C} \subseteq \mathcal{D}$  and  $P$  there are elements  $\delta$  of  $[\Delta](\mathcal{D}, (X:P))$  and  $f$  of  $[\phi_1 \supset \phi_2](\mathcal{D}, (X:P))$ .  $\eta_{\mathcal{B}, (\mathcal{D}, (X:P))}(\delta, f)$  is an element of  $[\phi](\mathcal{D}, (X:P))$ . This assignment is natural in  $(\mathcal{D}, (X:P))$ . By the induction hypothesis, we have  $\Vdash_{\mathcal{C}} \phi$ .

$\phi = \phi_1 \vee \phi_2$  By the definition of  $[-]$ , for all atoms  $p$  there exists a natural transformation  $\eta_{\mathcal{B}}$  from  $[\Gamma]^{W'}$  to  $[\phi_1 \vee \phi_2]^{W'}$  iff for all atoms  $p$  there exists a natural transformation  $\eta_{p, \mathcal{B}}$  from  $[\Gamma]^{W'}$  to  $[\phi_1 \supset p \supset (\phi_2 \supset p) \supset p]^{W'}$ . Hence the induction hypothesis yields  $\Gamma \Vdash_{\mathcal{B}} (\phi_1 \supset p) \supset (\phi_2 \supset p) \supset p$ . Hence, by definition,  $\Gamma \Vdash_{\mathcal{B}} \phi_1 \vee \phi_2$ .

$\Gamma = \Delta, \phi_1 \vee \phi_2$  By induction over the structure of  $\phi$ .

$\phi = p$  Suppose  $\Vdash_{\mathcal{B}} \Gamma$ . By algebraic soundness, there exists a natural transformation  $\eta': [\top]^{W'} \rightarrow [\Gamma]^{W'}$ . By definition  $\eta_{\mathcal{B}} \circ \eta'$  is a natural transformation from  $[\top]^{W'}$  to  $[p]^{W'}$ . As  $[\top](\mathcal{B}, (-: -))$  is non-empty,  $[p](\mathcal{B}, (-: -))$  is non-empty as well, and hence by definition of  $[-]$ , there exists a derivation  $\Vdash_{\mathcal{B}} p$ .

$\phi = \psi_1 \wedge \psi_2$  By definition, there exists natural transformations  $\eta_1$  and  $\eta_2$  such that  $\eta = \langle \eta_1, \eta_2 \rangle$ . By the induction hypothesis, there are derivations  $\Gamma \Vdash_{\mathcal{B}} \psi_1$  and  $\Gamma \Vdash_{\mathcal{B}} \psi_2$ . Therefore we also have  $\Gamma \Vdash_{\mathcal{B}} \psi_1 \wedge \psi_2$ .

$\phi = \psi_1 \supset \psi_2$  By definition, there exists also a natural transformation  $\eta'_{\mathcal{B}}$  from  $[\Gamma, \psi_1]^{W'}$  to  $[\psi_2]^{W'}$ . By the induction hypothesis, there exists a derivation  $\Gamma, \psi_1 \Vdash_{\mathcal{B}} \psi_2$ . Hence there is also a derivation  $\Gamma \Vdash_{\mathcal{B}} \psi_1 \supset \psi_2$ .

$\phi = \psi_1 \vee \psi_2$  Consider any atom  $p$ . By definition of  $[-]$ , there exists a natural transformation  $\eta'$  from  $[\Gamma]^{W'}$  to  $[(\psi_1 \supset p) \supset (\psi_2 \supset p) \supset p]^{W'}$ .

*By the induction hypothesis, there is a derivation  $\Gamma \Vdash_{\mathcal{B}} (\psi_1 \supset p) \supset (\psi_2 \supset p) \supset p$ . Hence there is also a derivation  $\Gamma \Vdash_{\mathcal{B}} \psi_1 \vee \psi_2$ .*

■

PROPOSITION 17. (Equivalence)  $\Gamma \Vdash \phi$  iff  $\Gamma \models \phi$ .

PROOF. *Direct consequence of Lemmas 15 and 16.*

■

## 4. Relation to Kripke, Presheaf, and Sheaf Models

The previous section has set out the details of a categorical account of Sandqvist’s proof-theoretic semantics. In this section, we situate that account with reference to classical Kripke models, to the algebraic treatment of intuitionistic logic via complete Heyting algebras, and to the categorical treatment of higher-order type theory via categories of presheaves and sheaves.

The treatment of conjunction and implication is standard. The treatment of disjunction, however, is not, and the close analysis of this exposes that there is a distinction between the proof-insensitive algebraic semantics and the proof-sensitive categorical semantics. In the algebraic semantics, the soundness of Sandqvist’s definition of disjunction can be established by exposing it as the natural disjunction in a sublocale (quotient complete Heyting algebra) of the locale obtained from the standard Kripke model (Definition 30, Lemma 33). The analogue in the categorical setting is sheaves for the corresponding (Grothendieck or Joyal-Tierney) topology. However, there is no real reason for the interpretations of atomic propositions to be sheaves, and the definition of disjunction is not the categorical coproduct.

The soundness of the interpretation instead comes from a structural induction, not a property of arbitrary sheaves (Lemmas 25, 15 and Proposition 27). This is unusual in categorical interpretations of type theory, and so noteworthy.

Classical Kripke models are grounded in a set of worlds,  $\mathcal{K}$ . For each atomic predicate  $p$ , we are told whether  $p$  holds in world  $w$ :  $w \Vdash p$ . When interpreting intuitionistic logic, the set of worlds can be thought of as possible states of knowledge. This set is partially ordered with the ordering representing increasing knowledge. This viewpoint gives rise to and is reflected in the monotonicity property: if  $w \leq w'$  and  $w \Vdash p$ , then  $w' \Vdash p$ .

The definition of validity is extended to arbitrary propositions by structural induction:

$$\begin{aligned}
(\supset) \quad w \Vdash \phi \supset \psi & \quad \text{iff} \quad \text{for all } w \leq w', \text{ if } w' \Vdash \phi, \text{ then } w' \Vdash \psi \\
(\wedge) \quad w \Vdash \phi \wedge \psi & \quad \text{iff} \quad w \Vdash \phi \text{ and } w \Vdash \psi \\
(\perp) \quad w \Vdash \perp & \quad \text{never} \\
(\vee) \quad w \Vdash \phi \vee \psi & \quad \text{iff} \quad w \Vdash \phi \text{ or } w \Vdash \psi
\end{aligned}$$

The monotonicity property then extends to all formulae. The quantification over extensions in the clause for  $(\supset)$  is required in order to ensure this.

Base-extension semantics relates closely to Kripke models. In the simplest formulation (ours is a little more complex), the set of worlds,  $\mathcal{K}$  is the set of bases being used, and the partial order is simply inclusion of bases. We define  $\mathcal{B} \Vdash p$  iff  $\vdash_{\mathcal{B}} p$ . This satisfies the monotonicity property because if  $\mathcal{B} \subseteq \mathcal{C}$  and  $\vdash_{\mathcal{B}} p$ , then  $\vdash_{\mathcal{C}} p$ . There are two main approaches to extending this definition for atomic formulas to general ones:

- the approach of Schroeder-Heister et al. (cf. [25]), which follows exactly the Kripke definitions
- the approach of Sandqvist [32], which follows the Kripke definitions for implication and conjunction, but not disjunction or false.

However, Kripke models also have a more algebraic interpretation.

LEMMA 18. In any Kripke model, the interpretation of an arbitrary proposition  $\phi$  satisfies the monotonicity property: if  $w \leq w'$  and  $w \Vdash \phi$ , then  $w' \Vdash \phi$ .

As a consequence, the interpretation of any proposition  $\phi$  also satisfies the monotonicity property and so is an upwards-closed subset of  $\mathcal{K}$ .

DEFINITION 19. Let  $\Omega$  be the set of upwards-closed subsets of  $\mathcal{K}$ ; that is,

$$\Omega = \{U \subseteq \mathcal{K} \mid \text{if } \mathcal{B} \in U \text{ and } \mathcal{B} \subseteq \mathcal{C}, \text{ then } \mathcal{C} \in U\}$$

$\Omega$  is partially ordered by inclusion.

LEMMA 20.  $\Omega$  is a (complete) Heyting algebra, and the interpretation of the logical connectives in that algebra coincides with that in the Kripke model.

Moreover, if  $w \in \mathcal{K}$ , then  $w^\uparrow = \{w' \in \mathcal{K} \mid w \leq w'\}$  is in  $\Omega$ . However, if  $w_0 \leq w_1$  then  $w_1^\uparrow \subseteq w_0^\uparrow$ . So it is  $\mathcal{K}^{op}$  that embeds in  $\Omega$ , not  $\mathcal{K}$ . Indeed,  $\Omega$  can be seen as a free completion under arbitrary disjunctions of  $\mathcal{K}^{op}$ . This means that Kripke models can be seen as simply a convenient way of presenting a complete Heyting algebra.

The standard Kripke account gives an interpretation of validity, but not really of proof, or justification. In order to obtain that, we need to move to a setting which can distinguish between proofs that are different. One way of doing that is to use presheaves.

DEFINITION 21. If  $\mathbf{C}$  is a category, then a *presheaf on  $\mathbf{C}$*  is a functor  $F : \mathbf{C}^{op} \rightarrow \mathbf{Set}$ , and a *morphism of presheaves* is a natural transformation between the functors.

Categories of presheaves generalize Kripke models. To cast a Kripke model in terms of presheaves, let  $\mathbf{C} = \mathcal{K}^{op}$ , so  $\mathbf{C}^{op} = \mathcal{K}$ . As a result the presheaf category corresponding to the Kripke model is actually  $\mathbf{Set}^{\mathcal{K}}$ . Here we take the standard interpretation of a partial order as a category: the elements of the partial order form the objects of the category, and the hom-set  $\mathcal{K}(w_0, w_1)$  contains a single element if  $w_0 \leq w_1$  and is empty otherwise.

In making the connection between Kripke and presheaves, a simple but important presheaf is the presheaf  $1$ , defined by  $1(w) = \{*\}$ . If  $\phi$  is a sub-presheaf of this, then each  $\phi(w)$  will either be  $\emptyset$  or  $\{*\}$ . And if  $w_0 \leq w_1$ , then there is a function  $\phi(w_0) \rightarrow \phi(w_1)$ . As a result, if  $\phi(w_0)$  is non-empty, then so is  $\phi(w_1)$ . Hence  $\phi$  corresponds to an upwards-closed subset of  $\mathcal{K}$ . Conversely, if  $U$  is an upwards-closed subset of  $\mathcal{K}$ , then we can define  $\chi_U : \mathcal{K} \rightarrow \mathbf{Set}$  by

- $\chi_U(w) = \{*\}$  if  $w \in U$
- $\chi_U(w) = \emptyset$  otherwise

$\chi_U$  is a subfunctor of  $1$ . For the case of a category of presheaves corresponding to a Kripke model, this demonstrates an order-preserving correspondence between the possible interpretations of propositions in the Kripke model and the subfunctors of  $1$  in the presheaf category. We have established:

LEMMA 22. There is an order-preserving isomorphism between

- $\Omega$ , The lattice of upwards-closed subsets of  $\mathcal{K}$ , and
- The lattice of subfunctors of  $1$  in the presheaf topos  $\mathbf{Set}^{\mathcal{K}}$ .

This means that either of these lattices can be used interchangeably as the target of a validity-based semantics for intuitionistic propositional logic.

However categories of presheaves include functors targeting sets with more than one element. They do not just provide models of intuitionistic propositional logic, they form models of simple intuitionistic type theory. They carry interpretations of type-forming operations including products and sums.

In the following, let  $F$  and  $G$  be two functors  $\mathbf{C}^{op} \rightarrow \mathbf{Set}$ :

- The product of  $F$  and  $G$  in the category of presheaves is the functor  $F \times G$  defined on objects by  $F \times G(A) = F(A) \times_{\mathbf{Set}} G(A)$ , where the action on morphisms is component-wise.
- The categorical function space  $[F \rightarrow G]$  is defined on objects as follows:  $[F \rightarrow G](A)$  is the set of natural transformations from  $F \times \mathbf{C}(-, A)$  to  $G$ . The action on morphisms is obtained via composition.
- The coproduct of  $F$  and  $G$  is the functor  $(F + G)(A) = F(A) +_{\mathbf{Set}} G(A)$ , where  $+_{\mathbf{Set}}$  is the coproduct (i.e., disjoint union) in  $\mathbf{Set}$ .

If  $\mathbf{C} = \mathcal{K}^{op}$  is a partial order, and  $F$  and  $G$  are subfunctors of  $1$ , then these definitions correspond to the interpretations of  $\wedge$ ,  $\supset$ , and  $\vee$  in the corresponding Kripke model.

Moreover, there are links to the categorical interpretation in Section 3. The constructions given there are internal constructions in the topos of presheaves:  $\mathcal{W}^{op} \rightarrow \mathbf{Set}$ . We recall from Definition 12, that the interpretation of an atom  $p$  is a functor  $\mathcal{W}^{op} \rightarrow \mathbf{Set}$ , and that for arbitrary propositions,  $\phi$  and  $\psi$ , the interpretations of  $\phi \wedge \psi$  and  $\phi \supset \psi$  are defined to be  $\llbracket \phi \rrbracket \times \llbracket \psi \rrbracket$  and  $\llbracket \phi \rrbracket \supset \llbracket \psi \rrbracket$  respectively.

LEMMA 23. For atoms,  $p$  and arbitrary propositions  $\phi$  and  $\psi$ , the interpretations defined in Definition 12 have the following properties:

- The interpretation of an atom  $p$  is a functor  $\llbracket p \rrbracket : \mathcal{W}^{op} \rightarrow \mathbf{Set}$
- The interpretation of  $\phi \wedge \psi$  is  $\llbracket \phi \rrbracket \times \llbracket \psi \rrbracket$
- The interpretation of  $\phi \supset \psi$  is  $\llbracket \phi \rrbracket \rightarrow \llbracket \psi \rrbracket$ .

However, the interpretation of  $\phi \vee \psi$  is not the coproduct  $\llbracket \phi \rrbracket + \llbracket \psi \rrbracket$  in the presheaf topos  $\mathcal{W}^{op} \rightarrow \mathbf{Set}$ . Instead it is the interpretation of the second order formula:

$$\forall p. (\llbracket \phi \rrbracket \rightarrow \llbracket p \rrbracket) \rightarrow (\llbracket \psi \rrbracket \rightarrow \llbracket p \rrbracket) \rightarrow \llbracket p \rrbracket$$

There are two equivalent ways of formalizing this intuition.

- We can take the collection of atomic propositions to be an external set, and the quantification as the conjunction of all the  $(\llbracket \phi \rrbracket \rightarrow \llbracket p \rrbracket) \rightarrow (\llbracket \psi \rrbracket \rightarrow \llbracket p \rrbracket) \rightarrow \llbracket p \rrbracket$ , using the fact that the lattice of propositions has infinite meets.
- We can internalize the collection of atomic propositions as the constant presheaf  $\Delta\mathbb{A}$ , where  $(\Delta\mathbb{A})(w) = \mathbb{A}$ . We interpret  $(\llbracket \phi \rrbracket \rightarrow \llbracket p \rrbracket) \rightarrow (\llbracket \psi \rrbracket \rightarrow \llbracket p \rrbracket) \rightarrow \llbracket p \rrbracket$  as an internal family of propositions indexed by  $\Delta\mathbb{A}$ , and

interpret the  $\forall p$  as the internal quantification in the logic of the topos, which in this case is the product:  $\prod_p . (\llbracket \phi \rrbracket \rightarrow \llbracket p \rrbracket) \rightarrow (\llbracket \psi \rrbracket \rightarrow \llbracket p \rrbracket) \rightarrow \llbracket p \rrbracket$ .

The second of these gives rise to the construction detailed in Section 3.

It follows that the structure we have defined in Section 3 is naturally a model of  $F_{at}$ , a restriction of the second-order lambda calculus with function types and polymorphism restricted to be over a collection of atomic types. The restriction of  $\forall$  to a predefined set means that  $F_{at}$  is predicative, in contrast to the full System  $F$  [14, 15], and it further means that it can be interpreted in a broadly set-theoretic setting where the universal types are interpreted as products over the set of atomic types. This is exactly what our categorical model is doing. There is therefore a strong link between the structures here and those investigated extensively by Ferreira et al. [5–9, 26]. However, Ferreira’s treatment is essentially syntactic, while ours is strongly semantic. In particular, the types of  $F_{at}$  exist in a wider context. However, our proof of Lemma 25 can be seen as a semantic equivalent of Ferreira’s ‘instantiation overflow’ for the case of disjunction.

This interpretation supports the introduction rules for disjunction: for any objects  $A$  and  $B$  of the topos there are morphisms  $A \rightarrow \forall p.(A \rightarrow \llbracket p \rrbracket) \rightarrow (B \rightarrow \llbracket p \rrbracket) \rightarrow \llbracket p \rrbracket$  and  $B \rightarrow \forall p.(A \rightarrow \llbracket p \rrbracket) \rightarrow (B \rightarrow \llbracket p \rrbracket) \rightarrow \llbracket p \rrbracket$ . But there is not a similarly general interpretation of the elimination rule. Instead, we get an interpretation where one of the variables is restricted to range over interpretations of formulae rather than arbitrary objects of the topos.

Consider the disjunction elimination rule:

$$\frac{\Gamma \vdash \phi \vee \psi \quad \Gamma, \phi \vdash \chi \quad \Gamma, \psi \vdash \chi}{\Gamma \vdash \chi}_{\vee E}$$

We want to encode this rule in terms of the structure of the topos. To do that, we consider an interpretation where the metavariables  $\phi$ ,  $\psi$ , and  $\chi$  represent objects of the topos rather than formulae. Replacing  $\phi$ ,  $\psi$ , and  $\chi$  by  $A$ ,  $B$ , and  $C$ , we see that we want a morphism

$$\sigma(A, B) \times C^A \times C^B \rightarrow C$$

where  $\sigma$  is a binary operation on objects of the topos giving the semantics of disjunction.

Specifically, Sandqvist’s semantics uses  $\sigma(A, B) = \forall p.(A \rightarrow \llbracket p \rrbracket) \rightarrow (B \rightarrow \llbracket p \rrbracket) \rightarrow \llbracket p \rrbracket$ , which is defined in any sheaf topos. For this interpretation, if  $\chi$  is interpreted by an arbitrary object, then the rule is not sound. However, if the object that interprets  $\chi$  is constructed propositionally from the

interpretations of atoms, and so is the interpretation of a proposition, then soundness is recovered.

DEFINITION 24. We say that a type constructor  $\sigma(A, B)$  supports disjunction elimination for an object  $C$  of the topos if there is a morphism

$$\sigma(A, B) \times [A \rightarrow C] \times [B \rightarrow C] \longrightarrow C$$

The objects for which a constructor  $\sigma$  supports disjunction elimination have good closure properties.

LEMMA 25. 1.  $\sigma$  always supports disjunction elimination for the terminal object  $1$ ; that is, when  $C$  is  $1$ , the constant functor taking value  $\{*\}$ .

2. If  $\sigma$  supports disjunction elimination for  $C_0$  and  $C_1$ , then  $\sigma$  supports disjunction elimination for  $C_0 \times C_1$ .
3. If  $\sigma$  supports disjunction elimination for  $C$ , then  $\sigma$  supports disjunction elimination for  $[D \rightarrow C]$ .
4. If  $\sigma$  supports disjunction elimination for any  $C_x$  where  $x \in X$ , then  $\sigma$  supports disjunction elimination for  $\prod_{x \in X} C_x$  (and a similar but more complex statement for internal products in the topos).

PROOF. *The proofs are all straightforward manipulation of the cartesian closed structure of the topos.*

1. We require a morphism  $\sigma(A, B) \times [A \rightarrow 1] \times [B \rightarrow 1] \rightarrow 1$ , which exists uniquely because  $1$  is terminal.
2. We require a morphism

$$\sigma(A, B) \times [A \rightarrow (C_0 \times C_1)] \times [B \rightarrow (C_0 \times C_1)] \rightarrow (C_0 \times C_1)$$

but  $[A \rightarrow (C_0 \times C_1)]$  is isomorphic to  $[A \rightarrow C_0] \times [A \rightarrow C_1]$ , and similarly for  $[B \rightarrow (C_0 \times C_1)]$ . This means that the required morphism can easily be constructed from the morphisms we are given

$$\sigma(A, B) \times [A \rightarrow C_0] \times [B \rightarrow C_0] \rightarrow C_0$$

and  $\sigma(A, B) \times [A \rightarrow C_1] \times [B \rightarrow C_1] \rightarrow C_1$ .

3. Similarly, we require a morphism

$$\sigma(A, B) \times [A \rightarrow [D \rightarrow C]] \times [B \rightarrow [D \rightarrow C]] \rightarrow [D \rightarrow C]$$

but this can be constructed from the exponential of the map

$$\sigma(A, B) \times [A \rightarrow C] \times [B \rightarrow C] \rightarrow C$$

4. This is essentially as 2. ■

LEMMA 26. Let  $\sigma(A, B) = \prod_p(A \rightarrow \llbracket p \rrbracket) \rightarrow (B \rightarrow \llbracket p \rrbracket) \rightarrow \llbracket p \rrbracket$ , then  $\sigma$  supports disjunction elimination for any object which is the semantics of a formula:  $\llbracket \phi \rrbracket$ .

PROOF. *The proof is by induction on the structure of  $\phi$ . The base case is  $\phi = q$  where  $q$  is an atom. The required morphism*

$$\left( \prod_p (A \rightarrow \llbracket p \rrbracket) \rightarrow (B \rightarrow \llbracket p \rrbracket) \rightarrow \llbracket p \rrbracket \right) \times (A \rightarrow \llbracket q \rrbracket) \times (B \rightarrow \llbracket q \rrbracket) \rightarrow \llbracket q \rrbracket$$

*is projection onto the  $q$  component followed by evaluation. The inductive cases for conjunction and implication follow immediately from Lemma 25 as the interpretations are given by product and exponentiation in the topos. The inductive case for disjunction also follows from the same lemma, since if  $\phi = \psi \vee \chi$ , then  $\llbracket \phi \rrbracket = \prod_p (\llbracket \psi \rrbracket \rightarrow \llbracket p \rrbracket) \rightarrow (\llbracket \chi \rrbracket \rightarrow \llbracket p \rrbracket) \rightarrow \llbracket p \rrbracket$ . ■*

It is not the case that arbitrary objects of the topos support disjunction elimination. Specifically, the object  $0$ , which corresponds to the constant functor with value the empty set does not do so. If  $A = B = C = 0$ , then  $\sigma(A, B) = \prod_p(A \rightarrow \llbracket p \rrbracket) \rightarrow (B \rightarrow \llbracket p \rrbracket) \rightarrow \llbracket p \rrbracket$  is isomorphic to  $\prod_p \llbracket p \rrbracket$ , while  $[A \rightarrow C]$  and  $[B \rightarrow C]$  are both isomorphic to  $1$ , but  $\prod_p \llbracket p \rrbracket$  may be non-empty and hence there may be no morphism  $\prod_p \llbracket p \rrbracket \rightarrow 0$ .

Soundness of the interpretation is now immediate:

PROPOSITION 27. Given a proof in intuitionistic propositional logic of  $\Gamma \vdash \phi$  then we can inductively define a morphism in the presheaf topos (a natural transformation between functors)  $[\Gamma] \rightarrow [\phi]$ . The induction is based on both the structure of the proof, and the syntactic structure of the formulae within it.

This soundness result sits in contrast to the soundness of the traditional interpretation of intuitionistic proof theory in topoi. In that case, the result allows proofs including propositional variables, which are to be interpreted as arbitrary objects of the topos. In our case, we are restricted to proofs with formulae built from a collection of atoms, each of which has a defined semantics as a particular object of the topos.

There is, however a way of escaping this restriction. We will do this in two stages. First we consider a semantics based on validity, and then one which is proof relevant.

For the validity-based semantics we take a semantics derived from our current semantics, but we equate all the elements of each  $\llbracket \phi \rrbracket w$ .

DEFINITION 28. We define the validity-based semantics as follows

$$\{\llbracket \phi \rrbracket\}w = \{ * \mid \exists x. x \in \llbracket \phi \rrbracket w \}$$



- LEMMA 29. 1.  $\{\phi\}$  is a subfunctor of the functor  $1$ , where  $1w = \{*\}$ .
2.  $\{\phi \wedge \psi\} = \{\phi\} \times \{\psi\}$
  3.  $\{\phi \supset \psi\} = \{\phi\} \rightarrow \{\psi\}$
  4.  $\{\phi\}w = \{*\}$  iff  $w \Vdash \phi$  in the sense of Sandqvist [32].

Now consider the map on objects of the topos, sending  $U$  to  $\prod_p(U \rightarrow \{p\}) \rightarrow \{p\}$ . If  $U$  is a subfunctor of  $1$ , then  $\prod_p(U \rightarrow \{p\}) \rightarrow \{p\}$  is itself isomorphic to a subfunctor of  $1$ .

DEFINITION 30. If  $U$  is a subfunctor of  $1$ , then let  $KU$  be the subfunctor of  $1$  isomorphic to  $\prod_p(U \rightarrow \{p\}) \rightarrow \{p\}$ .  $K$  is an operator on subfunctors of  $1$ .

LEMMA 31. For all subfunctors  $U$  and  $V$  of  $1$ :

1.  $U$  is a subfunctor of  $KU$ .
2.  $KKU$  is a subfunctor of  $KU$ , and hence  $K$  is idempotent ( $K^2 = K$ ).
3.  $K(U \times V) = KU \times KV$ .

It follows that  $K$  is a nucleus on the locale of subfunctors of  $1$  ordered by inclusion, see Johnstone [18], and hence that its fixpoints define a sublocale.

DEFINITION 32. A subfunctor  $U$  of  $1$  is said to be *closed* if  $KU = U$ . We write  $\Omega_K$  for the set of closed subfunctors of  $1$  ordered by pointwise inclusion.

LEMMA 33. 1. For any atomic proposition  $p$ ,  $\{p\}$  is closed.

2.  $1$  is closed.
3. If  $U$  and  $V$  are closed, then so is  $U \times V$ .
4. If  $U$  is closed and  $W$  is any subfunctor of  $1$ , then  $W \rightarrow U$  is closed.
5. An arbitrary product of closed subfunctors of  $1$  is closed.
6. If  $U$  and  $V$  are closed, then the least closed subfunctor of  $1$  containing both  $U$  and  $V$  is  $\prod_p(U \rightarrow \{p\}) \rightarrow (V \rightarrow \{p\}) \rightarrow \{p\}$ , and this is therefore the join of  $U$  and  $V$  in  $\Omega_K$ .
7.  $\Omega_K$  is a complete Heyting algebra (and hence an algebra supporting the interpretation of Intuitionistic Propositional Logic).

PROOF. (2–5) hold for arbitrary kernels, and follow from the properties of 31. (1) and (6) are specific to  $K$ , but follow from lemma 31 and basic properties of types and intuitionistic logic.

Specifically, (1) follows from lemma 31 (1) and the fact that if  $\forall p.(\{q\} \rightarrow \{p\}) \rightarrow \{p\}$ , then in particular  $(\{q\} \rightarrow \{q\}) \rightarrow \{q\}$ , and since  $\{q\} \rightarrow \{q\}$  always holds, then  $\{q\}$ . This establishes that  $K\{q\}$  is a subfunctor of  $\{q\}$  as well as conversely, and hence is equal to it.

Similarly for (6) we suppose  $U \leq W$ ,  $V \leq W$  and that  $W$  is closed. We show that

$$\prod_p (U \rightarrow \{p\}) \rightarrow (V \rightarrow \{p\}) \rightarrow \{p\} \leq \prod_p (W \rightarrow \{p\}) \rightarrow \{p\} = W$$

The proof is componentwise, for any  $p$ , we show that

$$(U \rightarrow \{p\}) \rightarrow (V \rightarrow \{p\}) \rightarrow \{p\} \leq (W \rightarrow \{p\}) \rightarrow \{p\}$$

This follows by monotonicity. Since  $U \leq W$ ,  $W \rightarrow \{p\} \leq U \rightarrow \{p\}$ , and since  $V \leq W$ ,  $W \rightarrow \{p\} \leq V \rightarrow \{p\}$ . Hence  $(U \rightarrow \{p\}) \rightarrow (V \rightarrow \{p\}) \rightarrow \{p\}$  is contained in  $(W \rightarrow \{p\}) \rightarrow (W \rightarrow \{p\}) \rightarrow \{p\}$  which is equal to  $(W \rightarrow \{p\}) \rightarrow \{p\}$ . ■

$\Omega_K$  therefore provides a model of intuitionistic propositional logic, with the connectives interpreted as standard (meet, join, heyting implication, etc). Moreover there is an immediate connection with Sandqvist's interpretation of Intuitionistic Propositional Logic. The interpretations of the connectives are exactly as in Sandqvist's definitions, and for any formula  $\phi$ , we have  $\{\phi\}w = \{*\}$  if and only if  $w \Vdash \phi$ . Sandqvist's interpretation of disjunction is the join operator in  $\Omega_K$ .

This can be further extended to a proof-sensitive account. The nucleus  $K$  on subfunctors of 1 internalizes to an endomorphism of the object  $\Omega$  of truth values for the topos of presheaves:

$$k = \lambda\omega. \forall p. (\omega \rightarrow \{p\}) \rightarrow \{p\} : \Omega \longrightarrow \Omega$$

The properties that make  $K$  a nucleus, then become the properties for  $k$  to be a Joyal-Tierney topology (see [21]), and we can obtain a proof-relevant model valued in the topos of sheaves:  $\text{Sh}_k(\mathbf{Set}^{\mathcal{W}^{op}})$ . The subobjects of 1 in this sheaf model are precisely the closed subfunctors of 1 for the nucleus  $k$ .

The sheaves for a topology form a full subcategory of the presheaf topos, closed under products, exponentials, and limits generally. This gives the category of sheaves limits and exponentials. The equalizer of  $k$  and the identity on  $\Omega$  gives a subobject of  $\Omega$ , which acts as the subobject classifier for the category of sheaves. The category of sheaves therefore has coproducts. These, however, are not the same as the coproducts in the category of presheaves. This structure can be used to give a proof-relevant model.

However, critically, the interpretation of atomic propositions,  $\llbracket p \rrbracket$ , that we have given does not in general form sheaves, and hence the model that we have given is not actually a sheaf-theoretic one.

To see why not, let's suppose we restrict to a setting in which we have two atomic propositions,  $p$  and  $q$ . There are eight possible first-order rules for these propositions. Four of them make sense from a proof-theoretic perspective:

$$\Rightarrow p, \quad \Rightarrow q, \quad q \Rightarrow p, \quad p \Rightarrow q$$

The remaining four are valid in format, but do not enable us to derive any new information:

$$p \Rightarrow p, \quad q \Rightarrow q, \quad p, q \Rightarrow p, \quad p, q \Rightarrow q$$

A base is given by a subset of these, and there are therefore  $2^8 = 256$  bases. The minimal bases in which both  $p$  and  $q$  are valid are:

$$\begin{aligned} &\Rightarrow p, \quad \Rightarrow q \\ &\Rightarrow p, \quad p \Rightarrow q \\ &\Rightarrow q, \quad q \Rightarrow p \end{aligned}$$

The base  $\{\Rightarrow p, \Rightarrow q\}$  generates one proof of  $p$ , the proof via  $\Rightarrow p$ . The base  $\{\Rightarrow q, q \Rightarrow p\}$  generates a different proof of  $p$ . Both of these bases are contained in  $\{\Rightarrow p, \Rightarrow q, q \Rightarrow p\}$ , which contains both proofs. As a result, it is impossible to choose a single proof of  $p$  that is valid for all bases that prove both  $p$  and  $q$ . However, this is precisely what we would get if  $\llbracket p \rrbracket$  were a sheaf in this example.

Those familiar with program semantics will recognise the construct  $((\ ) \rightarrow \{p\}) \rightarrow \{p\}$  as an instance of  $((\ ) \rightarrow R) \rightarrow R$ , and so a form of continuation semantics. In [29], John Reynolds quotes Christopher Wadsworth as describing continuations as ‘the meaning of the rest of the program’. Program semantics faces the issue that the meaning of a program is intended to be derived structurally, and so in terms of the meaning of its parts. However, it is not possible to run part of a program, only a complete one. As a result, to give meaning to part of a program, you have to say what that part will do in the context of its completion by the rest of a program; that is, in terms of what it will do given a continuation. In the  $((\ ) \rightarrow R) \rightarrow R$  construct,  $R$  represents the end result of the program.  $(\ ) \rightarrow R$  represents the continuation, something that will convert the meaning of a fragment into a final result, and  $((\ ) \rightarrow R) \rightarrow R$  expresses that the meaning of a program fragment is supposed to be something that takes a continuation and produces a final result. However, there is a hole here: if the meaning of a program fragment

is to be  $(X \rightarrow R) \rightarrow R$ , then this suggests continuations should have type  $((X \rightarrow R) \rightarrow R) \rightarrow R$ , and program fragments should have another iteration, opening up an infinite loop; but, in intuitionistic type theory, there is a natural morphism

$$(((X \rightarrow R) \rightarrow R) \rightarrow R) \longrightarrow (X \rightarrow R)$$

allowing escape from the recursion. The analogy for us is direct. The meaning of the proof of a non-atomic formula  $\phi$  is taken to be a mechanism that, given any atomic formula  $p$  and the rest of a proof of  $p$  from  $\phi$ , gives a proof of  $p$ . So, the complete proofs that are the analogues of programs are the proofs of atomic formulas, not proofs of arbitrary ones.

## 5. Logical Metatheory

We now have all that is required to complete our categorical account of Sandqvist's proof-theoretic semantics for intuitionistic propositional logic.

In Section 3, we established the (algebraic) soundness and completeness of our categorical set-up with respect to Sandqvist's formulation of validity. In this section, we give the two key meta-theoretic results, namely soundness and completeness with respect to the natural deduction system NJ. We defer a discussion of the choices around the interpretation of disjunction to Section 6.

### 5.1. Soundness

We establish the soundness of NJ [27], as given in Figure 1, with respect to proof-theoretic validity. The proof proceeds by induction over the structure of proofs in NJ. While this result does not depend directly upon the soundness and completeness results for the algebraic interpretation, it nevertheless makes essential use of the characterization of validity in terms of natural transformations.

**THEOREM 34.** (Soundness) If  $\Gamma \vdash \phi$ , then  $\Gamma \models \phi$ .

**PROOF.** *The proof, which is very similar to that of Lemma 15, is by induction over the structure of the derivation of  $\Gamma \vdash \phi$ . The critical cases are disjunction introduction and elimination.*

$\forall I$  *The rules are*

$$\frac{\Gamma \vdash \phi}{\Gamma \vdash \phi \vee \psi} \quad \text{and} \quad \frac{\Gamma \vdash \psi}{\Gamma \vdash \phi \vee \psi}$$

We consider only the first rule; the proof for second rule is similar. By assumption, there is a natural transformation  $\eta_\phi: [\Gamma] \rightarrow [\phi]$ . We define, using an informal  $\lambda$ -calculus notation, a natural transformation  $\eta: [\Gamma] \rightarrow [\phi \vee \psi]$  by

$$\eta(\gamma) = \lambda p. \lambda F_1: [\phi] \supset [p]. \lambda F_2: [\psi] \supset [p]. F_1(\eta_\phi \gamma)$$

$\vee E$  The rule is

$$\frac{\Gamma \vdash \phi \vee \psi \quad \Gamma, \phi \vdash \chi \quad \Gamma, \psi \vdash \chi}{\Gamma \vdash \chi}$$

By the induction hypothesis, there are natural transformations  $\eta_\phi: [\Gamma, \phi] \rightarrow [\chi]$ ,  $\eta_\psi: [\Gamma, \psi] \rightarrow [\chi]$  and  $\eta_{\phi \vee \psi}: [\Gamma] \rightarrow [\phi \vee \psi]$ . For any natural transformation  $\eta: [\Gamma] \times [\chi] \rightarrow [\sigma]$ , we write  $Cur(\eta): [\Gamma] \rightarrow [\chi] \supset [\sigma]$  for the natural transformation obtained by applying the adjunction defining function spaces to  $\eta$ .

We use now an induction over  $\chi$ .

$\chi = p, p$  atom We define a natural transformation  $\eta: [\Gamma] \rightarrow [\chi]$  by  $\eta(\gamma) = \eta_{\phi \vee \psi} p (Cur(\eta_\phi) \gamma) (Cur(\eta_\psi) \gamma)$ .

$\chi = \chi_1 \wedge \chi_2$  We also have  $\Gamma, \phi \vdash \chi_i$  and  $\Gamma \vdash \chi_i$  for  $i = 1, 2$ . Hence, by the induction hypothesis, there are natural transformations  $\eta_i: [\Gamma] \rightarrow [\chi_i]$  for  $i = 1, 2$ . The natural transformation  $\langle \eta_1, \eta_2 \rangle$  is a natural transformation from  $[\Gamma]$  to  $[\chi_1 \wedge \chi_2]$ .

$\chi = \chi_1 \supset \chi_2$  We also have  $\Gamma, \chi_1, \phi \vdash \chi_2$ ,  $\Gamma, \chi_1, \psi \vdash \chi_2$  and  $\Gamma, \chi_1 \vdash \phi \vee \psi$ . By induction hypothesis there is a natural transformation  $\eta: [\Gamma] \times [\chi_1] \rightarrow [\chi_2]$ . The natural transformation  $Cur(\eta)$  is a natural transformation from  $[\Gamma]$  to  $[\chi_1 \supset \chi_2]$ .

$\chi = \chi_1 \vee \chi_2$  For any atom  $p$  and any  $F_1: [\chi_1] \supset [p]$ ,  $F_2: [\chi_2] \supset [p]$ , we define a natural transformation  $\mu^\phi: [\Gamma] \times [\phi] \rightarrow [p]$  by  $\mu^\phi(\gamma, t) = \eta_\phi(\gamma, t) p F_1 F_2$ . We similarly define a natural transformation  $\mu^\psi: [\Gamma] \times [\psi] \rightarrow [p]$  by  $\mu^\psi(\gamma, t) = \eta_\psi(\gamma, t) p F_1 F_2$ . Now we define, again using an informal  $\lambda$ -calculus notation, a natural transformation  $\eta: [\Gamma] \rightarrow [\chi_1 \vee \chi_2]$  by

$$\eta(\gamma) = \lambda p. \lambda F_1: [\chi_1] \supset [p]. \lambda F_2: [\chi_2] \supset [p]. \eta_{\phi \vee \psi} \gamma p (Cur(\mu^\phi) \gamma) (Cur(\mu^\psi) \gamma)$$

■

## 5.2. Completeness

We now establish the completeness of proof-theoretic validity with respect to NJ. The proof begins with a construction that is analogous to the construction of a term model, but one which is with respect to (proof-theoretic) validity. The completeness statement then follows by an appeal to naturality.

For completeness, we seek to establish that if  $\Gamma \Vdash \phi$ , then  $\Gamma \vdash \phi$ . We make essential use of Sandqvist's flattening and the associated base  $\mathcal{N}_-$ , as defined in Definition 7 of Section 2.

LEMMA 35. Let  $\phi$  be any formula. Let  $\Delta^b$  be the flattening of  $\phi$ . Let  $\mathcal{N}_{\Delta^b}$  be the special base given by Definition 7. Let  $\mathcal{B}$  be any base such that  $\mathcal{B} \supseteq \mathcal{N}_{\Delta^b}$ . For any set of atoms  $Q$ , let  $\mathcal{W}' = \mathcal{W}_{\mathcal{B},(Y:Q)}$ .

1. There exists a natural transformation from  $\llbracket \phi \rrbracket^{\mathcal{W}'}$  to  $\llbracket \phi^b \rrbracket^{\mathcal{W}'}$ .
2. There exists a natural transformation from  $\llbracket \phi^b \rrbracket^{\mathcal{W}'}$  to  $\llbracket \phi \rrbracket^{\mathcal{W}'}$ .

PROOF. We prove both statements simultaneously by induction over the structure of  $\phi$ .

1.  $\phi = p$  By definition,  $p^b = p$ , hence the identity function has the desired properties.

$\phi = \phi_1 \wedge \phi_2$  Let  $f$  be an element of  $\llbracket \phi_1 \wedge \phi_2 \rrbracket^{\mathcal{W}'}(\mathcal{C}, (X:P))$ . Hence  $\pi_1(f)$  and  $\pi_2(f)$  are elements of  $\llbracket \phi_1 \rrbracket^{\mathcal{W}'}(\mathcal{C}, (X:P))$  and  $\llbracket \phi_2 \rrbracket^{\mathcal{W}'}(\mathcal{C}, (X:P))$ , respectively, where  $\pi_1$  and  $\pi_2$  are the projections from  $\llbracket \phi_1 \wedge \phi_2 \rrbracket^{\mathcal{W}'}$  to  $\llbracket \phi_1 \rrbracket^{\mathcal{W}'}$  and  $\llbracket \phi_2 \rrbracket^{\mathcal{W}'}$ , respectively. By the induction hypothesis, there exist functions  $\eta_1 : \llbracket \phi_1 \rrbracket^{\mathcal{W}'} \rightarrow \llbracket \phi_1^b \rrbracket^{\mathcal{W}'}$  and  $\eta_2 : \llbracket \phi_2 \rrbracket^{\mathcal{W}'} \rightarrow \llbracket \phi_2^b \rrbracket^{\mathcal{W}'}$ . By definition,  $\eta_1(\pi_1(f))$  and  $\eta_2(\pi_2(f))$  are derivations  $P \vdash_{\mathcal{C}} \phi_1^b$  and  $P \vdash_{\mathcal{C}} \phi_2^b$ , respectively. Using the  $\wedge I$ -rule of  $\mathcal{N}_{\Delta^b}$ , there is also a derivation  $P \vdash_{\mathcal{C}} (\phi_1 \wedge \phi_2)^b$ .

$\phi = \phi_1 \supset \phi_2$  Let  $f$  be an element of  $\llbracket \phi_1 \supset \phi_2 \rrbracket^{\mathcal{W}'}(\mathcal{C}, (X:P))$ . We start by showing that  $P, \phi_1^b \vdash_{\mathcal{C}} \phi_2^b$ . Suppose that  $\vdash_{\mathcal{D}} P$  and  $\vdash_{\mathcal{D}} \phi_1^b$ , for  $\mathcal{D} \supseteq \mathcal{C}$ . Then we also have  $P \vdash_{\mathcal{D}} \phi_1^b$ . By the induction hypothesis, there exists a function  $\eta_1$  from  $\llbracket \phi_1^b \rrbracket^{\mathcal{W}'}(\mathcal{D}, (X:P))$  to  $\llbracket \phi_1 \rrbracket^{\mathcal{W}'}(\mathcal{D}, (X:P))$ . Hence there is also an element  $g$  of  $\llbracket \phi_1 \rrbracket^{\mathcal{W}'}(\mathcal{D}, (X:P))$ . Therefore,  $f(\iota, g)$ , where  $\iota$  is the reverse inclusion of  $\mathcal{C}$  into  $\mathcal{D}$ , is an element of  $\llbracket \phi_2 \rrbracket^{\mathcal{W}'}(\mathcal{D}, (X:P))$ . By the induction hypothesis, there is also a function  $\eta_2$  from  $\llbracket \phi_2 \rrbracket^{\mathcal{W}'}(\mathcal{D}, X:P)$  to  $\llbracket \phi_2^b \rrbracket^{\mathcal{W}'}(\mathcal{D}, (X:P))$ . Hence there is a derivation  $P \vdash_{\mathcal{D}} \phi_2^b$ , and therefore by cut, also a derivation  $\vdash_{\mathcal{D}} \phi_2^b$ . The  $\supset I$ -rule of  $\mathcal{N}_{\Delta^b}$  now produces a derivation  $P \vdash_{\mathcal{C}} (\phi_1 \supset \phi_2)^b$ .

$\phi = \phi_1 \vee \phi_2$  Let  $f$  be an element of  $\llbracket \phi_1 \vee \phi_2 \rrbracket^{\mathcal{W}'}(\mathcal{C}, (X:P))$ . By definition, there exists also an element of  $\llbracket (\phi_1 \supset (\phi_1 \vee \phi_2)^b) \supset (\phi_2 \supset (\phi_1 \vee \phi_2)^b) \supset (\phi_1 \vee \phi_2^b) \rrbracket^{\mathcal{W}'}(\mathcal{C}, (X:P))$ . By the induction hypothesis, there exists also a derivation  $P \vdash_{\mathcal{C}} (\phi_1^b \supset (\phi_1 \vee \phi_2)^b) \supset (\phi_2^b \supset (\phi_1 \vee \phi_2)^b) \supset (\phi_1 \vee \phi_2)^b$ . Now an application of the  $\vee I$ -rules of  $\mathcal{N}_{\Delta^b}$  yields  $P \vdash_{\mathcal{C}} (\phi_1 \vee \phi_2)^b$ .

2.  $\phi = p$  By definition,  $p^b = p$ , hence the identity function has the desired properties.

$\phi = \phi_1 \wedge \phi_2$  Suppose  $P \vdash_{\mathcal{C}} (\phi_1 \wedge \phi_2)^b$ . Using the  $\wedge E$ -rule of  $\mathcal{N}_{\Delta^b}$  we obtain derivations  $\Phi_1$  of  $P \vdash_{\mathcal{C}} \phi_1^b$  and  $\Phi_2$  of  $P \vdash_{\mathcal{C}} \phi_2^b$  respectively. By the induction hypothesis, there exist functions  $\eta_1 : \llbracket \phi_1^b \rrbracket^{\mathcal{W}'}(\mathcal{C}, (X:P)) \rightarrow \llbracket \phi_1 \rrbracket^{\mathcal{W}'}(\mathcal{C}, (X:P))$  and  $\eta_2 : \llbracket \phi_2^b \rrbracket^{\mathcal{W}'}(\mathcal{C}, (X:P)) \rightarrow \llbracket \phi_2 \rrbracket^{\mathcal{W}'}(\mathcal{C}, (X:P))$ . Hence  $(\eta_1(\Phi_1), \eta_2(\Phi_2))$  is an element of  $\llbracket \phi_1 \wedge \phi_2 \rrbracket^{\mathcal{W}'}(\mathcal{C}, (X:P))$ .

$\phi = \phi_1 \supset \phi_2$  By the definition of function spaces, it suffices to show the existence of a function from  $\llbracket (\phi_1 \supset \phi_2)^b \rrbracket^{\mathcal{W}'}(\mathcal{C}, (X:P))$ ,  $\llbracket \phi_1 \rrbracket^{\mathcal{W}'}(\mathcal{C}, (X:P))$  to  $\llbracket \phi_2 \rrbracket^{\mathcal{W}'}(\mathcal{C}, (X:P))$ . Let  $\Phi$  be a derivation  $P \vdash_{\mathcal{C}} (\phi_1 \supset \phi_2)^b$  and  $f$  be an element of  $\llbracket \phi_1 \rrbracket^{\mathcal{W}'}(\mathcal{C}, (X:P))$ . By the induction hypothesis, there exists a function  $\eta_1 : \llbracket \phi_1 \rrbracket^{\mathcal{W}'}(\mathcal{C}, (X:P)) \rightarrow \llbracket \phi_1^b \rrbracket^{\mathcal{W}'}(\mathcal{C}, (X:P))$ , and therefore also a derivation  $P \vdash_{\mathcal{C}} \phi_1^b$ . The  $\supset E$ -rule of  $\mathcal{N}_{\Delta^b}$  now yields a derivation of  $P \vdash_{\mathcal{C}} \phi_2^b$ . By the induction hypothesis, there exists a function  $\eta_2 : \llbracket \phi_2^b \rrbracket^{\mathcal{W}'}(\mathcal{C}, (X:P)) \rightarrow \llbracket \phi_2 \rrbracket^{\mathcal{W}'}(\mathcal{C}, (X:P))$ , and therefore also an element of  $\llbracket \phi_2 \rrbracket^{\mathcal{W}'}(\mathcal{C}, (X:P))$ .

$\phi = \phi_1 \vee \phi_2$  It suffices to show that, for all atoms  $p$ , there is a function from  $\llbracket (\phi_1 \vee \phi_2)^b \rrbracket^{\mathcal{W}'}(\mathcal{C}, (X:P))$ ,  $\llbracket \phi_1 \supset p \rrbracket^{\mathcal{W}'}(\mathcal{C}, (X:P))$ ,  $\llbracket \phi_2 \supset p \rrbracket^{\mathcal{W}'}(\mathcal{C}, (X:P))$  to  $\llbracket p \rrbracket^{\mathcal{W}'}(\mathcal{C}, (X:P))$ . Suppose  $P \vdash_{\mathcal{C}} (\phi_1 \vee \phi_2)^b$  and let  $f_1$  and  $f_2$  be elements of  $\llbracket \phi_1 \supset p \rrbracket^{\mathcal{W}'}(\mathcal{C}, (X:P))$  and  $\llbracket \phi_2 \supset p \rrbracket^{\mathcal{W}'}(\mathcal{C}, (X:P))$  respectively. By the induction hypothesis, there exists also derivations  $P \vdash_{\mathcal{C}} (\phi_1 \supset p)^b$  and  $P \vdash_{\mathcal{C}} (\phi_2 \supset p)^b$ . The  $\vee E$ -rule of  $\mathcal{N}_{\Delta^b}$  now yields a derivation of  $P \vdash_{\mathcal{C}} p$ .

This completes the proof. ■

**THEOREM 36.** (Completeness) If  $\Gamma \models \phi$ , then  $\Gamma \vdash \phi$ .

**PROOF.** Let  $\Delta^b$  be the flattening of  $\Gamma$  and  $\phi$ . Let  $\mathcal{N}_{\Delta^b}$  be the special base given by Definition 7 for this set of atoms. By Lemma 35, there exists a function  $\eta$  from  $\llbracket \Gamma^b \rrbracket(\mathcal{N}_{\Delta^b}, (X : \Gamma^b))$  to  $\llbracket \phi^b \rrbracket(\mathcal{N}_{\Delta^b}, (X : \Gamma^b))$ . By definition,  $\eta(id)$  is a derivation  $\Gamma^b \vdash_{\mathcal{N}_{\Delta^b}} \phi^b$ . Theorem 5.1 of Sandqvist now implies  $\Gamma \vdash \phi$  in NJ. ■

## 6. Disjunction and Its Interpretation

There is only one interpretation of implication in any given semantics. This is not the case for disjunction. Often there are several ways of interpreting disjunction for a particular semantics with sometimes significantly different properties. In this section, we discuss possible alternatives for interpreting disjunction and give a justification for the interpretation of disjunction used for base-extension semantics.

The validity of formulae is commonly defined by an induction over the structure of formulae. The clause for disjunction usually states — for example, in elementary Kripke-style semantics — that a disjunction  $\phi \vee \psi$  is valid iff  $\phi$  or  $\psi$  are valid. This clause is problematic for a notion of validity using a proof-theoretic semantics, as completeness does not hold, as shown in [24]: specifically, the authors show that with this definition of validity, Harrop’s rule is valid; however, Harrop’s rule is not derivable in intuitionistic logic.

From the inferentialist perspective, Kripke’s clause is too strong because it assumes that the suasive content of a disjunction is identical to that of its disjuncts. However, Sandqvist’s treatment, corresponding to the  $\vee$ -elimination rule of NJ for disjunction, expresses that whatever can be inferred from both disjuncts can be inferred from the disjunction. It is this correspondence that allows completeness to go through.

Categorically, disjunction is typically interpreted as a coproduct. But coproducts are defined as a left adjoint. From the logical perspective, this means they are determined by the elimination rule, and hence that a characterization in terms of the introduction rule does not fit with the categorical interpretation, absent some strong properties on the indecomposability of formulae.

Instead, then, the validity of a disjunction is given by the elimination rule: if  $\phi \vee \psi$  is valid, and for all formulae  $\chi$ , if  $\phi$  is valid implies  $\chi$  is valid and if  $\psi$  is valid implies  $\chi$  is valid, then  $\chi$  is valid. Sandqvist’s definition of validity for disjunction restricts  $\chi$  to atoms (cf. [37,38]). In this way, we obtain an inductive definition of validity. Our categorical interpretation models this definition of validity. Completeness can be obtained by using a basis  $\mathcal{N}$ , which has a separate atom denoting each formula. It follows that the quantification over atoms captures quantification over all formulae.

We can also use the categorical definition of validity — as set up in Definition 13 — to show that the interpretation of disjunction by coproducts leads to incompleteness by proving that completeness would imply that the



rule

$$\frac{p \supset q \vee r}{(p \supset q) \vee (p \supset r)}$$

was derivable. This rule is a strong disjunction property and it is not derivable in IPC. The argument in the proof of the theorem is a reformulation of an argument by Sandqvist [32]:

**THEOREM 37.** Suppose in Definition 12, the definitional clause for  $\llbracket \phi \vee \psi \rrbracket$  is replaced by

–  $\llbracket \phi \vee \psi \rrbracket$  is the coproduct  $\llbracket \phi \rrbracket + \llbracket \psi \rrbracket$ .

Then there is a natural transformation from  $\llbracket p \supset (q \vee r) \rrbracket$  to  $\llbracket (p \supset q) \vee (p \supset r) \rrbracket$ .

**PROOF.** Consider the rule

$$\frac{p \supset q \vee r}{(p \supset q) \vee (p \supset r)}$$

Now consider any  $(\mathcal{B}, (X : P))$  that satisfies  $p \supset (q \vee r)$ , which means  $\llbracket p \supset (q \vee r) \rrbracket(\mathcal{B}, (X : P))$  is non-empty. Hence there is a natural transformation  $\eta$  from  $\mathcal{W}(-, (\mathcal{B}, (X : P))) \times \llbracket p \rrbracket$  to  $\llbracket q \vee r \rrbracket$ . Let  $\mathcal{C}$  be the base that is obtained by adding the rule  $\Rightarrow p$  to  $\mathcal{B}$ . Let  $\Phi$  the derivation of  $P \vdash_{\mathcal{C}} p$  in  $\mathcal{C}$  obtained by applying this rule. Let  $\iota$  be the reverse inclusion map from  $(\mathcal{C}, P)$  to  $(\mathcal{B}, P)$ . Hence,  $\eta_{(\mathcal{C}, P)}(\iota, \Phi)$  is an element of  $\llbracket q \vee r \rrbracket(\mathcal{C}, (X : P)) = \llbracket q \rrbracket(\mathcal{C}, (X : P)) + \llbracket r \rrbracket(\mathcal{C}, (X : P))$ . Therefore,  $\eta_{(\mathcal{C}, P)}(\iota, \Phi)$  is an element of  $\llbracket q \rrbracket(\mathcal{C}, (X : P))$  or  $\llbracket r \rrbracket(\mathcal{C}, (X : P))$ .

Now consider the first case of this disjunction; that is, in which there is a derivation of  $X : P \vdash_{\mathcal{C}} q$ . We must show that there is an element of  $\llbracket p \supset q \rrbracket(\mathcal{B}, (X : P)) = \text{Nat}(\mathcal{W}(-, (\mathcal{B}, (X : P))) \times \llbracket p \rrbracket, \llbracket q \rrbracket)$ . Consider any base  $\mathcal{D} \supseteq \mathcal{B}$  and set of atoms  $Q$  such that  $\Phi : Y : Q \vdash_{\mathcal{D}} P$  and there exists a derivation  $\Psi : Y : Q \vdash_{\mathcal{D}} p$ .

Now we show by induction over the derivation of  $Z : S, X : P \vdash_{\mathcal{C}} s$  that there exists a derivation  $Z : S, Y : Q \vdash_{\mathcal{D}} s$ .

For any rule other than  $\Rightarrow p$ , the substitution of  $\Phi$  for  $X$  yields a derivation  $Z : S, Y : Q \vdash_{\mathcal{D}} s$ . Now suppose the last rule is  $Z : S, X : P \Rightarrow p$ . By assumption, there is also a derivation of  $Z : S, Y : Q \vdash_{\mathcal{D}} p$ .

By a similar argument, we show that in the second case there is an element of  $\llbracket p \supset r \rrbracket(\mathcal{B}, (X : P))$ . It therefore follows that there is also an element of

$$(\llbracket p \supset q \rrbracket \vee \llbracket p \supset r \rrbracket)(\mathcal{B}, (X : P))$$

■

It follows that completeness cannot hold if disjunction is interpreted by coproducts (cf. the distinction between coproducts in presheaves and sheaves, as discussed in Section 4). This proof can be generalized by replacing the atom  $p$  by any formula  $\phi = p_n \supset \cdots \supset p_1 \supset p_0$ , where all  $p_i$ s are atoms. The modification of the proof consists of replacing  $\llbracket p \rrbracket$  by  $\llbracket p_n \supset \cdots \supset p_1 \supset p_0 \rrbracket$  and using this natural transformation instead of the rule  $\Rightarrow p$  to construct a derivation of  $Z : S, Y : Q \vdash_{\mathcal{D}} s$ .

The proof exploits the fact that the extensions in a base extension semantics have more structure. In particular, for each base  $\mathcal{B}$  and for each formula  $\phi$  as specified above one can construct a rule that corresponds to adding a derivation of  $\phi$ . Furthermore, there is a particular extension  $\mathcal{C}$  of the base  $\mathcal{B}$  (namely the one that just adds this rule) such that whenever an atom is derivable in  $\mathcal{C}$ , it is derivable in any extension of  $\mathcal{B}$ , which satisfies the formula  $\phi$ . If disjunction is modelled by coproducts, this implies a strong disjunction property: that there is a morphism from  $\llbracket \phi \supset (q \vee r) \rrbracket(\mathcal{B}, P)$  to  $\llbracket (\phi \supset q) \vee (\phi \supset r) \rrbracket(\mathcal{B}, P)$ .

This particular extension  $\mathcal{C}$  does not necessarily exist in a Kripke semantics, hence this argument does not hold for Kripke semantics.

**Acknowledgements.** We are grateful to the anonymous reviewers and to Alexander Gheorghiu and Tao Gu for helpful comments on this work.

**Open Access.** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

**Publisher's Note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

## Appendix: A Remarks on Consequence Relations

It will turn out that the significance of the base  $\mathcal{N}_-$  goes way beyond its use in Sandqvist's completeness theorem. Specifically, the algebraic interpretation that we give in Section 3 makes essential use of the canonical proofs of

atomic propositions of the form  $\phi^b$  available in  $\mathcal{N}_-$  in order to define the natural transformations that we needed in our interpretation.

As we noted, one of the key issues for proof-theoretic semantics is its completeness, or lack of it [23,24]. However, this issue is more delicate than it at first seems.

To begin with, we need to be clear about the judgements being used. Different forms of judgement give different answers to questions of soundness and completeness. At the most basic level, we have validity: the basic judgement is whether a formula is valid. Soundness means that theorems (provable formulae) should be valid in the semantics and completeness that any formula which is always valid in the semantics should be a theorem. This, however, is a fairly weak notion, and the judgements used are often strengthened to sequents. In this case, the soundness and completeness correspond to preservation and reflection of the validity of logical consequences, and there is more need to be specific how things are set up. This occurs even when we are just considering which consequences are valid. There is a further level of complexity when we come to consider possible representations of validations.

The common standard formulation is that a consequence relation is a relation expressing that a formula is a consequence of a set of other formulae. This concept is abstractly captured in the form of a reflexivity property, a monotonicity property, and a compositionality property, the last corresponding to cut.

**DEFINITION 38.** A (finitary) consequence relation is a relation  $\Gamma \triangleright p$  between a (finite) set of formulae and a formula, such that:

- $\phi \triangleright \phi$
- if  $\Gamma \triangleright \phi$ , then  $\Gamma, \psi \triangleright \phi$
- if  $\Gamma \triangleright \phi$  and  $\Delta, \phi \triangleright \psi$ , then  $\Gamma, \Delta \triangleright \psi$ .

There is a standard way of getting a consequence relation from a definition of validity. We write  $\blacktriangleright \phi$  to mean that  $\phi$  is valid.

**DEFINITION 39.** The *consequence relation generated from a validity* is defined by  $\Gamma \triangleright \psi$  iff if  $\blacktriangleright \gamma$ , for all  $\gamma \in \Gamma$ , then  $\blacktriangleright \psi$ .

Note that we can recover validity from consequence:

$$\blacktriangleright \psi \text{ iff } \emptyset \triangleright \psi$$

However, different consequence relations can give rise to the same validity. We therefore need to be careful about which consequence relations we are

using and why. But the consequence relation defined above is canonical in the following sense:

LEMMA 40. The finitary consequence relation generated from a validity is the largest finitary consequence relation that corresponds to that validity.

This property does not hold if we allow consequences  $\Gamma \triangleright \psi$  where  $\Gamma$  is infinite (the largest consequence relation corresponding to a validity is  $\Gamma \triangleright \psi$  iff  $\Gamma$  is finite and  $\psi$  is a consequence of  $\Gamma$  as per the finite case, or  $\Gamma$  is infinite and  $\psi$  is arbitrary). Moreover, the structures do not work well for substructural logic, and must be changed to a relation between an element of an algebra of formulae and a formula.

Definition 39 links strongly to the notion of an admissible rule. A rule is said to be admissible if its application preserves validity: if, in an instance, all hypotheses are valid, then so is the conclusion. But proof systems generate a different natural notion of consequence: the relation that expresses that a formula can be proved from a set of hypotheses.

## References

- [1] BELLIN, G., H. MARTIN, and P. EDMUND, Robinson, and Christian Urban Categorical proof theory of classical propositional calculus, *Theoretical Computer Science*, 364:146–165, 2006.
- [2] BLACKBURN, P., M. DE RIJKE, and Y. VENEMA, *Modal Logic*, Cambridge University Press, 2001.
- [3] VAN DALEN, D., *Logic and Structure*, Springer, Cham, 5th edition, 2012.
- [4] DUMMETT, M., *The Logical Basis of Metaphysics*, Harvard University Press, 1993.
- [5] ESPIRITO SANTO, J., and G. FERREIRA, A refined interpretation of intuitionistic logic by means of atomic polymorphism, *Studia Logica* 108:477–507, 2020.
- [6] FERREIRA, F., Comments on predicative logic, *Journal of Philosophical Logic* 35(1):1–8, 2006.
- [7] FERREIRA, F., and G. FERREIRA, The faithfulness of atomic polymorphism, in A. Indrzejczak, J. Kaczmarek, and M. Zawidzki, (eds.), *Proceedings of Trends in Logic XIII*, Łódź University Press, Łódź, 2014, pp. 55–65.
- [8] FERREIRA, F., and G. FERREIRA, Atomic polymorphism. *The Journal of Symbolic Logic* 78(1):260–274, 2013.
- [9] FERREIRA, F., and G. FERREIRA, The faithfulness of fat: a proof-theoretic proof, *Studia Logica* 103(6): 1301, 2015.
- [10] FÜHRMANN, C., and D. PYM, Order-enriched categorical models of the classical sequent calculus, *Journal of Pure and Applied Algebra* 204(1):21–78, 2006.
- [11] GENTZEN, G., Untersuchungen Über Das Logische Schließen. II. *Mathematische Zeitschrift* 39:405–431, 1935.
- [12] GHEORGHIU, A. V., and D. J. PYM, From Proof-theoretic Validity to Base-extension Semantics for Intuitionistic Propositional Logic. <https://arxiv.org/pdf/2210.05344>, 2022. Submitted.

- [13] GIRARD, J.Y., Interprétation fonctionnelle et élimination des coupures de l'arithmétique d'ordre supérieur. Thèse de Doctorat d'Etat, Université Paris VII, 1972.
- [14] GIRARD, J.-Y., Y. LAFONT, and P. TAYLOR. *Proofs and Types*, Cambridge University Press, 1989.
- [15] GIRARD, J. Y., Une Extension de l'Interpretation de Gödel à l'Analyse, et son Application à l'Élimination des Coupures dans l'Analyse et la Théorie des Types, in J.E. Fenstad, (ed.), *Proceedings of the Second Scandinavian Logic Symposium*, North-Holland, Amsterdam, 1971, pp. 63–92.
- [16] GOLDFARB, W., On Dummett's "proof-theoretic justifications of logical laws", in T. Piecha, and P. Schroeder-Heister, (eds.), *Advances in Proof-theoretic Semantics*, vol. 43 of Trends in Logic, Springer, 2016, pp. 195–210.
- [17] JACOBS, B.P.F., *Categorical Logic and Type Theory*, vol. 141 of *Studies in logic and the foundations of mathematics*, North-Holland, 2001.
- [18] JOHNSTONE, P.T., *Stone Spaces*, vol. 3 of *Cambridge Studies in Advanced Mathematics*, Cambridge University Press, 1982.
- [19] KRIPKE, S.A., Semantical Analysis of Intuitionistic Logic I, in J.N. Crossley, and M.A.E. Dummett, (eds.), *Formal Systems and Recursive Functions*, vol. 40 of *Studies in Logic and the Foundations of Mathematics*, Elsevier, 1965, pp. 92–130.
- [20] LAMBEK, J., and P. SCOTT, *Introduction to higher order categorical logic*, Cambridge University Press, 1986.
- [21] MAC LANE, S., and I. MOERDIJK, *Sheaves in Geometry and Logic: A First Introduction to Topos Theory*, Springer-Verlag, 1992.
- [22] NEGRI, S., and J. VON PLATO, *Structural Proof Theory*, Cambridge University Press, Cambridge 2008.
- [23] PIECHA, T., Completeness in proof-theoretic semantics, in T. Piecha, and P. Schroeder-Heister, (eds.), *Advances in Proof-Theoretic Semantics*, vol. 43 of *Trends in Logic*, Springer, 2016, pp. 231–251.
- [24] PIECHA, T., W. DE CAMPOS SANZ, and P. SCHROEDER-HEISTER, Failure of Completeness in Proof-theoretic Semantics, *Journal of Philosophical Logic* 44(3):321–335, 2015.
- [25] PIECHA, T., and P. SCHROEDER-HEISTER, Incompleteness of intuitionistic propositional logic with respect to proof-theoretic semantics, *Studia Logica* 107(1):233–246, 2019.
- [26] PISTONE, P., L. TRANCHINI, and M. PETROLO, The naturality of natural deduction (II). On atomic polymorphism and generalized propositional connectives, *Studia Logica* 110:545–592, 2022.
- [27] PRAWITZ, D., *Natural Deduction: A Proof-Theoretical Study*, Dover Publications, 1965.
- [28] PYM, D.J., E. RITTER, and E. ROBINSON, Proof-theoretic Semantics in Sheaves (Extended Abstract), in *Proceedings of the Eleventh Scandinavian Logic Symposium — SLSS 11*, 2022, pp. 36–38.
- [29] REYNOLDS, J.C., The discoveries of continuations, *Lisp and Symbolic Computation*, 6(3):233–247, 1993.

- [30] ROBINSON, E.P., Proof nets for classical logic, *Journal of Logic and Computation* 13(5):777–797, 2003.
- [31] SANDQVIST, T., *A note on definability of logical operators in second-order logic*, 2008.
- [32] SANDQVIST, T., Base-extension Semantics for Intuitionistic Sentential Logic, *Logic Journal of the IGPL* 23(5):719–731, 2015.
- [33] SCHROEDER-HEISTER, P., Validity concepts in proof-theoretic semantics, *Synthese* 148(3):525–571, 2006.
- [34] SCHROEDER-HEISTER, P., Proof-Theoretic versus Model-Theoretic Consequence, in M. Peliš, (ed.), *The Logica Yearbook 2007*, FILOSOFIA, Prague, 2008.
- [35] SEELY, R., Hyperdoctrines and natural deduction, *Zeitschrift für Mathematische Logik und Grundlagen der Mathematik* 29:505–542, 1983.
- [36] STAFFORD, W., and NASCIMENTO, V., Following all the rules: intuitionistic completeness for generalized proof-theoretic validity, *Analysis* 100, 07 2023.
- [37] TENNANT, N., Entailment and Proofs, *Proceedings of the Aristotelian Society* 79:167–189, 1978.
- [38] TENNANT, N., *Core Logic*, Oxford University Press, Oxford 2017.
- [39] TROELSTRA, A.S., and H. SCHWICHTENBERG, Basic Proof Theory, vol. 43 of *Cambridge Tracts in Theoretical Computer Science*, 2nd ed., Cambridge University Press, Cambridge, 2000.

D. PYM  
Computer Science and Philosophy  
University College London  
Gower Street  
London WC1E 6BT  
UK

and

Institute of Philosophy, School of Advanced Study  
University of London  
Malet Street  
London WC1E 7HU  
UK  
`d.pym@ucl.ac.uk`

E. RITTER  
Department of Computer Science  
University of Birmingham  
Edgbaston  
Birmingham B15 2TT  
UK  
`e.ritter@bham.ac.uk`

E. ROBINSON  
School of Electronic Engineering and Computer Science  
Queen Mary University of London  
Mile End Road  
London E1 4NS  
UK  
`e.p.robinson@qmul.ac.uk`