

Architecture of North Atlantic contourite drifts modified by transient circulation of the Icelandic mantle plume

Parnell-Turner, Ross; White, Nicholas; McCave, I. Nick; Henstock, Timothy J.; Murton, Bramley; Jones, Stephen

DOI:
[10.1002/2015GC005947](https://doi.org/10.1002/2015GC005947)

License:
None: All rights reserved

Document Version
Publisher's PDF, also known as Version of record

Citation for published version (Harvard):
Parnell-Turner, R, White, N, McCave, IN, Henstock, TJ, Murton, B & Jones, S 2015, 'Architecture of North Atlantic contourite drifts modified by transient circulation of the Icelandic mantle plume', *Geochemistry Geophysics Geosystems*, vol. 16, no. 10, pp. 3414-3435. <https://doi.org/10.1002/2015GC005947>

[Link to publication on Research at Birmingham portal](#)

Publisher Rights Statement:

AGU allows authors to deposit their journal articles if the version is the final published citable version of record, the AGU copyright statement is clearly visible on the posting, and the posting is made 6 months after official publication by the AGU

VC 2015. American Geophysical Union. All Rights Reserved.

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

1 **Supplementary Information for “Architecture of**
2 **North Atlantic Contourite Drifts Modified by**
3 **Transient Circulation of the Icelandic Mantle Plume”**

4
Ross Parnell-Turner^{1,4}, Nicky White¹, I. Nick McCave¹, Tim Henstock²,
Bramley Murton², Stephen M. Jones³

Corresponding authors: R. Parnell-Turner and N. White, Bullard Laboratories, Department of Earth Sciences, University of Cambridge, Madingley Road, Cambridge, CB3 0EZ, UK. (rparnellturner@whoi.edu, njw10@cam.ac.uk)

¹Department of Earth Sciences, University of Cambridge, Cambridge, CB3 0EZ, UK

²National Oceanography Centre Southampton, University of Southampton, European Way, Southampton SO14 3ZH

³School of Geography, Earth and Environmental Sciences, University of Birmingham, B15 2TT, UK

⁴Now at: Woods Hole Oceanographic Institution, Woods Hole, MA, USA

5 Contents of this file

6 1. Description of seismic reflection profiles

7 2. Description of Hatton Drift

8 3. Compaction parameter analysis

9 4. Figures:

10 – Figure S1: Seismic profiles 1 and 2

11 – Figure S2: Portion of profile 2 from Hatton Drift

12 – Figure S3: Portion of profile 1 from Eirik Drift

13 – Figure S4: Portion of profile 1b from Eirik Drift

14 – Figure S5: Detailed seismic images from Gardar drift

15 – Figure S6: Compaction parameter estimates from well-log data

16

1. Seismic reflection profiles

17 In this supplementary material, we present complete versions of seismic reflection pro-
18 files 1 and 2 (Figure S1). These profiles show the gross distribution and thickness of
19 sedimentary drifts in the Iceland and Irminger Basins (e.g. Björn, Gardar, Eirik and Hat-
20 ton Drifts). Figure S2 shows the detailed architecture of Hatton Drift imaged on profile
21 2. Figure S3 shows the westernmost portion of profile 1, where a section of Eirik Drift
22 is imaged. Figure S4 shows Eirik Drift imaged along profile 1b, which connects profiles
23 1 and 2 along the Greenland continental margin. Figure S5 shows detailed portions of
24 seismic images over Gardar Drift from profiles 1 and 2.

2. Hatton Drift

25 Hatton Drift sits on the lower slope of Hatton Bank which is located on the north-
26 west side of Rockall Plateau (Figures 1, S1 and S2). Maury Channel separates Hatton
27 and Gardar Drifts. Turbidity currents periodically transport volcanoclastic sediments
28 through this channel southward from Iceland [Ruddiman, 1972]. Sediments from Hatton
29 Drift have been recovered at various DSDP/ODP drilling sites (e.g. 403, 404, 552, 553,
30 555; Figure 1b). These sediments consist of carbonate oozes with occasional interbedded
31 terrigenous material within the upper few tens of meters [Stow and Holbrook, 1984]. Sed-
32 iments recovered in piston core V27-111 have an average silt-plus-sand content of 35%
33 increasing to $\sim 70\%$ toward the top. The carbonate-free coarse fraction is arkosic (80%
34 quartz and feldspar) and was presumably sourced from the continental shelf of the nearby
35 Rockall Plateau at glacial low sea levels [McCave et al., 1980]. Rippled sand dunes are
36 visible on photographs and sidescan sonar images acquired during deep camera tows on
37 Cruise KN-51 at the foot of Hatton Drift [see Figure 5 of McCave et al., 1980]. The pre-
38 dominantly foraminiferal sand has ripple and dune structures. It originates by winnowing
39 beneath a northward-flowing boundary current [McCave et al., 1980; McCartney, 1992].

40 The oldest material recovered from Hatton Drift consists of Late Paleocene volcanoclastic
41 and terrigenous sediments at Site 555. These deposits lie on top of igneous basement
42 [Stow and Holbrook, 1984]. A widespread non-depositional event occurred through much
43 of Eocene and Oligocene times and is visible at most Hatton Drift drilling sites. This
44 hiatus has been attributed to a global change in bottom-water circulation, triggered by a
45 combination of pronounced Antarctic cooling and development of strong Antarctic Bottom
46 Water flow into the Atlantic Ocean and formation of Labrador Sea Water [Schnitker, 1980;

47 *Woodruff and Savin, 1989; Zachos et al., 1993*]. Major growth of Hatton Drift probably
48 commenced during mid-Miocene times, with deposition of biogenic sediments continuing
49 into Holocene times [*Stow and Holbrook, 1984*].

50 Profile 2 traverses part of Hatton Drift on the northeast side of Rockall Plateau where
51 it is ~ 32 km wide (Figure S2). Drift sediments lie on the shoulder of Hatton Bank which
52 has a basement slope of $\sim 2^\circ$. They can be divided into four units named H-I to H-IV. The
53 youngest unit of Hatton Drift, H-I, consists of a 60 m thick continuous series of conformable
54 reflections (Figure S2d). This unit represents the most recent sedimentary drape deposited
55 by modern current activity. A package characterized by disrupted reflections, H-II, lies
56 directly underneath. This unit is up to 85 m thick and is acoustically transparent in
57 places. The base of H-II is marked by an erosional surface that truncates the continuous
58 reflections of H-III beneath. Unit H-II could be a debris flow or a mass-transport complex,
59 perhaps related to sediment supplied from the upslope direction on Hatton Bank. Unit
60 H-III is up to 230 m thick and comprises the main body of Hatton Drift on Profile 2.
61 Continuous reflections within unit H-III are easily identified but they are often truncated
62 by localized erosional surfaces and by minor buried channels (Figure S2b). Stratigraphic
63 onlap occurs at the base of unit H-III which lies unconformably above H-IV, the lowest
64 mappable unit. H-IV is probably the pre-drift sedimentary pile. Normal faulting occurs
65 at the sediment-basement interface and occasional faults reach the sea bed. Stratigraphic
66 growth occurs within H-IV but there is little evidence for growth within the Hatton Drift
67 itself.

2.1. Compaction parameters

68 We analyzed porosity-depth relationships from Sites 646 (Eirik Drift) and 984 (Björn
69 Drift) to determine the values of the initial porosity, ϕ_0 , and the compaction decay length,
70 λ (Figure S6). These parameters were used to determine solid accumulation rates for drift
71 deposits.

References

- 72 Schnitker, D. (1980), Global paleoceanography and its deep water linkage to the Antarctic
73 glaciation, *Earth-Science Reviews*, *16*, 1–20.
- 74 Stow, D. A. V., and J. A. Holbrook (1984), Hatton Drift contourites, Northeast Atlantic,
75 in *Init. Rep. Deep Sea Drill. Proj. 81*, edited by D. G. Roberts, D. Schnitker, et al., pp.
76 695–699, U.S. Govt. Printing Office, Washington.
- 77 Woodruff, F., and S. M. Savin (1989), Miocene deepwater oceanography, *Paleoceanogra-*
78 *phy*, *4*(1), 87–140.
- 79 Zachos, J. C., K. C. Lohmann, J. C. G. Walker, and S. W. Wise (1993), Abrupt Climate
80 Change and Transient Climates during the Paleogene: A Marine Perspective, *Journal*
81 *of Geology*, *101*, 191–213.




Figure S1: Time-migrated seismic reflection profiles 1 and 2 (see Figure 1 for location). a) Profile 2. Boxes show location of Figures S2 and S3. b) Interpretation. Solid lines = sedimentary drifts; yellow shading = sedimentary cover; dashed line = thin sediments; m = seabed multiple. c) Profile 1. d) Geologic interpretation. Note thin sediments are distributed along the western flank of the Reykjanes Ridge, however modern day body of Snorri Drift does not extend as far south as profile 2 (i.e. 62°N).




Figure S2: (a) Portion of profile 2 which crosses Hatton Drift (see Figures S1 and 1c for location). (b) and (c) Insets showing details of buried channel and onlapping stratigraphy, respectively. (d) Interpretation. Colored lines/intervals = interpreted horizons/units: yellow = H-I; green = H-II; blue = H-III; orange = H-IV; black line = basement; black lines = normal faults.




Figure S3: (a) Portion of profile 2 which crosses Eirik Drift (see Figure 1c for location). (b) Interpretation. Colored lines = interpreted horizons with ages noted; colored intervals = stratigraphic units; upper/lower black lines = seabed/basement reflections, respectively; black lines = faults; arrows = bounding channels.




Figure S4: (a) Portion of profile 1b located parallel to Greenland margin (see Figure 1c for location). (b) Interpretation. Colored lines = dated horizons; red dotted lines = interpretation of R2; colored intervals = stratigraphic units; upper/lower black lines = seabed/ basement reflections, respectively; black lines = faults. Subset of sedimentary units from Eirik Drift are seen. Two modern channels visible at ranges of 35 and 105 km. These channels initiated at R3 time (7.5 Ma), and have since migrated ~10 km southwards. Buried channel located at range 85 km. Channel initiated post-R5 times (10–12 Ma) and since southward by ~15 km until R3 times (7.5 Ma).




Figure S5: Detailed images of Gardar drift (see Figure 9 for location). (a) Detailed section of profile 1 at range of 80 km. (b) Interpretation showing normal faults and minor stratigraphic growth features. Thick lines = faults; thin lines = stratigraphic interpretation. (c) Detailed section of profile 2 at range of 89 km. (d) Interpretation showing two unconformity surfaces (bold lines) where older unit is truncated by overlying strata. (e) Detailed section of profile 2 at range of 920 km. (f) Interpretation showing westward-migrating mudwaves on eastern slope of drift. Mudwaves at seabed are ~ 25 m in amplitude, ~ 500 m in wavelength.




Figure S6: Porosity-depth relationship for Eirik and Björn Drifts from downhole log data. (a) Site 646 (Eirik Drift). Thin gray line = compensated neutron porosity. Noise at depths <220 m due to logging through drill pipe; black line = porosity calculated using best-fitting compaction parameters given in (c). (b) Site 984 (Björn Drift). (c) Misfit plot for compaction parameters ϕ_0 and λ at Site 646. Cross = misfit minimum. (d) Misfit plot for compaction parameters ϕ_0 and λ at Site 984.