

X-simple image eigencones of tropical matrices

Plavka, Jan; Sergeev, Sergey

DOI:

[10.1016/j.laa.2016.06.013](https://doi.org/10.1016/j.laa.2016.06.013)

License:

Creative Commons: Attribution-NonCommercial-NoDerivs (CC BY-NC-ND)

Document Version

Peer reviewed version

Citation for published version (Harvard):

Plavka, J & Sergeev, S 2016, 'X-simple image eigencones of tropical matrices', *Linear Algebra and its Applications*, vol. 507, pp. 169-190. <https://doi.org/10.1016/j.laa.2016.06.013>

[Link to publication on Research at Birmingham portal](#)

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

Accepted Manuscript

X -simple image eigencones of tropical matrices

Ján Plavka, Sergeĭ Sergeev


PII: S0024-3795(16)30230-0
DOI: <http://dx.doi.org/10.1016/j.laa.2016.06.013>
Reference: LAA 13747

To appear in: *Linear Algebra and its Applications*

Received date: 2 November 2015
Accepted date: 6 June 2016

Please cite this article in press as: J. Plavka, S. Sergeev, X -simple image eigencones of tropical matrices, *Linear Algebra Appl.* (2016), <http://dx.doi.org/10.1016/j.laa.2016.06.013>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



\mathbf{X} -simple image eigencones of tropical matrices

Ján Plavka^{a,1}, Sergej Sergeev^{b,*}

^a*Department of Mathematics and Theoretical Informatics, Technical University,
B. Němcovej 32, 04200 Košice, Slovakia*

^b*University of Birmingham, School of Mathematics, Edgbaston B15 2TT, UK*

Abstract

We investigate max-algebraic (tropical) one-sided systems $A \otimes x = b$ where b is an eigenvector and x lies in an interval \mathbf{X} . A matrix A is said to have \mathbf{X} -simple image eigencone associated with an eigenvalue λ , if any eigenvector x associated with λ and belonging to the interval \mathbf{X} is the unique solution of the system $A \otimes y = \lambda x$ in \mathbf{X} . We characterize matrices with \mathbf{X} -simple image eigencone geometrically and combinatorially, and for some special cases, derive criteria that can be efficiently checked in practice.

Keywords: Max algebra, one-sided system, weakly robust, interval analysis
AMS classification: 15A18, 15A80, 93C55

1. Introduction

1.1. Problem statement and main results

In this paper, by max algebra we mean the set of nonnegative numbers \mathbb{R}_+ equipped with usual multiplication $a \otimes b := a \cdot b$ and idempotent addition $a \oplus b := \max(a, b)$. Algebraically, \mathbb{R}_+ equipped with these operations forms a semifield. The operations of max algebra are then extended to matrices and vectors in the usual way, giving rise to an analogue of nonnegative linear algebra.

Max-algebraic one-sided systems $A \otimes x = b$ and max-algebraic eigenproblem $A \otimes x = \lambda x$ are two fundamental problems of max algebra whose

*Corresponding author. Email: sergiej@gmail.com

Email addresses: jan.plavka@tuke.sk (Ján Plavka), sergiej@gmail.com (Sergej Sergeev)

¹Supported by APVV grant 04-04-12.

solution goes back to the works of Cuninghame-Green [9, 11], Vorobyev [21] and Zimmermann [22] and these two topics are thoroughly discussed in any textbook of the max-plus (tropical) linear algebra [2, 3, 13]. Our intention is to consider the situation when the right-hand side of $A \otimes x = b$ is an eigenvector, and also when the solution has to lie in some interval of \mathbb{R}_+^n .

By an interval of \mathbb{R}_+^n we mean a subset of \mathbb{R}_+^n of the form $\mathbf{X} = \times_{i=1}^n \mathbf{X}_i$, where each \mathbf{X}_i is an arbitrary interval belonging to \mathbb{R}_+ , with its upper end possibly equal to $+\infty$ (and lower end possibly equal to 0). In particular, \mathbb{R}_+^n is an interval of itself. For each i we denote $\underline{x}_i := \inf \mathbf{X}_i$ and $\bar{x}_i = \sup \mathbf{X}_i$. Then we also have $\underline{x} := (\underline{x}_i)_{i=1}^n = \inf \mathbf{X}$ and $\bar{x} := (\bar{x}_i)_{i=1}^n = \sup \mathbf{X}$.

The notion of \mathbf{X} -simple image eigencone, which we introduce next, is related to the concept of simple image set [4]. By definition, simple image set of A is the set of vectors b such that the system $A \otimes x = b$ has a unique solution. If the only solution of the system $A \otimes x = b$ is $x = b$, then b is called a *simple image eigenvector*.

A matrix A is said to have \mathbf{X} -simple image eigencone associated with a (fixed) eigenvalue λ , if any eigenvector x associated with the eigenvalue λ and belonging to the interval \mathbf{X} is the unique solution in \mathbf{X} for the system $A \otimes y = \lambda x$. The characterization of a matrix with \mathbf{X} -simple image eigencone is described as the main result of the paper in Section 4.

Let us now give more details on the organization of the paper and on the results obtained there. Section 2 is devoted to basic notions of max algebra and its connections to the theory of digraphs and max-algebraic (tropical) convexity. In particular, we revisit here the spectral theory, focusing on the eigencone associated with an arbitrary eigenvalue, the generating matrix and the critical graph. Some aspects of the diagonal similarity scaling are also briefly discussed.

Section 3 starts by discussing the problem of covering the node set of a digraph by ingoing edges. We proceed with the theory of one-sided systems $A \otimes x = b$ where we describe the solution set to such systems and start analysing the case when b is an eigenvector of A . The main result of that section is Theorem 3.10, which characterizes matrices that have at least one simple image eigenvector corresponding to an eigenvalue λ .

In the beginning of Section 4 we first discuss the relation between \mathbf{X} -simple image eigencone and \mathbf{X} -weak robustness of a matrix. We then develop an interval version of theory of one-sided systems $A \otimes x = b$, i.e., when x has to belong to an interval \mathbf{X} . The second part of the section contains the main results of the paper: Theorem 4.16 and Theorem 4.17. More precisely,

Theorem 4.16 characterizes when A has \mathbf{X} -simple image eigencone in general, and Theorem 4.17 focuses on the case when \mathbf{X} is of a certain special type.

In the end of the paper we formulate some conclusions and discuss some directions for further research.

1.2. Motivations

In the literature, max algebra often appears as max-plus semiring developed over the set $\mathbb{R} \cup \{-\infty\}$ equipped with operations $a \otimes b := a + b$ and $a \oplus b := \max(a, b)$. However, this semiring is isomorphic to the semiring defined above, via a logarithmic transform. Max-plus algebra plays the crucial role in the study of discrete-event dynamic systems connected with the optimization problems such as scheduling or project management in which the objective function depends on the operations maximum and plus. The main principle of discrete-event dynamic systems consisting of n entities (machines [9, 11], processors [7], computers, etc.) is that the entities work interactively, i.e., a given entity must wait before proceeding to its next event until certain others have completed their current events. Cuninghame-Green [9] and Butkovič [3] discussed a hypothetical industrial discrete-event dynamic system and a multiprocessor interactive system, respectively, which can be described by the interferences using recurrence relations

$$x_i(r+1) = \max(x_1(r) + a_{1i}, x_2(r) + a_{2i}, \dots, x_n(r) + a_{ni}), \quad i \in \{1, 2, \dots, n\}.$$

The formula expresses the fact that entity i must wait with its $r+1$ st cycle until entities $j = 1, \dots, n$ have finished their r th cycle. The symbol $x_i(r)$ denotes the starting time of the r th cycle of entity i , and a_{ij} is the corresponding activity duration at which entity e_j prepares the outputs (products, components, data, etc.) for entity e_i . The steady states of such systems correspond to eigenvectors of max-plus matrices, therefore the investigation of properties of eigenvectors is important for the above mentioned applications.

In max-plus algebra the matrices for which the steady states of the systems are reached with any nontrivial starting vector are called robust. Such matrices have been studied in [3], [19]. The matrices for which the steady states of the systems are reached only if a nontrivial starting vector is an eigenvector of the matrix are called weakly robust. Efficient characterizations of such matrices are described in [6].

In practice, the values of starting vector are not exact numbers and usually they are rather contained in some intervals. Considering matrices and vectors with interval entries is therefore of practical importance.

See [12, 14, 15, 16, 18] for some of the recent developments. In particular, the weak robustness of an interval matrix is studied in [17].

The aim of this paper is to characterize the weak \mathbf{X} -robustness, i.e., the weak robustness of matrices with initial times confined in an interval vector \mathbf{X} , using \mathbf{X} -simplicity of image eigenecone.

2. Preliminaries

2.1. Matrices and graphs

Many problems of max algebra can be described and resolved in terms of digraphs (i.e., directed graphs). Let us give some of the relevant definitions here.

Definition 2.1 (Associated digraphs). The *weighted digraph associated with* $A \in \mathbb{R}_+^{n \times n}$ is the digraph $\mathcal{G}(A) = (N, E)$ with the node set $N := \{1, \dots, n\}$ and the edge set E such that $(i, j) \in E$ (edge from i to j) if and only if $a_{ij} > 0$. The number a_{ij} is called the *weight* of (i, j) .

Definition 2.2 (Paths). A *path* in the digraph $\mathcal{G}(A) = (N, E)$ is a sequence of nodes $p = (i_1, \dots, i_{k+1})$ such that $(i_j, i_{j+1}) \in E$ for $j = 1, \dots, k$. A path p is closed if $i_1 = i_{k+1}$, elementary if all nodes are distinct, and a cycle if it is closed and elementary. The number k is the *length* of the path p and is denoted by $l(p)$.

Definition 2.3 (Strongly connected components). By a *strongly connected component* (for brevity s.c.c.) of $\mathcal{G}(A) = (N, E)$ we mean a subdigraph $\mathcal{G}' = (N', E')$ with $N' \subseteq N$ and $E' \subseteq E$, such that any two distinct nodes $i, j \in N'$ are contained in a common cycle and N' is a maximal subset of N with that property. Particularly, $\mathcal{G}(A)$ is strongly connected if $\mathcal{G}' = (N', E')$ is strongly connected component of $\mathcal{G}(A)$ with $N' = N$ and $E' = E$.

Powers of max algebraic matrices are closely related to optimization on digraphs. Observe that the i, j th entry of the power $A^{\otimes k} := \underbrace{A \otimes \dots \otimes A}_k$ is

the biggest weight among all paths of length k connecting i to j .

If we define the formal series $A^+ = \bigoplus_{k=1}^{\infty} A^{\otimes k}$ then the i, j th entry of A^+ (possibly diverging to $+\infty$) equals to the greatest weight among all paths connecting i to j . Such weight is guaranteed to be finite if the weight of any cycle in $\mathcal{G}(A)$ does not exceed 1.

Definition 2.4 (Irreducibility). A matrix $A \in \mathbb{R}_+^{n \times n}$ is called *irreducible* if $\mathcal{G}(A)$ is strongly connected, and *reducible* otherwise.

Definition 2.5 (Graph restrictions). For arbitrary $K \subseteq N$, we denote by $\mathcal{G}(A)|_K$ the subgraph of $\mathcal{G}(A)$ consisting of all nodes of K and all edges of $\mathcal{G}(A)$ between the nodes of K .

2.2. Geometry

Max algebra also gives rise to the max-algebraic (tropical) analogue of convexity.

Definition 2.6 (Max cone). A subset $\mathcal{K} \subseteq \mathbb{R}_+^n$ is called a *max cone* if we have

- 1) $\lambda x \in \mathcal{K}$ for any $\lambda \geq 0$ and $x \in \mathcal{K}$,
- 2) $x \oplus y \in \mathcal{K}$ for any $x, y \in \mathcal{K}$.

The name “max cone” was suggested in [6]. In the literature this object also appears as tropical cone or max-plus linear space.

Definition 2.7 (Column span). For $A \in \mathbb{R}_+^{m \times n}$ define its *max-algebraic column span* as

$$\text{span}_{\oplus}(A) := \left\{ \bigoplus_{j=1}^n A_{.j} x_j : x_j \geq 0 \forall j \right\}, \quad (1)$$

where $A_{.j}$, for $j = 1, \dots, n$ denotes the j -th column of A .

The set of all positive vectors in $\text{span}_{\oplus}(A)$ will be denoted by $\text{span}_{\oplus}^+(A)$.

It is easily shown that $\text{span}_{\oplus}(A)$ is a max cone. Furthermore, $\text{span}_{\oplus}(A)$ is always closed in the Euclidean topology [6].

Consider now the following operator.

Definition 2.8 (Projector). Let \mathcal{W} be a closed max cone. Define

$$P_{\mathcal{W}}(x) := \max\{y \in \mathcal{W} : y \leq x\}. \quad (2)$$

In the case $\mathcal{W} = \text{span}_{\oplus}(A)$ we will write P_A instead of $P_{\text{span}_{\oplus}(A)}$, for brevity.

$P_{\mathcal{W}}$ is a nonlinear projector on the max cone \mathcal{W} . It is homogeneous ($P_{\mathcal{W}}(\lambda x) = \lambda P_{\mathcal{W}}x$) and isotone ($x \leq y \Rightarrow P_{\mathcal{W}}x \leq P_{\mathcal{W}}y$.) These operators are crucial for tropical convexity: see, e.g., [8].

2.3. Eigenvalues and eigenvectors

Definition 2.9 (Eigencone). The set

$$V(A, \lambda) = \{x: A \otimes x = \lambda x\}, \quad (3)$$

where $A \in \mathbb{R}_+^{n \times n}$ and $\lambda \geq 0$, is called the (max-algebraic) *eigencone* of A associated with λ . The nonzero vectors of $V(A, \lambda)$ are (max-algebraic) *eigenvectors* of A associated with λ .

The set of all positive vectors in $V(A, \lambda)$ will be denoted by $V^+(A, \lambda)$.

Note that $V(A, \lambda)$ consists of the eigenvectors associated with λ and vector $\mathbf{0}$. Obviously, $V(A, \lambda)$ is a max cone.

Definition 2.10 (Maximum cycle geometric mean). Let $A = (a_{ij}) \in \mathbb{R}_+^{n \times n}$. For any $i_1, \dots, i_k \in N$, the *geometric mean* of the cycle (i_1, i_2, \dots, i_k) is defined as $\sqrt[k]{a_{i_1 i_2} \cdots a_{i_k i_1}}$. The *maximum cycle geometric mean* of $A \in \mathbb{R}_+^{n \times n}$ equals to

$$\lambda(A) := \max_{k=1}^n \max_{1 \leq i_1, \dots, i_k \leq n} \sqrt[k]{a_{i_1 i_2} \cdots a_{i_k i_1}}. \quad (4)$$

$\lambda(A)$ is the greatest max-algebraic eigenvalue of A , for any $A \in \mathbb{R}_+^{n \times n}$. If A is irreducible then $\lambda(A)$ is the only max-algebraic eigenvalue of A (e.g. [3], Theorem 4.4.8).

Reducible $A \in \mathbb{R}_+^{n \times n}$ may have up to n max-algebraic eigenvalues, in general. We next give some elements of the spectral theory of reducible matrices. Although that theory is usually developed in terms of the Frobenius normal form and spectral classes [3] following a similar development of the spectral theory of nonnegative matrices, we choose not to do so, since in this paper we only need 1) the relation between the critical graph and the saturation graph, 2) the generating matrix of $V(A, \lambda)$.

Denote by $\Lambda(A)$ the set of (max-algebraic) eigenvalues of A . General $\lambda \in \Lambda(A)$ can be characterized as maximum cycle geometric mean of a certain subgraph of $\mathcal{G}(A)$.

Definition 2.11 (Support). For $x \in \mathbb{R}_+^n$ the set $\text{supp}(x) = \{i: x_i > 0\}$ is called the support of x .

The proof of the following statement is standard, but we give it for the reader's convenience.

Proposition 2.12. *Let $\lambda \in \Lambda(A)$ and $x \in V(A, \lambda)$ be nonzero. Then λ is the maximum cycle geometric mean of $\mathcal{G}(A)|_{\text{supp}(x)}$.*

PROOF: Take any cycle (i_1, \dots, i_k) with all indices belonging to $\text{supp}(x)$. Multiplying the inequalities $a_{ii_{l+1}}x_{i_{l+1}} \leq \lambda x_{i_l}$ for $l = 1, \dots, k-1$ and $a_{i_k i_1}x_{i_1} \leq \lambda x_{i_k}$, and cancelling all the coordinates of x we obtain that the cycle mean of (i_1, \dots, i_k) does not exceed λ .

Now, start with any $j_0 \in \text{supp}(x)$ and find j_1 such that $a_{j_0 j_1}x_{j_1} = \lambda x_{j_0}$. We again have $j_1 \in \text{supp}(x)$. Proceeding this way we obtain a cycle (j_t, \dots, j_{t+l}) (for some t and l) with the cycle mean equal to λ . \square

Let us also recall a useful link between the support of an eigenvector of A and the zero-nonzero pattern of A .

Let $A \in \mathbb{R}_+^{n \times n}$ and $J, L \subseteq N$. $A_{J,L}$ denotes the submatrix of A with row index set J and column index set L .

Proposition 2.13 (e.g. [3], p.96). *Let $A \in \mathbb{R}_+^{n \times n}$, $x \in V(A, \lambda)$ and $N' := \text{supp}(x)$. Then $A_{N \setminus N', N'} = \mathbf{0}$.*

Definition 2.14 (Critical graph $\mathcal{G}_c(A, x, \lambda)$). *For $x \in V(A, \lambda)$, define the critical graph $\mathcal{G}_c(A, x, \lambda)$ as the subgraph of $\mathcal{G}(A)$ consisting of all nodes and edges belonging to the cycles of $\mathcal{G}(A)|_{\text{supp}(x)}$ whose geometric mean is equal to λ .*

Definition 2.15 (Saturation graph). *For $x \in V(A, \lambda)$, the saturation graph $\text{Sat}(A, x, \lambda)$ is the subgraph of $\mathcal{G}(A)$ with set of nodes N and set of edges*

$$E_{\text{Sat}} = \{(i, j) : a_{ij}x_j = \lambda x_i \neq 0\}. \quad (5)$$

Proposition 2.16 ([2], Theorems 3.96 and 3.98). *For any $x \in V(A, \lambda)$,*

- (i) *Every node $i \in N$ such that $x_i \neq 0$ has an outgoing edge in $\text{Sat}(A, x, \lambda)$;*
- (ii) *Any cycle in $\text{Sat}(A, x, \lambda)$ belongs to $\mathcal{G}_c(A, x, \lambda)$;*
- (iii) *$\mathcal{G}_c(A, x, \lambda)$ is a subgraph of $\text{Sat}(A, x, \lambda)$;*

Definition 2.17 (Critical graph $\mathcal{G}_c(A, \lambda)$). *The critical graph $\mathcal{G}_c(A, \lambda)$ associated with λ and the set of nodes N^λ associated with λ :*

$$\begin{aligned} \mathcal{G}_c(A, \lambda) &:= \mathcal{G}_c(A, x', \lambda), \quad N^\lambda := \text{supp}(x'), \quad \text{where} \\ &\text{for } x' \in V(A, \lambda) \text{ such that } \text{supp}(x) \subseteq \text{supp}(x') \forall x \in V(A, \lambda). \end{aligned} \quad (6)$$

Each $\mathcal{G}_c(A, x, \lambda)$ consists of several strongly connected components isolated from each other.

In the following proposition let us collect some facts about the relation between $\mathcal{G}_c(A, x, \lambda)$ and $\mathcal{G}_c(A, \lambda)$.

Proposition 2.18. *Let $A \in \mathbb{R}_+^{n \times n}$, $\lambda \in \Lambda(A)$ and $x \in V(A, \lambda)$. Then*

- (i) $\mathcal{G}_c(A, x, \lambda) \subseteq \mathcal{G}_c(A, \lambda)$;
- (ii) $\mathcal{G}_c(A, x, \lambda) = \mathcal{G}_c(A, \lambda)|_{\text{supp}(x)}$;
- (iii) $\mathcal{G}_c(A, x, \lambda)$ consists of entire strongly connected components of $\mathcal{G}_c(A, \lambda)$.

PROOF: We only prove (ii) and (iii) since (i) follows from any of them.

(ii): Observe that both $\mathcal{G}_c(A, x, \lambda)$ and $\mathcal{G}_c(A, \lambda)|_{\text{supp}(x)}$ consist of the nodes and edges of the cycles on $\text{supp}(x)$ that have the cycle geometric mean λ , hence $\mathcal{G}_c(A, x, \lambda) = \mathcal{G}_c(A, \lambda)|_{\text{supp}(x)}$.

(iii): Since $\mathcal{G}_c(A, x, \lambda)$ is defined as a subgraph consisting of all nodes and edges of some critical cycles, it consists of several isolated strongly connected components, and each of these components is a subgraph of a component of $\mathcal{G}_c(A, \lambda)$. It remains to prove that none of these subgraphs is proper.

By the contrary, suppose that one of these components is a proper subgraph of a component of $\mathcal{G}_c(A, \lambda)$. Since $\mathcal{G}_c(A, x, \lambda) = \mathcal{G}_c(A, \lambda)|_{\text{supp}(x)}$, the component of $\mathcal{G}_c(A, \lambda)$ should contain a node in $\text{supp}(x)$ and a node not in $\text{supp}(x)$, otherwise it coincides with the component of $\mathcal{G}_c(A, x, \lambda)$. However, this contradicts with Proposition 2.13. Hence the claim. \square

We further define a generating matrix of $V(A, \lambda)$. For that, first define the matrix A_λ with the columns

$$(A_\lambda)_i = \begin{cases} A_{\cdot i} / \lambda & \text{if } i \in N^\lambda, \\ \mathbf{0}, & \text{otherwise.} \end{cases} \quad (7)$$

Here N^λ is as in (6). A_λ^+ is finite, since the weight of any cycle in A_λ does not exceed 1.

Definition 2.19 (Generating Matrix). *Let $N_c^{\lambda,1}, \dots, N_c^{\lambda,k}$ be the node sets of the strongly connected components of $\mathcal{G}_c(A, \lambda)$, and let j_1, \dots, j_k be the first*

indices in those components. Define the generating matrix of $V(A, \lambda)$ as the matrix resulting from stacking the columns $(A_\lambda^+)_{\cdot j_1}, \dots, (A_\lambda^+)_{\cdot j_k}$ together:

$$G_{A,\lambda} = [(A_\lambda^+)_{\cdot j_1}, \dots, (A_\lambda^+)_{\cdot j_k}] \quad (8)$$

Proposition 2.20 ([3] **Coro. 4.6.2**). *For any nonzero $\lambda \in \Lambda(A)$*

$$V(A, \lambda) = V(A_\lambda, 1) = \text{span}_\oplus(G_{A,\lambda}). \quad (9)$$

2.4. Invertible matrices and diagonal similarity scaling

The class of invertible matrices in max algebra is quite thin. In fact it coincides with that in nonnegative algebra, consisting of all products of positive diagonal and permutation matrices. The positive diagonal matrices will be especially interesting to us since they give rise to a particularly useful *visualization scaling*, also known as a Fiedler-Pták scaling. For a positive vector $x \in \mathbb{R}_+^n$, denote by $\text{diag}(x)$ matrix $X \in \mathbb{R}_+^{n \times n}$ whose i th diagonal entry is x_i and all off-diagonal entries are 0.

Proposition 2.21 ([20], **Theorem 3.7**). *Let $A \in \mathbb{R}_+^{n \times n}$. For a positive $x \in \mathbb{R}_+^n$, let $X = \text{diag}(x)$ and $\tilde{A} = X^{-1}AX$ with entries \tilde{a}_{ij} for $i, j = 1, \dots, n$.*

- (i) *If x satisfies $A \otimes x \leq x$ then $\tilde{a}_{ij} \leq 1$ and, in particular, $\tilde{a}_{ij} = 1$ for $(i, j) \in E_c(A, 1)$ (**visualization scaling**).*
- (ii) *There exists a positive x satisfying $A \otimes x \leq x$ such that $\tilde{a}_{ij} \leq 1$, and $\tilde{a}_{ij} = 1$ if and only if $(i, j) \in E_c(A, 1)$ (**strict visualization scaling**).*

Definition 2.22 (Visualization). *A matrix $A = (a_{ij}) \in \mathbb{R}_+^{n \times n}$ is called visualized if $a_{ij} \leq \lambda(A)$ for all $i, j \in N$ and strictly visualized if it is visualized and $a_{ij} = \lambda(A)$ holds only for $(i, j) \in \mathcal{G}_c(A, \lambda)$.*

We will also use the following observation about the diagonal similarity scaling, where by $\text{Sis}V(A, \lambda)$ we denote the set of vectors in $V(A, \lambda)$ that belong to the simple image set of A .

Lemma 2.23. *Let $A \in \mathbb{R}_+^{n \times n}$, let $X \in \mathbb{R}_+^{n \times n}$ be a positive diagonal matrix. and $\tilde{A} := X^{-1}AX$. Then*

- (i) $\Lambda(A) = \Lambda(\tilde{A})$, and $\mathcal{G}_c(A, \lambda) = \mathcal{G}_c(\tilde{A}, \lambda)$ for every $\lambda \in \Lambda(A)$;
- (ii) $y \in V(A, \lambda) \Leftrightarrow X^{-1}y \in V(\tilde{A}, \lambda)$;

(iii) $y \in \text{Sis } V(A, \lambda) \Leftrightarrow X^{-1}y \in \text{Sis } V(\tilde{A}, \lambda)$.

PROOF: The facts described in part (i) are well-known. Parts (ii) and (iii) follow from the observation that

$$A \otimes y = b \Leftrightarrow X^{-1}A \otimes y = X^{-1}b \Leftrightarrow X^{-1}AX \otimes (X^{-1}y) = X^{-1}b$$

for all $y, b \in \mathbb{R}_+^{n \times n}$. \square

Let us recall some properties of A_λ^+ and $G_{A,\lambda}$ when A is visualized. By E we denote a matrix of the same dimensions as A whose every element is 1.

Proposition 2.24 ([20], **Proposition 4.1**). *Let $A \in \mathbb{R}_+^{n \times n}$ be visualized and let $\lambda = \lambda(A)$. Let N_c^1, \dots, N_c^k be the node sets of the components of $\mathcal{G}_c(A, \lambda)$, and let $K = \{j_1, \dots, j_k\}$ be the index set of the columns of A_λ^+ forming $G_{A,\lambda}$.*

- (i) *For each $r, s \in K$ there exists $\alpha_{rs} \leq 1$ such that $(A_\lambda^+)_{N_c^r N_c^s} = \alpha_{rs} E_{N_c^r N_c^s}$ and $(G_{A,\lambda})_{N_c^r, s} = \alpha_{rs} E_{N_c^r, j_s}$.*
- (ii) *If $r = s$ then $\alpha_{rs} = 1$.*
- (iii) *There exists s such that $\alpha_{rs} < 1$ for all $r \neq s$.*

PROOF: Parts (i) and (ii) follow from [20] Proposition 4.1, part 2. For part (iii), note that if it does not hold, then there exist indices i_1, \dots, i_l belonging to K such that $\alpha_{i_1 i_2} = 1, \dots, \alpha_{i_l i_1} = 1$. This implies existence of a cycle in $\mathcal{G}_c(A, \lambda)$ going through different strongly connected components, contradicting the fact that they are isolated. \square

3. One-sided systems and simple image eigenvectors

3.1. Solving max-algebraic one-sided systems

In this section we shall suppose that $A \in \mathbb{R}_+^{m \times n}$ is a given matrix and recall the crucial results concerning a system of linear equations $A \otimes x = b$. Our notation is similar to that introduced in [3], [22]. However, unlike for example in [3], Section 3.1, we do not assume that b has full support (i.e., is positive), or even that every row and column of A contains a nonzero element. Denote

$$S(A, b) = \{x \in \mathbb{R}_+^n : A \otimes x = b\}. \quad (10)$$

For any $j \in N$ denote

$$\begin{aligned}\gamma_j^*(A, b) &= \max\{\alpha \in \mathbb{R}_+ \cup \{+\infty\} : \alpha A_{.j} \leq b\} \\ &= \min_{i: a_{ij} \neq 0} b_i a_{ij}^{-1},\end{aligned}\tag{11}$$

assuming that $0 \cdot (+\infty) = (+\infty) \cdot 0 = 0$, $M = \{1, \dots, m\}$ and

$$M_j(A, b) = \{i \in M : a_{ij} \gamma_j^*(A, b) = b_i \neq 0\}.\tag{12}$$

Vector $\gamma^*(A, b) = (\gamma_j^*(A, b))_{j=1}^n$ is closely related to the projection: $P_A(b) = A \otimes \gamma^*(A, b)$.

$\gamma^*(A, b)$ can have a $+\infty$ component in general, but only in the case when the corresponding column is zero: $A_{.j} = \mathbf{0}$.

The following lemma is crucial for the theory of $A \otimes x = b$.

Lemma 3.1 (e.g. [3] Theorem 3.1.1). $x \in S(A, b)$ if and only if $x \leq \gamma^*(A, b)$ and $\cup_{j: x_j = \gamma_j^*(A, b)} M_j(A, b) = \text{supp}(b)$.

We now give a description of the solution set of $A \otimes x = b$ in terms of minimal coverings.

Theorem 3.2. Let Ω be a collection of minimal subsets $N' \subseteq N$ such that $\cup_{j \in N'} M_j(A, b) = \text{supp}(b)$. Then

$$S(A, b) = \bigcup_{N' \in \Omega} S_{N'}(A, b),\tag{13}$$

where

$$\begin{aligned}S_{N'}(A, b) &= \{x \in \mathbb{R}_+^n : x_j = \gamma_j^*(A, b) \text{ for } j \in N', \\ &\quad x_j \leq \gamma_j^*(A, b) \text{ for } j \in N \setminus N'\}.\end{aligned}\tag{14}$$

PROOF: We first show that $S(A, b) \subseteq \cup_{N' \in \Omega} S_{N'}(A, b)$. If $x \in S(A, b)$ then by Lemma 3.1 we have $x \leq \gamma^*(A, b)$ and $\cup_{j: x_j = \gamma_j^*(A, b)} M_j(A, b) = \text{supp}(b)$. Hence $x \in S_{N'}(A, b)$ for some minimal $N' \subseteq \{j : x_j = \gamma_j^*(A, b)\}$ such that $\cup_{j \in N'} M_j(A, b) = \text{supp}(b)$.

Let $x \in S_{N'}(A, b)$. Then $\bigoplus_{j \in N'} A_{.j} x_j = b$ and $\bigoplus_{j \in N \setminus N'} A_{.j} x_j \leq b$, hence $A \otimes x = b$. \square

Corollary 3.3. *Let us have $\cup_{j \in N'} M_j(A, b) = \text{supp}(b)$ for some proper $N' \subseteq N$, and let $i \notin N \setminus N'$. Then for each $\alpha \leq \gamma_i^*(A, b)$ there exists $x \in S_{N'}(A, b) \subseteq S(A, b)$ with $x_i = \alpha$.*

PROOF: We can assume without loss of generality that N' is a minimal subset such that $\cup_{j \in N'} M_j(A, b) = \text{supp}(b)$. The claim then follows from (14) and (13). \square

Let us now formulate, without proof, conditions for existence and uniqueness of a finite solution to $A \otimes x = b$. Here $\bar{S}(A, b) := \{x \in (\mathbb{R} \cup \{+\infty\})^n : A \otimes x = b\}$.

Proposition 3.4 (e.g., [3], Coro. 3.1.2). *Let $A \in \mathbb{R}_+^{m \times n}$ and $b \in \mathbb{R}_+^m$. Then the following conditions are equivalent:*

- (i) $S(A, b) \neq \emptyset$,
- (ii) $\gamma^*(A, b) \in \bar{S}(A, b)$,
- (iii) $\bigcup_{j \in N} M_j(A, b) = \text{supp}(b)$.

Proposition 3.5 (e.g., [3], Coro. 3.1.3). *Let $A \in \mathbb{R}_+^{m \times n}$ and $b \in \mathbb{R}_+^m$, and let the solution to $A \otimes x = b$ exist. Then the following are equivalent:*

- (i) $A \otimes x = b$ has unique solution;
- (ii) $S(A, b) = \{\gamma^*(A, b)\}$;
- (iii) $\cup_{j \neq i} M_j(A, b) \neq \text{supp}(b) \quad \forall i: \gamma_i^*(A, b) \neq 0$.

3.2. *Digraph coverings and systems with eigenvector on the right-hand side*

Let $\mathcal{G} = (N, E)$ be a strongly connected digraph. Define

$$M_j(\mathcal{G}) = \{i \in N : (i, j) \in E\} \quad (15)$$

Theorem 3.6. *Let $\mathcal{G} = (N, E)$ be a strongly connected digraph. Then, $i \in N$ with the property $\cup_{j \in N \setminus \{i\}} M_j(\mathcal{G}) = N$ exists if and only if \mathcal{G} is not a cycle.*

PROOF: Observe first that if \mathcal{G} is a cycle then there are no nodes with such property.

To prove the converse, observe that if \mathcal{G} is not a cycle, then it contains two intersecting cycles, and one of the nodes in the intersection will have at least two ingoing edges.

Take a node with at least two incoming edges, and number this node as 1. Put $N_0 = \emptyset$ and $N_1 := \{1\}$. For each $l \geq 1$ let

$$N_{l+1} = N_l \cup \bigcup_{j \in N_l} M_j(\mathcal{G}). \quad (16)$$

Observe that since $\bigcup_{j \in N_{l-1}} M_j(\mathcal{G}) \subseteq N_l$, we can replace (16) with

$$N_{l+1} = N_l \cup \bigcup_{j \in N_l \setminus N_{l-1}} M_j(\mathcal{G}). \quad (17)$$

Hence for each $l \geq 1$ we have

$$N_{l+1} \setminus N_l \subseteq \bigcup_{j \in N_l \setminus N_{l-1}} M_j(\mathcal{G}). \quad (18)$$

Since \mathcal{G} is finite, there is $t > 1$ that satisfies $N_{t+1} = N_t$ and $N_{t-1} \neq N_t$. Observe that $N_t = N$, for if N_t is a proper subset of N , then this contradicts the assumption that \mathcal{G} is strongly connected.

Consider the case when $|N_t \setminus N_{t-1}| > 1$. Observe that a covering of N can be built by taking all nodes in N_{t-1} and a node in $N_t \setminus N_{t-1}$ that has an ingoing edge from 1, or just the nodes in N_{t-1} if one of the nodes of N_{t-1} has an ingoing edge from 1. Therefore in both of these cases there exists $i \in N_t \setminus N_{t-1}$ with $\bigcup_{j \in N \setminus \{i\}} M_j(\mathcal{G}) = N$.

Consider the remaining case when $|N_t \setminus N_{t-1}| = 1$ and when the node forming $N_t \setminus N_{t-1}$ is the only node that has an ingoing edge from 1. Since $|N_2 \setminus N_1| > 1$ there exists an $s < t$ with $|N_{s+1} \setminus N_s| < |N_s \setminus N_{s-1}|$. Since $N_s \setminus N_{s-1}$ does not contain a node to which 1 is connected with an edge, for some node $i \in N_s \setminus N_{s-1}$ we have

$$N_{s+1} \setminus N_s = \bigcup_{j \in N_s \setminus N_{s-1}, j \neq i} M_j(\mathcal{G}). \quad (19)$$

Combining this with (18) for all other $l \neq s$ we obtain that

$$N = \bigcup_{j \neq i} M_j(\mathcal{G}).$$

This completes the proof. \square

Corollary 3.7. *For each $x \in V(A, \lambda)$, an i with the property $\cup_{j \neq i} M_j(\mathcal{G}_c(A, \lambda, x)) = N_c(A, x, \lambda)$ exists if and only if $\mathcal{G}_c(A, x, \lambda)$ is not a union of disjoint cycles.*

We now briefly examine the link to one-sided systems with eigenvector on the right-hand side.

Proposition 3.8. *Let $x \in V(A, \lambda)$, and let $j \in N$ be such that there exists l with $(l, j) \in \text{Sat}(A, x, \lambda)$. Then*

- (i) $x \leq \gamma^*(A, \lambda x)$;
- (ii) $\gamma_j^*(A, x) = x_j$ and $M_j(A, \lambda x) = M_j(\text{Sat}(A, x, \lambda))$;
- (iii) $M_j(\mathcal{G}_c(A, \lambda)) \subseteq M_j(A, \lambda x)$.

PROOF: (i): follows by Lemma 3.1. (ii): Since $x \in V(A, \lambda)$, we have $a_{kj}x_j \leq \lambda x_k$ for all k , so $\gamma_j^*(A, \lambda x) = \min_{i \in N} \{\lambda x_i (a_{ij})^{-1}\} = \lambda x_l (a_{lj})^{-1} = x_j$. $M_j(A, \lambda x) = \{i : a_{ij}x_j = \lambda x_i\}$ follows from the definition of these sets, substituting $\gamma_j^*(A, \lambda x) = x_j$.

(iii): By (ii) we have $M_j(A, \lambda x) = M_j(\text{Sat}(A, x, \lambda))$, and also

$$M_j(\mathcal{G}_c(A, x, \lambda)) \subseteq M_j(\text{Sat}(A, x, \lambda)) = M_j(A, \lambda x)$$

since $\mathcal{G}_c(A, x, \lambda) \subseteq \text{Sat}(A, x, \lambda)$ by Proposition 2.16 part (iii). Graph $\mathcal{G}_c(A, x, \lambda)$ consists of entire components of $\mathcal{G}_c(A, \lambda)$ (Proposition 2.18 part (iii)), hence

$$M_j(\mathcal{G}_c(A, \lambda)) = M_j(\mathcal{G}_c(A, x, \lambda)) \subseteq M_j(\text{Sat}(A, x, \lambda)) = M_j(A, \lambda x).$$

This concludes the proof. \square

3.3. Simple image eigenvectors

Denote by $A^{(i)}$ the matrix which remains after the i th column of A is removed.

Lemma 3.9 (e.g. [3] Theorems 3.1.5 and 3.1.6). *Let $A \in \mathbb{R}_+^{m \times n}$ and $b \in \mathbb{R}_+^m$. Then, b belongs to the simple image set of A if and only if there is no i for which $\gamma_i^*(A, b) \neq 0$ and $b \in \text{span}_{\oplus}(A^{(i)})$.*

PROOF: By Proposition 3.4 $b \in \text{span}_{\oplus}(A^{(i)})$ if and only if $\cup_{j \in N \setminus \{i\}} M_j(A, b) = \text{supp}(b)$. By Proposition 3.5 the non-uniqueness of solution to $A \otimes x = b$ is equivalent to the existence of i with $\cup_{j \in N \setminus \{i\}} M_j(A, b) = \text{supp}(b)$ and $\gamma_i^*(A, b) \neq 0$. \square

Thus, x is not a simple image eigenvector if and only if there exists an i (and λ) such that $x \in \text{span}_{\oplus}(A^{(i)}) \cap V(A, \lambda)$ and $\gamma_i^*(A, \lambda x) > 0$.

Theorem 3.10. $V(A, \lambda)$ contains simple image vectors if and only if there exists a subset $N' \subseteq N$ with the following properties:

- (i) There exists $x \in V(A, \lambda)$ with $\text{supp}(x) = N'$;
- (ii) $\gamma_i^*(A, x) = 0$ for all $i \notin N'$;
- (iii) $N' = N_c(A, x, \lambda)$;
- (iv) All components of $\mathcal{G}_c(A, x, \lambda) = \mathcal{G}_c(A, \lambda)|_{N'}$ are cycles.

PROOF: By Proposition 2.20 we can assume without loss of generality that $\lambda = 1$.

“Only if”: Let us first argue that (i) and (ii) are necessary. Let x be a simple image vector in $V(A, \lambda)$, and take $N' = \text{supp}(x)$. Condition (i) is immediate, and for (ii) we observe that that $\gamma_i^*(A, x) > 0$ for some $i \notin N'$ would imply that x is not the only solution of the system $A \otimes y = x$, since $\gamma^*(A, x) \neq x$.

As condition (ii) depends only on the support of x and on the zero-nonzero pattern of A , it also holds for *arbitrary* $x \in V(A, 1)$ with $\text{supp}(x) = N'$. That condition also implies

$$\text{supp}(y) \subseteq N' \text{ for all } y \text{ such that } A \otimes y = x. \quad (20)$$

Moreover, we can further consider the system $A_{N'N'} \otimes y_{N'} = x_{N'}$ with $x_{N'} \in V(A_{N'N'})$, since we have the following correspondence:

$$\begin{aligned} x_{N'} \in V^+(A_{N'N'}, 1) &\leftrightarrow (x \in V(A, 1) \ \& \ \text{supp}(x) = N'), \\ y_{N'} \in S(A, x_{N'}) &\leftrightarrow y \in S(A, x). \end{aligned} \quad (21)$$

In that correspondence, x and y arise from $x_{N'}$ and $y_{N'}$ by setting $x_{N \setminus N'} = y_{N \setminus N'} = \mathbf{0}$, and $x_{N'}$ and $y_{N'}$ are formed as the usual subvectors of x and y

with indices in N' . The first part of that correspondence follows from Proposition 2.13, and the second part follows from (20).

Therefore, to show that (iii) and (iv) hold we can further assume that x is positive, that is, $N' = N$.

Assume for the contrary that (iii) does not hold. If $N_c(A, 1) \neq N$, consider the indices in $N \setminus N_c(A, 1)$. Every node with index in $N \setminus N_c(A, 1)$ has an outgoing edge in $\text{Sat}(A, x, 1)$. It cannot be that all of these edges also end in $N \setminus N_c(A, 1)$, because then there would be critical cycles and hence critical components in $N \setminus N_c(A, 1)$, a contradiction. Therefore there exists $(i, j) \in E_{\text{Sat}}(A, x, 1)$ with $i \notin N_c(A, 1)$ and $j \in N_c(A, 1)$ implying that

$$N_c(A, 1) \cup \{i\} \subseteq \bigcup_{j \in N_c(A, 1)} M_j(\text{Sat}(A, x, 1)). \quad (22)$$

For any $i' \notin N_c(A, 1)$ there exists j' such that $i' \in M_{j'}(\text{Sat}(A, x, 1))$, and therefore

$$N_c(A, 1) \cup \{i\} \cup \{i'\} \subseteq \bigcup_{j \in N_c(A, 1) \cup \{j'\}} M_j(\text{Sat}(A, x, 1)).$$

Thus adding indices to the left hand side of (22) one by one, we obtain that there exists $k \notin N_c(A, 1)$ such that $N = \cup_{j \neq k} M_j(\text{Sat}(A, x, 1))$, i.e., $N = \cup_{j \neq k} M_j(A, x)$, a contradiction.

Now assume that (iii) holds but (iv) does not hold, i.e., one of the components of $\mathcal{G}_c(A)$ is not a cycle. In this case by Corollary 3.7 there exists i such that $\cup_{j \neq i} M_j(\mathcal{G}_c(A, x, 1)) = N$. However, as $\mathcal{G}_c(A, x, 1) \subseteq \text{Sat}(A, x, 1)$ by Proposition 2.16, we also have $M_j(\mathcal{G}_c(A, x, 1)) \subseteq M_j(A, x)$ for all $j \in N$, implying $\cup_{j \neq i} M_j(A, x) = N$, a contradiction.

“If”: Let us now prove that (i)-(iv) are sufficient. Due to bijection (21) we can assume that $N' = N$. Then, by the main result of [20], there exists a diagonal matrix X such that $\tilde{A} := X^{-1}AX$ is strictly visualised, that is $\tilde{a}_{ij} \leq 1$, with the equality $\tilde{a}_{ij} = 1$ if and only if (i, j) is a critical edge, that is, if and only if (i, j) belongs to one of the disjoint critical cycles. Let u be the vector whose every component is 1. For this vector we obtain $\text{Sat}(\tilde{A}, u, 1) = \mathcal{G}_c(\tilde{A}, u, 1) = \mathcal{G}_c(\tilde{A}, 1)$ and hence $M_j(\tilde{A}, u) = M_j(\mathcal{G}_c(\tilde{A}, 1))$ for all $j \in N$. Since $\mathcal{G}_c(A)$ consists of disjoint cycles only, by Corollary 3.7 we have $\cup_{j \neq i} M_j(\mathcal{G}_c(\tilde{A}, 1)) \neq N$ for any $i \in N$, and since $M_j(\tilde{A}, u) = M_j(\mathcal{G}_c(\tilde{A}, 1))$ for all $j \in N$ we obtain that u is a simple image eigenvector of \tilde{A} . By Lemma 2.23, Xu is a simple image eigenvector of A .

□

Remark 3.11. Condition (i) of Theorem 3.10 can be expressed in terms of the Frobenius normal form of A , see for instance [10] Ch. IV Theorem 2.2.4. Condition (ii) of Theorem 3.10 is equivalent to the following:

$$\forall i \notin N' \exists l \notin N': a_{li} \neq 0. \quad (23)$$

The next result is mostly needed for Theorem 4.17, but it is also of independent interest. We formulate it for positive vectors only, for the sake of simplicity.

Theorem 3.12. *Let $N_c(A, \lambda) = N$ and let all the s.c.c of $\mathcal{G}_c(A, \lambda)$ be cycles and $K = \{j_1, \dots, j_k\}$ be the index set of the columns of A_λ^+ forming $G_{A, \lambda}$. Then*

$$\left(\bigcup_{i \in N} \text{span}_\oplus(A^{(i)}) \cap V^+(A, \lambda) \right) = \left(\bigcup_{s \in K} \text{span}_\oplus^+(G_{A, \lambda}^{(s)}) \right). \quad (24)$$

PROOF: Assume $\lambda = 1$.

For any $x \in V(A, 1)$ we have $x \in \text{span}_\oplus(A)$ and $x \in \text{span}_\oplus(G_{A, 1})$, and if x is positive then we also have coverings $\cup_{i \in N} M_i(A, x) = N$ and $\cup_{r \in K} M_r(G_{A, 1}, x) = N$. The claim of the theorem, reducing to $\exists i: x \in \text{span}_\oplus(A^{(i)}) \Leftrightarrow \exists s: x \in \text{span}_\oplus(G_{A, 1}^{(s)})$ for positive $x \in V(A, 1)$, then amounts to equivalence between the following statements:

- (a) $\exists i \in N: M_i(A, x) \subseteq \bigcup_{j \neq i} M_j(A, x)$ and
- (b) $\exists s \in K: M_s(G_{A, 1}, x) \subseteq \bigcup_{r \neq s} M_r(G_{A, 1}, x)$.

Let $X = \text{diag}(x)$ and consider the matrix $\tilde{A} := X^{-1}AX$. Matrix $G_{\tilde{A}, 1}$ then generates the cone $V(\tilde{A}, 1)$ which is the same as $\{y: Xy \in V(A, 1)\}$, by Lemma 2.23 part (ii). Proposition 3.8 part (ii) implies that $M_i(A, x) = M_i(\text{Sat}(A, x, 1))$ for all i , since all nodes are critical and $\mathcal{G}_c(A, 1) \subseteq \text{Sat}(A, x, 1)$.

By Proposition 2.21 $\tilde{A} =: (\tilde{a}_{ij})_{i, j=1}^n$ has the property $\tilde{a}_{ij} \leq 1$ with the equality if and only if $(i, j) \in \text{Sat}(A, x)$, that is, if and only if $i \in M_j(A, x)$. In \tilde{A}^+ , all columns with indices belonging to the same component of $\mathcal{G}_c(A)$ are equal to each other (Proposition 2.24).

Denote the entries of $G_{A, 1}$ and $G_{\tilde{A}, 1}$ by g_{is} and \tilde{g}_{is} respectively.

Assume (a). It means that there exists $i \in N_c^s$ for some $s \in K$ such that for each $i' \in M_i(A, x) = M_i(\text{Sat}(A, x))$ there is an index $l(i')$ with $l(i') \in N_c^{s'}$ with $s' \neq s$, such that $i' \in M_{l(i')}(A, x) = M_{l(i')}(\text{Sat}(A, x))$. In terms of matrix \tilde{A} it means that

$$\tilde{a}_{i'i} = 1 \Rightarrow \tilde{a}_{i'l(i')} = 1 \text{ with } l(i') \in N_c^{s'}, i \in N_c^s \text{ and } s \neq s'. \quad (25)$$

We will show that for $s \in K$ as above and for every $e \in N$ such that $\tilde{g}_{es} = 1$ we can find s' with $\tilde{g}_{es'} = 1$. Firstly if $\tilde{g}_{es} = 1$ then $\tilde{a}_{ejs}^+ = 1$ and $\tilde{a}_{ei}^+ = 1$ since i and j_s are in the same cycle of $\mathcal{G}_c(A)$. This implies that for some i' there is a path P of weight 1 connecting e to i' , and edge (i', i) of weight 1, in digraph $\mathcal{G}(\tilde{A})$. By (25) there exists index $l(i')$ such that $\tilde{a}_{i'l(i')} = 1$ with $l(i') \in N_c^{s'}$ and $s' \neq s$. Concatenating P with edge $(i', l(i'))$ we obtain a path of weight 1 such that $\tilde{a}_{el(i')}^+ = 1$ and hence $\tilde{g}_{es'} = 1$ with $s' \neq s$.

Assume (b). By Proposition 2.24 part (i) we have $\tilde{g}_{es} = 1$ for any $s \in K$ and $e \in N_c^s$, and statement (b) implies that there also exist e and $r \in K$ such that $\tilde{g}_{er} = 1$ and $e \in N_c^s$ with $s \neq r$. Moreover, by Proposition 2.24 part (ii) we can also assume that $\tilde{g}_{is} < 1$ for all $i \notin N_c^s$.

As $\tilde{g}_{er} = 1$ for all $e \in N_c^s$, we have $\tilde{a}_{ej}^+ = 1$ for all $e \in N_c^s$ and $j \in N_c^r$. This implies that there exist $i_1 \in N_c^s$, $i' \in N_c^{s'}$ such that $\tilde{a}_{i_1 i'} = 1$, where $s \neq s'$. But as $i_1 \in N_c^s$, there also exists $i_2 \in N_c^s$ such that $\tilde{a}_{i_1 i_2} = 1$.

As $\tilde{g}_{es} < 1$ for all $e \notin N_c^s$, we have $\tilde{a}_{ei_2}^+ < 1$ implying also that $\tilde{a}_{ei_2} < 1$ for all $e \notin N_c^s$. Since each component of $\mathcal{G}_c(A, 1)$ is a cycle, we conclude that $M_{i_2}(\text{Sat}(A, x)) = \{i_1\}$, and it is covered by $M_{i'}(\text{Sat}(A, x))$, implying (a). \square

4. Interval problems

4.1. Weak \mathbf{X} -robustness and \mathbf{X} -simple image eigencone

In this section we consider an interval extension of weak robustness and its connection to \mathbf{X} -simplicity, the main notion studied in this paper.

Definition 4.1 (Weak \mathbf{X} -robustness). Let $A \in \mathbb{R}_+^{n \times n}$ and $\mathbf{X} \subseteq \mathbb{R}_+^n$ be an interval.

- (i) $\text{attr}(A, \lambda) := \{x: A^{\otimes t} \otimes x \in V(A, \lambda) \text{ for some } t\}$.
- (ii) A is called weakly (\mathbf{X}, λ) -robust if $\text{attr}(A, \lambda) \cap \mathbf{X} \subseteq V(A, \lambda)$.

If $\mathbf{X} = \mathbb{R}^n$, then the notion of weak robustness can be described in terms of simple image eigenvectors. The proof is omitted.

Proposition 4.2. $A \in \mathbb{R}_+^{n \times n}$. The following are equivalent:

- (i) A is weakly λ -robust;
- (ii) $|S(A, x)| = 1$ for all $x \in V(A, \lambda)$;

(iii) Each $x \in V(A, \lambda)$ is a simple image eigenvector.

Definition 4.3 (Invariance). Let $A \in \mathbb{R}_+^{n \times n}$ and $\mathbf{X} \subseteq \mathbb{R}_+^n$ be an interval. We say that \mathbf{X} is invariant under A if $x \in \mathbf{X}$ implies $A \otimes x \in \mathbf{X}$.

Theorem 4.4. Let $A \in \mathbb{R}_+^{n \times n}$ and \mathbf{X} be an interval.

- (i) If A is weakly (\mathbf{X}, λ) -robust then A has \mathbf{X} -simple image eigencone associated with λ .
- (ii) If A has \mathbf{X} -simple image eigencone associated with λ and if \mathbf{X} is invariant under A then A is weakly (\mathbf{X}, λ) -robust.

PROOF: (i) Suppose that A is weakly (\mathbf{X}, λ) -robust and $x \in V(A, \lambda) \cap \mathbf{X}$. If the system $A \otimes y = \lambda x$ has a solution $y \neq x$ in \mathbf{X} , then y is not an eigenvector but belongs to $\text{attr}(A, \lambda) \cap \mathbf{X}$, which contradicts the weak \mathbf{X} -robustness.

(ii) Assume that A has \mathbf{X} -simple image eigencone and x is an arbitrary element of $\text{attr}(A, \lambda) \cap \mathbf{X}$. As \mathbf{X} is invariant under A , we have that $A^{\otimes k} \otimes x \in \mathbf{X}$ for all k . Moreover from the definition of \mathbf{X} -simple image eigencone we get $A \otimes x \in V(A, \lambda) \Rightarrow x \in V(A, \lambda)$. Then $A^{\otimes k} \otimes x \in V(A, \lambda)$ for some k implies $A^{\otimes(k-1)} \otimes x \in V(A, \lambda), \dots, x \in V(A, \lambda)$. \square

Thus the \mathbf{X} -simplicity is a necessary condition for weak \mathbf{X} -robustness. It is also sufficient if \mathbf{X} is invariant under A .

Observe that A is order-preserving ($x \leq y \Rightarrow A \otimes x \leq A \otimes y$), which is due to the following arithmetic properties:

$$\begin{aligned} x \leq y &\Rightarrow \alpha x \leq \alpha y \quad \forall \alpha, x, y \in \mathbb{R}_+ \\ \alpha_1 \leq \alpha_2, \beta_1 \leq \beta_2 &\Rightarrow \alpha_1 \oplus \beta_1 \leq \alpha_2 \oplus \beta_2 \quad \forall \alpha_1, \alpha_2, \beta_1, \beta_2 \in \mathbb{R}_+. \end{aligned}$$

Since A is order-preserving the invariance of \mathbf{X} under A admits the following simple characterization:

Proposition 4.5. Let \mathbf{X} be closed. Then \mathbf{X} is invariant under A if and only if $\underline{x} \leq A \otimes \underline{x} \leq A \otimes \bar{x} \leq \bar{x}$.

4.2. \mathbf{X} -simple image of a matrix

Let us first introduce the following bits of notation:

Definition 4.6.

$$c(\underline{x}) = \{i \in N : \underline{x}_i \in \mathbf{X}_i\}, \quad o(\underline{x}) = N \setminus c(\underline{x}) \quad (26)$$

Definition 4.7.

$$\begin{aligned}\mathbf{X}^\dagger &= \times_{i=1}^n \mathbf{X}_i^\dagger, \text{ where} \\ \mathbf{X}_i^\dagger &= \mathbf{X}_i \cup [\text{sup}(\mathbf{X}_i), +\infty).\end{aligned}\quad (27)$$

We illustrate the definitions by the following example.

Example 4.8. Suppose that $\mathbf{X} = [1, 2] \times [3, 5] \times (7, 9]$ be given. Then we have $\underline{x} = (1, 3, 7)$, $c(\underline{x}) = (1, 2)$ and $\mathbf{X}^\dagger = [1, \infty) \times [3, \infty) \times (7, \infty)$.

Lemma 4.9. Let $A \in \mathbb{R}_+^{m \times n}$ and $b \in \mathbb{R}_+^m$. A solution to $A \otimes x = b$ in \mathbf{X} exists if and only if

$$\gamma^*(A, b) \in \mathbf{X}^\dagger \ \& \ \bigcup_{j: \gamma_j^*(A, b) \in \mathbf{X}_j} M_j(A, b) = \text{supp}(b). \quad (28)$$

PROOF: “Only if”. Assume that \mathbf{X} contains a solution to $A \otimes x = b$, but (28) does not hold. If $\gamma^*(A, b) \notin \mathbf{X}^\dagger$, then since every solution to $A \otimes x = b$ satisfies $x \leq \gamma^*(A, b)$, we have $x \notin \mathbf{X}^\dagger$ and hence $x \notin \mathbf{X}$ for every solution. Let $\gamma^*(A, b) \in \mathbf{X}^\dagger$, $\bigcup_{j: \gamma_j^*(A, b) \in \mathbf{X}_j} M_j(A, b) \neq \text{supp}(b)$. Since $x \in \mathbf{X}$ we have $x_j < \gamma_j^*(A, b)$ for all j such that $\gamma_j^*(A, b) \notin \mathbf{X}_j$. This implies that $A \otimes x \neq b$, a contradiction.

“If”. Assume that (28) holds. Then by Corollary 3.3 any x with $x_j = \gamma_j^*(A, b)$ for $j: \gamma_j^*(A, b) \in \mathbf{X}_j$, and $x_j \leq \gamma_j^*(A, b)$ for $j: \gamma_j^*(A, b) \in \mathbf{X}_j^\dagger \setminus \mathbf{X}_j$ is a solution to $A \otimes x = b$. Hence $A \otimes x = b$ has a solution in \mathbf{X} . \square

Theorem 4.10. Let $A \otimes x = b$ have a solution in \mathbf{X} . That solution is unique in \mathbf{X} if and only if the following condition is satisfied for each i :

$$\bigcup_{j \neq i} M_j(A, b) = \text{supp}(b) \Rightarrow [i \in c(\underline{x}) \ \& \ \underline{x}_i = \min(\bar{x}_i, \gamma_i^*(A, b))] \quad (29)$$

PROOF: As $A \otimes x = b$ has a solution in \mathbf{X} , condition (28) holds, and by Corollary 3.3 we have

$$\forall \alpha \in \mathbf{X}_i \text{ s.t. } \alpha \leq \gamma_i^*(A, b) \ \exists x \in S(A, b) \cap \mathbf{X}: x_i = \alpha. \quad (30)$$

In particular if $\gamma_i^*(A, b) \in \mathbf{X}_i^\dagger \setminus \mathbf{X}_i$ then

$$\forall \alpha \in \mathbf{X}_i \ \exists x \in S(A, b) \cap \mathbf{X}: x_i = \alpha. \quad (31)$$

“Only if”: Suppose that there is a unique solution belonging to \mathbf{X} . By (31) if \mathbf{X}_i does not reduce to one point, then $S(A, b)$ also contains more than one vector. Thus, $\underline{x}_i = \bar{x}_i$ for all i such that $\gamma_i^*(A, b) \in \mathbf{X}_i^\uparrow \setminus \mathbf{X}_i$, which satisfies (29).

If we assume that (29) does not hold for some i with $\gamma_i^*(A, b) \in \mathbf{X}_i$ then $\bigcup_{j \neq i} M_j(A, b) = \text{supp } b$ and (31) implies that the solution is non-unique since the interval $\mathbf{X}_i \cap \{\alpha : \alpha \leq \gamma_i^*(A, b)\}$ contains more than one point.

“If”: Assume that (29) holds and, by contradiction, that there is more than one solution to $A \otimes x = b$ belonging to \mathbf{X} .

Since the solution is non-unique, it follows that there exists a proper subset N' of N such that $\bigcup_{j \in N'} M_j(A, b) = \text{supp}(b)$. Assume that N' is a minimal such subset, with respect to inclusion.

We have $\bigcup_{j \neq i} M_j(A, b) = \text{supp}(b)$ for all $i \in N \setminus N'$. By (29) $N \setminus N' \subseteq c(\underline{x})$, and $\underline{x}_i = \min(\bar{x}_i, \gamma_i^*(A, b))$ for all $i \in N \setminus N'$. This condition implies that there exists only one N' -solution to $A \otimes x = b$ belonging to \mathbf{X} , and it has coordinates

$$x_i = \begin{cases} \underline{x}_i, & \text{if } \underline{x}_i = \bar{x}_i, \\ \gamma_i^*(A, b), & \text{otherwise.} \end{cases}$$

As this solution is the same for any minimal subset N' , system $A \otimes x = b$ has a unique solution, contradicting the non-uniqueness of it. \square

Corollary 4.11. *Let $x \in V(A) \cap \mathbf{X}$. Then x is an \mathbf{X} -simple image eigenvector if and only if*

$$\bigcup_{j \neq i} M_j(A, b) = \text{supp}(x) \Rightarrow [i \in c(\underline{x}) \ \& \ \underline{x}_i = \min(\bar{x}_i, \gamma_i^*(A, b))] \quad (32)$$

If $N_c = N$ then this condition can be replaced with the following one:

$$\bigcup_{j \neq i} M_j(A, b) = \text{supp}(x) \Rightarrow [i \in c(\underline{x}) \ \& \ \underline{x}_i = \min(\bar{x}_i, x_i)] \quad (33)$$

PROOF: As $x \in V(A) \cap \mathbf{X}$, system $A \otimes y = x$ has a solution in \mathbf{X} , which is $y = x$. So we can apply Theorem 4.10 yielding (32) for the uniqueness of this solution. Further if all nodes are critical, then $\gamma_i^*(A, x) = x_i$ for all x , so (32) gets replaced with (33). \square

4.3. Characterizing matrices with \mathbf{X} -simple image eigencone

We begin with the following definition and key lemma of geometric kind.

Definition 4.12. An interval $\mathbf{X} \subseteq \mathbb{R}_+^n$ is called \underline{x} -open if $o(\underline{x}) = N$. It is called \bar{x} -closed if it $\bar{x} \in \mathbf{X}$.

Lemma 4.13. Let \mathbf{X} be an \bar{x} -closed interval and let $A \in \mathbb{R}^{m \times n}$. Then $\mathbf{X} \cap \text{span}_{\oplus}(A) \neq \emptyset$ if and only if $P_A \bar{x} \in \mathbf{X}$.

PROOF: If $P_A \bar{x} \in \mathbf{X}$ then $\mathbf{X} \cap \text{span}_{\oplus}(A) \neq \emptyset$ (since $P_A \bar{x} \in \text{span}_{\oplus}(A)$).

If $\mathbf{X} \cap \text{span}_{\oplus}(A) \neq \emptyset$, take $y \in \mathbf{X} \cap \text{span}_{\oplus}(A) \neq \emptyset$. Vector $z = y \oplus P_A \bar{x}$ belongs to $\text{span}_{\oplus}(A)$ and satisfies $y \leq z \leq \bar{x}$. It follows then that $z \in \mathbf{X}$. However, $P_A \bar{x} \leq z \leq \bar{x}$ while $P_A \bar{x}$ is the greatest vector of $\text{span}_{\oplus}(A)$ bounded from above by \bar{x} . This implies that $z = P_A \bar{x}$ and $P_A \bar{x} \in \mathbf{X}$. \square

Definition 4.14. For any $l, i \in N$, let

$$\bar{x}_i^{(l)} = \begin{cases} \underline{x}_l, & \text{if } i = l, \\ \bar{x}_i, & \text{otherwise,} \end{cases} \quad (34)$$

and let $\bar{x}^{(l)} = (\bar{x}_i^{(l)})_{i \in N}$.

Definition 4.15. For any $l \in N$, let

$$\begin{aligned} \mathbf{X}_{A,\lambda}^l &= \{x \in \mathbf{X} : x_l = \underline{x}_l, \quad \lambda x_j > a_{jl} \underline{x}_l \quad \forall j\}, \\ \mathbf{X}^{(l)} &= \{x \in \mathbf{X} : x_l > \underline{x}_l\}. \end{aligned} \quad (35)$$

Observe that if \mathbf{X} is \bar{x} -closed then \mathbf{X}_A^l is $\bar{x}^{(l)}$ -closed (if it is nonempty) for every l . Also note that $a_{ll} < \lambda$ is a necessary condition for $\mathbf{X}_{A,\lambda}^l$ to be non-empty.

Theorem 4.16. Let $\lambda > 0$ and $V(A, \lambda) \cap \mathbf{X} \neq \emptyset$.

(i) A has \mathbf{X} -simple image eigencone corresponding to λ if and only if

$$\begin{aligned} \text{span}_{\oplus}(A^{(i)}) \cap V(A, \lambda) \cap \mathbf{X}^{(i)} &= \emptyset \quad \forall i, \\ V(A, \lambda) \cap \mathbf{X}_{A,\lambda}^l &= \emptyset \quad \forall l \in c(\underline{x}) \setminus N_c(A, \lambda) \text{ s.t. } \underline{x}_l < \bar{x}_l. \end{aligned} \quad (36)$$

(ii) If \mathbf{X} is \bar{x} -closed then (36) is equivalent to

$$\begin{aligned} P_{\mathcal{W}^i} \bar{x} &\notin \mathbf{X}^{(i)} \quad \forall i \in N \text{ where } \mathcal{W}^{(i)} = \text{span}_{\oplus}(A^{(i)}) \cap V(A, \lambda), \\ P_{V(A, \lambda)} \bar{x}^{(l)} &\notin \mathbf{X}_{A, \lambda}^l, \quad \forall l \in c(\underline{x}) \setminus N_c(A, \lambda) \text{ s.t. } \underline{x}_l < \bar{x}_l. \end{aligned} \quad (37)$$

PROOF: Assume without loss of generality that $\lambda = 1$.

(i): Let $x \in V(A, 1) \cap \mathbf{X}$. In general, $x \leq \gamma^*(A, x)$. More precisely, for $\gamma_i^*(A, x)$ we may have $\gamma_i^*(A, x) = x_i$ (for all $i \in N_c(A, 1)$ and some other nodes), or $\gamma_i^*(A, x) > x_i$ (for at least one node in $N \setminus N_c(A, 1)$ if $N \neq N_c(A, 1)$).

“Only if”: Let us show that the conditions (36) are necessary. For this assume by contradiction that one of these conditions is violated but A has \mathbf{X} -simple image eigencone.

(a) If the first condition does not hold then take $x' \in \mathbf{X}^{(i)} \cap V(A, 1) \cap \text{span}_{\oplus}(A^{(i)})$. Then x' satisfies $x'_i > \underline{x}_i$ and also $\cup_{j \neq i} M_j(A, x') = \text{supp}(x')$, hence x' is not an \mathbf{X} -simple image eigenvector by Corollary 4.11: either $i \in o(\underline{x})$ and $\cup_{j \neq i} M_j(A, x') = \text{supp}(x')$, or $i \in c(\underline{x})$, $\cup_{j \neq i} M_j(A, x') = \text{supp}(x')$ and $\underline{x}_i < \min(\gamma_i^*(A, x'), \bar{x}_i)$.

(b) If the second condition does not hold then take $x'' \in V(A, 1) \cap \mathbf{X}_A^l$ for some $l \in c(\underline{x}) \setminus N_c(A, 1)$ such that $\underline{x}_l < \bar{x}_l$. We have $x''_l = \underline{x}_l$, $\underline{x}_l < \bar{x}_l$ and $\gamma_l^*(A, x'') > x''_l$ (equivalent with the condition $x''_j > a_{jl} x''_l \quad \forall j$ from (35)). The inequality $\gamma_l^*(A, x'') > x''_l$ implies that $\cup_{j \neq l} M_j(A, x'') = \text{supp}(x'')$, and we also have $\underline{x}_l < \min(\gamma_l^*(A, x''), \bar{x}_l)$ and $l \in c(\underline{x})$. By Corollary 4.11, this shows that x is not an \mathbf{X} -simple image eigenvector.

“If”: By contradiction, suppose that the conditions hold but A does not have simple image eigencone. Let x be an \mathbf{X} -simple image eigenvector. Then either $x_i > \underline{x}_i$ and $\cup_{j \neq i} M_j(A, x) = \text{supp}(x)$ for some i , which implies $x \in \mathbf{X}^{(i)} \cap V(A) \cap \text{span}_{\oplus}(A^{(i)})$, or $x_l = \underline{x}_l$, $\cup_{j \neq l} M_j(A, x) = \text{supp}(x)$ and $\underline{x}_l < \min(\gamma_l^*(A, x), \bar{x}_l)$ for some $l \in c(\underline{x})$. In this case necessarily $\underline{x}_l < \bar{x}_l$ and $x_l < \gamma_l^*(A, x)$, which is only possible for $l \in N \setminus N_c$.

This shows the sufficiency of (36).

(ii) By Lemma 4.13, $\mathcal{W}^i \cap \mathbf{X}^{(i)} \neq \emptyset$ if and only if $P_{\mathcal{W}^i} \bar{x} \in \mathbf{X}^{(i)}$, and $V(A, 1) \cap \mathbf{X}_A^l \neq \emptyset$ if and only if $P_{V(A, 1)} \bar{x}^{(l)} \in \mathbf{X}_A^l$. \square

In general, the basis of $\text{span}_{\oplus}(A^{(i)}) \cap V(A, \lambda)$ can be computed algorithmically, using the method of Butkovič, Hegedus [5] or the more recent and efficient methods of Allamigeon et al. [1]

Let us now examine the case when \mathbf{X} is \underline{x} -open.

Theorem 4.17. *Let \mathbf{X} be an \underline{x} -open interval and $V(A, \lambda) \cap \mathbf{X}$ be non-empty.*

(i) *A has \mathbf{X} -simple image eigencone corresponding to λ if and only if*

$$\text{span}_{\oplus}(A^{(i)}) \cap V(A, \lambda) \cap \mathbf{X} = \emptyset \quad \forall i = 1, \dots, n. \quad (38)$$

(ii) *In that case $N_c(A, \lambda) = N$ and $\mathcal{G}_c(A, \lambda)$ consists of disjoint cycles c_1, \dots, c_k for some k .*

(iii) *Condition (38) is equivalent to*

$$\text{span}_{\oplus}(G_{A^{(s)}, \lambda}) \cap \mathbf{X} = \emptyset \quad \forall s = 1, \dots, k. \quad (39)$$

(iv) *If \mathbf{X} is also \bar{x} -closed then (38) is also equivalent to*

$$P_{G_{A^{(s)}, \lambda}} \bar{x} \not\prec \underline{x} \quad \forall s = 1, \dots, k. \quad (40)$$

PROOF: Assume $\lambda = 1$. (i): Follows from Theorem 4.16 part (i), where we take into account that $c(\underline{x})$ is empty and $\mathbf{X}^{(i)} = \mathbf{X}$ for each i .

(ii): Note that each vector in $V(A, 1) \cap \mathbf{X}$ is positive. By Theorem 3.10, if $\mathcal{G}_c(A)$ does not consist of disjoint cycles or $N_c \neq N$ then each vector $x \in V(A, 1)$ belongs to $\text{span}_{\oplus}(A^{(i)})$ for some $i \in N$. Hence $\text{span}_{\oplus}(A^{(i)}) \cap V(A, 1) \cap \mathbf{X} \neq \emptyset$ for some i , a contradiction.

(iii): By Theorem 3.12 we have

$$\left(\bigcup_{i \in N} \text{span}_{\oplus}(A^{(i)}) \right) \cap V(A, 1) = \bigcup_{s=1}^k \text{span}_{\oplus}(G_{A^{(s)}, 1}).$$

Hence (38) and (39) are equivalent.

(iv): By applying Lemma 4.13 to (39), we obtain that (39) is equivalent to

$$P_{G_{A^{(s)}, 1}} \bar{x} \notin \mathbf{X} \quad \forall s = 1, \dots, k, \quad (41)$$

and that is the same as (40). \square

- [1] X. Allamigeon, É. Goubault and S. Gaubert. Computing the Vertices of Tropical Polyhedra using Directed Hypergraphs. *Discrete and Computational Geometry*, 49:2 (2013), 247–279.
- [2] F.L. Baccelli, G. Cohen, G.J. Olsder and J.P. Quadrat. *Synchronization and Linearity*. Wiley and Sons, 1992. Available online: <https://www.rocq.inria.fr/metalau/cohen/SED/book-online.html>
- [3] P. Butkovič. *Max-linear Systems: Theory and Algorithms*, Springer Monographs in Mathematics, Springer-Verlag 2010.
- [4] P. Butkovič. Simple image set of $(\max, +)$ linear mappings. *Discrete Appl. Math.* vol. 105 (2000), 73 – 86.
- [5] P. Butkovič and G. Hegedüs. An elimination method for finding all solutions of the system of linear equations over an extremal algebra *Ekonomicko-Matematický Obzor*, 20 (1984), pp. 203–214
- [6] P. Butkovič, H. Schneider and S. Sergeev. Generators, extremals and bases of max cones. *Linear Algebra Appl.* **421**, 2007, 394-406.
- [7] P. Butkovič, H. Schneider and S. Sergeev. Recognizing weakly stable matrices. *SIAM J. Control Optim.* **50** (5), 2012, 3029-3051.
- [8] G. Cohen, S. Gaubert, J.P. Quadrat and I. Singer. Max-plus convex sets and functions. In: G.L. Litvinov and V.P. Maslov (eds.), *Idempotent Mathematics and Mathematical Physics*, Cont. Math. **377** , AMS, 2005, pp. 105–129,
- [9] R. A. Cuninghame-Green. *Minimax Algebra*. Lecture Notes in Economics and Mathematical Systems **166**, Springer, Berlin, 1979.
- [10] S. Gaubert. *Théorie des systèmes linéaires dans les dioïdes*. PhD Thesis, L'École des Mines de Paris, 1992. Available online: <http://www.cmap.polytechnique.fr/~gaubert/PAPERS/ALL.pdf>
- [11] R. A. Cuninghame-Green. *Minimax algebra and applications*. Advances in Imaging and Electron Physics **90**, (1995) 1–121.
- [12] M. Gavalec, K. Zimmermann. Classification of solutions to systems of two-sided equations with interval coefficients. *Inter. J. of Pure and Applied Math.* **45** (2008), 533–542.

- [13] B. Heidergott, G.-J. Olsder, and J. van der Woude. *Max-plus at Work*. Princeton Univ. Press, 2005.
- [14] H. Myšková. Interval systems of max-separable linear equations. *Linear Algebra and Its Applications* 403 (2005) 263–272.
- [15] H. Myšková. Control solvability of interval systems of max-separable linear equations. *Linear Algebra and Its Applications* 416 (2006) 215–223.
- [16] J. Plavka. The weak robustness of interval matrices in max-plus algebra. *Discrete Appl. Math.* 173 (2014) 92–101.
- [17] J. Plavka. The weak robustness of interval matrices in max-plus algebra. *Discrete Applied Mathematics* 173 (2014) 92–101.
- [18] J. Rohn. Systems of Linear Interval Equations. *Linear Algebra and Its Applications* 126 (1989) 39–78.
- [19] S. Sergeev. Max-algebraic cones of nonnegative irreducible matrices, *Linear Algebra and its Applications* 435 (2011), 1736–1757.
- [20] S. Sergeev, H. Schneider and P. Butkovič. On visualization scaling, subeigenvectors and Kleene stars in max algebra. *Linear Algebra and Its Applications* 431 (2009) 2395–2406.
- [21] N.N. Vorobyev. Extremal algebra of positive matrices (in Russian). *Elektronische Informationsverarbeitung und Kybernetik*, 3 (1967) 39–71.
- [22] K. Zimmermann. *Extremální algebra* (in Czech), Ekon. ústav ČSAV Praha, 1976.