# UNIVERSITY<sup>OF</sup> BIRMINGHAM University of Birmingham Research at Birmingham

Evidence for Increased 5α-Reductase Activity During Early Childhood in Daughters of Women with Polycystic Ovary Syndrome

Arlt, Wiebke; Torchen, Laura; Idkowiak, Jan; Fogel, Naomi; O'Neil, Donna; Shackleton, Cedric; Dunaif, Andrea

DOI: 10.1210/jc.2015-3926#sthash.9Wp6MrQ4.dpuf

License: Creative Commons: Attribution (CC BY)

Document Version Publisher's PDF, also known as Version of record

Citation for published version (Harvard):

Arlt, W, Torchen, L, Idkowiak, J, Fogel, N, O'Neil, D, Shackleton, C & Dunaif, A 2016, 'Evidence for Increased 5α-Reductase Activity During Early Childhood in Daughters of Women with Polycystic Ovary Syndrome', *Journal of Clinical Endocrinology and Metabolism*, vol. 101, no. 5, pp. 2069-2075. https://doi.org/10.1210/jc.2015-3926#sthash.9Wp6MrQ4.dpuf

Link to publication on Research at Birmingham portal

#### **General rights**

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

•Users may freely distribute the URL that is used to identify this publication.

•Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.

•User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?) •Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

#### Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

# **Evidence for Increased 5***α***-Reductase Activity During Early Childhood in Daughters of Women With Polycystic Ovary Syndrome**

Laura C. Torchen, Jan Idkowiak, Naomi R. Fogel, Donna M. O'Neil, Cedric H. L. Shackleton, Wiebke Arlt,\* and Andrea Dunaif\*

Division of Endocrinology, Metabolism, and Molecular Medicine (A.D.), Feinberg School of Medicine, Northwestern University, Chicago, Illinois 60611; Division of Pediatric Endocrinology (L.C.T., N.R.F.), Ann & Robert H. Lurie Children's Hospital of Chicago, Feinberg School of Medicine, Northwestern University, Chicago, Illinois 60611; Institute of Metabolism and Systems Research (J.I., D.M.O., C.H.L.S., W.A.), University of Birmingham, Birmingham B15 2TT, UK; Centre for Endocrinology, Diabetes and Metabolism (J.I., W.A.), Birmingham Health Partners, Birmingham B15 2TT, UK

**Context:** Polycystic ovary syndrome (PCOS) is a heritable, complex genetic disease. Animal models suggest that androgen exposure at critical developmental stages contributes to disease pathogenesis. We hypothesized that genetic variation resulting in increased androgen production produces the phenotypic features of PCOS by programming during critical developmental periods. Although we have not found evidence for increased in utero androgen levels in cord blood in the daughters of women with PCOS (PCOS-d), target tissue androgen production may be amplified by increased  $5\alpha$ -reductase activity analogous to findings in adult affected women. It is possible to noninvasively test this hypothesis by examining urinary steroid metabolites.

**Objective:** We performed this study to investigate whether PCOS-d have altered androgen metabolism during early childhood.

**Design, Setting, and Participants:** Twenty-one PCOS-d, 1–3 years old, and 36 control girls of comparable age were studied at an academic medical center.

Main Outcome Measures: Urinary steroid metabolites were measured by gas chromatography/ mass spectrometry. Twenty-four hour steroid excretion rates and precursor to product ratios suggestive of  $5\alpha$ -reductase and  $11\beta$ -hydroxysteroid dehydrogenase activities were calculated.

**Results:** Age did not differ but weight for length Z-scores were higher in PCOS-d compared to control girls (P = .02). PCOS-d had increased  $5\alpha$ -tetrahydrocortisol:tetrahydrocortisol ratios (P = .04), suggesting increased global  $5\alpha$ -reductase activity. There was no evidence for differences in  $11\beta$ -hydroxysteroid dehydrogenase activity. Steroid metabolite excretion was not correlated with weight.

**Conclusions:** Our findings suggest that differences in androgen metabolism are present in early childhood in PCOS-d. Increased  $5\alpha$ -reductase activity could contribute to the development of PCOS by amplifying target tissue androgen action. (*J Clin Endocrinol Metab* 101: 2069–2075, 2016)

Polycystic ovary syndrome (PCOS) is a common, highly heritable disorder with substantial reproductive and metabolic morbidities (1). Hyperandrogenemia is the car-

First Published Online March 18, 2016

dinal reproductive phenotype in adult female first-degree relatives of women with PCOS (2). Further, hyperandrogenemia is present in the daughters (PCOS-d) of affected

doi: 10.1210/jc.2015-3926

J Clin Endocrinol Metab, May 2016, 101(5):2069–2075 press.endocrine.org/journal/jcem 2069

RIGHTSLINK()

ISSN Print 0021-972X ISSN Online 1945-7197 Printed in USA

This article has been published under the terms of the Creative Commons Attribution License (CC-BY; https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited. Copyright for this article is retained by the author(s). Received November 11, 2015. Accepted March 14, 2016.

<sup>\*</sup> W.A. and A.D. served as joint senior authors.

Abbreviations:  $5\alpha$ THF,  $5\alpha$ -reduced tetrahydrocortisol;  $5\alpha$ THF/THF,  $5\alpha$ THF/s  $5\beta$ -reduced metabolite; An, androsterone;  $11\beta$ HSD1,  $11\beta$ -hydroxysteroid dehydrogenase; BMI, body mass index; Et, etiocholanolone; PCOS, polycystic ovary syndrome; PCOS-d, daughters of women with PCOS; THE, tetrahydrocortisone.

women (3). We have found increased circulating testosterone (T) levels in PCOS-d as young as age 8 years (4). Although a number of susceptibility loci for PCOS have been reproducibly mapped (5, 6), these loci account for less than 5% of the approximately 70% heritability of PCOS (7). Epigenetic mechanisms may account for some of this so-called "missing heritability" that has been observed in PCOS and other complex genetic traits (8). Phenocopies of PCOS can be produced in animal models by exposure to androgens during gestation (9, 10), neonatally (11), or peripubertally (12). Therefore, the PCOS phenotype may result from the interactions of genetic variation resulting in hyperandrogenemia and the epigenetic actions of these androgens in utero as well as during critical postnatal developmental windows.

The source of androgens could be increased glandular androgen production. However, at birth, most studies (13, 14) suggest that cord blood androgen levels in PCOS-d are decreased. It remains possible that target tissue androgen action is amplified by upregulation of  $5\alpha$ reductase activity resulting in increased production of DHT from T (15). If such increases in  $5\alpha$ -reductase activity were the result of variation in PCOS susceptibility genes, target tissue androgen action would be enhanced in affected individuals when T was available.

Alterations in intracellular steroidogenic enzyme activity could also increase adrenal androgen production. Increased  $5\alpha$ -reductase or decreased  $11\beta$ -hydroxysteroid dehydrogenase type 1 ( $11\beta$ HSD1) activity would enhance cortisol metabolism resulting in a compensatory increase in ACTH secretion and stimulation of adrenal steroidogenesis (16, 17). Indeed, evidence for increased  $5\alpha$ -reductase and decreased  $11\beta$ HSD1 activity has been found in adult women with PCOS (16-21). It is possible to noninvasively assess global androgen metabolism by measuring urinary metabolites and precursor: product ratios. We tested the hypothesis that  $5\alpha$ -reductase activity is increased in PCOS-d, who are at genetically high risk for PCOS, by measuring urinary steroid metabolites.

## **Materials and Methods**

#### Study population

Twenty-one PCOS-d and 36 control girls aged 1 to 3 years were studied. We chose this age range to ensure that all subjects were at a similar developmental stage when the mini-puberty of infancy was completed, but the infants were still in diapers for ease of urine collection (22). Target sample sizes of 20 cases and 40 controls were calculated for the primary end point, the ratio of 5 $\alpha$ -reduced tetrahydrocortisol (5 $\alpha$ THF) to its 5 $\beta$ -reduced metabolite, THF (5 $\alpha$ THF/THF), a measure of 5 $\alpha$ -reductase activity (17, 23), assuming 80% power and alpha of 0.05 based on the

mean and SD of this ratio in previous studies comparing PCOS to control women (17, 23). Subjects were recruited by advertisements in local media and on-line, and by contacting women who have previously participated in our studies of PCOS and control adult women. PCOS-d had a mother who fulfilled National Institutes of Health criteria for PCOS (hyperandrogenism and oligo-anovulation with exclusion of other reproductive disorders (1) as confirmed before recruitment by us or by their personal physician. The mothers of control girls had regular menses every 27-35 days and no signs or symptoms of androgen excess by a validated questionnaire (2). All girls were in good health and not taking any medications known to alter steroid hormone metabolism. The Institutional Review Boards of the Feinberg School of Medicine, Northwestern University, and Ann & Robert H. Lurie Children's Hospital of Chicago approved this study. Written informed consent was obtained from the parent or legal guardian of all subjects.

#### Data collection

Age, height, weight, medical and reproductive histories were ascertained by validated questionnaires (2) in PCOS and control mothers. Timed urine samples were obtained in PCOS-d and control girls through collection of consecutive wet diapers for a minimum of 10 and maximum of 24 hours. Pediatric growth records for height and weight were available in 21 PCOS-d and 30 control girls. Weight for length Z score was calculated based on the US Centers for Disease Control and Prevention 2000 growth charts (24).

#### Assays

Urine was extracted from diapers as described previously (25) and urinary steroid metabolites measured by a quantitative gas chromatography-mass spectrometry (GC-MS) selected ionmonitoring method (18). Metabolite concentrations were corrected to reflect 24-hour excretion rates by applying the following formula: [(metabolite concentration ( $\mu$ g/L) × urine volume (L))/(24/urine collection time (h))]. Substrate-to-product metabolite ratios were calculated to reflect the in vivo net activity of steroidogenic enzymes. Specifically, the ratio of the glucocorticoid metabolite  $5\alpha$ THF to its  $5\beta$ -reduced metabolite, THF  $(5\alpha THF/THF)$ , was the primary ratio used to assess  $5\alpha$ -reductase activity because of the low concentrations of androgen and mineralocorticoid metabolites in this age group (Table 2) (26). The following ratios were used to assess 11<sup>β</sup>-hydroxysteroid dehydrogenase type 1 (11 $\beta$ -HSD1) activity: (5 $\alpha$ THF + THF)/tetrahydrocortisone (THE), and cortols/cortolones (27). Total cortisol production was quantified by the sum of cortisol metabolites: THF +  $5\alpha$ THF + cortisol + cortols (26). Total cortisone production was quantified by the sum cortisone metabolites: THE + cortisone + cortolones (26). Total glucocorticoid production was quantified by the sum of total cortisol and total cortisone production (26). Total androgen production was quantified by the sum of androsterone (An) + etiocholanolone (Et) (26).

#### **Statistical analysis**

Data were log-transformed when necessary to achieve homogeneity of variance. Differences between groups were assessed by two-tailed unpaired t tests or Mann-Whitney tests, if homogeneity of variance was not achieved. Because weight for length Z score was higher in the PCOS-d, Pearson or Spear-

The Endocrine Society. Downloaded from press.endocrine.org by [\$[individualUser.displayName]] on 24 June 2016. at 02:11 For personal use only. No other uses without permission. . All rights reserved

| Mothers  | PCOS Mothers<br>n = 21  | Control Mothers<br>n = 36  | P Value     |  |  |
|--|---|--|-------------|--|--|
| Age, y<br>BMI, kg/m <sup>2</sup>                 | 32 ± 4<br>31 ± 8  | 33 ± 3<br>26 ± 7   | .77<br>.01  |  |  |
| Daughters  | PCOS-d<br>n = 21  | Control Girls<br>n = 36  |             |  |  |
| Age, y<br>Weight for length Z-score <sup>a</sup> | $1.7 \pm 0.6$<br>$0.43 \pm 1.04$  | $1.9 \pm 0.5$<br>-0.29 ± 0.99  | 0.45<br>.02 |  |  |
| Length of urine collection, h                    | 6% White, non-Hispanic, 6% Hispanic,<br>6% Asian Indian, 3% Asian<br>17.8 $\pm$ 4.5 | 10% White, non-Hispanic, 10% Hispanic,<br>10% Black, non-Hispanic, 5% Asian Indian<br>16.5 $\pm$ 4.7 | .33         |  |  |

# **Table 1.**Baseline Characteristics

Data are presented as mean  $\pm$  sp with the exception of the categorical variable of race/ethnicity, where the percentage of subjects included in each group are noted. *P* values listed from *t* tests unless noted otherwise.

<sup>a</sup> Weight and length data available on 21 PCOS-d and 30 control girls.

<sup>b</sup> Categorical variables analyzed by Fisher's exact test.

man correlation, depending on the normality of the data, was performed on all endpoints to determine whether there was a significant correlation with weight for length Z score. Because there was no correlation of weight for length Z score with any endpoint, no statistical adjustment for weight was necessary. Statistical analyses were performed using SAS 9.4 (SAS Institute, Inc.). Categorical variables were compared by Fisher's exact test. Differences were considered to be significant at  $P \leq$ .05. Data are reported as the untransformed mean  $\pm$  SD.

Urinary Metabolite Excretion PCOS Daughters and Control Girls

# Results

The PCOS mothers had increased body mass index (BMI) compared to control mothers but their age did not differ (Table 1). Age and ethnicity of the PCOS-d and control girls were similar by design (Table 1). Weight for length Z score (P = .02, Table 1) was higher in the PCOS-d. The length of urine collection did not differ between the groups (P = .28, Table 1).

|                                  | PCOS-d<br>n = 21 | Control Girls<br>n = 36 | P Value | P Value Correlation<br>Weight-for-Length Z Score |  |
|----------------------------------|------------------|-------------------------|---------|--|--|
| Cortisol metabolites             |                  |                         |         |  |  |
| THF                              | 44.7 ± 41.3      | 39.0 ± 30.7             | .64     | .50  |  |
| $5\alpha$ -THF                   | 115.2 ± 130.6    | $68.8 \pm 48.0$         | .13     | .40  |  |
| $\alpha$ -cortol                 | $11.5 \pm 11.0$  | $7.4 \pm 5.5$           | .07     | .82  |  |
| $\beta$ -cortol                  | $24.7 \pm 25.9$  | $16.8 \pm 14.5$         | .27     | .44  |  |
| Cortisol <sup>a</sup>            | 9.4 ± 10.2       | 6.8 ± 9.1               | .14     | .87  |  |
| Cortisone metabolites            |                  |                         |         |  |  |
| THE                              | 192.4 ± 200.6    | 165.5 ± 132.9           | .69     | .56  |  |
| $\alpha$ -cortolone <sup>b</sup> | $40.4 \pm 32.6$  | 34.0 ± 28.6             | .64     | .73  |  |
| $\beta$ -cortolone <sup>b</sup>  | 37.1 ± 30.2      | 32.9 ± 24.2             | .80     | .84  |  |
| Cortisone                        | $6.2 \pm 5.2$    | $5.9 \pm 6.2$           | .51     | .85  |  |
| Androgen metabolites             |                  |                         |         |  |  |
| An                               | 2.9 ± 3.1        | $2.5 \pm 2.8$           | .42     | .45  |  |
| Et                               | $1.6 \pm 1.4$    | $1.6 \pm 1.7$           | .59     | .64  |  |
| Mineralocorticoid metabolites    |                  |                         |         |  |  |
| THDOC                            | $1.7 \pm 2.0$    | $1.2 \pm 2.0$           | .06     | .47  |  |
| 5α-THDOC                         | $2.3 \pm 2.2$    | $1.4 \pm 1.7$           | .05     | .69  |  |
| THA                              | $11.1 \pm 15.0$  | $8.0 \pm 6.5$           | .49     | .35  |  |
| 5α-THA                           | 12.9 ± 16.3      | $7.3 \pm 5.6$           | .08     | .25  |  |
| THB                              | 7.8 ± 11.9       | $4.5 \pm 4.2$           | .22     | .37  |  |
| $5\alpha$ -THB                   | 45.6 ± 74.1      | 19.8 ± 16.7             | .07     | .25  |  |

All metabolite concentrations reported in  $\mu$ g/24 h. The *t* tests were performed for two-group comparisons and Spearman correlations for correlation with weight for length Z-scores unless noted otherwise.

Abbreviations: THA, tetrahydro-11-dehydrocorticosterone; THB, tetrahydrocorticosterone; THDOC, tetrahydrodeoxycorticosterone; THF, tetrahydrocortisol.

 $^{\rm a}$  One outlier PCOS-d cortisol result excluded (result  ${>}4$  sp above mean).

<sup>b</sup> Nonparametric tests performed for  $\alpha$ -cortolone and  $\beta$ -cortolone because of a failure to achieve homogeneity of variance.

Table 2.

The Endocrine Society. Downloaded from press.endocrine.org by [\${individualUser.displayName}] on 24 June 2016. at 02:11 For personal use only. No other uses without permission. All rights reserved



**Figure 1.** Steroidogenesis pathway highlighting steroidogenic enzyme activities and urinary steroid metabolites. Urinary steroid metabolites are circled and steroidogenic enzymes are indicated by black boxes. \*The  $5\alpha$ -reduced metabolites. ALDO, aldosterone; B, corticosterone;  $3\beta$ -HSD,  $3\beta$ -hydroxysteroid dehydrogenase;  $11\beta$ -HSD,  $11\beta$ -hydroxysteroid dehydrogenase;  $17\beta$ -HSD,  $17\beta$ -hydroxysteroid dehydrogenase; D4, androstenedione; DOC, deoxycorticosterone; DHEA, dehydroepiandrosterone; 17HP, 17-OHpregnanolone; 18OHB, 18-hydrocxycorticosterone; 17OHP, 17-hydroxyprogesterone; PD, pregnanediol; 5PD, pregnanediol; PREG, pregnenolone; 17PREG, 17-hydroxypregnenolone; PROG, progesterone; PT, pregnanetriol; 5PT, 5-pregnenetriol; S, 11-deoxycortisol; THA, tetrahydro-11-dehydrocorticosterone; THB, tetrahydrocorticosterone; THDOC, tetrahydrodeoxycorticosterone; THF, tetrahydrocortisol; THS, tetrahydrodeoxycortisol.

Figure 1 depicts the steroidogenic enzyme and urinary steroid metabolites investigated. There were no differences between the groups in 24-hour excretion of any of the cortisol metabolites, although there was a trend toward increased  $\alpha$ -cortol in PCOS-d (P = .07, Table 2). Excretion of the cortisone metabolites and androgen metabolites was also similar (Table 2). Ex-



**Figure 2.** Metabolite ratios reflective of  $5\alpha$ -reductase activity and total cortisol and cortisone metabolite excretion. PCOS-d (black bars) had an increased  $5\alpha$ THF/THF ratio compared to control girls (open bars), reflecting global increases in  $5\alpha$ -reductase activity (\*P = .04, A). There was no difference in An/Et, a second ratio indicating  $5\alpha$ -reductase activity (P = .59, B). This discrepancy may be related to the very low concentrations of these androgen metabolites in this young cohort. There was no significant difference between the groups in total glucocorticoid metabolites (P = .30, C) or total androgen metabolites (P = .53, D). Data are mean and SEM.

cretion of the mineralocorticoid metabolites was similar between the groups, with the exception of  $5\alpha$ -tetrahydrodeoxycorticosterone, which was higher in the PCOS-d (P = .05, Table 2). Finally, excretion of total cortisol metabolites (211 ± 212 PCOS-d vs 139  $\pm$  99 control girls,  $\mu$ g/24 hours, P = .16), total cortisone metabolites (276  $\pm$  264 PCOS-d vs  $238 \pm 188$  control girls,  $\mu g/24$ hours, P = .55), total glucocorticoid metabolites (498  $\pm$  466 PCOS-d vs  $377 \pm 281$  control girls,  $\mu$ g/24 hours, P = .30, Figure 2) and total and rogen metabolites (4.5  $\pm$  4.0 PCOS-d vs  $4.1 \pm 4.3$  control girls,  $\mu$ g/24 hours, P = .53, Figure 2) did not differ between the groups.

PCOS-d had significantly higher in vivo  $5\alpha$ -reductase net activity compared to control girls based on

the  $5\alpha$ THF/THF ratio (2.6 ± 1.1 PCOS-d vs 2.0 ± 0.9 control girls, P = .04, Figure 2), the most reliable measure of  $5\alpha$ -reductase net activity in this age group. The An/Et ratio (2.1 ± 1.3 PCOS-d vs 1.8 ± 0.9 control girls, P = .59, Figure 2) and other ratios of  $5\alpha$ - to  $5\beta$ -reduced metabolites (Supplemental Table 1) did not differ between the two groups. The ratios reflective of 11 $\beta$ -

HSD1 activity were similar between the groups:  $(5\alpha \text{THF} + \text{THF})/\text{THE} (P = .17)$  and cortols/ cortolones (P = .33). Weight for length Z score did not correlate with  $5\alpha \text{THF}/\text{THF}$ , An/Et,  $(5\alpha \text{THF} + \text{THF})/\text{THE}$ , or cortols/cortolones.

## Discussion

PCOS-d had evidence for increased global  $5\alpha$ -reductase activity analogous to findings in adult women with PCOS (16, 17, 19–21).  $5\alpha$ -reductase converts testosterone to its more potent  $5\alpha$ -reduced metabolite, DHT (15). Intracellular conversion of testosterone to DHT is responsible for androgen actions in most target tissues (15). In women, increased  $5\alpha$ reductase activity has been associated with idiopathic hirsutism (28),

#### RIGHTSLINKA)

The Endocrine Society. Downloaded from press.endocrine.org by [\${individualUser.displayName}] on 24 June 2016. at 02:11 For personal use only. No other uses without permission. All rights reserved

androgenic alopecia (29), and acne (30). Increased  $5\alpha$ -reductase activity in specific tissues, such as the skin (31) and ovary (21), has been reported in women with PCOS. If genetic variation caused increased  $5\alpha$ -reductase activity in androgen target tissues of affected PCOS-d, it would result in the amplification of peripheral androgen action by increasing the intracellular production of DHT from T (32, 33). These findings are consistent with our hypothesis that increases in target tissue androgen production play a role in the development of PCOS.

Increased  $5\alpha$ -reductase activity would only be physiologically relevant when T was available. Although persistent increases in androgen secretion do not occur until the time of adrenarche (32), there are earlier developmental phases in which androgen secretion may be increased in girls. The fetal adrenal secretes DHEA, an androgen precursor that may undergo extraglandular metabolism to T (34). Although less is known about fetal ovarian androgen production, CYP17A1 is strongly expressed in fetal theca cells starting as early as 33 weeks, suggesting that intrauterine ovarian androgen production is possible (35). A "mini-puberty" of infancy occurs during the first 3-4 months of life during which time the hypothalamic-pituitary-gonadal axis is active, so it is biologically plausible that ovarian androgen production could be increased during this time (22). Following the first year of life, adrenal and ovarian androgen production remains quiescent until the time of adrenarche and subsequent puberty (32, 33). Detailed investigation of gonadal and adrenal steroid physiology during infancy and early childhood has been limited because of ethical and practical challenges of performing invasive studies in this age group (36). Analysis of urinary steroid metabolites provides a valuable means to noninvasively investigate sex steroid production in young children (18).

Androgen exposure in utero (9, 10), neonatally (11), or peripubertally (12) produces phenocopies of PCOS in several animal species (37). During gestation, robust placental aromatase activity likely protects the fetus from exposure to the increased circulating androgen levels in mothers with PCOS (38). Nevertheless, it remains possible that increases in fetal glandular or target tissue androgen production are present in females at risk for PCOS, such as those daughters of affected women who inherited PCOS susceptibility genes. No increase in cord blood androgen levels has been found in PCOS-d in studies using sensitive and specific testosterone assay methods (13, 14). However, increases in androgen production earlier in gestation or enhanced target tissue  $5\alpha$ -reductase activity resulting in local DHT increases could have escaped detection in studies using cord blood (13, 14). Alternatively, there could be increases in androgen production postnatally during other developmental windows, such as the "mini-puberty" of infancy (22).

In contrast to adult affected women, only the  $5\alpha$ THF/ THF ratio was increased in PCOS-d, whereas the An/Et ratio did not differ between the groups. However, the excretion of An and Et was near the limits of detection of the assay, limiting the utility of this ratio as a measure of  $5\alpha$ reductase activity in children of this age, in whom androgen generation is physiologically very low (32, 33). In adult women with PCOS,  $5\alpha$ -reductase was positively correlated with body weight (17, 23). Although body weight was significantly higher in PCOS-d compared to control girls, there was no correlation of body weight with any urinary steroid metabolites suggesting weight differences did not contribute to the differences in  $5\alpha$ -reductase activity we found in PCOS-d.

Increased hepatic  $5\alpha$ -reductase activity may also enhance cortisol inactivation and clearance, resulting in a compensatory increase in ACTH secretion to normalize circulating cortisol levels, at the expense of increased adrenal androgen production (16, 17). The excretion of glucocorticoid metabolites was increased in adult women with PCOS (16, 17, 23, 39), consistent with ACTH-mediated cortisol production. Further, decreased activity of 11 $\beta$ -HSD1, which reduces inactive cortisone to cortisol, has been reported in some (23, 39) but not all (17) studies of adult affected women. This change may also result lower cortisol levels and in compensatory increases in ACTH release (23, 39). In the present study, neither glucocorticoid metabolite excretion nor 11β-HSD1 activity differed in PCOS-d compared to control girls. These findings suggest that increases in zona fasciculata activity are not important in the early origins of PCOS.

Our study has some potential limitations. First, urinary steroid metabolite ratios can estimate global enzyme activity, but do not reflect intracellular steroidogenic enzyme activity in target tissues (28–30). Nevertheless, this non-invasive approach is a valuable tool for investigation of alterations in steroid metabolism in infants and young children. Second, the difference we observed in  $5\alpha$ THF/ THF achieved only modest statistical significance (P = .04, Figure 2). However, this difference was similar in magnitude to differences in  $5\alpha$ -reductase activity observed in lean adult women with PCOS compared to lean control women (17, 23). Further, because genetic factors likely contributed to the alterations in steroidogenesis, these effects would be diluted because the PCOS-d group contained both genetically affected and unaffected girls.

In conclusion, our findings suggest that increased  $5\alpha$ reductase activity is present in girls at risk for PCOS in early childhood. Hence, enzyme expression in target tissues may be primed toward androgen activation in genet-

RIGHTSLINK ()

The Endocrine Society. Downloaded from press.endocrine.org by [\$[individualUser.displayName]] on 24 June 2016. at 02:11 For personal use only. No other uses without permission. All rights reserved.

ically affected PCOS-d before the onset of increased glandular DHEA secretion during adrenarche (32, 40). These findings support our hypothesis that genetic variation leading to hyperandrogenemia, specifically increased target tissue production of DHT, contributes to the development of PCOS through epigenetic actions during critical developmental windows. Prospective cohort studies are needed to determine whether those PCOS-d with early childhood activation of  $5\alpha$ -reductase activity ultimately develop PCOS.

# Acknowledgments

Address all correspondence and requests for reprints to: Andrea Dunaif, MD, Division of Endocrinology, Metabolism, and Molecular Medicine, Feinberg School of Medicine, Northwestern University, 303 East Chicago Avenue, Tarry Building 15-709, Chicago, IL 60611. E-mail: a-dunaif@northwestern.edu.

This research was supported by P50 HD044405 (to A.D.) and K12 HD055884 (to L.T.) from the *Eunice Kennedy Shriver* National Institute of Child Health and Development, the Medical Research Council UK Research Training Fellowship G1001964 (to J.I.), and the Wellcome Trust Project grant 092283 (to W.A.).

Disclosure Summary: The authors report no potential conflicts of interest relevant to this article.

# References

- Diamanti-Kandarakis E, Dunaif A. Insulin resistance and the polycystic ovary syndrome revisited: an update on mechanisms and implications. *Endocr Rev.* 2012;33:981–1030.
- 2. Legro RS, Driscoll D, Strauss JF, 3rd, Fox J, Dunaif A. Evidence for a genetic basis for hyperandrogenemia in polycystic ovary syndrome. *Proc Natl Acad Sci U S A*. 1998;95:14956–14960.
- Sir-Petermann T, Codner E, Perez V, et al. Metabolic and reproductive features before and during puberty in daughters of women with polycystic ovary syndrome. *J Clin Endocrinol Metab.* 2009; 94:1923–1930.
- Torchen LC, Fogel NR, Brickman WJ, Paparodis R, Dunaif A. Persistent apparent pancreatic beta-cell defects in premenarchal PCOS relatives. J Clin Endocrinol Metab. 2014;99:3855–3862.
- Hayes MG, Urbanek M, Ehrmann DA, et al. Genome-wide association of polycystic ovary syndrome implicates alterations in gonadotropin secretion in European ancestry populations. *Nat Commun.* 2015;6:7502.
- Shi Y, Zhao H, Shi Y, et al. Genome-wide association study identifies eight new risk loci for polycystic ovary syndrome. *Nat Genet*. 2012; 44:1020–1025.
- 7. Vink JM, Sadrzadeh S, Lambalk CB, Boomsma DI. Heritability of polycystic ovary syndrome in a Dutch twin-family study. *J Clin Endocrinol Metab*. 2006;91:2100–2104.
- Maher B. Personal genomes: the case of the missing heritability. Nature. 2008;456:18–21.
- Dumesic DA, Abbott DH, Eisner JR, Goy RW. Prenatal exposure of female rhesus monkeys to testosterone propionate increases serum luteinizing hormone levels in adulthood. *Fertil Steril*. 1997;67:155– 163.
- 10. Eisner JR, Dumesic DA, Kemnitz JW, Abbott DH. Timing of prenatal androgen excess determines differential impairment in insulin

secretion and action in adult female rhesus monkeys. J Clin Endocrinol Metab. 2000;85:1206–1210.

- 11. Pinilla L, Trimino E, Garnelo P, et al. Changes in pituitary secretion during the early postnatal period and anovulatory syndrome induced by neonatal oestrogen or androgen in rats. *J Reprod Fertil*. 1993;97:13–20.
- McGee WK, Bishop CV, Bahar A, et al. Elevated androgens during puberty in female rhesus monkeys lead to increased neuronal drive to the reproductive axis: a possible component of polycystic ovary syndrome. *Hum Reprod.* 2012;27:531–540.
- Anderson H, Fogel N, Grebe SK, Singh RJ, Taylor RL, Dunaif A. Infants of women with polycystic ovary syndrome have lower cord blood androstenedione and estradiol levels. J Clin Endocrinol Metab. 2010;95:2180–2186.
- Maliqueo M, Lara HE, Sanchez F, Echiburu B, Crisosto N, Sir-Petermann T. Placental steroidogenesis in pregnant women with polycystic ovary syndrome. *Eur J Obstet Gynecol Reprod Biol.* 2013;166:151–155.
- Wilson JD, Griffin JE, Russell DW. Steroid 5 alpha-reductase 2 deficiency. *Endocr Rev.* 1993;14:577–593.
- Stewart PM, Shackleton CH, Beastall GH, Edwards CR. 5 alphareductase activity in polycystic ovary syndrome. *Lancet*. 1990;335: 431–433.
- Vassiliadi DA, Barber TM, Hughes BA, et al. Increased 5 alphareductase activity and adrenocortical drive in women with polycystic ovary syndrome. *J Clin Endocrinol Metab.* 2009;94:3558–3566.
- Shackleton C, Marcos J, Malunowicz EM, et al. Biochemical diagnosis of Antley-Bixler syndrome by steroid analysis. Am J Med Genet A. 2004;128A:223–231.
- 19. Chin D, Shackleton C, Prasad VK, et al. Increased 5alpha-reductase and normal 11beta-hydroxysteroid dehydrogenase metabolism of C19 and C21 steroids in a young population with polycystic ovarian syndrome. *J Pediatr Endocrinol Metab.* 2000;13:253–259.
- Fassnacht M, Schlenz N, Schneider SB, Wudy SA, Allolio B, Arlt W. Beyond adrenal and ovarian androgen generation: Increased peripheral 5 alpha-reductase activity in women with polycystic ovary syndrome. J Clin Endocrinol Metab. 2003;88:2760–2766.
- Jakimiuk AJ, Weitsman SR, Magoffin DA. 5alpha-reductase activity in women with polycystic ovary syndrome. J Clin Endocrinol Metab. 1999;84:2414–2418.
- Winter JS, Hughes IA, Reyes FI, Faiman C. Pituitary-gonadal relations in infancy: 2. Patterns of serum gonadal steroid concentrations in man from birth to two years of age. J Clin Endocrinol Metab. 1976;42:679–686.
- Tsilchorozidou T, Honour JW, Conway GS. Altered cortisol metabolism in polycystic ovary syndrome: insulin enhances 5alphareduction but not the elevated adrenal steroid production rates. *J Clin Endocrinol Metab.* 2003;88:5907–5913.
- Kuczmarski RJ, Ogden CL, Guo SS, et al. CDC growth charts for the United States: methods and development. *Vital Health Stat 11*. 2000;2002:1–190.
- 25. Rogers SL, Hughes BA, Jones CA, et al. Diminished 11beta-hydroxysteroid dehydrogenase type 2 activity is associated with decreased weight and weight gain across the first year of life. *J Clin Endocrinol Metab.* 2014;99:E821–E831.
- 26. Tomlinson JW, Finney J, Hughes BA, Hughes SV, Stewart PM. Reduced glucocorticoid production rate, decreased 5alpha-reductase activity, and adipose tissue insulin sensitization after weight loss. *Diabetes*. 2008;57:1536–1543.
- 27. Walker EA, Ahmed A, Lavery GG, et al. 11beta-Hydroxysteroid dehydrogenase type 1 regulation by intracellular glucose 6-phosphate provides evidence for a novel link between glucose metabolism and hypothalamo-pituitary-adrenal axis function. *J Biol Chem.* 2007;282:27030–27036.
- Gompel A, Wright F, Kuttenn F, Mauvais-Jarvis P. Contribution of plasma androstenedione to 5 alpha-androstanediol glucuronide in women with idiopathic hirsutism. J Clin Endocrinol Metab. 1986; 62:441–444.

- 29. Hoffmann R, Happle R. Current understanding of androgenetic alopecia. Part I: etiopathogenesis. *Eur J Dermatol*. 2000;10:319–327.
- Thiboutot D, Bayne E, Thorne J, Gilliland K, Flanagan J, Shao Q, Light J, Helm K. Immunolocalization of 5alpha-reductase isozymes in acne lesions and normal skin. *Arch Dermatol.* 2000;136:1125– 1129.
- 31. Skalba P, Dabkowska-Huc A, Kazimierczak W, Samojedny A, Samojedny MP, Chelmicki Z. Content of 5-alpha-reductase (type 1 and type 2) mRNA in dermal papillae from the lower abdominal region in women with hirsutism. *Clin Exp Dermatol*. 2006;31:564–570.
- 32. Auchus RJ. The physiology and biochemistry of adrenarche. *Endocr Dev*. 2011;20:20–27.
- Winter JS, Faiman C. Pituitary-gonadal relations in female children and adolescents. *Pediatr Res.* 1973;7:948–953.
- 34. Ishimoto H, Jaffe RB. Development and function of the human fetal adrenal cortex: a key component in the feto-placental unit. *Endocr Rev.* 2011;32:317–355.

- Cole B, Hensinger K, Maciel GA, Chang RJ, Erickson GF. Human fetal ovary development involves the spatiotemporal expression of p450c17 protein. J Clin Endocrinol Metab. 2006;91:3654–3661.
- 36. Levine RJ GM, Cuttler L, Becker DJ, Nieman L, Rosenfield RL. Overcoming burdens in the regulation of clinical research in children. Proceedings of a consensus conference, in historical context. *Int J Pediatr Endocrinol.* 2011;201:19.
- Dumesic DA, Abbott DH, Padmanabhan V. Polycystic ovary syndrome and its developmental origins. *Rev Endocr Metab Disord*. 2007;8:127–141.
- McClamrock HD, Adashi EY. Gestational hyperandrogenism. Fertil Steril. 1992;57:257–274.
- Rodin A, Thakkar H, Taylor N, Clayton R. Hyperandrogenism in polycystic ovary syndrome. Evidence of dysregulation of 11 betahydroxysteroid dehydrogenase. N Engl J Med. 1994;330:460–465.
- Maliqueo M, Sir-Petermann T, Perez V, et al. Adrenal function during childhood and puberty in daughters of women with polycystic ovary syndrome. J Clin Endocrinol Metab. 2009;94:3282–3288.

RIGHTSLINK4)

The Endocrine Society. Downloaded from press.endocrine.org by [\${individualUser.displayName}] on 24 June 2016. at 02:11 For personal use only. No other uses without permission. . All rights reserved