

## Somatosensory attention identifies both overt and covert awareness in disorders of consciousness

Gibson, Raechelle M; Chennu, Srivas; Fernandez-Espejo, Davinia; Naci, Lorina; Owen, Adrian M; Cruse, Damian

DOI:  
[10.1002/ana.24726](https://doi.org/10.1002/ana.24726)

License:  
None: All rights reserved

Document Version  
Peer reviewed version

Citation for published version (Harvard):  
Gibson, RM, Chennu, S, Fernandez-Espejo, D, Naci, L, Owen, AM & Cruse, D 2016, 'Somatosensory attention identifies both overt and covert awareness in disorders of consciousness', *Annals of Neurology*, vol. 80, no. 3, pp. 412-423. <https://doi.org/10.1002/ana.24726>

[Link to publication on Research at Birmingham portal](#)

### Publisher Rights Statement:

This is the peer reviewed version of the following article: Gibson, R. M., Chennu, S., Fernández-Espejo, D., Naci, L., Owen, A. M. and Cruse, D. (2016), Somatosensory attention identifies both overt and covert awareness in disorders of consciousness. *Ann Neurol.*, 80: 412–423, which has been published in final form at <http://dx.doi.org/10.1002/ana.24726>. This article may be used for non-commercial purposes in accordance with Wiley Terms and Conditions for Self-Archiving.

Checked 15/7/2016

### General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

### Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact [UBIRA@lists.bham.ac.uk](mailto:UBIRA@lists.bham.ac.uk) providing details and we will remove access to the work immediately and investigate.

**Somatosensory attention identifies both overt and covert awareness in disorders of consciousness**

Raechelle M. Gibson<sup>1,2\*</sup>, Srivas Chennu PhD<sup>3,4</sup>, Davinia Fernández-Espejo PhD<sup>5</sup>, Lorina Naci PhD<sup>1,2</sup>, Adrian M. Owen PhD<sup>1,2</sup>, and Damian Cruse PhD<sup>5</sup>

<sup>1</sup>Department of Psychology, University of Western Ontario, London, CAN

<sup>2</sup>The Brain and Mind Institute, University of Western Ontario, London, CAN

<sup>3</sup>School of Computing, University of Kent, Chatham Maritime, UK

<sup>4</sup>Department of Clinical Neurosciences, University of Cambridge, Cambridge, UK

<sup>5</sup>School of Psychology, University of Birmingham, Birmingham, UK

\*Correspondence to: Raechelle M. Gibson  
The Brain and Mind Institute  
Natural Sciences Centre Room 120K  
Western University  
London, ON, CAN N6A 3K7  
rgibso5@uwo.ca

## Abstract

*Objective:* Some patients diagnosed with disorders of consciousness retain sensory and cognitive abilities beyond those apparent from their overt behaviour. Characterising these covert abilities is crucial for diagnosis, prognosis, and medical ethics. This multimodal study investigates the relationship between electroencephalographic evidence for perceptual/cognitive preservation and both overt and covert markers of awareness.

*Methods:* Fourteen patients with severe brain injuries were evaluated with an electroencephalographic vibrotactile attention task designed to identify a hierarchy of residual somatosensory and cognitive abilities: 1) somatosensory steady-state evoked responses, 2) bottom-up attention orienting (P3a event-related potential), and 3) top-down attention (P3b event-related potential). Each patient was also assessed with a clinical behavioural scale and two functional magnetic resonance imaging assessments of covert command following.

*Results:* Six patients produced only sensory responses, with no evidence of cognitive event-related potentials. A further eight patients demonstrated reliable bottom-up attention orienting responses (P3a). No patient showed evidence of top-down attention (P3b). Only those patients who followed commands, whether overtly with behaviour or covertly with functional neuroimaging, also demonstrated event-related potential evidence of attentional orienting.

*Interpretation:* Somatosensory attentional orienting event-related potentials differentiated patients who could follow commands from those who could not. Crucially, this differentiation was irrespective of whether command following was evident through overt external behaviour, or through covert functional neuroimaging methods. Bedside electroencephalographic methods may corroborate more expensive and challenging methods such as functional neuroimaging, and thereby assist in the accurate diagnosis of awareness.

# 1   **Introduction**

2           Disorders of consciousness (DoC) are states that a person may enter when they emerge from  
3   coma following a severe brain injury. Patients in a vegetative state (VS) do not demonstrate purposeful  
4   behaviour and are considered to lack awareness<sup>1–3</sup>. In contrast, patients in a minimally conscious state  
5   (MCS) are considered to have fluctuating awareness and demonstrate variable, but reproducible,  
6   purposeful behaviour<sup>4</sup>. Furthermore, the MCS can be sub-divided into MCS *Plus* or *Minus* on the basis  
7   of the patient’s ability to follow commands<sup>5</sup>. Patients who demonstrate accurate communication and/or  
8   functional object use are considered *emergent* from a MCS (EMCS)<sup>4</sup>. However, the accurate  
9   identification of a patient’s diagnostic group comprises a considerable clinical challenge<sup>1–3,6–8</sup>.

10          To facilitate more accurate diagnosis of the DoC, researchers have developed brain imaging  
11   paradigms to assess volition and command following in the absence of outward responsiveness<sup>9–14</sup>.  
12   Patients who produce behaviour consistent with a VS, but who exhibit evidence of *covert* awareness  
13   with functional neuroimaging – such as imagining movements to command<sup>6,9,10,13,15–17</sup> – have been  
14   considered to exhibit a non-behavioural MCS<sup>18</sup>. However, in both behavioural and neuroimaging-based  
15   assessments, a patient may produce a false negative due to fatigue or insufficient cognitive resources to  
16   successfully complete the demanding diagnostic task<sup>8,19</sup>.

17          Researchers have developed assessments of brain function to place a patient along a hierarchy of  
18   increasingly complex attentional information processing<sup>20–24</sup>. However, there are inconsistencies in the  
19   prognostic value of the event-related potentials used in these hierarchical approaches; some investigators  
20   have reported positive prognostic value in these attentional markers<sup>25</sup>, while others have not<sup>26</sup>. These  
21   inconsistencies may have occurred because multimodal assessments were not used to identify patients in  
22   a non-behavioural MCS. Therefore, 15% of the patient sample considered to be VS may have possessed  
23   a non-behavioural MCS and consequently misrepresented the diagnostic category<sup>27</sup>. Similarly, most  
24   studies of patients with DoC employ auditory stimulation because many patients lack oculomotor  
25   control; however, this tendency limits the characterisation of a patient’s sensory abilities to the auditory  
26   domain.

27          We report a hierarchical cognitive assessment in a sample of fourteen patients with severe brain  
28   injuries using vibrotactile stimulation. The assessment employed an oddball paradigm to elicit steady-  
29   state evoked responses of sensory processing and event-related potential (ERP) markers of bottom-up  
30   and top-down attention (the P3a and P3b, respectively)<sup>28</sup>. As with previous hierarchical designs, this

approach discretizes a patient's sensory and cognitive abilities. A novel aspect of our method is the assessment of a patient's ability to sense and attend to touch. Importantly, patients were also evaluated using two previously established neuroimaging-based assessments of covert command following – mental imagery<sup>6,9,10,13,15–17</sup> and selective auditory attention<sup>29,30</sup> – and a clinical behavioural assessment<sup>31</sup>. By identifying patients with covert command following abilities, these additional assessments ensured a more accurate representation of each patient's level of awareness. Furthermore, we were in a position to test the divergence and convergence of these methods. It was expected that ERP markers of higher-order attention would be evident in patients who were aware, either expressed overtly in their behaviour, or covertly by wilful modulations of brain activity detected with neuroimaging.

## Materials and methods

### Participants

Fourteen patients [mean age 41 (range: 19 to 58) years] contributed sufficient data for inclusion in this investigation. Seven patients were diagnosed as VS<sup>3</sup>, four patients were diagnosed as MCS, two patients were diagnosed as EMCS<sup>4</sup>, and one patient was diagnosed with Locked-In Syndrome (LIS)<sup>32</sup>. Six patients had sustained traumatic brain injuries from motor vehicle accidents. The remaining eight patients had sustained non-traumatic brain injuries from different aetiologies including cardiac arrest (3 cases) and near-drowning (1 case; see Supplementary Table 1). Each patient's surrogate decision maker provided informed, written consent for the patient's participation in the study. Ethical approval was obtained from the University of Western Ontario's Health Sciences Research Ethics Board (London, Canada).

As a scientific control, a sample of fifteen healthy volunteers also participated in the somatosensory selective attention task. These participants ranged in age from 17 to 23 years (mean age 18 years). All healthy volunteers provided informed written consent and received course credit for their participation. The Psychology Research Ethics Board of the University of Western Ontario (London, Canada) provided ethical approval for the control study. Control studies of the other neuroimaging paradigms have been reported elsewhere<sup>15,30,33</sup>.

### Procedure

For each patient, participation in this study comprised assessments with: (1) electroencephalography (EEG) during their completion of a somatosensory selective attention paradigm;

(2) functional magnetic resonance imaging (fMRI) during their completion of a mental imagery paradigm<sup>6,9,10,13,15–17</sup>; (3) fMRI during their completion of an auditory selective attention paradigm<sup>29,30</sup>; and (4) the Coma Recovery Scale-Revised (CRS-R<sup>31</sup>; see Supplementary Tables 1 and 2). fMRI data from Patient EMCS2 could not be analysed due to excessive motion artefacts. However, this patient was included in this investigation because his ability to follow simple commands and communicate was evident from his overt behaviour. Similarly, the data for Patient VS7 from one fMRI session (selective auditory attention) were discarded due to excessive movement. This patient was included in the current investigation because useable data were obtained from this patient for the other three paradigms.

All patients completed the two fMRI paradigms within a two-day period. Ten patients completed the fMRI assessments within two days of their EEG assessments (see Supplementary Table 2). The other four patients completed the EEG assessments after the fMRI assessment with the following delay: 1.5-months (EMCS1); 7.5-months (MCS3); 1-year (VS3); and 3.5-years (VS7). Only Patient MCS3 demonstrated a clinical status change between assessments with EEG and fMRI (MCS- to MCS+). Given the aetiology, age, and time *post-ictus* of those patients with a year or more between assessments (Supplementary Table 2), it is unlikely (although not impossible) that either of these patients underwent a change in their conscious states between assessments<sup>1–3</sup>. Indeed, Patients VS3 and VS7 demonstrated overt behaviour consistent with a VS at all assessments.

### **Somatosensory selective attention paradigm**

Participants completed a short somatosensory selective attention task as their EEGs were recorded. One stimulator was affixed to each wrist and the upper back (three total). Each stimulator administered non-painful vibrotactile stimuli via a motor housed in a rubberized casing<sup>34</sup>. A similar paradigm has also been evaluated for patients with LIS<sup>35</sup>. The experiment comprised 14 blocks. Participants were presented with a series of vibrations alternating among their wrists (10% per wrist) and upper back (80%). A vibration occurred every 200ms and lasted for 50ms. The number of vibrations presented to each wrist in a block was selected on a random uniform interval from 28 to 32. There was always a minimum of three (maximum=21) upper back stimuli between wrist vibrations; on average, 49% (standard deviation=13%) of the wrist stimuli followed exactly three upper back stimuli. Participants were instructed to count the vibrations presented only to the target wrist. The experimenter touched the patient's target wrist after the instruction. The right wrist was always the target wrist for the first block and subsequently alternated between the left and right wrists. The healthy volunteers reported

their count at the end of each block; these participants reported the correct number of vibrations for 12/14 blocks on average (all reports were within  $\pm 3$  of the true number of targets). One block of trials lasted for approximately one minute.

### **Mental imagery paradigm**

During an fMRI scan, patients were asked to engage in two mental imagery paradigms<sup>6,9,10,13,15–17</sup>. In the motor imagery task, patients were instructed to imagine swinging their right arm to hit a tennis ball. In the spatial navigation task, patients were instructed to imagine walking from room to room in their house and visualise all objects they would encounter. Instructions were delivered with noise cancellation headphones (Silent ScanTM, Avotec Inc. for patients scanned in the Trio system, as well as Patient VS6 [first visit], and Sensimetrics S14 for the patients scanned in the Prisma system, including Patient VS6 [second visit]). Patients VS1, VS2, VS4, VS5, VS6 (second visit), MCS4, and EMCS1 completed two sessions of each task, while patients VS3, VS6 (first visit), VS7, MCS1, MCS2, MCS3, and LIS1 completed only one session due to scanner availability or patient fatigue.

### **Auditory selective attention paradigm**

The fMRI selective auditory attention paradigm has been previously described in healthy individuals<sup>30</sup> and patients with DoC<sup>29</sup>, and is designed to identify an ability to follow commands to selectively attend to stimuli – *i.e.*, top-down attention. On each trial, participants were instructed to either count a target word ('yes' or 'no') presented among pseudorandom distractors (spoken digits one to nine), or to relax. Each trial had an on/off design: sound (~22.5s) followed by silence (10s). The scan lasted five minutes, including instructions.

### **Replication data**

Each task alternated five 30-second blocks of mental imagery and five 30-second blocks of rest for a total of five minutes. Patients VS4, MCS3, and EMCS1 participated in second assessments with the somatosensory selective attention task and the CRS-R. These assessments occurred from 2- to 3.5-months following their initial participation. Patient VS6 completed a second assessment with all paradigms (CRS-R, fMRI, and EEG) 22-months after her initial assessment. All four patients maintained their clinical status at follow-up (Supplementary Table 2).

### **EEG data acquisition and pre-processing**

EEG data were recorded at sites FC1, Fz, FC2, C3, Cz, C4, CP1, CP2, Pz, Oz, PO7, and PO8 using an electrode cap with the g.Gamma active electrode system (g.tec Medical Engineering GmbH, Austria). This montage was selected following a previous study conducted in patients with LIS<sup>35</sup> and previous work concerning optimal P300 classification<sup>36</sup>. Data were sampled at 256 Hz and filtered between 0.5 and 30 Hz using a digital Butterworth filter. Stimuli were presented with the g.VIBROstim box (g.tec Medical Engineering GmbH, Austria) using a custom MATLAB® script for Simulink® (MathWorks, Inc., Natick, MA). The recordings were referenced to the right earlobe with a forehead (Fpz) ground. Impedances were kept below 5 kΩ. Data processing was conducted with EEGLAB<sup>37</sup>. The data were segmented into 1-second epochs with a 200ms pre-stimulus period, and linear detrending and baseline correction were applied to each epoch. For artefact correction, all trials containing data with voltages exceeding  $\pm 100 \mu\text{V}$  were rejected. In a second step, the kurtosis of the signal across all channels was calculated for each stimulus type separately, and all trials exceeding 2.5 standard deviations of the mean were rejected. Final trial numbers are reported in (Table 1).

#### **fMRI data acquisition and pre-processing**

The MRI data were acquired in a 3-Tesla Siemens scanner (Siemens, Erlangen, Germany) with a Siemens 32-channel head-coil at the Centre for Functional and Metabolic Mapping at Robarts Research Institute, Western University, Canada. The patients were recruited over 30-months, in which time the 3-T scanner was upgraded. Three patients (VS3, VS7, and MCS3) were scanned in a Magnetom Trio system. All other patients were scanned in a Magnetom Prisma system. Functional echo-planar images of 36 slices covering the whole brain were acquired (repetition time=2000ms, echo time=30ms, matrix size=420 x 420, slice thickness=3 mm, in-plane resolution=3x3 mm, flip angle=78°; for patients VS6 and LIS1 only, matrix size=384x384 and flip angle=75°). High-resolution T1-weighted 3D images were acquired in the same session (Trio system: repetition time=2300ms, echo time=2.98ms, inversion time=900ms, matrix size=256x240, voxel size  $1 \times 1 \times 1$  mm, flip angle=9°; Prisma system: repetition time=2300ms, echo time=2.32ms, inversion time=900ms, matrix size=256x256, flip angle=8°; for patients VS6 and LIS1 only, matrix size=240 x 256 and flip angle=9°). Data from the mental imagery paradigm were pre-processed using SPM8 (<http://www.fil.ion.ucl.ac.uk/spm>), as described elsewhere<sup>13</sup>. For the selective attention paradigm, pre-processing was performed with the AA software<sup>38</sup>.

#### **Statistical analyses**



## EEG responses

The EEG data were assessed for the presence of a steady-state evoked potential to the repetitive vibrotactile stimulation. As one vibration occurred every 200ms, an evoked response was considered present when the averaged peak of the frequency spectrum of the data at the stimulation rate (5 Hz) and its first harmonic (10 Hz) was significantly higher than the background noise<sup>39</sup>. A frequency spectrum was calculated with a discrete Fourier transform over the entire 1-second epoch from the average of all trials using data only from site Pz<sup>40,41</sup>. An  $F$  ratio ( $\alpha=.05$ ;  $F_{2,20} \geq 3.49$ ) was computed to compare the power at 5 and 10 Hz with the average power in the ten adjacent  $\sim 1$  Hz frequency bins (2-4 Hz, 6-9 Hz, and 11-13 Hz)<sup>39</sup>.

Two analyses of the EEG data were conducted to identify the attention-based event-related potentials. For the bottom-up attention effect (P3a), responses to wrist (deviant) and upper back (standard) stimuli were compared. A random subset of the standard stimuli (equal in number to the deviant stimuli) was selected because there were many more standard than deviant stimuli. For the top-down attention effect (P3b), responses to the target and non-target wrist stimuli were compared. Trial numbers were matched between the target and non-target trials. Data from 50 to 750ms post-stimulus were analysed using the cluster-mass procedure<sup>42</sup> of the MATLAB® toolbox FieldTrip<sup>43</sup>. This technique has been described in detail previously<sup>42,44</sup>. In the first step, data were compared at each time-point using a  $t$ -test. In the second step,  $t$ -values of adjacent spatiotemporal points with  $p < .05$  were clustered together by summing their  $t$ -values. The largest cluster was retained. This entire procedure was repeated 1000 times with recombination and randomized resampling of the ERP data. This Monte Carlo method generated a nonparametric estimate of the  $p$ -value representing the statistical significance of the originally identified cluster.

## Blood oxygen level-dependent (BOLD) mental imagery responses

Single subject fixed-effect analyses were performed for each patient. The analysis was based on the general linear model using the canonical hemodynamic response function<sup>45</sup> implemented with SPM8 (<http://www.fil.ion.ucl.ac.uk/spm>). The analysis pipeline was previously reported<sup>13</sup>. Linear contrasts were used to obtain subject-specific estimates, and results were thresholded at a voxel level, familywise error (FWE), whole-brain  $p < .05$ . When no significant activations were found at this level, the statistical threshold was reduced to an uncorrected  $p < .001$  because of the strong anatomical *a priori*

hypotheses<sup>6,9,10,13,15–17</sup>. This less conservative threshold excluded the possibility of failing to detect more subtle changes in the signal<sup>45,46</sup>.

### **BOLD auditory selective attention responses**

The general linear model (SPM8) was used to explore effects of interest. Two event types were defined corresponding to the on/off periods (count/relax; ~22.5s, or vice-versa). The silent period (10s) served as an implicit baseline for all trials. Events for these regressors were modelled by convolving boxcar functions with the canonical hemodynamic response function. Also included in the general linear model were the following nuisance variables: the movement parameters in the three directions of motion and three degrees of rotation, and the mean of each scan. Linear contrasts were used to obtain subject-specific estimates for the effect of interest. Clusters that survived the  $p < .05$  threshold after the FWE correction were reported as significant.

## **Results**

All patient outcomes are summarized in ( **Figure 1** ) and (Supplementary Table 3).

### **EEG responses**

A steady-state evoked potential was detected in the EEG data of all patients ( $n=14$ ) and all healthy volunteers ( $n=15$ ; **Figure 2**).

**Bottom-up attention effects** (deviant versus standard stimuli) were detected from eight patients and all of the healthy volunteers ( $n=15$ ; **Figure 3**). All patients who demonstrated a differential response to the deviant versus standard stimuli also demonstrated evidence of command following in either a behavioural or a neuroimaging-based assessment (**Figure 1** and Supplementary Table 3).

Top-down ERP attention effects (target versus non-target wrist vibrations) were not detected from any of the patients. However, this ERP effect was evident for healthy volunteers at the group level ( $n=15$ ) and at the single-subject level, albeit with a hit-rate of 67% (**Figure 4**). Hit-rates of at least 80% (12/15) and 100% (15/15) have been reported for fMRI-detected mental imagery and selective attention respectively<sup>30</sup>. Given the relatively lower sensitivity of the top-down attention ERP analysis (*i.e.*, 67%), additional *post-hoc* comparisons were conducted. While the number of trials available after artefact rejection did not differ across groups (Table 1;  $\chi^2(2)=0.21$ ,  $p=0.9$ ), some patients had many fewer trials available than healthy individuals. The single-subject ERP analyses for the healthy volunteers were thus

repeated in the *post-hoc* analyses using only a pseudorandom subset of trials equal in number to the minimum number of trials available in the single-subject analyses of the patient data (180 trials, in the case of Patient MCS2).

Bottom-up attentional ERP effects were detected at the single-subject level for all healthy volunteers when as few as 180 trials were included for each stimulus type. However, top-down attentional ERP effects were detected from only seven healthy volunteers. Subsequent analyses revealed that a minimum of 300 trials were required to detect the top-down attentional ERP effects from the same 10 healthy volunteers as in the *a priori* analyses. Four patients did not have enough trials available to meet this criterion. Overall, these analyses indicate that the top-down attentional ERP effect may not have been detected in some single-subject analyses due to low trial numbers. Nevertheless, the bottom-up attentional ERP effect was robust to data loss.

## **BOLD mental imagery responses**

In her first visit, Patient VS6 produced reliable, appropriate activation during the motor imagery task in the supplementary motor area and cerebellum bilaterally at an uncorrected  $p < .001$  (cluster level FWE-corrected  $p < .05$ ). In her second visit, Patient VS6 produced reliable, isolated clusters of activation during the motor imagery and spatial navigation tasks in the left precentral gyrus at an uncorrected  $p < .001$  (cluster level FWE-corrected  $p < .05$ ). The patient was thus reclassified as in a non-behavioural MCS<sup>18</sup>.

Patients VS7 showed high levels of motion requiring 37% and 37.5% of his data to be discarded (for motor imagery and spatial navigation respectively). The analysis of the remaining data revealed appropriate activation during the spatial navigation task only (*i.e.*, the left occipito-parietal junction at uncorrected  $p < .001$ ). The patient was thus reclassified as in a non-behavioural MCS<sup>18</sup>.

Patients MCS3, MCS4, EMCS1, and LIS1 showed reliable activation during the spatial navigation task only. This involved: bilateral occipito-parietal junction (uncorrected  $p < .001$ ) for MCS3; right temporo-occipito-parietal junction (FWE-corrected  $p < .05$ ), as well as right dorsal premotor cortex, right insular cortex, and right putamen (uncorrected  $p < .001$ ) for MCS4; right occipito-parietal junction, a region in the boundaries between right lingual gyrus/parahippocampal cortex, left precentral gyrus (comprising the supplementary and pre-supplementary motor areas), as well as some less typical areas such as the inferior frontal gyrus, the left superior temporal gyrus, and the left striatum (FWE-corrected

$p < .05$ ) for EMCS1; and supplementary motor area, right precentral gyrus, occipito-parietal junction, posterior temporo-occipital region, and the cerebellum (uncorrected  $p < .001$ ) for LIS1.

The remaining seven patients (VS1-5, MCS1, and MCS2) showed no activation at the conservative FWE-corrected statistical threshold, or at uncorrected  $p < .001$ .

## **BOLD auditory selective attention responses**

Of the patients diagnosed as in a VS, only Patient VS6 showed significantly more activation following the instruction to count than to relax. This patient showed significant activation in the temporal and parietal cortex bilaterally (FWE-corrected at  $p < .05$ ).

Patients MCS1-4 and LIS1 also showed significantly more activation following the instruction to count than to relax. Patient MCS1 showed significant activation in the frontotemporal and parietal cortex bilaterally. Patient MCS2 showed significant activation in the temporal cortex bilaterally (FWE-corrected at  $p < .05$ ). Patient MCS3 showed significant activation in the parietal cortex bilaterally. Patient MCS4 showed significant activation in the frontotemporal and parietal cortex bilaterally (FWE-corrected at  $p < .05$ ). Patient LIS1 produced significant brain activity in the frontotemporal cortex bilaterally (FWE-corrected at  $p < .05$ ).

Of note, Patient EMCS1 did not show significant differences in activation in the command following task even though she was able to follow commands with her overt behaviour immediately prior to her assessment. Patients VS7 and EMCS2 were excluded from this analysis because both patients moved excessively during their functional scans.

## **Correspondence between command following and EEG responses**

The main hypothesis in this investigation was that patients who were aware would exhibit concordant EEG markers of higher-order attention processing. While top-down processing (P3b) was not detected from any patients, an interesting observation from the current data is the relationship between a specific marker of awareness – command-following – and the bottom-up attention orienting ERP effect, the P3a. A patient was considered to have evidence of such awareness if they demonstrated evidence of command following in any one of the three non-EEG assessments (selective auditory attention, mental imagery, or a behavioural assessment with the CRS-R). This approach is consistent with clinical behavioural guidelines in which a diagnosis of awareness (MCS) is given if a patient follows commands on one occasion across multiple assessments. A Fisher's exact test revealed a

significant positive association between evidence for command following and evidence for the P3a ( $p=.007$ ; note  $p=.0047$  if the two observations of Patient VS6 are not included to maintain the assumption of independence). This relationship is summarised in (Figure 1).

## Replication data

The replication results are depicted in (Figure 5). All patients exhibited consistent effects across assessments with the exception of Patient VS6 for whom a P3a was significant only during her initial assessment.

## Discussion

We investigated a novel EEG method for the assessment of residual sensory and cognitive processing alongside two fMRI-based assessments of covert command following and one behavioural assessment of overt command following in a sample of fourteen patients with severe brain injuries. The primary novel finding of this work is the relationship between an ERP marker of bottom-up attention orienting (the P3a) and command following such that all patients with a P3a response demonstrated positive evidence of command following. Similarly, most patients who did not generate a P3a response also did not demonstrate evidence of command following (see Figure 1 and Supplementary Table 3).

Some investigators have reported positive prognostic value in the presence of a P300 following traumatic brain injury<sup>25</sup>. There have also been reports of correlations between cognitive ERPs and behavioural markers of awareness<sup>14,24</sup>, as well as the prediction of recovery from the DoC using cognitive ERPs<sup>47,26</sup>. Crucially, the current study included two neuroimaging-based assessments of covert command following. This step is important given that a recent meta-analysis estimates a 15% rate of covert awareness among patients diagnosed as in a VS<sup>27</sup>. Previous studies of the P300 in patients with DOC are likely to have included patients capable of covert command following, thus obscuring the relationship reported here. While the feasibility of routine neuroimaging assessments in clinical practice is limited by important health, safety, and financial factors, the findings of this work suggest that these assessments are necessary to elucidate the relationship between a patient's conscious state and their residual sensory and cognitive abilities.

It is curious that an ERP marker of unconscious (or preconscious) processing – *i.e.*, the P3a – is closely linked to awareness in this work. Indeed, the P3a can be elicited by unattended stimuli and during REM sleep and deep sedation<sup>28,48</sup>. We speculate that the correspondence between the P3a and

1 command following stems from the overlap of the neural networks that support attention, and those that  
2 are relatively more preserved in conscious patients<sup>49,50</sup>. Indeed, frontal lobe lesions have been associated  
3 with diminished P3a responses to auditory<sup>51</sup> and somatosensory<sup>52</sup> stimulation. Equally, this association  
4 suggests that a P3a response may be less informative for patients with specific frontal lobe injuries.  
5 Nevertheless, a P3a can be elicited without the explicit collaboration of the individual – *i.e.*, without  
6 following task instructions<sup>48</sup>. This feature is appealing, as it suggests that a passive assessment of  
7 attention orienting, which entails lower cognitive demands than active assessments of voluntary top-  
8 down attention, may be sufficient to identify patients with covert awareness.

9         The P3b marker of top-down attention in the current EEG task was not detected from any of the  
10 patients in this sample, as has been reported previously<sup>53</sup>. In fact, P3b responses in the current work  
11 were detected from only 67% (10/15) of the healthy volunteers. *Post-hoc* analyses of the ERP data  
12 indicated that this low sensitivity may be exacerbated by the fewer usable trials in the patient data, as  
13 this comparison was sensitive to a reduced signal-to-noise ratio. Additionally, time-variant levels of  
14 arousal and fatigue characteristic of the DoC may have led to inconsistent engagement in the counting  
15 task needed to generate the top-down ERP effect<sup>8,19</sup>. In contrast to the fMRI-based selective attention  
16 task, the selective attention manipulation in the EEG task may have placed higher cognitive demands on  
17 participants due to the longer duration of the EEG task. Participants were required to sustain attention  
18 for five minutes in ~22.5-second blocks for both fMRI tasks, whereas the EEG task involved fifteen  
19 minutes of attention in ~1-minute blocks. The EEG task was longer to ensure that a high EEG signal-to-  
20 noise ratio was achieved, and *post-hoc* analyses confirmed that the top-down ERP effect was sensitive to  
21 trial numbers. Unfortunately, increased task duration requires participants to sustain attention for an  
22 even longer period, making it unlikely that this manipulation would increase the sensitivity of the task.  
23 Some investigators use machine learning to circumvent these issues and address possible spatiotemporal  
24 variations in the electrocortical responses of patients with brain injuries<sup>54</sup>. For simplicity of  
25 interpretation and consistency with clinical methods, we employed a more traditional approach to  
26 comparing scalp voltages. While no false alarms were evident in the current sample, misses occurred  
27 with two patients – *i.e.*, patients demonstrated evidence of command following but no evidence of a P3a.  
28 As has been discussed elsewhere, signs of awareness in both behavioural and neuroimaging assessments  
29 may be missed due to fluctuating arousal<sup>13</sup>. Nevertheless, when a P3a is elicited, the current data suggest  
30 the sophisticated cognitive networks that underlie an ability to follow commands are also preserved.

1           The detection of awareness in the DoC is a clinical standard of care. In order to provide  
2 sufficient evidence to influence clinical practice, it is essential to compare novel assessments to existing  
3 techniques. The current investigation allowed for a comparison of two previously reported  
4 neuroimaging-based assessments of covert command following, based on mental imagery<sup>6,9,10,13,15-17</sup> and  
5 selective auditory attention<sup>29,30</sup>. The results of these assessments converged for nine of the twelve  
6 patients with useable data from both paradigms. Two patients demonstrated positive evidence of  
7 command following in only the selective auditory attention task, while one patient showed positive  
8 evidence of command following only in the mental imagery task. The behavioural profile of the DoC –  
9 that is, time-variant fatigue and arousal – always affords the possibility that a patient did not  
10 demonstrate positive evidence of covert command following due to lack of voluntary engagement in the  
11 task. Likewise, false negatives occur in assessments of healthy volunteers<sup>11,55</sup>. Nevertheless, the less  
12 than perfect correspondence of the two covert fMRI command following tasks may have occurred  
13 because the demands of one task were better suited to the patient. For example, some individuals find it  
14 difficult to engage in motor imagery<sup>56</sup>, and in some reports, brain-computer interfaces based on selective  
15 attention tasks are successfully operated by more users than those based on responses to motor  
16 imagery<sup>57,58</sup>. Accordingly, assessments of covert command following based on selective attention may  
17 be better suited to a general population. Overall, however, an optimal evaluation of a patient with a DoC  
18 should include multiple assessments to maximise the likelihood of detecting responses that are not  
19 evident from overt behaviour<sup>13</sup>. In the absence of unambiguous ground truth, an investigation of the  
20 concordance between assessments may be the best way to improve diagnostic and prognostic accuracy.

21           In summary, the brain responses of fourteen patients with severe brain injuries were assessed  
22 using an EEG-based somatosensory selective attention task, two fMRI-based assessments of covert  
23 command following, and one behavioural instrument. While limited by a relatively small sample of  
24 patients, the data tentatively suggest that the detection of a somatosensory bottom-up P3a effect in a  
25 patient correlates with an ability to follow commands, as evaluated by multimodal assessments. This  
26 provides evidence that a bedside somatosensory oddball procedure can improve diagnostic accuracy in  
27 the DoC and more accurately characterise the level of neurocognitive preservation. Overall, this work  
28 provides a valuable addition to neuroimaging batteries for the clinical assessment of patients with DoC  
29 and convergent, multimodal evidence for the utility of these techniques.

## Acknowledgements

We are grateful to Dr. Laura E. Gonzalez-Lara (Western University, London, Canada) for her assistance with our patient work. This work was supported by grants to A.M.O. from the DECODER project, the European Commission in the 7<sup>th</sup> Framework Programme (2007-2013), the James S. McDonnell Foundation, and the Canada Excellence Research Chairs Program through the Canadian Institutes of Health Research. D.C. and D.F.E. were each supported by fellowships from the Canadian Institutes of Health Research; S.C. was supported by the James S. McDonnell Foundation, the NIHR Brain Injury Healthcare Technology Co-operative, and the Evelyn Trust; and R.M.G. was supported by a Vanier Canada Graduate Scholarship from the Natural Sciences and Engineering Research Council of Canada. The views expressed are those of the authors and not necessarily those of the UK National Health Service, the NIHR or the UK Department of Health. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

## Author Contributions

R.M.G., D.F.E., L.N., and D.C. contributed to conception and design of the study. R.M.G., D.F.E., L.N., and D.C. contributed to data collection and analysis. R.M.G., D.C., S.C., and A.M.O contributed to writing the manuscript.

## Potential Conflicts of Interest

Nothing to report.



## References

1. Multi-Society Task Force on PVS. Medical aspects of the persistent vegetative state (first part). *N. Engl. J. Med.* 1994;330(21):1499 [PubMed](#) –1508.
2. Multi-Society Task Force on PVS. Medical aspects of the persistent vegetative state (second part). *N. Engl. J. Med.* 1994;330(22):1572 [PubMed](#) –1579.
3. Royal College of Physicians Working Group. The vegetative state: Guidance on diagnosis and management. *Clin. Med. (Northfield. Il).* 2003;3(3):249–254.
4. Giacino JT, Ashwal S, Childs N, et al. The minimally conscious state: Definition and diagnostic criteria. *Neurology* 2002;58(3):349 [PubMed](#) –353.
5. Bruno M-A, Vanhaudenhuyse A, Thibaut A, et al. From unresponsive wakefulness to minimally conscious PLUS and functional locked-in syndromes: Recent advances in our understanding of disorders of consciousness. *J. Neurol.* 2011;258(7):1373 [PubMed](#) –1384.
6. Fernández-Espejo D, Owen AM. Detecting awareness after severe brain injury. *Nat. Rev. Neurosci.* 2013;14:801–809.
7. Gosseries O, Di H, Laureys S, Boly M. Measuring consciousness in severely damaged brains. *Annu. Rev. Neurosci.* 2014;37:457– [PubMed](#) ;478.
8. Giacino JT, Fins JJ, Laureys S, Schiff ND. Disorders of consciousness after acquired brain injury: The state of the science. *Nat. Rev. Neurol.* 2014;10(2):99 [PubMed](#) –114.
9. Owen AM, Coleman MR, Boly M, et al. Detecting awareness in the vegetative state. *Science.* 2006;313(5792):1402.
10. Monti MM, Vanhaudenhuyse A, Coleman MR, et al. Willful modulation of brain activity in disorders of consciousness. *N. Engl. J. Med.* 2010;362(7):579 [PubMed](#) –589.
11. Cruse D, Chennu S, Chatelle C, et al. Bedside detection of awareness in the vegetative state: A cohort study. *Lancet* 2011;378(9809):2088 [PubMed](#) –2094.
12. Goldfine AM, Victor JD, Conte MM, et al. Determination of awareness in patients with severe brain injury using EEG power spectral analysis. *Clin. Neurophysiol.* 2011;122(11):2157–2168.
13. Gibson RM, Fernández-Espejo D, Gonzalez-Lara LE, et al. Multiple tasks and neuroimaging

modalities increase the likelihood of detecting covert awareness in patients with disorders of consciousness. *Front. Hum. Neurosci.* 2014;8:00950.

14. Schnakers C, Perrin F, Schabus M, et al. Voluntary brain processing in disorders of consciousness. *Neurology* 2008;71(20):1614 [PubMed](#) –1620.

15. Boly M, Coleman MR, Davis MH, et al. When thoughts become action: An fMRI paradigm to study volitional brain activity in non-communicative brain injured patients. *Neuroimage* 2007;36(3):979 [PubMed](#) –992.

16. Bardin JC, Fins JJ, Katz DI, et al. Dissociations between behavioural and functional magnetic resonance imaging-based evaluations of cognitive function after brain injury. *Brain* 2011;134(3):769 [PubMed](#) –782.

17. Stender J, Gosseries O, Bruno M-A, et al. Diagnostic precision of PET imaging and functional MRI in disorders of consciousness: a clinical validation study. *Lancet* 2014;384(9942):514 [PubMed](#) –522.

18. Gosseries O, Zasler ND, Laureys S. Recent advances in disorders of consciousness: Focus on the diagnosis. *Brain Inj.* 2014;28(9):1141 [PubMed](#) –1150.

19. Whyte J, Nordenbo AM, Kalmar K, et al. Medical complications during inpatient rehabilitation among patients with traumatic disorders of consciousness. *Arch. Phys. Med. Rehabil.* 2013;94(10):1877–1883.

20. Bekinschtein TA, Dehaene S, Rohaut B, et al. Neural signature of the conscious processing of auditory regularities. *Proc. Natl. Acad. Sci. U. S. A.* 2009;106(5):1672–1677.

21. Fischer C, Luaute J, Morlet D. Event-related potentials (MMN and novelty P3) in permanent vegetative or minimally conscious states. *Clin. Neurophysiol.* 2010;121(7):1032 [PubMed](#) –1042.

22. Chennu S, Finoia P, Kamau E, et al. Dissociable endogenous and exogenous attention in disorders of consciousness. *NeuroImage. Clin.* 2013;3:450– [PubMed](#) ;461.

23. Faugeras F, Rohaut B, Weiss N, et al. Event related potentials elicited by violations of auditory regularities in patients with impaired consciousness. *Neuropsychologia* 2012;50(3):403 [PubMed](#) –418.

24. Kotchoubey B, Lang S, Mezger G, et al. Information processing in severe disorders of

- consciousness: Vegetative state and minimally conscious state. *Clin. Neurophysiol.* 2005;116(10):2441–2453.
25. Lew HL, Dikmen S, Slimp J, et al. Use of somatosensory-evoked potentials and cognitive event-related potentials in predicting outcomes of patients with severe traumatic brain injury. *Am. J. Phys. Med. Rehabil.* 2003;82(1):53 [PubMed](#) –61.
26. Steppacher I, Eickhoff S, Jordanov T, et al. N400 predicts recovery from disorders of consciousness. *Ann. Neurol.* 2013;73(5):594 [PubMed](#) –602.
27. Kondziella D, Friberg CK, Frokjaer VG, et al. Preserved consciousness in vegetative and minimal conscious states: Systematic review and meta-analysis. *J. Neurol. Neurosurg. Psychiatry* 2016;87(5):485 [PubMed](#) -492.
28. Polich J. Updating P300: an integrative theory of P3a and P3b. *Clin. Neurophysiol.* 2007;118(10):2128–2148.
29. Naci L, Owen AM. Making every word count for nonresponsive patients. *JAMA Neurol.* 2013;70(10):1235 [PubMed](#) -1241.
30. Naci L, Cusack R, Jia VZ, Owen AM. The brain’s silent messenger: Using selective attention to decode human thought for brain-based communication. *J. Neurosci.* 2013;33(22):9385–9393.
31. Kalmar K, Giacino JT. The JFK Coma Recovery Scale—Revised. *Neuropsychol. Rehabil.* 2005;15(3/4):454–460.
32. Smith E, Delargy M. Locked-in syndrome. *Br. Med. J.* 2005;330(7488):406-409.
33. Gabriel D, Henriques J, Comte A, et al. Substitute or complement? Defining the relative place of EEG and fMRI in the detection of voluntary brain reactions. *Neuroscience* 2015;290:435–[PubMed](#) ;444.
34. Ortner R, Prückl R, Guger C. A tactile P300-based BCI for communication and detection of awareness. *Biomed. Technol.* 2013;58:1– [PubMed](#) ;2.
35. Lugo ZR, Rodriguez J, Lechner A, et al. A vibrotactile P300-based Brain-Computer Interface for consciousness detection and communication. *Clin. EEG Neurosci.* 2014;45(1):14–21.
36. Krusienski DJ, Sellers EW, Cabestaing F, et al. A comparison of classification techniques for the P300 Speller. *J. Neural Eng.* 2006;3(4):299 [PubMed](#) –305.

- 1 37. Delorme A, Makeig S. EEGLAB: An open source toolbox for analysis of single-trial EEG  
2 dynamics including independent component analysis. *J. Neurosci. Methods*  
3 2004;134(1):9 [PubMed](#) –21.
- 4 38. Cusack R, Vicente-Grabovetsky A, Mitchell DJ, et al. Automatic analysis (AA): Efficient  
5 neuroimaging workflows and parallel processing using Matlab and XML. *Front. Neuroinform.*  
6 2014;8:90.
- 7 39. Dobie RA, Wilson MJ. A comparison of t test, F test, and coherence methods of detecting steady-  
8 state auditory-evoked potentials, distortion-product otoacoustic emissions, or other sinusoids. *J.*  
9 *Acoust. Soc. Am.* 1996;100(4 [PubMed](#) Pt 1):2236–2246.
- 10 40. Mouraux A, Iannetti GD, Colon E, et al. Nociceptive steady-state evoked potentials elicited by  
11 rapid periodic thermal stimulation of cutaneous nociceptors. *J. Neurosci.* 2011;31(16):6079–6087.
- 12 41. Tobimatsu S, Zhang YM, Kato M. Steady-state vibration somatosensory evoked potentials:  
13 Physiological characteristics and tuning function. *Clin. Neurophysiol.* 1999;110(11):1953–1958.
- 14 42. Maris E, Oostenveld R. Nonparametric statistical testing of EEG- and MEG-data. *J. Neurosci.*  
15 *Methods* 2007;164(1):177 [PubMed](#) –190.
- 16 43. Oostenveld R, Fries P, Maris E, Schoffelen J-M. FieldTrip: Open source software for advanced  
17 analysis of MEG, EEG, and invasive electrophysiological data. *Comput. Intell. Neurosci.*  
18 2011;2011:1–9.
- 19 44. Cruse D, Beukema S, Chennu S, et al. The reliability of the N400 in single subjects: Implications  
20 for patients with disorders of consciousness. *NeuroImage. Clin.* 2014;4:788–799.
- 21 45. Friston KJ, Holmes A, Poline JB, et al. Detecting activations in PET and fMRI: Levels of  
22 inference and power. *Neuroimage* 1996;4(3 [PubMed](#) Pt 1):223–235.
- 23 46. Fernández-Espejo D, Junque C, Cruse D, et al. Combination of diffusion tensor and functional  
24 magnetic resonance imaging during recovery from the vegetative state. *BMC Neurol.* 2010;10:77.
- 25 47. Wijnen VJM, van Boxtel GJM, Eilander HJ, de Gelder B. Mismatch negativity predicts recovery  
26 from the vegetative state. *Clin. Neurophysiol.* 2007;118(3):597 [PubMed](#) –605.
- 27 48. Chennu S, Bekinschtein TA. Arousal modulates auditory attention and awareness: Insights from  
28 sleep, sedation, and disorders of consciousness. *Front. Psychol.* 2012;3:65.

- 1 49. Fernández-Espejo D, Soddu A, Cruse D, et al. A role for the default mode network in the bases of  
2 disorders of consciousness. *Ann. Neurol.* 2012;72(3):335 **PubMed** –343.
- 3 50. Thibaut A, Bruno MA, Chatelle C, et al. Metabolic activity in external and internal awareness  
4 networks in severely brain-damaged patients. *J. Rehabil. Med.* 2012;44(6):487–494.
- 5 51. Knight RT. Decreased response to novel stimuli after prefrontal lesions in man.  
6 *Electroencephalogr. Clin. Neurophysiol. Evoked Potentials* 1984;59(1):9 **PubMed** –20.
- 7 52. Yamaguchi S, Knight RT. Anterior and posterior association cortex contributions to the  
8 somatosensory P300. *J. Neurosci.* 1991;11(7):2039 **PubMed** –2054.
- 9 53. Pokorny C, Klobassa DS, Pichler G, et al. The auditory P300-based single-switch brain-computer  
10 interface: Paradigm transition from healthy subjects to minimally conscious patients. *Artif. Intell.*  
11 *Med.* 2013;59(2):81 **PubMed** –90.
- 12 54. King JR, Faugeras F, Gramfort a., et al. Single-trial decoding of auditory novelty responses  
13 facilitates the detection of residual consciousness. *Neuroimage* 2013;83:726– **PubMed** ;738.
- 14 55. Fernández-Espejo D, Norton L, Owen AM. The clinical utility of fMRI for identifying covert  
15 awareness in the Vegetative State: A comparison of sensitivity between 3T and 1.5T. *PLoS One*  
16 2014;9(4): **PubMed** e95082.
- 17 56. Hammer EM, Halder S, Blankertz B, et al. Psychological predictors of SMR-BCI performance.  
18 *Biol. Psychol.* 2012;89(1):80 **PubMed** –86.
- 19 57. Guger C, Daban S, Sellers E, et al. How many people are able to control a P300-based brain-  
20 computer interface (BCI)? *Neurosci. Lett.* 2009;462(1):94 **PubMed** –98.
- 21 58. Guger C, Edlinger G, Harkam W, et al. How many people are able to operate an EEG-based  
22 brain-computer interface (BCI)? *IEEE Trans. Neural Syst. Rehabil. Eng.* 2003;11(2):145–147.
- 23 59. Bruno M-A, Vanhaudenhuyse A, Schnakers C, et al. Visual fixation in the vegetative state: An  
24 observational case series PET study. *BMC Neurol.* 2010;10:35.
- 25
- 26
- 27

## 1 **Figure Captions**

2 **Figure 1 . Summary of the relationship between command following and outcomes on the selective**  
3 **somatosensory attention task.**

4 The summary depicts the number of patients and healthy volunteers who generated each of the three  
5 possible outcomes on the somatosensory selective attention task.

6 VS=vegetative state; MCS=minimally conscious state; EMCS=emergent from a minimally conscious  
7 state; LIS=Locked-In Syndrome.

8 **Figure 2 . Steady-state evoked responses to the repetitive vibrotactile stimulation.**

9 Power spectra (top panels) and averaged EEG responses (bottom panels) calculated over a period  
10 of 1-second. Analyses were conducted using the data recorded from site Pz only; each waveform  
11 (bottom panels) is depicted with  $\pm 1$  standard error of the mean.

12 EEG=electroencephalography; \*\*= $p < 0.01$ ; \*\*\*= $p < .001$ ; VS=vegetative state; MCS=minimally  
13 conscious state; EMCS=emergent from a minimally conscious state; LIS=Locked-In Syndrome.

14 **Figure 3 . Bottom-up attention event-related potentials to the standard and deviant vibrotactile**  
15 **stimulation.**

16 Spatiotemporal clusters were calculated across all twelve electrodes and are depicted with  $\pm 1$   
17 standard error of the mean in colour-matched shading. The electrodes included in the significant  
18 spatiotemporal cluster are enclosed with a black line on each topographic plot. The temporal  
19 boundaries and the probability value of each cluster are indicated with shading and inset text. (A)  
20 depicts the grand-averaged ERP effect for the healthy volunteers, (B) depicts the single-subject  
21 ERP effects for the healthy volunteers ( $p < 9.9E-03$  in all cases), and (C) depicts the single-subject  
22 ERP effects for the patients with statistically significant results.

23 VS=vegetative state; MCS=minimally conscious state; EMCS=emergent from a minimally  
24 conscious state; LIS=Locked-In Syndrome.

25 **Figure 4 . Top-down attention event-related potentials to the target and non-target vibrotactile**  
26 **stimulation for the healthy volunteers.**

Spatiotemporal clusters were calculated across all twelve electrodes with each waveform depicted with  $\pm 1$  standard error of the mean. The electrodes included in the significant spatiotemporal cluster are enclosed with a black outline on each topographic plot. The temporal boundaries and the probability value of each cluster are indicated with shading and inset text. The grand-averaged result ( $n=15$ ) is depicted in (A). For the single subject results (B), only results from participants with statistically significant clusters are shown.

**Figure 5 .** Replication data from the four patients with whom follow-up investigations were conducted.

Data are depicted for the initial and follow up tests of Patients VS4, MCS3, EMCS1, and VS6, as labelled. For the steady-state evoked potentials, power spectra (top left panels within each cell) and averaged EEG data (bottom left panels within each cell) were calculated over a period of 1-second. Analyses were conducted using the data recorded from site Pz only; each waveform is depicted with  $\pm 1$  standard error of the mean. For the bottom-up attention ERP effects (right panels within each cell), spatiotemporal clusters were calculated across all twelve electrodes and are depicted with  $\pm 1$  standard error of the mean. The electrodes included in the significant spatiotemporal cluster are enclosed with a black line on each topographic plot. The temporal boundaries and the probability value of each cluster are indicated with shading and inset text. For Patient VS6 only, two separate fMRI assessments were conducted at each testing session. For the fMRI mental imagery paradigm, significant task-related fMRI activation is depicted (Imagery>Rest), and results are thresholded at an uncorrected  $p<.001$ . For the fMRI selective auditory attention task, only activation clusters within the attention network (Count>Relax) that survived the familywise error correction threshold of  $p<.05$  at the whole-brain level are displayed. The fMRI results are rendered on the patient's T1 anatomical MRI image, and scales depicting the  $t$ -value statistical maps are inset.

\*= $p<0.05$ ; \*\*= $p<0.01$ ; \*\*\*= $p<.001$ ; n.s.=not statistically significant; VS=vegetative state; MCS=minimally conscious state; EMCS=emergent from a minimally conscious state.

**Table 1. Number of trials available for the analyses of the EEG data from the somatosensory selective attention paradigm following artefact rejection.**

	Stimulus Type <sup>a</sup>			
	<i>M (MIN-MAX)</i>			
	Upper Back	Target Wrist	Non-Target Wrist	Trials Rejected (%)
Patients ( <i>n</i> =14)	2614 (1591-3246)	313 (188-384)	311 (180-388)	35 (20-59)
Controls ( <i>n</i> =15)	2890 (2718-5026)	345 (327-363)	345 (321-359)	25 (20-32)
<p><i>Notes.</i> M=mean; MIN=minimum; MAX=maximum.</p> <p><sup>a</sup>A 2x3 Chi-square goodness of fit test indicated that the minimum number of trials in each of the three stimulus types did not significantly differ between the controls and patients, <math>\chi^2(2)=0.21</math>, <math>p=0.9</math>.</p>				