

Sunitinib treatment enhances metastasis of innately drug resistant breast tumors

Wragg, Joseph; Heath, Victoria; Bicknell, Roy

DOI:

[10.1158/0008-5472.CAN-16-1982](https://doi.org/10.1158/0008-5472.CAN-16-1982)

License:

None: All rights reserved

Document Version

Peer reviewed version

Citation for published version (Harvard):

Wragg, J, Heath, V & Bicknell, R 2017, 'Sunitinib treatment enhances metastasis of innately drug resistant breast tumors', *Cancer Research*, vol. 77, no. 4, pp. 1008-1020. <https://doi.org/10.1158/0008-5472.CAN-16-1982>

[Link to publication on Research at Birmingham portal](#)

Publisher Rights Statement:

<http://cancerres.aacrjournals.org/content/77/4/1008>

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

Supplementary materials and methods.

Sunitinib resistance inheritance experiment

4T1 tumours displaying acquired (responsive cohort) or innate (non-responsive cohort) resistance to sunitinib treatment *in vivo* were removed at 1300 mm³ in size, minced, and tumour derived cells cultured in high glucose Dulbecco's modified eagle medium (DMEM) (Sigma, Gillingham, UK), supplemented with 10% FCS (PAA, Cambridge, UK) and 2mM L-glutamine (Gibco, Paisley, UK), for 2 weeks. 2.5×10^5 4T1-Luc cells, from these cultures, suspended in Optimem (Gibco) and in a volume of 100 μ L, were injected into the third mammary fat pad of anaesthetised 8-week-old female Balb/C mice. After a 1-week tumour establishment period, mice received either 40 mg sunitinib (Selleck Chemicals, USA) per kg body weight in phosphate buffered saline (PBS) (Sigma) and 3.72% DMSO (Sigma) or the drug vehicle only. The tumour growth was measured by daily caliper measurements.