

The generic character table of a Sylow p -subgroup of a finite Chevalley group of type D_4

Magaard, Kay; Goodwin, Simon; Le, Tung

DOI:

[10.1080/00927872.2017.1298771](https://doi.org/10.1080/00927872.2017.1298771)

Document Version

Peer reviewed version

Citation for published version (Harvard):

Magaard, K, Goodwin, S & Le, T 2017, 'The generic character table of a Sylow p -subgroup of a finite Chevalley group of type D_4 ', *Communications in Algebra*. <https://doi.org/10.1080/00927872.2017.1298771>

[Link to publication on Research at Birmingham portal](#)

Publisher Rights Statement:

Checked for eligibility: 16/02/2017

"This is an Accepted Manuscript of an article published by Taylor & Francis in *Communications in Algebra* on 2nd March 2017, available online: <http://www.tandfonline.com/10.1080/00927872.2017.1298771>

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

THE GENERIC CHARACTER TABLE OF A SYLOW p -SUBGROUP OF A FINITE CHEVALLEY GROUP OF TYPE D_4

SIMON M. GOODWIN, TUNG LE AND KAY MAGAARD

ABSTRACT. Let U be a Sylow p -subgroup of the finite Chevalley group of type D_4 over the field of q elements, where q is a power of a prime p . We describe a construction of the generic character table of U .

1. INTRODUCTION

Let p be a prime and let q be a power of p . Let G be a finite reductive group over the field of q elements, and let U be a Sylow p -subgroup of G . The aim of this paper is to describe a construction of the *generic character table* of U for the case where G is split and of type D_4 . This generic character table can be determined immediately from the sequence of propositions and lemmas in Section 5. The construction is “uniform” over primes $p > 2$, but we observe differences for the bad prime $p = 2$.

There has been much research into constructing generic character tables of G . For G of small rank these have been programmed into the computing system CHEVIE, [10]. More recently there has been interest in generic character tables for Borel and parabolic subgroups of G , see for example [17], [19], [21], [22] and [25], where many cases for G of \mathbb{F}_q -rank 3 or less are considered. These papers and similar methods have led to applications in determining decomposition numbers in the modular character theory of G , see for example, [1], [18], [20], [23], [26], [34] and [41].

There has been a great deal of research interest in the character theory of U , with motivation coming from understanding how this character theory varies with q .

Interest in properties of U for $G = \mathrm{GL}_n(q)$ goes back to G. Higman [16], where the conjugacy classes of U are considered. In [31], G. Lehrer determined how discrete series characters decompose when restricted to U for $G = \mathrm{GL}_n(q)$. This led to a study of the irreducible characters of U and in particular it was conjectured that the degrees of the irreducible characters are always a power of q , and that the number of characters of degree q^d , for $d \in \mathbb{Z}_{\geq 0}$ is given by a polynomial in q with integer coefficients; this refines a well known conjecture attributed to Higman. There has been further work in this direction, for $G = \mathrm{GL}_n(q)$, by I. M. Isaacs, and more generally on the character theory of algebra groups, see [27] and [28]. In particular, Isaacs verified that the degree of any irreducible character of an algebra group over \mathbb{F}_q is a power of q . For other recent developments we refer, for example, to the work of A. Evseev [8], E. Marberg [32], and I. Pak and A. Soffer [35]; in particular, we remark that the results of [35] suggest that the aforementioned conjecture on the number of characters being given by a polynomial in q is false. An interesting aspect of the character theory of U is the theory of supercharacters, which was first studied by André, see for example [2]; this theory was fully developed by Diaconis and Isaacs in [6].

There has been much interest on the complex irreducible characters of Sylow p -subgroups of finite reductive groups of other types. For instance, it has been proved that for G of type B , C or D the degrees of all irreducible characters of $U(q)$ are powers of q if and only if $p \neq 2$, see [36], [37] and [40]. There has been much further interest in determining certain families of irreducible character, see for example [15], [33], [38] and [39]. Recently, a notion of supercharacters of U for G of types B , C and D was introduced by André and Neto in [3]. For p larger than the Coxeter number of G , the Kirillov orbit method gives a bijection between the coadjoint orbits of U on its Lie algebra and the irreducible characters of U ; in [13] an algorithm is given, which is used to calculate a parametrization of these coadjoint orbits when G is split and of rank less than or equal to 8, excluding type E_8 . We refer the reader also to [30] for other results on the parametrization of irreducible characters of U .

From now on we suppose G is a finite Chevalley group of type D_4 . The two key ingredients that enable us to calculate the generic table of U are the papers [24] and [14], which provide a parametrization of the irreducible characters and the conjugacy classes of U ; we also use [4] for the conjugacy classes for the case $p = 2$. Our construction of this generic character table is given in such a way that it is straightforward to convert it to a computer file that can be used in future calculations.

Our parametrization of the characters and conjugacy classes of U partitions them into families that are independent of q , as can be observed in Tables 3 and 4. We have refined the parameterizations from [14] and [24], so that they are stable under the automorphisms of G coming from the graph automorphisms of the Dynkin diagram; this is likely to be of significant benefit for applications. Our calculations are based on developing the methods from [24]. A key additional ingredient is that we modify those constructions making them more explicit by realizing the irreducible characters of U as induced characters of linear characters of certain subgroups.

By the *generic character table* we mean the table with rows labelled by the families of the characters, and columns labelled by the families of representatives of conjugacy classes. The values in the table are given by expressions depending on the parameters for the families and on q . This can be viewed as an extension of the ideas discussed above regarding how the character theory of U varies with q .

We observe that the characters and their values are uniform for $p > 2$, whereas for $p = 2$ there are differences. More explicitly the parametrization of the family of irreducible characters $\mathcal{F}_{8,9,10}$ is different for $p = 2$, and although the parametrization of the characters in the families \mathcal{F}_{11} and \mathcal{F}_{12} is not different for $p = 2$, the values of these characters do differ. These differences give some explanation of why $p = 2$ is a bad prime for D_4 .

We summarize our results in the following theorem.

Theorem 1.1. *Let G be a finite Chevalley group of type D_4 over \mathbb{F}_q and let U be a Sylow p -subgroup of G . The generic character table of U is determined from: the parametrization of the conjugacy classes given in Section 3; the parametrization of the irreducible characters given in Section 4; and the character values given in Section 5.*

The methods in this paper serve as a model, which can be applied to determine generic character tables of Sylow p -subgroups of finite groups of Lie type, where p is the defining characteristic.

We outline the structure of paper. First we recall the required theory and notation regarding the finite Chevalley group of type D_4 in Section 2. Then in Sections 3 and 4, we

recall the results from [4], [12] and [24] giving the parametrization of the conjugacy classes and irreducible characters of U . The majority of the work is done in Section 5, where we determine the character values.


Acknowledgements: We would like to thank the referee for many helpful suggestions that led to improvements in the paper. We also thank A. Paolini for pointing out some corrections. Part of this research was completed during a visit of the second author to the University of Birmingham; we thank the LMS for a grant to cover his expenses.

2. NOTATION AND SETUP

Let p be a prime, let q be a power of p , and denote the finite field with q elements by \mathbb{F}_q . We let \mathbf{G} denote a simply connected simple algebraic group of type D_4 defined and split over \mathbb{F}_p ; in other words \mathbf{G} is Spin_8 defined over \mathbb{F}_p . Let \mathbf{B} be a Borel subgroup of \mathbf{G} defined over \mathbb{F}_p , let \mathbf{U} be the unipotent radical of \mathbf{B} , and $\mathbf{T} \subseteq \mathbf{B}$ be a maximal torus defined over \mathbb{F}_p . For a subgroup \mathbf{H} of \mathbf{G} defined over \mathbb{F}_p , we write $H = \mathbf{H}(q)$ for the group of \mathbb{F}_q -rational points of \mathbf{H} . So $G = \mathbf{G}(q)$ is a finite Chevalley group of type D_4 and $U = \mathbf{U}(q)$ is a Sylow p -subgroup of G . We cite [7, Sec. 3], as a reference for the theory of algebraic groups over finite fields, and for the terminology used here.

We note that $G = \text{Spin}_8^+(q)$ is not a simple group, but that $G/Z(G) \cong P\Omega_8^+(q)$ is a simple group. As we are mainly interested in U in this paper, which is isomorphic to its image in $G/Z(G)$, this distinction causes us no harm.

Let Φ be a root system of \mathbf{G} with respect to \mathbf{T} and let Φ^+ be the system of positive roots determined by \mathbf{B} . We write $\{\alpha_1, \alpha_2, \alpha_3, \alpha_4\}$ for the corresponding base of simple roots ordered so that the Dynkin diagram of Φ is



For $\alpha = \sum_{i=1}^4 a_i \alpha_i \in \Phi$ we write $\text{ht}(\alpha) = \sum_{i=1}^4 a_i$ for the height of α . We use the notation $\frac{1}{1} \ 2 \ 1$ for the root $\alpha_1 + \alpha_2 + 2\alpha_3 + \alpha_4 \in \Phi^+$ and we use a similar notation for the remaining positive roots. We enumerate the roots in Φ^+ as shown in Table 1 below.

The triality automorphism of the Dynkin diagram can be lifted to an automorphism of \mathbf{G} that stabilizes \mathbf{T} and \mathbf{B} , and restricts to an automorphism of G . We refer the reader to [5, Section 4.2] for more information and let τ denote this automorphism of G . In a similar way we can define three automorphisms of G corresponding to the involutory automorphisms of the Dynkin diagram given by switching two of the outer vertices. We denote these by σ_{12} , σ_{14} and σ_{24} in the obvious way.

The group G is generated by the root elements $x_\alpha(t)$ for $\alpha \in \Phi$ and $t \in \mathbb{F}_q$. We may choose these root elements so that, for $\alpha, \beta \in \Phi$ such that $\alpha + \beta \in \Phi$, we have $[x_\alpha(t), x_\beta(u)] = x_{\alpha+\beta}(\pm tu)$ and the signs are invariant under τ and the σ_{ij} . For positive roots, we use the abbreviation $x_i(t) := x_{\alpha_i}(t)$ and let $X_i = \{x_i(t) \mid t \in \mathbb{F}_q\}$, for $i = 1, 2, \dots, 12$. We have $\tau(x_i(t)) = x_{i\rho}(t)$ where ρ is the permutation $\rho = (1, 4, 2)(5, 7, 6)(8, 9, 10)$; and

| Height | Roots | | | |
|--------|---|--|---|------------|
| 1 | α_1 | α_2 | α_3 | α_4 |
| 2 | $\alpha_5 := \begin{smallmatrix} 1 & 1 & 0 \\ 0 & & \end{smallmatrix}$ | $\alpha_6 := \begin{smallmatrix} 0 & 1 & 0 \\ 1 & & \end{smallmatrix}$ | $\alpha_7 := \begin{smallmatrix} 0 & 1 & 1 \\ 0 & & \end{smallmatrix}$ | |
| 3 | $\alpha_8 := \begin{smallmatrix} 1 & 1 & 0 \\ 1 & & \end{smallmatrix}$ | $\alpha_9 := \begin{smallmatrix} 1 & 1 & 1 \\ 0 & & \end{smallmatrix}$ | $\alpha_{10} := \begin{smallmatrix} 0 & 1 & 1 \\ 1 & & \end{smallmatrix}$ | |
| 4 | $\alpha_{11} := \begin{smallmatrix} 1 & 1 & 1 \\ 1 & & \end{smallmatrix}$ | | | |
| 5 | $\alpha_{12} := \begin{smallmatrix} 1 & 2 & 1 \\ 1 & & \end{smallmatrix}$ | | | |

TABLE 1. Positive roots of the root system Φ of type D_4 .

have a similar description for the automorphisms σ_{ij} . The commutators $[x_i(t), x_j(u)] = x_i(t)^{-1}x_j(u)^{-1}x_i(t)x_j(u)$ are given in Table 2. All $[x_i(t), x_j(u)]$ not listed in this table are equal to 1. We refer the reader to [5, Chapter 4] for details on root elements and commutator relations.

$$\begin{array}{ll}
[x_1(t), x_3(u)] &= x_5(tu), & [x_1(t), x_6(u)] &= x_8(tu), \\
[x_1(t), x_7(u)] &= x_9(tu), & [x_1(t), x_{10}(u)] &= x_{11}(tu), \\
[x_2(t), x_3(u)] &= x_6(tu), & [x_2(t), x_5(u)] &= x_8(tu), \\
[x_2(t), x_7(u)] &= x_{10}(tu), & [x_2(t), x_9(u)] &= x_{11}(tu), \\
[x_3(t), x_4(u)] &= x_7(-tu), & [x_3(t), x_{11}(u)] &= x_{12}(tu), \\
[x_4(t), x_5(u)] &= x_9(tu), & [x_4(t), x_6(u)] &= x_{10}(tu), \\
[x_4(t), x_8(u)] &= x_{11}(tu), & [x_5(t), x_{10}(u)] &= x_{12}(-tu), \\
[x_6(t), x_9(u)] &= x_{12}(-tu), & [x_7(t), x_8(u)] &= x_{12}(-tu).
\end{array}$$

TABLE 2. Commutator relations for type D_4 .

We have that U is the subgroup of G generated by the elements $x_i(t_i)$ for $i = 1, 2, \dots, 12$ and $t_i \in \mathbb{F}_q$. In fact we have that $U = \prod_{i=1}^{12} X_i$. It turns out that it is convenient for us to write our elements of U with an element from X_3 at the front, and we use the notation

$$(2.1) \quad x(\mathbf{t}) = x_3(t_3)x_1(t_1)x_2(t_2)x_4(t_4)x_5(t_5) \cdots x_{12}(t_{12}),$$

where $\mathbf{t} = (t_1, \dots, t_{12})$.

There is a further piece of notation that we use frequently in the sequel, we define the normal subgroups M_i of U for $i = 1, \dots, 13$ by

$$\begin{aligned}
(2.2) \quad M_1 &= U, \\
M_2 &= X_1 X_2 \prod_{j=4}^{12} X_j, \\
M_3 &= X_2 \prod_{j=4}^{12} X_j \text{ and}
\end{aligned}$$

$$(2.3) \quad M_i = \prod_{j=i}^{12} X_j \quad \text{for } i = 4, \dots, 13.$$

3. CONJUGACY CLASSES AND THEIR REPRESENTATIVES

In [14], an algorithm for calculating the conjugacy classes of Sylow p -subgroups of finite Chevalley groups in good characteristic is described; so we assume $p > 2$ for the first part of this section. This algorithm was implemented in GAP4 [9] and, in particular, used to determine the conjugacy classes of $U = U(q)$. The conjugacy class representatives are given as *minimal representatives* as defined in [11] and discussed below.

(For ease of exposition we temporarily let $x_1 = x_{\alpha_3}$, $x_2 = x_{\alpha_1}$ and $x_3 = x_{\alpha_2}$ in the next paragraph.)

The idea behind the calculation of minimal representatives is to recursively determine the orbits of U on U/M_{i+1} for $i = 1, \dots, 12$; recall that M_{i+1} is defined in (2.2). To explain this we consider the set $A(x, i) = \{xx_i(t)M_{i+1} \mid t \in \mathbb{F}_q\} \subseteq U/M_{i+1}$ for $x \in U$. The centralizer $C_U(xM_i)$ acts on $A(x, i)$ by conjugation. The key results for the theory of minimal representatives are [11, Lem. 5.1 and Prop. 6.2], which imply the following dichotomy:

- (I) all elements of $A(x, i)$ are conjugate under $C_U(xM_i)$; or
- (R) no two elements of $A(x, i)$ are conjugate under $C_U(xM_i)$.

We say that i is an *inert point* of x if (I) holds and i is a *ramification point* of x if (R) holds. We say $x(\mathbf{t})$ as defined in (2.1) is the *minimal representative* of its conjugacy class if $t_i = 0$ whenever i is an inert point of $x(\mathbf{t})$. Thanks to [11, Prop. 5.4 and Prop. 6.2], the minimal representatives of conjugacy classes give a complete set of representatives of the conjugacy classes in U .

In Table 3, we present the minimal representatives of the conjugacy classes in $U(q)$. Starting with the minimal representatives calculated as in [14], we have parameterized the conjugacy classes giving representatives partitioned into families. This has been done to reduce the number of families required to what appears to be an optimal number.

The first column of Table 3 gives a label for the family, then in the second column a parametrization of the elements of the family is given, where the notational convention is that an a_i is an element of $\mathbb{F}_q^\times = \mathbb{F}_q \setminus \{0\}$, a b_i is any element of \mathbb{F}_q , and for a sequence of the form c_{i_1}, \dots, c_{i_r} , the c_{i_j} are elements of \mathbb{F}_q , which are not all zero. These parameters have to satisfy the conditions given in the third column. In the fourth column we give the number of elements in the family. The families are constructed so that the size of the centralizer of an element in the family does not depend on the choice, and the final column gives this centralizer size.

The families of representatives have been chosen, so that they are permuted by the automorphisms τ , σ_{12} , σ_{14} and σ_{24} . Thus we can read off the action of these automorphisms on the conjugacy classes. However, for the family, $\mathcal{C}_{1,2,4,q^6}$, it is not possible for $p = 3$ to get this stability under the triality automorphism and we have to distinguish the cases $p = 3$ and $p > 3$; we note that the family given for $p = 3$ is actually also valid for $p > 3$.

To end this section we explain the differences in the parametrization of the conjugacy classes for $p = 2$, where the theory of minimal representatives fails. First we note that there are a number of families, where there is a relation between parameters involving a sum of two

| Label | Representative | Conditions | Number | Centralizer |
|-----------------------------------|--|-----------------------------------|------------------|-------------|
| $\mathcal{C}_{1,2,3,4}$ | $x_3(a_3)x_1(a_1)x_2(a_2)x_4(a_4)$ | | $(q-1)^4$ | q^4 |
| $\mathcal{C}_{1,2,3}$ | $x_3(a_3)x_1(a_1)x_2(a_2)x_9(b_9)x_{10}(b_{10})$ | $b_9/a_1 + b_{10}/a_2 = 0$ | $(q-1)^3q$ | q^5 |
| $\mathcal{C}_{1,3,4}$ | $x_3(a_3)x_1(a_1)x_4(a_4)x_8(b_8)x_{10}(b_{10})$ | $b_8/a_1 + b_{10}/a_4 = 0$ | $(q-1)^3q$ | q^5 |
| $\mathcal{C}_{2,3,4}$ | $x_3(a_3)x_2(a_1)x_4(a_4)x_8(b_8)x_9(b_9)$ | $b_8/a_2 + b_9/a_3 = 0$ | $(q-1)^3q$ | q^5 |
| $\mathcal{C}_{1,3}$ | $x_3(a_3)x_1(a_1)x_{10}(b_{10})$ | | $(q-1)^2q$ | q^5 |
| $\mathcal{C}_{2,3}$ | $x_3(a_3)x_2(a_2)x_9(b_9)$ | | $(q-1)^2q$ | q^5 |
| $\mathcal{C}_{3,4}$ | $x_3(a_3)x_4(a_4)x_8(b_8)$ | | $(q-1)^2q$ | q^5 |
| $\mathcal{C}_{1,2,4,q^6} (p > 3)$ | $x_1(a_1)x_2(a_2)x_4(a_4)x_5(c_5)x_6(c_6)x_7(c_7)$ | $c_5/a_1 + c_6/a_2 + c_7/a_4 = 0$ | $(q-1)^3(q^2-1)$ | q^6 |
| $\mathcal{C}_{1,2,4,q^6} (p = 3)$ | $x_1(a_1)x_2(a_2)x_4(a_4)x_6(c_6)x_7(c_7)$ | | $(q-1)^3(q^2-1)$ | q^6 |
| $\mathcal{C}_{1,2,4,q^7}$ | $x_1(a_1)x_2(a_2)x_4(a_4)x_{12}(b_{12})$ | | $(q-1)^3q$ | q^7 |
| $\mathcal{C}_{1,2,q^6}$ | $x_1(a_1)x_2(a_2)x_5(c_5)x_6(c_6)x_7(c_7)$ | $c_5/a_1 + c_6/a_2 = 0$ | $(q-1)^2(q^2-1)$ | q^6 |
| $\mathcal{C}_{1,2,q^7}$ | $x_1(a_1)x_2(a_2)x_9(a_9)x_{10}(a_{10})$ | $a_9/a_1 + a_{10}/a_2 = 0$ | $(q-1)^3$ | q^7 |
| $\mathcal{C}_{1,2,q^8}$ | $x_1(a_1)x_2(a_2)x_{12}(b_{12})$ | | $(q-1)^2q$ | q^8 |
| $\mathcal{C}_{1,4,q^6}$ | $x_1(a_1)x_4(a_4)x_5(c_5)x_6(c_6)x_7(c_7)$ | $c_5/a_1 + c_7/a_4 = 0$ | $(q-1)^2(q^2-1)$ | q^6 |
| $\mathcal{C}_{1,4,q^7}$ | $x_1(a_1)x_4(a_4)x_8(a_8)x_{10}(a_{10})$ | $a_8/a_1 + a_{10}/a_4 = 0$ | $(q-1)^3$ | q^7 |
| $\mathcal{C}_{1,4,q^8}$ | $x_1(a_1)x_4(a_4)x_{12}(b_{12})$ | | $(q-1)^2q$ | q^8 |
| $\mathcal{C}_{2,4,q^6}$ | $x_2(a_2)x_4(a_4)x_5(c_5)x_6(c_6)x_7(c_7)$ | $c_6/a_2 + c_7/a_4 = 0$ | $(q-1)^2(q^2-1)$ | q^6 |
| $\mathcal{C}_{2,4,q^7}$ | $x_2(a_2)x_4(a_4)x_8(a_8)x_9(a_9)$ | $a_8/a_2 + a_9/a_4 = 0$ | $(q-1)^3$ | q^7 |
| $\mathcal{C}_{2,4,q^8}$ | $x_2(a_2)x_4(a_4)x_{12}(b_{12})$ | | $(q-1)^2q$ | q^8 |
| \mathcal{C}_{1,q^6} | $x_1(a_1)x_6(c_6)x_7(c_7)$ | | $(q-1)(q^2-1)$ | q^6 |
| \mathcal{C}_{1,q^7} | $x_1(a_1)x_{10}(a_{10})$ | | $(q-1)^2$ | q^7 |
| \mathcal{C}_{1,q^8} | $x_1(a_1)x_{12}(b_{12})$ | | $(q-1)q$ | q^8 |
| \mathcal{C}_{2,q^6} | $x_2(a_2)x_5(c_5)x_7(c_7)$ | | $(q-1)(q^2-1)$ | q^6 |
| \mathcal{C}_{2,q^7} | $x_2(a_2)x_9(a_9)$ | | $(q-1)^2$ | q^7 |
| \mathcal{C}_{2,q^8} | $x_2(a_2)x_{12}(b_{12})$ | | $(q-1)q$ | q^8 |
| \mathcal{C}_{4,q^6} | $x_4(a_4)x_5(c_5)x_6(c_6)$ | | $(q-1)(q^2-1)$ | q^6 |
| \mathcal{C}_{4,q^7} | $x_4(a_4)x_8(a_8)$ | | $(q-1)^2$ | q^7 |
| \mathcal{C}_{4,q^8} | $x_4(a_4)x_{12}(b_{12})$ | | $(q-1)q$ | q^8 |
| \mathcal{C}_3 | $x_3(a_3)x_8(b_8)x_9(b_9)x_{10}(b_{10})x_{11}(b_{11})$ | | $(q-1)q^4$ | q^8 |
| $\mathcal{C}_{5,6,7}$ | $x_5(a_5)x_6(a_6)x_7(a_7)x_{11}(b_{11})$ | | $(q-1)^3q$ | q^8 |
| $\mathcal{C}_{5,6,q^8}$ | $x_5(a_5)x_6(a_6)x_9(a_9)x_{10}(a_{10})$ | $a_9/a_5 + a_{10}/a_6 = 0$ | $(q-1)^3$ | q^8 |
| $\mathcal{C}_{5,6,q^9}$ | $x_5(a_5)x_6(a_6)x_{11}(b_{11})$ | | $(q-1)^2q$ | q^9 |
| $\mathcal{C}_{5,7,q^8}$ | $x_5(a_5)x_7(a_7)x_8(a_8)x_{10}(a_{10})$ | $a_8/a_5 + a_{10}/a_7 = 0$ | $(q-1)^3$ | q^8 |
| $\mathcal{C}_{5,7,q^9}$ | $x_5(a_5)x_7(a_7)x_{11}(b_{11})$ | | $(q-1)^2q$ | q^9 |
| $\mathcal{C}_{6,7,q^8}$ | $x_6(a_6)x_7(a_7)x_8(a_8)x_9(a_9)$ | $a_8/a_6 + a_9/a_7 = 0$ | $(q-1)^3$ | q^8 |
| $\mathcal{C}_{6,7,q^9}$ | $x_6(a_6)x_7(a_7)x_{11}(b_{11})$ | | $(q-1)^2q$ | q^9 |
| \mathcal{C}_{5,q^8} | $x_5(a_5)x_{10}(a_{10})$ | | $(q-1)^2$ | q^8 |
| \mathcal{C}_{5,q^9} | $x_5(a_5)x_{11}(b_{11})$ | | $(q-1)q$ | q^9 |
| \mathcal{C}_{6,q^8} | $x_6(a_6)x_9(a_9)$ | | $(q-1)^2$ | q^8 |
| \mathcal{C}_{6,q^9} | $x_6(a_6)x_{11}(b_{11})$ | | $(q-1)q$ | q^9 |
| \mathcal{C}_{7,q^8} | $x_7(a_7)x_8(a_8)$ | | $(q-1)^2$ | q^8 |
| \mathcal{C}_{7,q^9} | $x_7(a_7)x_{11}(b_{11})$ | | $(q-1)q$ | q^9 |
| $\mathcal{C}_{8,9,10}$ | $x_8(c_8)x_9(c_9)x_{10}(c_{10})$ | | $q^3 - 1$ | q^{10} |
| \mathcal{C}_{11} | $x_{11}(a_{11})$ | | $q - 1$ | q^{11} |
| \mathcal{C}_{12} | $x_{12}(b_{12})$ | | q | q^{12} |

TABLE 3. Representatives of conjugacy classes in $U(q)$ for $p > 2$

terms and this has to be replaced by just setting one of the parameters equal to zero. For example in $\mathcal{C}_{1,2,3}$, we have to replace the representatives simply with $x_3(a_3)x_1(a_1)x_2(a_2)x_{10}(b_{10})$ and there are no conditions. This does lead to the families of representatives not being stable under the automorphisms, but this cannot be helped.

The more significant differences are as follows.

- The family $\mathcal{C}_{1,2,3,4}$ is replaced by the family $\mathcal{C}_{1,2,3,4}^{p=2}$. The representatives are of the form $x_3(a_3)x_1(a_1)x_2(a_2)x_4(a_4)x_{10}(d_{10})$, where d_{10} is either 0, or an element of \mathbb{F}_q not in the image of the map $t \mapsto a_3a_2a_4(t^2 + t)$. So there are $2(q-1)^4$ representatives in this family. Their centralizers have order $2q^4$.
- The family $\mathcal{C}_{1,2,4,q^6}$ should be included as in the $p = 3$ case in the table above.
- The family $\mathcal{C}_{1,2,4,q^7}$ is replaced by two families.
 - $\mathcal{C}_{1,2,4,2q^7}^{p=2}$. The representatives are of the form $x_1(a_1)x_2(a_2)x_4(a_4)x_{10}(a_{10})x_{12}(d_{12})$, where d_{12} is either 0, or an element of \mathbb{F}_q not in the image of the map $t \mapsto a_1a_2a_4t^2 + a_1a_{10}t$. So there are $2(q-1)^4$ representatives in this family. Their centralizers have order $2q^7$.
 - $\mathcal{C}_{1,2,4,q^7}^{p=2}$. The representatives are of the form $x_1(a_1)x_2(a_2)x_4(a_4)$. There are $(q-1)^3$ representatives in this family and their centralizers have order q^7 .
- The family $\mathcal{C}_{5,6,7}$ is replaced by two new families namely.
 - $\mathcal{C}_{5,6,7,2q^8}^{p=2}$. The representatives are of the form $x_5(a_5)x_6(a_6)x_7(a_7)x_{10}(a_{10})x_{11}(d_{11})$, where d_{11} is either 0, or an element of \mathbb{F}_q not in the image of the map $t \mapsto a_5a_6a_7t^2 + a_5a_{10}t$. There are $2(q-1)^4$ representatives in this family and their centralizers have order $2q^8$.
 - $\mathcal{C}_{5,6,7,q^8}^{p=2}$. The representatives are of the form $x_5(a_5)x_6(a_6)x_7(a_7)$. The number of representatives in this family is $(q-1)^3$ and their centralizers have order q^8 .

4. THE IRREDUCIBLE CHARACTERS OF U

In [24], the irreducible characters of U were determined giving a partition into families, which are given in Table 4 for the case $p > 2$. We have modified these constructions to make them more explicit, so that we can realize them all as induced characters from linear characters of certain subgroups of U . In the next section we calculate the values of the characters, and in doing so we explain their construction.

We explain the contents of Table 4. The first column gives the name of the family and the second column gives the names of the characters in that family. The notational convention is that parameters denoted by a_i range over $\mathbb{F}_q^\times = \mathbb{F}_q \setminus \{0\}$, and parameters denoted by b_i range over \mathbb{F}_q . The subscripts in the notation give an idea of “where these parameters act”, as can be understood from the descriptions of how the characters are constructed given in the next section. For some families there is a relation between some of the parameters and this is given in the third column. The fourth column records the number of characters in that family. The families are constructed in such a way that the degree of the characters in a family is constant, and the fifth column gives this degree.

The triality automorphism τ of G , permutes the families of characters by acting on the subscripts of the families by the permutation ρ ; similarly the automorphisms σ_{12} , σ_{14} and σ_{24} act on the families. As mentioned earlier, we have updated the parametrization from [24, §4], so that we can read off the action of these automorphisms on the characters. Though it is not possible to do this for the family $\mathcal{F}_{5,6,7}$ for $p = 3$.

We end this section by noting that for $p = 2$, the parametrization of irreducible characters is slightly different. As explained in [24], the family denoted $\mathcal{F}_{8,9,10}$ is more complicated and has to be replaced by the family $\mathcal{F}_{8,9,10}^{p=2}$, which consists of two types of characters as explained below and in more detail at the end of §5.4.

| Family | Notation | Conditions | Number | Degree |
|-------------------------------|--|-----------------------------------|--------------|--------|
| \mathcal{F}_{12} | $\chi_{12}^{a_{12}, b_1, b_2, b_4}$ | | $q^3(q-1)$ | q^4 |
| \mathcal{F}_{11} | $\chi_{11}^{a_{11}, b_5, b_6, b_7, b_3}$ | | $q^4(q-1)$ | q^3 |
| $\mathcal{F}_{8,9,10}$ | $\chi_{8,9,10}^{a_8, a_9, a_{10}, b_3}$ | | $q(q-1)^3$ | q^3 |
| $\mathcal{F}_{8,9}$ | $\chi_{8,9,q^3}^{a_8, a_9, a_6, a_7}$ | $a_6/a_8 + a_7/a_9 = 0$ | $(q-1)^3$ | q^3 |
| | $\chi_{8,9,q^2}^{a_8, a_9, b_2, b_3, b_4}$ | $b_2/a_8 + b_4/a_9 = 0$ | $q^2(q-1)^2$ | q^2 |
| $\mathcal{F}_{8,10}$ | $\chi_{8,10,q^3}^{a_8, a_{10}, a_5, a_7}$ | $a_5/a_8 + a_7/a_{10} = 0$ | $(q-1)^3$ | q^3 |
| | $\chi_{8,10,q^2}^{a_8, a_{10}, b_1, b_3, b_4}$ | $b_1/a_8 + b_4/a_{10} = 0$ | $q^2(q-1)^2$ | q^2 |
| $\mathcal{F}_{9,10}$ | $\chi_{9,10,q^3}^{a_9, a_{10}, a_5, a_6}$ | $a_5/a_9 + a_6/a_{10} = 0$ | $(q-1)^3$ | q^3 |
| | $\chi_{9,10,q^2}^{a_9, a_{10}, b_1, b_2, b_3}$ | $b_1/a_9 + b_2/a_{10} = 0$ | $q^2(q-1)^2$ | q^2 |
| \mathcal{F}_8 | $\chi_{8,q^3}^{a_8, a_7}$ | | $(q-1)^2$ | q^3 |
| | $\chi_{8,q^2}^{a_8, b_3, b_4}$ | | $q^2(q-1)$ | q^2 |
| \mathcal{F}_9 | $\chi_{9,q^3}^{a_9, a_6}$ | | $(q-1)^2$ | q^3 |
| | $\chi_{9,q^2}^{a_9, b_2, b_3}$ | | $q^2(q-1)$ | q^2 |
| \mathcal{F}_{10} | $\chi_{10,q^3}^{a_{10}, a_5}$ | | $(q-1)^2$ | q^3 |
| | $\chi_{10,q^2}^{a_{10}, b_1, b_2}$ | | $q^2(q-1)$ | q^2 |
| $\mathcal{F}_{5,6,7} (p > 3)$ | $\chi_{5,6,7}^{a_5, a_6, a_7, b_1, b_2, b_4}$ | $b_1/a_5 + b_2/a_6 + b_4/a_7 = 0$ | $q^2(q-1)^3$ | q |
| $\mathcal{F}_{5,6,7} (p = 3)$ | $\chi_{5,6,7}^{a_5, a_6, a_7, b_2, b_4}$ | | $q^2(q-1)^3$ | q |
| $\mathcal{F}_{5,6}$ | $\chi_{5,6}^{a_5, a_6, b_1, b_2, b_4}$ | $b_1/a_5 + b_2/a_6 = 0$ | $q^2(q-1)^2$ | q |
| $\mathcal{F}_{5,7}$ | $\chi_{5,7}^{a_5, a_7, b_1, b_2, b_4}$ | $b_1/a_5 + b_4/a_7 = 0$ | $q^2(q-1)^2$ | q |
| $\mathcal{F}_{6,7}$ | $\chi_{6,7}^{a_6, a_7, b_1, b_2, b_4}$ | $b_2/a_6 + b_4/a_7 = 0$ | $q^2(q-1)^2$ | q |
| \mathcal{F}_5 | $\chi_5^{a_5, b_2, b_4}$ | | $q^2(q-1)$ | q |
| \mathcal{F}_6 | $\chi_6^{a_6, b_1, b_4}$ | | $q^2(q-1)$ | q |
| \mathcal{F}_7 | $\chi_7^{a_7, b_1, b_2}$ | | $q^2(q-1)$ | q |
| \mathcal{F}_{lin} | $\chi_{\text{lin}}^{b_1, b_2, b_3, b_4}$ | | q^4 | 1 |

TABLE 4. The irreducible characters of U for $p > 2$.

- The characters $\chi_{8,9,10,q^3}^{a_8, a_9, a_{10}}$. There are $(q-1)^3$ of these and they have degree q^3 .

- The characters $\chi_{8,9,10,\frac{q^3}{2}}^{a_8,a_9,a_{10},a_{5,6,7},d_{1,2,4},d_3}$. There are $4(q-1)^4$ of these and their degree is $\frac{q^3}{2}$. The parameters $d_{1,2,4}$ and d_3 can each take one of two values as explained at the end of §5.4.

Also for $p = 2$ the parametrization of characters where there is a relation between parameters involving the sum of two terms has to be updated in a similar way to the corresponding situation for conjugacy class representatives. For example, in the family $\mathcal{F}_{5,6}$, we just have characters $\chi_{5,6}^{a_5,a_6,b_2}$ (so we have set $b_1 = 0$ and b_2 ranges over \mathbb{F}_q).

5. DETERMINING THE CHARACTER VALUES

In the following subsections we give the construction of the irreducible characters and determine the character values. We split up our determination of the character values according to the families given in Table 4, and in each of these subsections we calculate the values of the irreducible characters on elements of U . For the families considered in §5.1, §5.2 and §5.3 we determine the character values on general elements of U , whereas for the other characters we take advantage of the representatives of the conjugacy classes from Table 3.

Before we embark on calculating the values of the irreducible characters of U , we give some notation that we use.

Denote by $\text{Tr} : \mathbb{F}_q \rightarrow \mathbb{F}_p$ the trace map, and define $\phi : \mathbb{F}_q \rightarrow \mathbb{C}^\times$ by $\phi(x) = e^{\frac{i2\pi \text{Tr}(x)}{p}}$, so that ϕ is a nontrivial character from the additive group of \mathbb{F}_q to the multiplicative group \mathbb{C}^\times .

We record two important elementary observations about ϕ that we require frequently in the sequel. First we note that, for $t \in \mathbb{F}_q$, we have $\phi(st) = 1$ for all $s \in \mathbb{F}_q$ if and only if $t = 0$, second we note that $\sum_{s \in \mathbb{F}_q} \phi(s) = 0$.

Let H be a finite group. We write $Z(H)$ for the centre of H . Let K be a subgroup of H , let $x \in H$, $h \in H$ and let $\psi : K \rightarrow \mathbb{C}$. We write $\dot{\psi} : H \rightarrow \mathbb{C}$ for the function defined by $\dot{\psi}(x) = \psi(x)$ if $x \in K$ and $\dot{\psi}(x) = 0$ if $x \notin K$. We denote conjugation by $x^h = h^{-1}xh$ and ${}^h\psi : H \rightarrow \mathbb{C}$ is defined by ${}^h\psi(x) = \dot{\psi}(x^h)$. If ψ is a class function on K , then we write ψ^H for the induced class function on H , which is given by the standard formula $\psi^H(h) = \frac{1}{|K|} \sum_{x \in H} \dot{\psi}(x^h)$.

We use the notation $x(\mathbf{t})$ from (2.1), where we recall that $x_3(t_3)$ is on the left. Frequently, we consider quotients of U by the normal subgroups M_i defined in (2.2). When we do this we often want to identify the root subgroup X_j with its image in U/M_i for $j < i$, and we will do this without further explanation. We consider subgroups V of U/M_i of the form $\prod_{j \in I} X_j$, where I is a subset of $\{1, \dots, i\}$. For such a subgroup V we use the notation $x_V(\mathbf{t}) = \prod_{j \in I} x_j(t_j)$ to denote a general element of V , where the factors are ordered as in $x(\mathbf{t})$, so that $x_3(t_3)$ is on the left (if $3 \in I$).

We often use the notation $\delta_{\mathbf{s},0}$, where $\mathbf{s} \in \mathbb{F}_q^m$ for some m , which is defined as usual by $\delta_{\mathbf{s},0} = 1$ if $\mathbf{s} = 0$ and $\delta_{\mathbf{s},0} = 0$ if $\mathbf{s} \neq 0$.

As mentioned earlier, we have adapted the construction of the characters from [24] so that they are more explicit. In particular, we realise each of the irreducible characters as an induced character of a certain subgroup, which leads to a more precise parametrization. Given these differences, we briefly explain why our constructions line up with those in [24].

For each irreducible character χ that we construct, we give a subgroup V of a quotient U/M of U and a linear character λ of V . Then χ is equal to the induced character $\lambda^{U/M}$

inflated to U possibly tensored with a linear character of U . In each case there is a subgroup X of U/M , which is a transversal of V in U/M . In each of subsequent subsections, where we calculate the characters values, we state what V and X are. We also justify that the induced characters that we obtain are indeed irreducible, and that the characters as given in Table 4 are distinct. The main tool for achieving this is the Clifford theory for characters of special groups of type q^{1+2m} , as is also the case in [24]. Now we can count the number of characters that we have constructed in each family, and verify that this gives the same numbers in [24]. Therefore, we can deduce that we have covered all of the irreducible characters of U .

5.1. The family \mathcal{F}_{lin} . Here we state the value of $\chi_{\text{lin}}^{b_1, b_2, b_3, b_4}$ on all elements of U . We note that M_5 is the commutator subgroup of U , and the characters $\chi_{\text{lin}}^{b_1, b_2, b_3, b_4}$ are precisely the characters of U/M_5 inflated to U ; they are all the linear characters of U . The following formula, therefore, defines these characters.

$$\chi_{\text{lin}}^{b_1, b_2, b_3, b_4}(x(\mathbf{t})) = \phi(b_1 t_1 + b_2 t_2 + b_3 t_3 + b_4 t_4).$$

5.2. The family $\mathcal{F}_{5,6,7}^*$. The characters in the families $\mathcal{F}_5, \mathcal{F}_6, \mathcal{F}_7, \mathcal{F}_{5,6}, \mathcal{F}_{5,7}, \mathcal{F}_{6,7}$ and $\mathcal{F}_{5,6,7}$ can all be constructed in essentially the same way. Thus we combine these families and denote their union by $\mathcal{F}_{5,6,7}^*$.

We begin by giving the construction of the characters $\chi_{5,6,7}^{c_5, c_6, c_7, b_1, b_2, b_4}$, where $c_5, c_6, c_7 \in \mathbb{F}_q$ are not all zero and $b_1, b_2, b_4 \in \mathbb{F}_q$. From these characters we obtain all of the characters in $\mathcal{F}_{5,6,7}^*$. First we note that M_8 is in the kernel of $\chi_{5,6,7}^{c_5, c_6, c_7, b_1, b_2, b_4}$. As explained at the start of the section we identify X_i with its image in U/M_8 for $i = 1, 2, \dots, 7$.

We have that $U/M_8 = X_3 X_1 X_2 X_4 X_5 X_6 X_7$, and that $V = X_1 X_2 X_4 X_5 X_6 X_7 \cong \mathbb{F}_q^6$ is an elementary abelian subgroup of U/M_8 of order q^6 , and $X = X_3$ is a transversal of U in U/M_8 . We define the linear character λ^{c_5, c_6, c_7} of V by $\lambda^{c_5, c_6, c_7}(x_V(\mathbf{t})) = \phi(c_5 t_5 + c_6 t_6 + c_7 t_7)$.

Let $\chi_{5,6,7}^{c_5, c_6, c_7}$ be the character of U/M_8 we obtain by inducing λ^{c_5, c_6, c_7} ; by a mild abuse of notation we also write $\chi_{5,6,7}^{c_5, c_6, c_7}$ for the character of U given by inflation. The character $\chi_{5,6,7}^{c_5, c_6, c_7, b_1, b_2, b_4}$ is defined by tensoring $\chi_{5,6,7}^{c_5, c_6, c_7}$ with the linear character $\chi_{\text{lin}}^{b_1, b_2, 0, b_4}$.

The following proposition gives the character values of $\chi_{5,6,7}^{c_5, c_6, c_7, b_1, b_2, b_4}$. It is clear from the calculations in the proof below that $x \lambda^{c_5, c_6, c_7} \neq \lambda^{c_5, c_6, c_7}$ for all $x \in X$ with $x \neq 1$, so that $\chi_{5,6,7}^{c_5, c_6, c_7, b_1, b_2, b_4}$ is irreducible by an application of Clifford theory.

Proposition 5.1.

$$\chi_{5,6,7}^{c_5, c_6, c_7, b_1, b_2, b_4}(x(\mathbf{t})) = q \delta_{(t_3, c_5 t_1 + c_6 t_2 + c_7 t_4), 0} \phi(b_1 t_1 + b_2 t_2 + b_4 t_4 + c_5 t_5 + c_6 t_6 + c_7 t_7).$$

Proof. We write λ for λ^{c_5, c_6, c_7} and χ for $\chi_{5,6,7}^{c_5, c_6, c_7}$. We have that $X = X_3$ is a transversal of V in U , and also that $Z(U/M_8) = X_5 X_6 X_7$. From the induction formula we easily obtain that $\chi(x(\mathbf{t})) = 0$ if $t_3 \neq 0$, and also that $\chi(x(\mathbf{t})) = \delta_{t_3, 0} \chi(x_1(t_1)x_2(t_2)x_4(t_4))\lambda(x_5(t_5)x_6(t_6)x_7(t_7))$. Thus it suffices to calculate $\chi(x_1(t_1)x_2(t_2)x_4(t_4))$. We have

$$\begin{aligned} \chi(x_1(t_1)x_2(t_2)x_4(t_4)) &= \sum_{s \in \mathbb{F}_q} x_3(s) \lambda(x_1(t_1)x_2(t_2)x_4(t_4)) \\ &= \sum_{s \in \mathbb{F}_q} \lambda((x_1(t_1)x_2(t_2)x_4(t_4))^{x_3(s)}) \end{aligned}$$

$$\begin{aligned}
&= \sum_{s \in \mathbb{F}_q} \lambda([x_1(t_1), x_3(s)][x_2(t_2), x_3(s)][x_4(t_4), x_3(s)]) \\
&= \sum_{s \in \mathbb{F}_q} \lambda(x_5(st_1)x_6(st_2)x_7(st_4)) \\
&= \sum_{s \in \mathbb{F}_q} \phi(s(c_5t_1 + c_6t_2 + c_7t_4)) \\
&= q\delta_{c_5t_1+c_6t_2+c_7t_4,0}.
\end{aligned}$$

From this we can deduce the proposition. \square

We note that the characters $\chi_{5,6,7}^{c_5, c_6, c_7, b_1, b_2, b_4}$ can be equal for different values of b_i . A fairly easy calculation shows that $\chi_{5,6,7}^{c_5, c_6, c_7, b_1, b_2, b_4} = \chi_{5,6,7}^{c_5, c_6, c_7, b'_1, b'_2, b'_4}$ if and only if $(b_1 - b'_1, b_2 - b'_2, b_4 - b'_4)$ is a multiple of (c_5, c_6, c_7) . Thus we choose representatives as follows for $p \neq 2, 3$.

- If $c_5 \neq 0$ and $c_6, c_7 = 0$, then we take $b_1 = 0$.
- If $c_5, c_6 \neq 0$ and $c_7 = 0$, then we take b_1, b_2 such that $c_6b_1 + c_5b_2 = 0$.
- If $c_5, c_6, c_7 \neq 0$, then we take b_1, b_2, b_4 such that $c_6c_7b_1 + c_5c_7b_2 + c_5c_6b_4 = 0$.

We deal with the other cases symmetrically with respect to the triality. This gives the characters as in Table 4, and we can give their character values from Proposition 5.1.

We note that for $p = 2$ or $p = 3$ we have to choose our representatives slightly differently. For $p = 3$ the only difference is when $c_5, c_6, c_7 \neq 0$ and this is shown in Table 4. In the case $p = 2$, and we have $c_5, c_6 \neq 0$ and $c_7 = 0$, then we take $b_1 = 0$, and we deal with the other cases symmetrically with respect to the triality.

5.3. The Families $\mathcal{F}_8, \mathcal{F}_9, \mathcal{F}_{10}, \mathcal{F}_{8,9}, \mathcal{F}_{8,10}$ and $\mathcal{F}_{9,10}$. The characters in the families $\mathcal{F}_8, \mathcal{F}_9, \mathcal{F}_{10}, \mathcal{F}_{8,9}, \mathcal{F}_{8,10}$ and $\mathcal{F}_{9,10}$ are constructed in a similar way, so we deal with them together. We include the details for the characters in the family $\mathcal{F}_{8,9}$ and then remark that the calculations for the family \mathcal{F}_8 are entirely similar. The character in the families $\mathcal{F}_9, \mathcal{F}_{10}, \mathcal{F}_{8,10}$ and $\mathcal{F}_{9,10}$ are defined from characters in \mathcal{F}_8 or $\mathcal{F}_{8,9}$ through the triality τ . Thus the values of these characters can be immediately deduced.

From Table 4 we see that the characters in $\mathcal{F}_{8,9}$ fall into two subfamilies determined by degree. We start with the characters of degree q^3 and explain the construction of the characters in this family.

First we note that M_{10} lies in the kernel of the characters in $\mathcal{F}_{8,9}$. We have that $U/M_{10} = \prod_{i=1}^9 X_i$ and we define the normal subgroup

$$V = \prod_{i \neq 1, 2, 4} X_i$$

of U/M_{10} . We have that V is isomorphic to \mathbb{F}_q^6 and so is elementary abelian, and $X = X_1X_2X_4$ is a transversal of V in U/M_{10} . Therefore, we have linear characters $\lambda^{a_8, a_9, a_6, a_7}$ of V defined by

$$\lambda^{a_8, a_9, a_6, a_7}(x_V(\mathbf{t})) = \phi(a_6t_6 + a_7t_7 + a_8t_8 + a_9t_9),$$

where $a_8, a_9, a_6, a_7 \in \mathbb{F}_q^\times$ and we assume that (a_6, a_7) is not a multiple of (a_8, a_9) . Then we define $\chi_{8,9,q^3}^{a_8, a_9, a_6, a_7}$ to be the character we obtain by inducing $\lambda^{a_8, a_9, a_6, a_7}$ to U/M_{10} .

From the calculations of the character values in Proposition 5.2 below, we observe that $x\lambda^{a_8,a_9,a_6,a_7} \neq \lambda^{a_8,a_9,a_6,a_7}$ for all $x \in X$ with $x \neq 1$. Therefore, an application of Clifford theory implies that each $\chi_{8,9,q^3}^{a_8,a_9,a_6,a_7}$ is irreducible.

We proceed to consider the characters in $\mathcal{F}_{8,9}$ of degree q^2 . Again we have that M_{10} is in the kernel of these characters. We consider the subgroup

$$W = \prod_{i \neq 1,5} X_i$$

of U/M_{10} . We note that $M = X_6X_7$ is a normal subgroup of W , that W/M is isomorphic to \mathbb{F}_q^5 and that $X = X_1X_5$ is a transversal of W in U/M_{10} . Therefore, we have linear characters μ^{a_8,a_9} of W defined by

$$\mu^{a_8,a_9}(x_W(\mathbf{t})) = \phi(a_8t_8 + a_9t_9),$$

where $a_8, a_9 \in \mathbb{F}_q^\times$. Then we define $\chi_{8,9,q^2}^{a_8,a_9}$ to be the character we obtain by inducing μ^{a_8,a_9} to U/M_{10} . The characters $\chi_{8,9,q^2}^{a_8,a_9,b_2,b_3,b_4}$ are given by tensoring $\chi_{8,9,q^2}^{a_8,a_9}$ with $\chi_{\text{lin}}^{0,b_2,b_3,b_4}$.

To see that $\chi_{8,9,q^2}^{a_8,a_9}$ is irreducible we can consider the restriction μ' of μ^{a_8,a_9} to the subgroup $V' = X_2X_6X_8X_9$ of V . Then we let $\chi' = \mu'^{U'}$, where U' is the subgroup $X_1X_2X_5X_6X_8X_9$ of U/M_{10} . Now V' is normal in U' and it is an easy calculation to check that ${}^x\mu' \neq \mu'$ for all $x \in X$ with $x \neq 1$. Now we deduce that χ' is irreducible by an application of Clifford theory. From this it follows that $\chi_{8,9,q^2}^{a_8,a_9}$ is irreducible and thus also $\chi_{8,9,q^2}^{a_8,a_9,b_2,b_3,b_4}$ is irreducible.

We now state our proposition giving the values of the characters in $\mathcal{F}_{8,9}$.

Proposition 5.2. (a)

$$\chi_{8,9,q^3}^{a_8,a_9,a_6,a_7}(x(\mathbf{t})) = q^3\delta_{(t_1,t_2,t_3,t_4,t_5,a_8t_6+a_9t_7),0}\phi(a_6t_6 + a_7t_7 + a_8t_8 + a_9t_9).$$

(b) If $t_3 = 0$, then

$$\chi_{8,9,q^2}^{a_8,a_9,b_2,b_3,b_4}(x(\mathbf{t})) = q^2\delta_{(t_1,t_5,a_8t_2+a_9t_4,a_8t_6+a_9t_7),0}\phi(b_2t_2 + b_4t_4 + a_8t_8 + a_9t_9).$$

If $t_3 \neq 0$, then

$$\begin{aligned} \chi_{8,9,q^2}^{a_8,a_9,b_2,b_3,b_4}(x(\mathbf{t})) = \\ q\delta_{(t_1,a_8t_2+a_9t_4),0}\phi(b_2t_2 + b_3t_3 + b_4t_4 + a_8t_8 + a_9t_9 + t_5(a_8t_2 + a_9t_4 - (a_8t_6 + a_9t_7)/t_3)). \end{aligned}$$

Proof. (a) We write $\lambda = \lambda^{a_8,a_9,a_6,a_7}$ and $\chi = \chi_{8,9,q^2}^{a_8,a_9,a_6,a_7}$. We consider the series of normal subgroups of U/M_{10}

$$V_0 = U/M_{10}, \quad V_1 = \prod_{i \neq 1} X_i, \quad V_2 = \prod_{i \neq 1,2} X_i, \quad V_3 = V.$$

First we calculate the values of λ^{V_2} . We have $Z(V_2) = X_6X_7X_8X_9$, so that

$$\lambda^{V_2}(x_{V_2}(\mathbf{t})) = \lambda^{V_2}(x_3(t_3)x_4(t_4)x_5(t_5))\lambda(x_6(t_6)x_7(t_7)x_8(t_8)x_9(t_9)).$$

Also $\lambda^{V_2}(x_{V_2}(\mathbf{t})) = 0$ if $t_4 \neq 0$. Thus it suffices to determine $\lambda^{V_2}(x_3(t_3)x_5(t_5))$. Now for $s \in \mathbb{F}_q$, we have $(x_3(t_3)x_5(t_5))^{x_4(s)} = x_3(t_3)x_5(t_5)x_7(-st_3)x_9(-st_5)$ in U/M_{10} . Therefore,

$$\lambda^{V_2}(x_3(t_3)x_5(t_5)) = \sum_{s \in \mathbb{F}_q} \lambda(x_3(t_3)x_5(t_5)x_7(-st_3)x_9(-st_5))$$

$$\begin{aligned}
&= \sum_{s \in \mathbb{F}_q} \phi(-a_7 s t_3 - a_9 s t_5) \\
&= \sum_{s \in \mathbb{F}_q} \phi(-s(a_7 t_3 + a_9 t_5)) \\
&= q \delta_{a_7 t_3 + a_9 t_5, 0}.
\end{aligned}$$

and thus

$$\lambda^{V_2}(x_{V_2}(\mathbf{t})) = q \delta_{(t_4, a_7 t_3 + a_9 t_5), 0} \phi(a_6 t_6 + a_7 t_7 + a_8 t_8 + a_9 t_9).$$

Next we induce λ^{V_2} to V_1 ; this requires a very similar calculation, which we omit. We obtain

$$\lambda^{V_1}(x_{V_1}(\mathbf{t})) = q^2 \delta_{(t_2, t_4, a_7 t_3 + a_9 t_5, a_6 t_3 + a_8 t_5), 0} \phi(a_6 t_6 + a_7 t_7 + a_8 t_8 + a_9 t_9).$$

Our assumption that (a_6, a_7) is not a multiple of (a_8, a_9) implies that for the above to be nonzero we require $t_3, t_5 = 0$. Thus we deduce that

$$\lambda^{V_1}(x_{V_1}(\mathbf{t})) = q^2 \delta_{(t_2, t_3, t_4, t_5), 0} \phi(a_6 t_6 + a_7 t_7 + a_8 t_8 + a_9 t_9).$$

Now we do the final induction up to U/M_{10} to obtain χ . First we observe that $Z(V_0) = X_8 X_9$ and it is easy to check that $\chi(x(\mathbf{t})) = 0$ if any of t_1, t_2, t_3, t_4 or t_5 is equal to 0. Thus it suffices to calculate $\chi(x_6(t_6)x_7(t_7))$, and we obtain

$$\begin{aligned}
\chi(x_6(t_6)x_7(t_7)) &= \sum_{s \in \mathbb{F}_q} \lambda^{V_1}(x_6(t_6)x_7(t_7)x_8(-st_6)x_9(-st_7)) \\
&= q^2 \sum_{s \in \mathbb{F}_q} \phi(a_6 t_7 + a_7 t_7 - a_8 s t_6 - a_9 s t_7) \\
&= q^3 \delta_{0, a_8 t_6 + a_9 t_7} \phi(a_6 t_6 + a_7 t_7).
\end{aligned}$$

Putting this all together we obtain the stated value of $\chi_{8,9,q^3}^{a_8, a_9, a_6, a_7}$.

(b) We write $\mu = \mu^{a_8, a_9}$ and $\chi = \chi_{8,9,q^2}^{a_8, a_9}$. In order to determine the value of χ we first note that $Z(U/M_{10}) = X_8 X_9$ and that we have $\chi(x(\mathbf{t})) = 0$ if $t_1 \neq 0$. Thus it suffices to calculate $\chi(x_3(t_3)x_2(t_2)x_4(t_4)x_5(t_5)x_6(t_6)x_7(t_7))$. Next we calculate that

$$\begin{aligned}
&(x_3(t_3)x_2(t_2)x_4(t_4)x_5(t_5)x_6(t_6)x_7(t_7))^{x_5(s_5)x_1(s_1)} = \\
&x_3(t_3)x_2(t_2)x_4(t_4)x_5(t_5 - s_1 t_3)x_6(t_6)x_7(t_7)x_8(-s_1 t_6 + s_5 t_2 + s_1 t_2 t_3)x_9(-s_1 t_7 + s_5 t_4 + s_1 t_3 t_4).
\end{aligned}$$

Thus, $\chi(x_3(t_3)x_2(t_2)x_4(t_4)x_5(t_5)x_6(t_6)x_7(t_7))$ is equal to

$$\begin{aligned}
&\sum_{s_1, s_5 \in \mathbb{F}_q} \mu(x_3(t_3)x_2(t_2)x_4(t_4)x_5(t_5)x_6(t_6)x_7(t_7))^{x_5(s_5)x_1(s_1)}) \\
&= \sum_{s_1, s_5 \in \mathbb{F}_q} \delta_{t_5 - s_1 t_3, 0} \phi(a_8(-s_1 t_6 + s_5 t_2 + s_1 t_2 t_3) + a_9(-s_1 t_7 + s_5 t_4 + s_1 t_3 t_4)).
\end{aligned}$$

If $t_3 = 0$, then we see that this is equal to

$$\delta_{t_5, 0} \sum_{s_1, s_5 \in \mathbb{F}_q} \phi(a_8(-s_1 t_6 + s_5 t_2) + a_9(-s_1 t_7 + s_5 t_4)) = q^2 \delta_{(t_5, a_8 t_2 + a_9 t_4, a_8 t_6 + a_9 t_7), 0}.$$

Whereas for $t_3 \neq 0$, in the sum over s_1 we only get a nonzero contribution for $s_1 = t_5/t_3$, thus we get

$$\sum_{s_5 \in \mathbb{F}_q} \phi(a_8(-t_5 t_6/t_3 + s_5 t_2 + t_2 t_5) + a_9(-t_5 t_7/t_3 + s_5 t_4 + t_4 t_5)) =$$

$$q \delta_{a_8 t_2 + a_9 t_4, 0} \phi(t_5(a_8 t_2 + a_9 t_4 - (a_8 t_6 + a_9 t_7)/t_3)).$$

From the formulas above we deduce the stated values of $\chi_{8,9,q^2}^{a_8,a_9,b_2,b_3,b_4}$. \square

We note the characters $\chi_{8,9,q^3}^{a_8,a_9,a_6,a_7}$ can be equal for different values of a_i . From the character values we observe that $\chi_{8,9,q^3}^{a_8,a_9,a_6,a_7} = \chi_{8,9,q^3}^{a'_8,a'_9,a'_6,a'_7}$ if and only if (a'_8, a'_9, a'_6, a'_7) is obtained from (a_8, a_9, a_6, a_7) by adding a multiple of (a_8, a_9) to (a_6, a_7) . So we choose (a_6, a_7) such that $a_6 a_9 + a_7 a_8 = 0$, except when $p = 2$ where we just put $a_6 = 0$.

Also we observe that $\chi_{8,9,q^3}^{a_8,a_9,b_2,b_3,b_4} = \chi_{8,9,q^3}^{a'_8,a'_9,b'_2,b'_3,b'_4}$ if and only if $(a'_8, a'_9, b'_2, b'_3, b'_4)$ is obtained from $(a_8, a_9, b_2, b_3, b_4)$ by adding a multiple of (a_8, a_9) to (b_2, b_4) . So we choose a coset representative such that $b_2 a_9 + b_4 a_8 = 0$ (except in the case $p = 2$ where we just take $b_2 = 0$).

We move on to briefly discuss the characters in the family \mathcal{F}_8 . These can be defined in essentially the same way as those in $\mathcal{F}_{8,9}$. The characters of degree q^3 are denoted $\chi_{8,q^3}^{a_8,a_7}$ and are defined in exactly the same way as $\chi_{8,9,q^3}^{a_8,a_9,a_6,a_7}$ except we set $a_9 = 0$ and $a_6 = 0$. The characters $\chi_{8,q^2}^{a_8,b_3,b_4}$ are defined in the same way as $\chi_{8,9,q^2}^{a_8,a_9,b_2,b_3,b_4}$ except we set $a_9 = 0$ and $b_2 = 0$. Thus we can immediately deduce the values of these characters given in the proposition below.

Proposition 5.3. (a)

$$\chi_{8,q^3}^{a_8,a_7}(x(\mathbf{t})) = q^3 \delta_{(t_1,t_2,t_3,t_4,t_5,t_6),0} \phi(a_7 t_7 + a_8 t_8).$$

(b) If $t_3 = 0$, then

$$\chi_{8,q^2}^{a_8,b_3,b_4}(x(\mathbf{t})) = q^2 \delta_{(t_1,t_2,t_5,t_6),0} \phi(b_4 t_4 + a_8 t_8).$$

If $t_3 \neq 0$, then

$$\chi_{8,q^2}^{a_8,b_3,b_4}(x(\mathbf{t})) = q \delta_{(t_1,t_2),0} \phi(b_3 t_3 + b_4 t_4 + a_8(t_8 - t_5 t_6/t_3)).$$

5.4. The Family $\mathcal{F}_{8,9,10}$. We begin with the assumption that $p > 2$, as the situation for $p = 2$ is slightly different and is discussed at the end of the section.

First we note that M_{11} lies in the kernel of the characters in $\mathcal{F}_{8,9,10}$. We have that $U/M_{11} = \prod_{i=1}^{10} X_i$ and we define the normal subgroup

$$V = \prod_{i \neq 1,2,4} X_i$$

of U/M_{11} . We have that V is isomorphic to \mathbb{F}_q^7 and that $X = X_1 X_2 X_4$ is a transversal of V in U/M_{11} . We define the character $\lambda^{a_8,a_9,a_{10}}$ of V by

$$\lambda^{a_8,a_9,a_{10}}(x_V(\mathbf{t})) = \phi(a_8 t_8 + a_9 t_9 + a_{10} t_{10})$$

for $a_8, a_9, a_{10} \in \mathbb{F}_q^\times$. Then we define $\chi_{8,9,10}^{a_8,a_9,a_{10}}$ to be the character obtained by inducing $\lambda^{a_8,a_9,a_{10}}$ to U/M_{11} . The irreducible characters $\chi_{8,9,10}^{a_8,a_9,a_{10},b_3}$ in the family $\mathcal{F}_{8,9,10}$ are then obtained by tensoring $\chi_{8,9,10}^{a_8,a_9,a_{10}}$ with the linear character $\chi_{\text{lin}}^{0,0,b_3,0}$. The construction of these characters is the same as in [24, §4], where they are shown to be irreducible and distinct.

In order to obtain the values of χ we take advantage of the conjugacy class representatives in Table 3. First note that since $\chi(x(\mathbf{t})) = 0$ for $x(\mathbf{t}) \notin V$, so we do not need to consider

any conjugacy classes in Table 3 above the family \mathcal{C}_3 . We split up our calculations for the remaining elements by considering elements in the family \mathcal{C}_3 , and those lying in any of the other classes.

Proposition 5.4. *Suppose $p > 2$.*

- (a) *Suppose that at least one of t_1, t_2 or t_4 is nonzero. Then $\chi_{8,9,10}^{a_8,a_9,a_{10},b_3}(x(\mathbf{t})) = 0$.*
- (b) *Let $x(\mathbf{t}) = x_3(t_3)x_8(t_8)x_9(t_9)x_{10}(t_{10})$, where $t_3 \neq 0$. Then*

$$\chi_{8,9,10}^{a_8,a_9,a_{10},b_3}(x(\mathbf{t})) = q\phi(b_3t_3 + a_8t_8 + a_9t_9 + a_{10}t_{10}) \sum_{s \in \mathbb{F}_q} \phi(-a_8a_9a_{10}t_3s^2).$$

- (c) *Let $x(\mathbf{t}) = x_5(t_5)x_6(t_6)x_7(t_7)x_8(t_8)x_9(t_9)x_{10}(t_{10})$. Then*

$$\chi_{8,9,10}^{a_8,a_9,a_{10},b_3}(x(\mathbf{t})) = q^3\delta_{(t_5,t_6,t_7),0}\phi(a_8t_8 + a_9t_9 + a_{10}t_{10}).$$

Remark 5.5. Before we proceed to the proof, we remark on the term $\sum_{t \in \mathbb{F}_q} \phi(-a_8a_9a_{10}t_3t^2)$ in (b) (for the case $t_3 \neq 0$). Clearly this can be deduced from the value of $\sum_{t \in \mathbb{F}_q} \phi(t^2)$: if $-a_8a_9a_{10}t_3 \in \mathbb{F}_q^\times$ is a square, then $\sum_{t \in \mathbb{F}_q} \phi(-a_8a_9a_{10}t_3t^2) = \sum_{t \in \mathbb{F}_q} \phi(t^2)$ and if $-a_8a_9a_{10}t_3 \in \mathbb{F}_q^\times$ is not a square, then $\sum_{t \in \mathbb{F}_q} \phi(-a_8a_9a_{10}t_3t^2) = -\sum_{t \in \mathbb{F}_q} \phi(t^2)$. The sum $\sum_{t \in \mathbb{F}_q} \phi(t^2)$ is referred to as a *quadratic Gauss sum*. For $q = p$ the value of this Gauss sum is well known: if $p \equiv 1 \pmod{4}$, then $\sum_{t \in \mathbb{F}_p} \phi(t^2) = \sqrt{p}$, and if $p \equiv -1 \pmod{4}$, then $\sum_{t \in \mathbb{F}_p} \phi(t^2) = i\sqrt{p}$. However, for general q the situation is more complicated: it is known that the absolute value is \sqrt{q} , see for example [29, Proposition 11.5], but a formula giving the exact value appears not to be known in general.

Proof. We begin by noting that it suffices to consider the case $b_3 = 0$. We write $\lambda = \lambda^{a_8,a_9,a_{10}}$ and $\chi = \chi_{8,9,10}^{a_8,a_9,a_{10}}$.

- (a) We dealt with this case in the discussion before the statement of the proposition.
- (b) The first step is to calculate in U/M_{11} that

$$x_3(t_3)^{x_1(s_1)x_2(s_2)x_4(s_4)} = x_3(t_3)x_5(-s_1t_3)x_6(-s_2t_3)x_7(-s_4t_3)x_8(s_1s_2t_3)x_9(s_1s_4t_3)x_{10}(s_2s_4t_3).$$

Therefore, we have

$$\begin{aligned} \chi(x_3(t_3)) &= \sum_{s_1, s_2, s_4 \in \mathbb{F}_q} \phi(a_8s_1s_2t_3 + a_9s_1s_4t_3 + a_{10}s_2s_4t_3) \\ &= \sum_{s_2, s_4 \in \mathbb{F}_q} \phi(a_{10}s_2s_4t_3) \sum_{s_1 \in \mathbb{F}_q} \phi(s_1(a_8s_2 + a_9s_4)t_3) \\ &= \sum_{s_2, s_4 \in \mathbb{F}_q} \phi(a_{10}t_3s_2s_4)q\delta_{0,a_8s_2+a_9s_4} \\ &= q \sum_{s_4 \in \mathbb{F}_q} \phi(-a_{10}a_9s_4^2t_3/a_8) \\ &= q \sum_{s \in \mathbb{F}_q} \phi(-a_8a_9a_{10}t_3s^2). \end{aligned}$$

For the last equality we just substitute $s_4 = a_8s$. Now using that $Z(U/M_{11}) = X_8X_9X_{10}$, the value of χ given in (b) can be deduced.

- (c) In this case the starting point is to calculate in U/M_{11} that

$$(x_5(t_5)x_6(t_6)x_7(t_7))^{x_1(s_1)x_2(s_2)x_4(s_4)} = x_5(t_5)x_6(t_6)x_7(t_7)x_8(-s_1t_6 - s_2t_5)x_9(-s_1t_7 - s_4t_5)x_{10}(-s_2t_7 - s_4t_6).$$

Therefore, $\chi(x_5(t_5)x_6(t_6)x_7(t_7))$ is equal to

$$\begin{aligned} & \sum_{s_1 \in \mathbb{F}_q} \phi(-s_1(a_8t_6 + a_9t_7)) \sum_{s_2 \in \mathbb{F}_q} \phi(-s_2(a_8t_5 + a_{10}t_7)) \sum_{s_4 \in \mathbb{F}_q} \phi(-s_4(a_9t_5 + a_{10}t_6)) \\ &= q^3 \delta_{(a_8t_6 + a_9t_7, a_8t_5 + a_{10}t_7, a_9t_5 + a_{10}t_6), 0}. \end{aligned}$$

Now the linear system

$$\begin{aligned} a_8t_6 &+ a_9t_7 &= 0 \\ a_8t_5 &+ a_{10}t_7 &= 0 \\ a_9t_5 &+ a_{10}t_6 &= 0 \end{aligned}$$

has only the trivial solution $t_5 = t_6 = t_7 = 0$, as we are assuming $p > 2$. Thus we get $\chi(x_5(t_5)x_6(t_6)x_7(t_7)) = q^3 \delta_{(t_5, t_6, t_7), 0}$. Now using that $Z(U/M_{11}) = X_8X_9X_{10}$, the value of $\chi_{8,9,10}^{a_8, a_9, a_{10}, b_3}$ given in (c) can be deduced. \square

To end this section we consider the case $p = 2$, where the situation is more complicated. The characters $\chi_{8,9,10,q^3}^{a_8, a_9, a_{10}}$ are defined in the same way as the characters $\chi_{8,9,10}^{a_8, a_9, a_{10}}$ for $p > 3$. There are also characters denoted $\chi_{8,9,10,q^3/2}^{a_8, a_9, a_{10}, a_5, 6, 7, d_1, 2, 4, d_3}$, which are defined as follows.

Again we let $V = \prod_{i \neq 1, 2, 4} X_i$ and let $a = a_5, 6, 7, a_8, a_9, a_{10} \in \mathbb{F}_q^\times$. We define the linear character $\lambda = \lambda^{a_8, a_9, a_{10}, a}$ of V by

$$\lambda(x_V(\mathbf{t})) = \phi(a_8a_9at_5 + a_8a_{10}at_6 + a_9a_{10}at_7 + a_8t_8 + a_9t_9 + a_{10}t_{10}).$$

Now let $Y_{124} = \{1, x_1(a_{10}a)x_2(a_9a)x_4(a_8a)\}$ and $\bar{V} = Y_{124}V$. We can extend λ to \bar{V} in two ways, namely $\bar{\lambda}^{d_1, 2, 4}$, where $\bar{\lambda}^{d_1, 2, 4}(x_1(a_{10}a)x_2(a_9a)x_4(a_8a)) = (-1)^{d_1, 2, 4}$ for $d_1, 2, 4 \in \mathbb{F}_2$. We then induce $\bar{\lambda}^{d_1, 2, 4}$ to U/M_{11} to obtain a character $\psi^{d_1, 2, 4}$, which is irreducible as shown in [24, §4]. Now let d_3 be either 0 or an element of \mathbb{F}_q , which is not in the image of the map $s \mapsto aa_8a_9a_{10}s + a_8a_9a_{10}s^2$. Then we define $\chi_{8,9,10,q^3/2}^{a_8, a_9, a_{10}, a_5, 6, 7, d_1, 2, 4, d_3}$ by tensoring $\psi^{d_1, 2, 4}$ with $\chi_{\text{lin}}^{0, 0, d_3, 0}$.

The proposition below states the values of these characters, these can be deduced from [30, Thm. 2.3]. Note that with our specific choice of ϕ , we have $\ker(\phi) = \{t^p - t : t \in \mathbb{F}_q\}$. Thus for each $a \in \mathbb{F}_q^\times$, we have that $\ker(\phi) = a^{-p}\mathbb{T}_a$, where $\mathbb{T}_a = \{t^p - a^{p-1}t \mid t \in \mathbb{F}_q\}$ as defined in [30, Definition 1.2]. In turn this implies that a_ϕ as defined in [30, Definition 1.4] is equal to a^{-p} . We note that the values for $\chi_{8,9,10,q^3}^{a_8, a_9, a_{10}}$ look different to those for $p > 2$, which is explained by the fact that the quadratic Gauss sums for $p = 2$ are clearly zero.

Proposition 5.6. *Let $p = 2$.*

(a)

$$\chi_{8,9,10,q^3}^{a_8, a_9, a_{10}}(x(\mathbf{t})) = \delta_{(t_1, t_2, t_3, t_4, a_8t_5 + a_{10}t_7, a_8t_6 + a_9t_7), 0} q^3 \phi(a_8t_8 + a_9t_9 + a_{10}t_{10}).$$

(b) *Let $a = a_5, 6, 7$. If $(t_1, t_2, t_4) \notin \{(0, 0, 0), (a_{10}a, a_9a, a_8a)\}$, then*

$$\chi_{8,9,10,q^3/2}^{a_8, a_9, a_{10}, a_5, 6, 7, d_1, 2, 4, d_3}(x(\mathbf{t})) = 0.$$

If $x(\mathbf{t}) = x_1(a_{10}r)x_2(a_9r)x_4(a_8r)x_5(a_{10}t)x_6(a_9t)x_7(a_8t)x_8(t_8)x_9(t_9)x_{10}(t_{10})$, where $r \in \{0, a\}$ and $t, t_8, t_9, t_{10} \in \mathbb{F}_q$, then

$$\chi_{8,9,10,q^3/2}^{a_8,a_9,a_{10},a_5,6,7,d_1,2,4,d_3}(x(\mathbf{t})) = \frac{q^3}{2}\phi(d_{1,2,4}r + a_8a_9a_{10}at + a_8t_8 + a_9t_9 + a_{10}t_{10}).$$

For $r \in \{0, a\}$ and $x(\mathbf{t}) = x_3(t_3)x_1(a_{10}r)x_2(a_9r)x_4(a_8r)x_5(t_5)x_6(t_6)x_7(t_7)x_8(t_8)x_9(t_9)x_{10}(t_{10})$,

$$\begin{aligned} \chi_{8,9,10,q^3/2}^{a_8,a_9,a_{10},a_5,6,7,d_1,2,4,d_3}(x(\mathbf{t})) &= \frac{q^2}{2}\delta_{t_3, \frac{a-p}{a_8a_9a_{10}}}\phi(d_{1,2,4}r + d_3t_3 + a_8a_9a_{10}a\frac{t_7}{a_8} + \\ &\quad \frac{(a_8a_9a_{10})^2}{a^{-p}}(\frac{t_5}{a_{10}} + \frac{t_7}{a_8})(\frac{t_6}{a_9} + \frac{t_7}{a_8}) + a_8t_8 + a_9t_9 + a_{10}t_{10}). \end{aligned}$$

5.5. The Family \mathcal{F}_{11} . First we note that M_{12} lies in the kernel of the characters in \mathcal{F}_{11} . We have that $U/M_{12} = \prod_{i=1}^{11} X_i$ and we define the normal subgroup

$$V = \prod_{i \neq 1,2,4} X_i$$

of U/M_{12} . We have that V is isomorphic to \mathbb{F}_q^8 and $X = X_1X_2X_4$ is a transversal of V in U/M_{12} . We define the character $\lambda^{a_{11},b_5,b_6,b_7}$ of V by

$$\lambda^{a_{11},b_5,b_6,b_7}(x_V(\mathbf{t})) = \phi(b_5t_5 + b_6t_6 + b_7t_7 + a_{11}t_{11})$$

for $a_{11} \in \mathbb{F}_q^\times$ and $b_5, b_6, b_7 \in \mathbb{F}_q$.

We define $\chi_{11}^{a_{11},b_5,b_6,b_7}$ to be the character obtained by inducing $\lambda^{a_{11},b_5,b_6,b_7}$ to U/M_{12} . The irreducible characters $\chi_{11}^{a_{11},b_5,b_6,b_7,b_3}$ in the family \mathcal{F}_{11} are then obtained by tensoring $\chi_{11}^{a_{11},b_5,b_6,b_7}$ with the linear character $\chi_{\text{lin}}^{0,0,b_3,0}$.

Using [30, Lemma 1.5] with $Z = X_{11}$, $Y = X_8X_9X_{10}$, $X = X_1X_2X_4$ and $M = X_3X_5X_6X_7$, we obtain that each $\chi_{11}^{a_{11},b_5,b_6,b_7,b_3}$ is irreducible, and that $\chi_{11}^{a_{11},b_5,b_6,b_7,b_3} = \chi_{11}^{a'_{11},b'_5,b'_6,b'_7,b'_3}$ if and only if $(a_{11}, b_5, b_6, b_7, b_3) = (a'_{11}, b'_5, b'_6, b'_7, b'_3)$. The fact that the hypothesis of that lemma holds follows from the fact that XYZ forms a subgroup of V which is special of type q^{1+6} .

In order to obtain the values of $\chi_{11}^{a_{11},b_5,b_6,b_7,b_3}$ in Proposition 5.7 below, we take advantage of the conjugacy class representatives in Table 3. We assume $p > 2$ at first, as the situation is slightly different for $p = 2$, which we discuss after. First note that since $\chi_{11}^{a_{11},b_5,b_6,b_7,b_3}(x(\mathbf{t})) = 0$ for $x(\mathbf{t}) \notin V$, we can deal with any conjugacy classes in Table 3 above the family \mathcal{C}_3 quickly in (a). Dealing with elements in the family \mathcal{C}_3 in (b) is the most complicated part of the proposition. Part (c) covers all of those in the families $\mathcal{C}_{5,6,7}$, $\mathcal{C}_{5,6}$ and \mathcal{C}_5 , and then we can then use the triality automorphism τ to determine the values on representatives in the classes $\mathcal{C}_{5,7}$, $\mathcal{C}_{6,7}$, \mathcal{C}_6 and \mathcal{C}_7 . We finish by considering the remaining classes together in (d), so that we have determined the value of $\chi_{11}^{a_{11},b_5,b_6,b_7,b_3}$.

Proposition 5.7. *Suppose $p > 2$.*

(a) *Suppose that at least one of t_1 , t_2 or t_4 is nonzero. Then*

$$\chi_{11}^{a_{11},b_5,b_6,b_7,b_3}(x(\mathbf{t})) = 0.$$

(b) *Let $x(\mathbf{t}) = x_3(t_3)x_8(t_8)x_9(t_9)x_{10}(t_{10})x_{11}(t_{11})$, where $t_3 \neq 0$. Then*

$$\chi_{11}^{a_{11},b_5,b_6,b_7,b_3}(x(\mathbf{t})) = q\phi(b_3t_3 + a_{11}t_{11})(q\delta_{(b_6t_3+a_{11}t_9, b_7t_3+a_{11}t_8), 0})$$

$$\sum_{s \in \mathbb{F}_q^\times} \phi(-(b_5 t_3 + a_{11} t_{10})s + (b_6 t_3 + a_{11} t_9)(b_7 t_3 + a_{11} t_8)(a_{11} t_3)^{-1} s^{-1})).$$

(c)

(i) Let $x(\mathbf{t}) = x_5(t_5)x_6(t_6)x_7(t_7)x_{11}(t_{11})$, where $t_5, t_6, t_7 \neq 0$. Then

$$\chi_{11}^{a_{11}, b_5, b_6, b_7, b_3}(x(\mathbf{t})) = q\phi(b_5 t_5 + b_6 t_6 + b_7 t_7 + a_{11} t_{11}) \sum_{s \in \mathbb{F}_q} \phi(-a_{11} t_5 t_6 t_7 s^2).$$

(ii) Let $x(\mathbf{t}) = x_5(t_5)x_6(t_6)x_9(t_9)x_{10}(t_{10})x_{11}(t_{11})$, where $t_5 \neq 0$. Then

$$\chi_{11}^{a_{11}, b_5, b_6, b_7, b_3}(x(\mathbf{t})) = q^2 \delta_{t_6 t_9, t_5 t_{10}} \phi(b_5 t_5 + b_6 t_6 + a_{11} t_{11}).$$

(d) Let $x(\mathbf{t}) = x_8(t_8)x_9(t_9)x_{10}(t_{10})x_{11}(t_{11})$. Then

$$\chi_{11}^{a_{11}, b_5, b_6, b_7, b_3} = q^3 \delta_{(t_8, t_9, t_{10}), 0} \phi(a_{11} t_{11}).$$

Remark 5.8. Before the proof, we comment on the term of the form $\sum_{s \in \mathbb{F}_q} \phi(As + \frac{B}{s})$, where $A, B \in \mathbb{F}_q^\times$ in (b). It seems that a general formula for this is not known.

Proof. We note that it suffices to deal with the case $b_3 = 0$ and we let $\lambda = \lambda^{a_{11}, b_5, b_6, b_7}$ and $\chi = \chi^{a_{11}, b_5, b_6, b_7}$.

(a) We dealt with this case in the discussion before the statement of the proposition.

(b) First we note that $Z(U/M_{12}) = X_{11}$. Therefore, it suffices to calculate the values of $\chi(x_3(t_3)x_8(t_8)x_9(t_9)x_{10}(t_{10}))$. We calculate

$$\begin{aligned} (x_3(t_3)x_8(t_8)x_9(t_9)x_{10}(t_{10}))^{x_1(s_1)x_2(s_2)x_4(s_4)} = \\ x_3(t_3)x_5(-t_3 s_1)x_6(-t_3 s_2)x_7(-t_3 s_4)x_8(t_8 + t_3 s_1 s_2) \\ x_9(t_9 + t_3 s_1 s_4)x_{10}(t_{10} + t_3 s_2 s_4)x_{11}(-t_9 s_2 - t_{10} s_1 - t_8 s_4 - t_3 s_1 s_2 s_4). \end{aligned}$$

Thus we see that $\chi(x_3(t_3)x_8(t_8)x_9(t_9)x_{10}(t_{10}))$ is equal to

$$\begin{aligned} & \sum_{s_1, s_2, s_4 \in \mathbb{F}_q} \phi(-b_5 t_3 s_1 - b_6 t_3 s_2 - b_7 t_3 s_4 + a_{11}(-t_9 s_2 - t_{10} s_1 - t_8 s_4 - t_3 s_1 s_2 s_4)) \\ &= \sum_{s_1, s_2 \in \mathbb{F}_q} \phi(-b_5 t_3 s_1 - b_6 t_3 s_2 + a_{11}(-t_9 s_2 - t_{10} s_1)) \sum_{s_4 \in \mathbb{F}_q} \phi(-(b_7 t_3 + a_{11} t_8 + a_{11} t_3 s_1 s_2) s_4) \\ &= \sum_{s_1, s_2 \in \mathbb{F}_q} \phi(-b_5 t_3 s_1 - b_6 t_3 s_2 + a_{11}(-t_9 s_2 - t_{10} s_1)) q \delta_{b_7 t_3 + a_{11} t_8 + a_{11} t_3 s_1 s_2, 0} \\ &= q \left(\sum_{s_2 \in \mathbb{F}_q} \phi((-b_6 t_3 - a_{11} t_9) s_2) \delta_{b_7 t_3 + a_{11} t_8, 0} \right. \\ & \quad \left. + \sum_{s_1 \in \mathbb{F}_q^\times} \phi(-b_5 t_3 s_1 - a_{11} t_{10} s_1) \sum_{s_2 \in \mathbb{F}_q} \phi((-b_6 t_3 - a_{11} t_9) s_2) \delta_{b_7 t_3 + a_{11} t_8 + a_{11} t_3 s_1 s_2, 0} \right) \\ &= q \left(q \delta_{(b_7 t_3 + a_{11} t_8, b_6 t_3 + a_{11} t_9), 0} \right. \end{aligned}$$

$$\begin{aligned}
& + \sum_{s_1 \in \mathbb{F}_q^\times} \phi(-(b_5 t_3 + a_{11} t_{10}) s_1) \phi((b_6 t_3 + a_{11} t_9)(b_7 t_3 + a_{11} t_8)(a_{11} t_3)^{-1} s_1^{-1}) \Big) \\
& = q(q\delta_{(b_7 t_3 + a_{11} t_8, b_6 t_3 + a_{11} t_9), 0} \\
& + \sum_{s \in \mathbb{F}_q^\times} \phi(-(b_5 t_3 + a_{11} t_{10}) s + (b_6 t_3 + a_{11} t_9)(b_7 t_3 + a_{11} t_8)(a_{11} t_3)^{-1} s^{-1})).
\end{aligned}$$

This gives the formula in (b).

(c) We move on to consider the case where $t_3 = 0$ and $t_5 \neq 0$. Consider the normal subgroup $W = \prod_{i \neq 1} X_i$ of U/M_{12} . A calculation shows that

$$\begin{aligned}
\lambda^W(x_5(t_5)x_6(t_6)x_7(t_7)x_8(t_8)x_9(t_9)x_{10}(t_{10})x_{11}(t_{11})) = \\
q\phi(b_5 t_5 + b_6 t_6 + b_7 t_7 + a_{11}(-t_5^{-1} t_8 t_9 + t_{11})).
\end{aligned}$$

To obtain $\chi(x_5(t_5)x_6(t_6)x_7(t_7)x_8(t_8)x_9(t_9)x_{10}(t_{10})x_{11}(t_{11}))$ we just have to induce this formula over X_1 to U/M_{12} . First we consider elements of the conjugacy class $\mathcal{C}_{5,6,7}$, and we get

$$\begin{aligned}
& \chi(x_5(t_5)x_6(t_6)x_7(t_7)x_{11}(t_{11})) \\
& = q\phi(b_5 t_5 + b_6 t_6 + b_7 t_7 + a_{11} t_{11}) \sum_{s_1 \in \mathbb{F}_q} \phi(-a_{11} t_5^{-1} (-s_1 t_6) (-s_1 t_7)) \\
& = q\phi(b_5 t_5 + b_6 t_6 + b_7 t_7 + a_{11} t_{11}) \sum_{s \in \mathbb{F}_q} \phi(-a_{11} t_5 t_6 t_7 s^2)
\end{aligned}$$

as stated in (i).

Next we consider elements in the class $\mathcal{C}_{5,6}$ or \mathcal{C}_5 and we get

$$\begin{aligned}
& \chi(x_5(t_5)x_6(t_6)x_9(t_9)x_{10}(t_{10})x_{11}(t_{11})) \\
& = q\phi(b_5 t_5 + b_6 t_6 + a_{11} t_{11}) \sum_{s_1 \in \mathbb{F}_q} \phi(-a_{11} t_5^{-1} ((-s_1 t_6) t_9 - s_1 t_{10})) \\
& = q\phi(b_5 t_5 + b_6 t_6 + b_7 t_7 + a_{11} t_{11}) \sum_{s \in \mathbb{F}_q} \phi(a_{11} t_5^{-1} s_1 (t_6 t_9 + t_{10})) \\
& = q^2 \delta_{t_6 t_9 + t_{10}, 0} \phi(b_5 t_5 + b_6 t_6 + b_7 t_7 + a_{11} t_{11})
\end{aligned}$$

as stated in (ii).

(d) This is an easy calculation, which we omit. \square

In the case $p = 2$, the only difference is that we now have to consider elements of the form $x_5(t_5)x_6(t_6)x_7(t_7)x_{10}(t_{10})x_{11}(t_{11})$ even when $t_5, t_6, t_7 \neq 0$, to cover the family of conjugacy classes $\mathcal{C}_{5,6,7,2q^8}^{p=2}$; the value given below is in a form where we do not have a closed formula. Also the value for the family $\mathcal{C}_{5,6,7,q^8}^{p=2}$ is simplified as the quadratic Gauss sum is equal to 0.

Lemma 5.9. *Let $p = 2$.*

(a) *Let $x(\mathbf{t}) = x_5(t_5)x_6(t_6)x_7(t_7)x_{10}(t_{10})x_{11}(t_{11})$, where $t_5, t_6, t_7, t_{10} \in \mathbb{F}_q^\times$ and $t_{11} \in \mathbb{F}_q$. Then*

$$\chi_{11}^{a_{11}, b_5, b_6, b_7, b_3}(x(\mathbf{t})) = q\phi(b_5 t_5 + b_6 t_6 + b_7 t_7 + a_{11} t_{11}) \sum_{s \in \mathbb{F}_q} \phi(a_{11}(t_5 t_6 t_7 s^2 + t_5 t_{10} s)).$$

(b) Let $x(\mathbf{t}) = x_5(t_5)x_6(t_6)x_7(t_7)$, where $t_5, t_6, t_7 \in \mathbb{F}_q^\times$. Then

$$\chi_{11}^{a_{11}, b_5, b_6, b_7, b_3}(x(\mathbf{t})) = 0.$$

Proof. (a) We can proceed as in the case $p > 2$ and we obtain the formula

$$\begin{aligned} \lambda^W(x_5(t_5)x_6(t_6)x_7(t_7)x_8(t_8)x_9(t_9)x_{10}(t_{10})x_{11}(t_{11})) = \\ q\phi(b_5t_5 + b_6t_6 + b_7t_7 + a_{11}(t_5^{-1}t_8t_9 + t_{11})). \end{aligned}$$

Now we induce over X_1 to obtain

$$\begin{aligned} \chi(x_5(t_5)x_6(t_6)x_7(t_7)x_{10}(t_{10})x_{11}(t_{11})) = \\ \sum_{s_1 \in \mathbb{F}_q} \lambda^W(x_5(t_5)x_6(t_6)x_7(t_7)x_8(s_1t_6)x_9(s_1t_7)x_{10}(t_{10})x_{11}(s_1t_{10} + t_{11})) \\ = \sum_{s_1 \in \mathbb{F}_q} q\phi(b_5t_5 + b_6t_6 + b_7t_7 + a_{11}(t_5^{-1}s_1^2t_6t_7 + s_1t_{10} + t_{11})) \\ = q\phi(b_5t_5 + b_6t_6 + b_7t_7 + a_{11}t_{11}) \sum_{s \in \mathbb{F}_q} \phi(a_{11}(t_5t_6t_7s^2 + t_5t_{10}s)). \end{aligned}$$

(b) This is obtained directly from the calculation for $p > 2$ by noting that the sum there is equal to 0. \square

5.6. The Family \mathcal{F}_{12} . Let $V = \prod_{i \neq 3,5,6,7} X_i$ and $W = X_8X_9X_{10}X_{11}$. Then W is a normal subgroup of V and the quotient V/W is elementary abelian isomorphic to \mathbb{F}_q^4 . Also $X = X_3X_5X_6X_7$ is a transversal of V in U . Given $a_{12} \in \mathbb{F}_q^\times$, we define the linear character $\lambda^{a_{12}} : V \rightarrow \mathbb{C}$ by $\lambda^{a_{12}}(x_V(\mathbf{t})) = \phi(a_{12}t_{12})$. Then we define $\chi_{12}^{a_{12}} = (\lambda^{a_{12}})^U$ and $\chi_{12}^{a_{12}, b_1, b_2, b_4} = \chi_{12}^{a_{12}} \chi_{\text{lin}}^{b_1, b_2, 0, b_4}$.

Using [30, Lemma 1.5] with $Z = X_{12}$, $Y = X_8X_9X_{10}X_{11}$, $X = X_3X_5X_6X_7$ and $M = X_1X_2X_4$, we obtain that each $\chi_{11}^{a_{12}, b_1, b_2, b_4}$ is irreducible, and that $\chi_{11}^{a_{12}, b_1, b_2, b_4} = \chi_{11}^{a'_{12}, b'_1, b'_2, b'_4}$ if and only if $(a_{12}, b_1, b_2, b_4) = (a'_{12}, b'_1, b'_2, b'_4)$. The hypothesis of that lemma holds due to the fact that XYZ forms a subgroup of V which is special of type q^{1+8} .

In the next proposition we state the values of $\chi_{12}^{a_{12}, b_1, b_2, b_4}$ by considering a number of cases, which cover all of the conjugacy class representatives in Table 3. We note that in (c) we only deal with conjugacy classes up to the action of τ . We first restrict to the case $p > 2$, and then afterwards we describe some differences that occur for the case $p = 2$.

Proposition 5.10. *Suppose that $p > 2$.*

(a) *Suppose that $t_3 \neq 0$. Then*

$$\chi_{12}^{a_{12}, b_1, b_2, b_4}(x(\mathbf{t})) = 0.$$

(b) *Suppose $x(\mathbf{t})$ lies in one of the families $\mathcal{C}_{1,2,4,q^6}$, $\mathcal{C}_{1,2,q^6}$, $\mathcal{C}_{1,4,q^6}$, $\mathcal{C}_{2,4,q^6}$, \mathcal{C}_{1,q^6} , \mathcal{C}_{2,q^6} or \mathcal{C}_{4,q^6} . Then*

$$\chi_{12}^{a_{12}, b_1, b_2, b_4}(x(\mathbf{t})) = 0.$$

(c) *The values of $\chi_{12}^{a_{12}, b_1, b_2, b_4}$ on the families of conjugacy classes $\mathcal{C}_{1,2,4,q^7}$, $\mathcal{C}_{1,2,q^7}$, $\mathcal{C}_{1,2,q^8}$, \mathcal{C}_{1,q^7} and \mathcal{C}_{1,q^8} are given as follows.*

$$\begin{aligned} \mathcal{C}_{1,2,4,q^7}: \quad & \chi_{12}^{a_{12},b_1,b_2,b_4}(x_1(t_1)x_2(t_2)x_4(t_4)x_{12}(t_{12})) \\ & = q\phi(b_1t_1 + b_2t_2 + b_4t_4 + a_{12}t_{12}) \sum_{s \in \mathbb{F}_q} \phi(-a_{12}t_1t_2t_4s^2), \text{ for } t_1, t_2, t_4 \in \mathbb{F}_q^\times \text{ and } t_{12} \in \mathbb{F}_q. \end{aligned}$$

$$\mathcal{C}_{1,2,q^7}: \quad \chi_{12}^{a_{12},b_1,b_2,b_4}(x_1(t_1)x_2(t_2)x_9(t_9)x_{10}(t_{10})) = q^2\delta_{0,t_{10}t_1-t_2t_9}\phi(b_1t_1 + b_2t_2),$$

for $t_1, t_2, t_9, t_{10} \in \mathbb{F}_q^\times$.

$$\mathcal{C}_{1,2,q^8}: \quad \chi_{12}^{a_{12},b_1,b_2,b_4}(x_1(t_1)x_2(t_2)x_{12}(t_{12})) = q^2\phi(b_1t_1 + b_2t_2 + a_{12}t_{12}),$$

for $t_1, t_2 \in \mathbb{F}_q^\times$ and $t_{12} \in \mathbb{F}_q$.

$$\mathcal{C}_{1,q^7}: \quad \chi_{12}^{a_{12},b_1,b_2,b_4}(x_1(t_1)x_{10}(t_{10})) = 0, \text{ for } t_1, t_{10} \in \mathbb{F}_q^\times.$$

$$\mathcal{C}_{1,q^8}: \quad \chi_{12}^{a_{12},b_1,b_2,b_4}(x_1(t_1)x_{12}(t_{12})) = q^2\phi(b_1t_1 + a_{12}t_{12}), \text{ for } t_1 \in \mathbb{F}_q^\times \text{ and } t_{12} \in \mathbb{F}_q.$$

(d) Let $x(\mathbf{t}) = \prod_{i \geq 5} x_i(t_i)$. Then

$$\chi_{12}^{a_{12},b_1,b_2,b_4}(x(\mathbf{t})) = q^4\delta_{0,(t_5,t_6,t_7,t_8,t_9,t_{10},t_{11})}\phi(a_{12}t_{12}).$$

Proof. It is clear that it suffices to calculate the values of $\chi_{12}^{a_{12}}$. We write $\lambda = \lambda^{a_{12}}$ and $\chi = \chi_{12}^{a_{12}}$.

In order to explain our calculations, we introduce some intermediate subgroups. We let $V_1 = VX_7$, $V_2 = VX_6X_7$ and $V_3 = VX_5X_6X_7$. To abbreviate notation we write $\lambda_i = \lambda^{V_i}$ for each $i = 1, 2, 3$. By direct calculation we find

$$\lambda_1(x_{V_1}(\mathbf{t})) = q\delta_{(t_7,t_8),0}\phi(a_{12}t_{12}),$$

and

$$(5.1) \quad \lambda_2(x_{V_2}(\mathbf{t})) = \begin{cases} q\delta_{0,(t_6,t_7)}\phi(a_{12}(t_{12} - \frac{t_9t_8}{t_1})) & \text{if } t_1 \neq 0 \\ q^2\delta_{0,(t_6,t_7,t_8,t_9)}\phi(a_{12}t_{12}) & \text{if } t_1 = 0. \end{cases}$$

We do not calculate λ_3 fully here, but do note that

$$(5.2) \quad \lambda_3(x_{V_3}(\mathbf{t})) = \delta_{0,(t_5,t_6,t_7)}\lambda_3(x_{V_3}(\mathbf{t})).$$

(a) It is clear that $\chi(x(\mathbf{t})) = 0$, if $t_3 \neq 0$.

(b) Let $x = x_1(c_1)x_2(c_2)x_4(c_4)x_5(e_5)x_6(e_6)x_7(e_7)$ lie in the family $\mathcal{C}_{1,2,4,q^6}$ and consider $y = x^{x_3(s_3)}$ for $s_3 \in \mathbb{F}_q$. The theory of minimal representatives outlined in Section 3 implies that the coefficient of one of x_5 , x_6 or x_7 in y must be nonzero. Therefore, $\chi(x) = \sum_{s \in \mathbb{F}_q} \dot{\psi}_3(x^{x_3(s)}) = 0$ by (5.2).

Similar arguments deal with the other families in the statement.

(c) First we show that if one of t_1 , t_2 or t_4 is nonzero, then

$$(5.3) \quad \chi\left(\prod_{i \neq 3,5,6,7} x_i(t_i)\right) = \lambda_3\left(\prod_{i \neq 3,5,6,7} x_i(t_i)\right).$$

We just deal with the case $t_1 \neq 0$, as the other cases can be dealt with similarly. Consider $y = \prod_{i \neq 3,5,6,7} x_i(t_i)^{x_3(s_3)}$, for $s_3 \in \mathbb{F}_q$. We calculate the coefficient of x_5 in y to be t_1s_3 . Therefore, by (5.2), the only nonzero term in the right side of

$$\chi\left(\prod_{i \neq 3,5,6,7} x_i(t_i)\right) = \sum_{s_3 \in \mathbb{F}_q} \dot{\lambda}_3\left(\prod_{i \neq 3,5,6,7} x_i(t_i)^{x_3(s_3)}\right)$$

is when $s_3 = 0$.

We now give the calculation for $\mathcal{C}_{1,2,4,q^7}$

$$\begin{aligned}
& \chi(x_1(t_1)x_2(t_2)x_4(t_4)x_{12}(t_{12})) \\
&= \lambda_3(x_1(t_1)x_2(t_2)x_4(t_4)x_{12}(t_{12})) \\
&= \sum_{s_5 \in \mathbb{F}_q} \dot{\lambda}_2((x_1(t_1)x_2(t_2)x_4(t_4)x_{12}(t_{12}))^{x_5(s_5)}) \\
&= \sum_{s_5 \in \mathbb{F}_q} \dot{\lambda}_2(x_1(t_1)x_2(t_2)x_8(t_2s_5)x_4(t_4)x_9(t_4s_5)x_{12}(t_{12})) \\
&= \sum_{s_5 \in \mathbb{F}_q} \dot{\lambda}_2(x_1(t_1)x_2(t_2)x_4(t_4)x_8(t_2s_5)x_9(t_4s_5)x_{11}(-t_2t_4s_5)x_{12}(t_{12})) \\
&= q\phi(a_{12}t_{12}) \sum_{s \in \mathbb{F}_q} \phi(-a_{12}t_1t_2t_4s^2).
\end{aligned}$$

The first equality is given by (5.3), and the last uses (5.1) and substitution $s = t_1s_5$.

We omit the calculations for the other conjugacy classes in (c), as they are very similar.

(d) This is a straightforward calculation, which we omit. \square

For the case $p = 2$, we just have to deal with the classes $\mathcal{C}_{1,2,4,2q^7}^{p=2}$ and $\mathcal{C}_{1,2,4,q^7}^{p=2}$, which we do in the following lemma. The calculations involved are a minor modification to those for the class $\mathcal{C}_{1,2,4}$ for $p > 2$, so we omit the proof.

Lemma 5.11. *Let $p = 2$. The values of on the classes $\mathcal{C}_{1,2,4,2q^7}^{p=2}$ and $\mathcal{C}_{1,2,4,q^7}^{p=2}$ are given as follows.*

$$\begin{aligned}
\mathcal{C}_{1,2,4,2q^7}^{p=2}: \quad & \chi_{12}^{a_{12},b_1,b_2,b_4}(x_1(t_1)x_2(t_2)x_4(t_4)x_{10}(t_{10})x_{12}(t_{12})) \\
&= q\phi(b_1t_1 + b_2t_2 + b_4t_4 + a_{12}t_{12}) \sum_{s \in \mathbb{F}_q} \phi(a_{12}(st_1t_{10} + t_1t_2t_4s^2)), \text{ for } t_1, t_2, t_4, t_{10} \in \mathbb{F}_q^\times \\
&\text{ and } t_{12} \in \mathbb{F}_q. \\
\mathcal{C}_{1,2,4,q^7}^{p=2}: \quad & \chi_{12}^{a_{12},b_1,b_2,b_4}(x_1(t_1)x_2(t_2)x_4(t_4)x_{12}(t_{12})) = 0, \text{ for } t_1, t_2, t_4 \in \mathbb{F}_q^\times \text{ and } t_{12} \in \mathbb{F}_q.
\end{aligned}$$

REFERENCES

- [1] J. An and G. Hiss, *Restricting unipotent characters in finite symplectic groups*, Comm. Algebra **39** (2011), no. 3, 1104–1130.
- [2] C. A. M. André, *The basic character table of the unitriangular group* J. Algebra **241** (2001), no. 1, 437–471.
- [3] C. André and A. Neto, *A supercharacter theory for the Sylow p -subgroups of the finite symplectic and orthogonal groups*, J. Algebra **322** (2009), 1273–1294.
- [4] J. D. Bradley and S. M. Goodwin, *Conjugacy classes in Sylow p -subgroups of finite Chevalley groups in bad characteristic*, Comm. Algebra **42** (2014), no. 8, 3245–3258.
- [5] R. W. Carter, *Simple groups of Lie type*, A Wiley-Interscience publication, London, 1972.
- [6] P. Diaconis, I.M. Isaacs, *Supercharacters and superclasses for algebra groups*, Trans. Amer. Math. Soc. **360** (2008), no. 5, 2359–2392.
- [7] F. Digne and J. Michel, *Representations of finite groups of Lie type*, London Mathematical Society Student Texts **21**, Cambridge University Press, Cambridge, 1991.
- [8] A. Evseev, *Reduction for characters of finite algebra groups*, J. Algebra **325** (2011), 321–351.
- [9] The GAP Group, *GAP – Groups, Algorithms, and Programming, Version 4.4.12* (2008), <http://www.gap-system.org>.

- [10] M. Geck, G. Hiss, F. Lübeck, G. Malle and G. Pfeiffer, *CHEVIE A system for computing and processing generic character tables for finite groups of Lie type, Weyl groups and Hecke algebras*, Appl. Algebra Engrg. Comm. Comput. **7** (1996), 175–210.
- [11] S. M. Goodwin, *On the conjugacy classes in maximal unipotent subgroups of simple algebraic groups*, Transform. Groups **11** (2006), no. 1, 51–76.
- [12] S. M. Goodwin, P. Mosch and G. Röhrle, *Calculating conjugacy classes in Sylow p -subgroups of finite Chevalley groups of rank six and seven*, LMS J. Comput. Math. **17** (2014), no. 1, 109–122.
- [13] ———, *On the coadjoint orbits of maximal unipotent subgroups of reductive groups*, Transform. Groups **21** (2016), no. 2, 399–426.
- [14] S. M. Goodwin and G. Röhrle, *Calculating conjugacy classes in Sylow p -subgroups of finite Chevalley groups*, J. Algebra **321** (2009), 3321–3334.
- [15] R. Gow, M. Marjoram and A. Previtali *On the irreducible characters of a Sylow 2-subgroup of the finite symplectic group in characteristic 2*, J. Algebra **241** (2001), 393–409.
- [16] G. Higman, *Enumerating p -groups. I: Inequalities*, Proc. London Math. Soc. **10** (1960), 24–30.
- [17] F. Himstedt, *Character tables of parabolic subgroups of Steinberg’s triality groups*, J. Algebra **281** (2004), 774–822.
- [18] ———, *On the 2-decomposition numbers of Steinberg’s triality groups ${}^3D_4(q)$, q odd*, J. Algebra **309** (2007), 569–593.
- [19] ———, *Character tables of parabolic subgroups of Steinberg’s triality groups ${}^3D_4(2^n)$* , J. Algebra **316** (2007), 254–283.
- [20] ———, *On the decomposition numbers of the Ree groups ${}^2F_4(q^2)$ in non-defining characteristic*, J. Algebra **325** (2011), 364–403.
- [21] F. Himstedt and S. Huang, *Character table of a Borel subgroup of the Ree groups ${}^2F_4(q^2)$* , LMS J. Comput. Math. **12** (2009), 1–53.
- [22] ———, *Character tables of the maximal parabolic subgroups of the Ree groups ${}^2F_4(q^2)$* , LMS J. Comput. Math. **13** (2010), 90–110.
- [23] ———, *On the decomposition numbers of Steinberg’s triality groups ${}^3D_4(2^n)$ in odd characteristics*, Comm. Algebra **41** (2013), no. 4, 1484–1498.
- [24] F. Himstedt, T. Le and K. Magaard *Characters of the Sylow p -subgroups of the Chevalley groups $D_4(p^n)$* , J. Algebra **332** (2011), 414–427.
- [25] F. Himstedt and F. Noeske, *Decomposition numbers of $SO_7(q)$ and $Sp_6(q)$* , J. Algebra **413** (2014), 15–40.
- [26] ———, *Restricting unipotent characters in special orthogonal groups*, LMS J. Comput. Math. **18** (2015), 456–488.
- [27] I. M. Isaacs, *Characters of groups associated with finite algebras*, J. Algebra, **177** (1995), 708–730.
- [28] ———, *Counting characters of upper triangular groups*, J. Algebra **315** (2007), no. 2, 698–719.
- [29] H. Iwaniec and E. Kowalski, *Analytic number theory*, American Mathematical Society Colloquium Publications **53**. American Mathematical Society, Providence, RI, 2004.
- [30] T. Le and K. Magaard, *On the character degrees of Sylow p -subgroups of Chevalley groups $G(p^f)$ of type E* , Forum Math. **27** (2015), no. 1, 1–55.
- [31] G. Lehrer, *Discrete series and the unipotent subgroup*, Compos. Math., **28** (1974), 9–19.
- [32] E. Marberg, *Combinatorial methods for character enumeration for the unitriangular groups*, J. Algebra **345** (2011), 295–323.
- [33] M. Marjoram, *Irreducible characters of a Sylow p -subgroup of the orthogonal group*, Comm. Algebra **27** (1999), 1171–1195.
- [34] T. Okuyama and K. Waki, *Decomposition numbers of $Sp(4, q)$* , J. Algebra **199** (1998), 544–555.
- [35] I. Pak and A. Soffer, *On Higman’s $k(U_n(\mathbb{F}_q))$ conjecture*, preprint, arxiv:1507.00411.
- [36] A. Previtali, *On a conjecture concerning character degrees of some p -groups*, Arch. Math. **65** (1995) 375–378.
- [37] J. Sangroniz, *Character degrees of the Sylow p -subgroups of classical groups*, Groups St. Andrews 2001 in Oxford. Vol. II, 487–493, London Math. Soc. Lecture Note Ser., 305, Cambridge Univ. Press, Cambridge, 2003.

- [38] ———, *Irreducible characters of large degree of Sylow p -subgroups of classical groups*, J. Algebra **321** (2009), 1480–1496.
- [39] F. Szechtman, *Irreducible characters of Sylow subgroups of symplectic and unitary groups*, J. Algebra **303** (2006), 722–730.
- [40] B. Szegedy, *Characters of the Borel and Sylow subgroups of classical groups*, J. Algebra **267** (2003), 130–136.
- [41] K. Waki, *A note on decomposition numbers of $G_2(2^n)$* , J. Algebra **274** (2004), 602–606.

SCHOOL OF MATHEMATICS, UNIVERSITY OF BIRMINGHAM, BIRMINGHAM, B15 2TT, U.K.

E-mail address: s.m.goodwin@bham.ac.uk

E-mail address: k.magaard@bham.ac.uk

DEPARTMENT OF MATHEMATICS AND APPLIED MATHEMATICS, UNIVERSITY OF PRETORIA, PRETORIA 0002, SOUTH AFRICA

E-mail address: lttung96@yahoo.com