

# A glucagon-like peptide-1 receptor agonist reduces intracranial pressure in a rat model of hydrocephalus

Botfield, Hannah F; Uldall, Maria S; Westgate, Connor S J; Mitchell, James L; Hagen, Snorre M; Gonzalez, Ana Maria; Hodson, David J; Jensen, Rigmor H; Sinclair, Alexandra J

DOI:

[10.1126/scitranslmed.aan0972](https://doi.org/10.1126/scitranslmed.aan0972)

License:

Other (please specify with Rights Statement)

*Document Version*

Peer reviewed version

*Citation for published version (Harvard):*

Botfield, HF, Uldall, MS, Westgate, CSJ, Mitchell, JL, Hagen, SM, Gonzalez, AM, Hodson, DJ, Jensen, RH & Sinclair, AJ 2017, 'A glucagon-like peptide-1 receptor agonist reduces intracranial pressure in a rat model of hydrocephalus', *Science Translational Medicine*, vol. 9, no. 404, ean0972.  
<https://doi.org/10.1126/scitranslmed.aan0972>

[Link to publication on Research at Birmingham portal](#)

## **Publisher Rights Statement:**

This is the author's version of the work. It is posted here by permission of the AAAS for personal use, not for redistribution. The definitive version was published in *Science Translational Medicine*, Volume 9 on 23rd August 2017, DOI: 10.1126/scitranslmed.aan0972

## **General rights**

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

## **Take down policy**

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact [UBIRA@lists.bham.ac.uk](mailto:UBIRA@lists.bham.ac.uk) providing details and we will remove access to the work immediately and investigate.

**A glucagon-like peptide-1 receptor agonist reduces intracranial pressure in a rat model of hydrocephalus**

**Authors:** Hannah F. Botfield<sup>1,2,†</sup>, Maria S. Uldall<sup>3,†</sup>, Connar S.J. Westgate<sup>1,2</sup>, James L. Mitchell<sup>1,2,4</sup>, Snorre M. Hagen<sup>3</sup>, Ana Maria Gonzalez<sup>5</sup> David J. Hodson<sup>1,6</sup>, Rigmor H. Jensen<sup>3</sup>, Alexandra J. Sinclair<sup>1,2,4,\*</sup>

**Affiliations:**

<sup>1</sup>Institute of Metabolism and Systems Research, University of Birmingham, Edgbaston, B15 2TT, UK.

<sup>2</sup>Centre for Endocrinology, Diabetes and Metabolism, Birmingham Health Partners, B15 2TH, United Kingdom

<sup>3</sup>Danish Headache Center, Clinic of Neurology, Rigshospitalet-Glostrup, University of Copenhagen, Nordre Ringvej 69, 2600 Glostrup, Denmark

<sup>4</sup>Department of Neurology, University Hospitals Birmingham NHS Foundation Trust, B15 2TH, United Kingdom

<sup>5</sup>Institute of Inflammation and Ageing, University of Birmingham, Edgbaston, B15 2TT, UK.

<sup>6</sup>COMPARE University of Birmingham and University of Nottingham, Midlands, UK.

\*Corresponding author: Dr Alexandra Sinclair, [a.b.sinclair@bham.ac.uk](mailto:a.b.sinclair@bham.ac.uk)

<sup>†</sup>Joint first authors

**Overline:**

26

27 **One sentence summary:**

28 GLP-1R agonists show promise as a therapeutic agent to lower intracranial pressure in rodents.

29

30 **Abstract**

31 Current therapies for reducing raised intracranial pressure (ICP) under conditions such as  
32 idiopathic intracranial hypertension or hydrocephalus have limited efficacy and tolerability.  
33 Thus, there is a pressing need to identify alternative drugs. Glucagon-like peptide-1 receptor  
34 (GLP-1R) agonists are used to treat diabetes and promote weight loss but have also been shown  
35 to affect fluid homeostasis in the kidney. Here, we investigated whether exendin-4, a GLP-1R  
36 agonist, is able to modulate cerebrospinal fluid (CSF) secretion at the choroid plexus and  
37 subsequently reduce ICP in rats. We used tissue sections and cell cultures to demonstrate  
38 expression of GLP-1R in the choroid plexus and its activation by exendin-4, an effect blocked  
39 by the GLP-1R antagonist exendin 9-39. Acute treatment with exendin-4 reduced  $\text{Na}^+ \text{K}^+$   
40 ATPase activity, a key regulator of CSF secretion, in cell cultures. Finally, we demonstrated  
41 that administration of exendin-4 to female rats with raised ICP (hydrocephalic) resulted in a  
42 GLP-1R-mediated reduction in ICP. These findings suggest that GLP-1R agonists can reduce  
43 ICP in rodents. Repurposing existing GLP-1R agonist drugs may be a useful therapeutic  
44 strategy for treating raised ICP.

45

46

47

48

49 **Introduction**

Elevated intracranial pressure (ICP) is caused by alterations in the volume of either cerebral blood, cerebrospinal fluid (CSF) or brain tissue. CSF volume is tightly regulated and depends on the balance between CSF secretion, which is modulated predominantly by the choroid plexus, and drainage through the arachnoid granulations and lymphatic (1). Reducing CSF volume, by either CSF drainage or decreasing CSF secretion is used therapeutically to lower ICP (2, 3) in conditions characterized by raised ICP such as idiopathic intracranial hypertension and hydrocephalus.

In the choroid plexus, CSF is secreted by the choroid plexus epithelial (CPE) cells, and is driven by net movement of sodium ions ( $\text{Na}^+$ ) from the blood into the cerebral ventricles. This creates an osmotic gradient, which drives water transport into the cerebral ventricles. There are numerous ion channels involved in this process, but the apical  $\text{Na}^+ \text{K}^+$  ATPase that pumps  $\text{Na}^+$  into the ventricles is the most important of these channels and represents the rate limiting step (4, 5). Specific inhibition of the  $\text{Na}^+ \text{K}^+$  ATPase with ouabain, reduces CSF secretion by 70-80% (6). As such, the CPE cells function akin to inverted renal proximal tubule epithelial cells with an analogous mechanism of fluid transport (7, 8).

The incretin glucagon-like peptide-1 (GLP-1), is a gut peptide secreted by the distal small intestine in response to food intake (9). GLP-1 stimulates glucose-dependent insulin secretion and inhibits glucagon release, lowering blood glucose (10). In addition, GLP-1 is synthesized in neurons of the nucleus tractus solitarius, which project to the hypothalamus (11) and promote satiety and weight loss (12-14). GLP-1 signals through the GLP-1 receptor (GLP-1R), a class-B G protein-coupled receptor expressed in selected cell types within the central nervous system including the hypothalamus, hippocampus, olfactory cortex, circumventricular organs, hindbrain and choroid plexus (15-17).

GLP-1 also has effects on renal proximal tubule  $\text{Na}^+$  secretion, reducing  $\text{Na}^+$  reabsorption and increasing diuresis (18). Here, GLP-1R activation stimulates the conversion

of adenosine triphosphate (ATP) to cyclic adenosine monophosphate (cAMP) by adenylate cyclase. cAMP activates protein kinase A (PKA), which inhibits the  $\text{Na}^+ \text{H}^+$  exchanger, thereby preventing  $\text{Na}^+$  reabsorption into the bloodstream (18). The diuretic actions of incretins have led to investigation of their use as antihypertensive agents (19). Similar to its activity in the kidney, we hypothesize that GLP-1 also modulates  $\text{Na}^+$  transport and subsequently fluid movement at the choroid plexus. We propose that GLP-1R activation may inhibit the basal  $\text{Na}^+ \text{H}^+$  exchanger through cAMP-dependent PKA activation, thus impeding the  $\text{Na}^+ \text{K}^+$  ATPase-dependent secretion of CSF. Stabilized GLP-1 mimetics are widely used to treat diabetes and obesity, and therefore could be repurposed for treating raised ICP.

In the present study, we used tissue sections and CPe cell cultures to assess the localization and distribution of the GLP-1R in rat and human choroid plexus and determined the effects of GLP-1R stimulation on CSF secretion. Furthermore, we conducted in vivo studies to evaluate the effects of GLP-1R agonists on ICP in a hydrocephalus rat model with raised ICP.

## Results

### *GLP-1R expression in human choroid plexus tissue*

Immunohistochemical analysis using haematoxylin and eosin staining confirmed that human donor tissue comprised the choroid plexus, demonstrating choroid plexus morphology including a cuboidal CPe cell monolayer resting on a basement membrane, the underlying interstitial tissue and capillary vessels (**Fig. 1A**). *GLP-1R* mRNA expression in five human choroid plexus samples was compared to known commercially available GLP-1R-positive tissues (pooled samples; see methods for source details). Human pancreas had the highest expression of *GLP-1R* mRNA, with heart and ovary having the least. Human choroid plexus

showed *GLP-1R* mRNA expression (**Fig. 1B**). To determine the localization of the receptor protein, paraffin embedded human choroid plexus sections were immunostained with a specific monoclonal antibody to human GLP-1R previously validated in human and monkey tissue (20, 21). Based on the morphology of the choroid plexus, GLP-1R positive staining was detected in CPe cells (**Fig. 1C-F**). Together, these studies demonstrate that the human choroid plexus expresses GLP-1R mRNA and protein.

#### *Exendin-4 treatment of whole rat choroid plexus in vitro*

Given the lack of validated antibodies against rodent GLP-1R, we instead incubated whole rat choroid plexus in vitro with a fluorescently tagged GLP-1R agonist, exendin-4 (FLEX), to demonstrate the presence of the receptor in the choroid plexus. After 15 minutes of 1 $\mu$ M FLEX incubation, only a few CPe cells were positive for FLEX within the cytoplasm (**Fig. 2A**). However this increased by 30 minutes (**Fig. 2A**). In both cases, GLP-1R appeared to localize predominantly in the cytoplasm, consistent with agonist-induced receptor internalization and trafficking, most likely via endosomes (22). The GLP-1R antagonist exendin 9-39 (1 $\mu$ M) reduced the number of FLEX-positive cells within the choroid plexus (**Fig. 2A**), suggesting specific binding of the FLEX ligand to GLP-1R.

Next, we determined *Glp-1r* mRNA expression in whole rat choroid plexus tissue after incubation with 100nM exendin-4. Incubation of the rat choroid plexus with exendin-4 for 3 hours showed an increase in *Glp-1r* mRNA compared to artificial CSF ( $3.21 \pm 0.70$  fold,  $P < 0.01$ ), with a return to baseline at 6 hours ( $0.78 \pm 0.12$  fold) (**Fig 2B**). There was also a small but detectable increase in *Na<sup>+</sup> K<sup>+</sup> atpase* mRNA expression after 3 hours of exendin-4 treatment compared to incubation with artificial aCSF ( $1.82 \pm 0.28$  fold;  $P < 0.05$ ), which again returned to baseline at 6 hours ( $0.97 \pm 0.21$  fold) (**Fig. 2C**). The expression of other channels

and transporters involved in CSF secretion, including the water channel aquaporin 1 (*Aqp1*) and the  $\text{Na}^+ \text{H}^+$  exchanger (*Nhe1*), were not altered after exendin-4 treatment (**Fig. 2D, E**).

#### *Exendin-4 treatment increases cAMP and reduces $\text{Na}^+ \text{K}^+$ ATPase activity*

To explore further the effects of exendin-4 on the choroid plexus, monolayers of rat CPe cells were grown in culture. These CPe cells were characterized using antibodies against specific identity markers and were shown to be similar to their in vivo counterparts (**Fig. S1A**), including the expression of *Glp-1r* mRNA (**Fig. S1B**). To determine the effect of exendin-4 on GLP-1R signaling, cAMP generation was assessed using two enzyme immunoassay systems. Treatment of CPe cells with exendin-4 increased cAMP compared to control ( $2.14 \pm 0.61$  fold,  $P < 0.01$ ) (**Fig. 3A**) in a concentration-dependent manner, and this could be inhibited by exendin 9-39 (**Fig. 3B**). Forskolin, an adenylate cyclase activator, was used as a positive control to maximally stimulate cAMP production (**Fig 3A-B**) ( $5.30 \pm 0.74$  fold compared to control).

The role of GLP-1R signaling in CSF secretion was assessed in rat CPe cell cultures by measuring  $\text{Na}^+ \text{K}^+$  ATPase activity (proposed as a marker of CSF secretion from the choroid plexus) (6). Exendin-4 treatment reduced  $\text{Na}^+ \text{K}^+$  ATPase specific phosphate production compared to control ( $39.3 \pm 9.4\%$ ,  $P < 0.05$ ) (**Fig. 3C**). In addition, inhibition of PKA with PKI-16-22-amide (PKI) abolished the exendin-4-induced reduction in  $\text{Na}^+ \text{K}^+$  ATPase activity ( $95.4 \pm 17.6\%$ ,  $P < 0.05$ ) (**Fig. 3C**).

#### *Exendin-4 treatment reduces ICP in conscious rats*

To establish whether exendin-4 was able to modulate ICP in vivo, healthy female adult rats were implanted with an ICP monitor (Day 0) before receiving daily subcutaneous (SC) injections of either saline or 20  $\mu\text{g/kg}$  exendin-4 for 5 days (day 2-6). ICP was measured before and after the SC injection on days 2, 4 and 6 (**Fig. 4A**). Examples of the ICP traces are shown

in **Fig. 4B**. On the first day of treatment (day 2), exendin-4 significantly reduced ICP 10 minutes after the SC injection; by 30 minutes ICP was  $65.2 \pm 6.6\%$  of baseline compared to  $91.0 \pm 3.9\%$  of baseline in saline-treated rats ( $P < 0.01$ ) (**Fig. 4C**). A similar drop in ICP was observed on day 4 ( $50.4 \pm 6.9\%$  of baseline;  $P < 0.001$ ) and day 6 ( $54.5 \pm 8.2\%$  of baseline;  $P < 0.001$ ), 30 minutes after exendin-4 administration (**Fig. 4D-E**).

In addition to reducing ICP immediately after treatment, exendin-4 had a cumulative effect on reducing ICP. Exendin-4 caused a significant reduction in ICP measured pre-dose on day 2 (baseline, 100%) to day 4 ( $79.3 \pm 7.3\%$ ;  $P < 0.05$ ) and day 6 ( $72.5 \pm 5.6\%$ ;  $P < 0.01$ ) (**Fig. 4F**), which was not observed in saline-treated rats (day 2, baseline 100%; day 4,  $95.5 \pm 13.6\%$ ; day 6,  $105.3 \pm 12.5\%$ ; **Fig. 4G**).

As there is evidence that weight loss can alter ICP (23), weights were monitored over the treatment period. Whilst both saline- and exendin-4-treated rats lost weight during treatment ( $P < 0.05$ ), there was no significant difference between the groups at any time point (**Fig. 4H**). In the saline group, weight change correlated with alterations in ICP ( $r = 0.710$ ,  $P = 0.032$ ), although no relationship was detected for the exendin-4 treatment group ( $r = -0.300$ ,  $P = 0.552$ ) (**Fig. 4I**).

The effect of SC administration of 20  $\mu\text{g/kg}$  exendin-4 on blood and CSF pH and CSF electrolytes was analyzed 60 minutes post-treatment. Exendin-4 maintained normal blood pH ( $7.35 \pm 0.01$ ; **Fig. 4J**), however, it caused a reduction in CSF pH ( $7.41 \pm 0.03$ ;  $P < 0.05$ ) compared to saline (blood pH  $7.35 \pm 0.03$ , CSF pH  $7.61 \pm 0.07$ ) (**Fig. 4K**). CSF  $\text{Na}^+$  concentration remained unaltered (saline,  $150.3 \pm 0.9$ ; exendin-4,  $150.3 \pm 0.6$ ) (**Fig. 4L**), whereas the concentration of  $\text{Cl}^-$  ions in the CSF was reduced in the exendin-4 group ( $117 \pm 0.5$  mmol/L;  $P < 0.05$ ) compared to the saline group ( $123.8 \pm 0.9$  mmol/L) (**Fig. 4M**). Exendin-4 treatment also increased the concentration of  $\text{Ca}^{2+}$  ions in the CSF ( $1.09 \pm 0.01$ ,  $P > 0.05$ ) compared to saline ( $1.03 \pm 0.02$ ) (**Fig. 4N**).



*Exendin-4 acts via GLP-1R in the brain to reduce ICP in rats*

To assess whether the reduction in ICP was specific to the brain, exendin-4 was injected into the lateral ventricle through an intracerebroventricular (ICV) cannula in anesthetized rats. ICV delivery of exendin-4 reduced ICP over time, which was significantly different from baseline at 15 minutes ( $68.9 \pm 6.4\%$ ,  $P < 0.05$ ). ICV delivery of saline also reduced ICP over time (technical effect due to the ICV cannula itself reducing ICP), and this was significantly different from baseline at 50 minutes ( $74.5 \pm 7.9\%$ ,  $P < 0.05$ ). Over the 60 minutes of ICP measurement, ICV delivery of exendin-4 significantly reduced the area-under-the-curve (AUC) of ICP compared to saline delivered via the same route ( $3852 \pm 397$  versus  $4974 \pm 262$  AUC,  $P < 0.05$ ) (**Fig. 4O**). To determine if the effects of exendin-4 on ICP were mediated by the GLP-1R, the antagonist exendin 9-39 was continuously infused ( $4 \mu\text{g}/\text{hour}$ ) into the lateral ventricle for 2 days prior to SC administration of either saline or  $20 \mu\text{g}/\text{kg}$  exendin-4. SC injection of exendin-4 (ICV saline + SC exendin-4) lowered ICP ( $P < 0.0001$ ) compared to a SC injection of saline with ICV delivery of exendin 9-39 (ICV exendin 9-39 + SC saline; **Fig. 4P**). **Central ICV exendin 9-39 infusion decreased the ICP-lowering effect of SC exendin-4 at 5 minutes (ICV exendin 9-39 + SC exendin-4  $96.7 \pm 13.7\%$  vs ICV saline + SC exendin-4  $65.7 \pm 8.3\%$ ,  $P < 0.001$ ) (Fig. 4P).** These data suggest that exendin-4 in part exerts its effects on ICP via the GLP-1R signaling pathway in the brain.

*Exendin-4 reduces ICP in a dose-dependent manner and the effects last for 24 hours*

Rats were treated subcutaneously with 1, 3 and  $5 \mu\text{g}/\text{kg}$  exendin-4 to determine whether exendin-4 reduces ICP at lower concentrations. At 60 minutes 1, 3 and  $5 \mu\text{g}/\text{kg}$  exendin-4 significantly reduced ICP to  $79.0 \pm 7.0\%$  of baseline ( $P < 0.05$ ),  $69.9 \pm 8.8\%$  of baseline ( $P < 0.0001$ ) and  $60.6 \pm 3.6\%$  of baseline ( $P < 0.0001$ ), respectively, compared to saline ( $97.2 \pm$

2.5% of baseline) (**Fig. 5A-B**). Five  $\mu\text{g/kg}$  exendin-4 showed the greatest reduction in ICP and the effect was still present 3 hours after the treatment ( $P<0.001$ ). Conversely, in 1 and 3  $\mu\text{g/kg}$  exendin-4 groups ICP had returned to baseline by 3 hours (**Fig. 5C**).

Alterations in mRNA and protein expression of GLP-1R and molecules involved in CSF secretion were assessed in the choroid plexus of rats 3 hours after treatment with 1, 3 and 5  $\mu\text{g/kg}$  exendin-4. *Glp-1R* and  $\text{Na}^+ \text{K}^+ \text{atpase}$  mRNA expression was not altered by exendin-4 treatment (**Fig. 5D-E**). Conversely, there was a 2-fold increase in the amount of *Aqp1* mRNA in the 5  $\mu\text{g/kg}$  exendin-4 treatment group ( $P<0.05$ ) (**Fig. 5F**), and a 4-fold increase in the amount of *Nhe1* mRNA expression in the 1  $\mu\text{g/kg}$  exendin-4 treatment group ( $P<0.05$ ) (**Fig. 5G**). Although no significant changes were observed in  $\text{Na}^+ \text{K}^+ \text{atpase}$  mRNA expression, there was a small increase in  $\text{Na}^+ \text{K}^+ \text{ATPase}$  protein in the 5  $\mu\text{g/kg}$  exendin-4 treatment group ( $2.16 \pm 0.22 \text{ AU}$ ,  $P<0.05$ ) (**Fig 5H-I**). Two bands were observed for the water channel aquaporin 1 (AQP1), representing the glycosylated (top band) and non-glycosylated (bottom band) forms of AQP1 (**Fig. 5H**). The total amount of AQP1 protein was slightly reduced by 1 and 3  $\mu\text{g/kg}$  exendin-4 treatment but not with the higher 5  $\mu\text{g/kg}$  exendin-4 dose (1  $\mu\text{g/kg}$ ,  $1.96 \pm 0.17 \text{ AU}$ ,  $P<0.05$ , 3  $\mu\text{g/kg}$ ,  $1.75 \pm 0.08 \text{ AU}$ ,  $P<0.01$ , 5  $\mu\text{g/kg}$ ,  $2.75 \pm 0.30 \text{ AU}$ ) (**Fig. 5J**). The ratio of glycosylated AQP1 to non-glycosylated AQP1 was increased after 1 and 3  $\mu\text{g/kg}$  exendin-4 treatment but not after 5  $\mu\text{g/kg}$  exendin-4 treatment (1  $\mu\text{g/kg}$ ,  $0.97 \pm 0.06 \text{ AU}$ ,  $P<0.05$ , 3  $\mu\text{g/kg}$ ,  $1.08 \pm 0.06 \text{ AU}$ ,  $P<0.01$ , 5  $\mu\text{g/kg}$ ,  $0.81 \pm 0.08 \text{ AU}$ ) (**Fig. 5K**). Glycosylation is important for intracellular trafficking and protein stability, making proteins more resistant to proteolysis (24), therefore these data suggest that exendin-4 may lower AQP1 through enhanced degradation of the non-glycosylated AQP1.

The effect of 5  $\mu\text{g/kg}$  exendin-4 was monitored for 24 hours in healthy rats to determine its duration of action. A single SC injection of 5  $\mu\text{g/kg}$  exendin-4 maintained lower ICP compared to saline over 24 hours and returned to the pre-dose ICP baseline at 24 hours (1 hour,  $60.2 \pm$

3.5%,  $P<0.0001$ , 3 hours,  $71.3 \pm 3.7\%$ ,  $P<0.001$ , 6 hours,  $70.3 \pm 4.0\%$ ,  $P<0.0001$ , 12 hours,  $88.9 \pm 16.6\%$ ,  $P<0.01$ , 24 hours,  $100.3 \pm 14.3\%$ ,  $P<0.01$ ) (**Fig. 6A**). Effects on weight and food and water intake were also noted in relation to changes in ICP over 24 hours. Although exendin-4 caused a greater reduction in weight at 3 and 6 hours (**Fig. 6B**), there were no differences between food or water intake at any time point between exendin-4-treated and saline-treated rats (**Fig. 6C-D**). *Glp-1R*,  $Na^+ K^+ atpase$  and *Nhe1* mRNA expression did not change over the 24 hour period (**Fig. 6E-F,H**). As shown previously, 5  $\mu\text{g/kg}$  exendin-4 increased *Aqp1* mRNA expression at 3 hours compared to saline, although this was not observed at any other time point (**Fig. 6G**). There were also no significant changes in the amount of  $Na^+ K^+$  ATPase or AQP1 protein over the 24 hour time period (**Fig. 6I-L**).

#### *Exendin-4 treatment reduces ICP in a rodent model of raised ICP*

To determine the efficacy of exendin-4 to reduce ICP under conditions of raised ICP, a well-characterized kaolin model of hydrocephalus in rats was used. Kaolin, an aluminium silicate, acts as an irritant, inducing an inflammatory response with concomitant deposition of collagen and dense fibrosis in areas of the subarachnoid space close to the injection site, which leads to raised ICP (25, 26). Kaolin was injected into the cisterna magna, leading to development of hydrocephalus, before implantation of the ICP monitor. ICP was recorded before and after a SC injection of either saline or 20  $\mu\text{g/kg}$  exendin-4 (**Fig. 7A**). The injection of kaolin significantly increased baseline ICP ( $11.1 \pm 1.3$  mmHg;  $P<0.0001$ ) compared to that of normal rats ( $5.5 \pm 0.4$  mmHg) (**Fig. 7B**). Exendin-4 treatment significantly reduced ICP almost immediately after the SC injection, and at 30 minutes was  $62.6 \pm 5.1\%$  of baseline ( $P<0.0001$ ) compared to  $105.0 \pm 4.6\%$  of baseline in saline-treated rats (**Fig. 7C**). Eight rats in the kaolin group had baseline ICP values of greater than 10 mmHg and had an average baseline ICP of  $13.7 \pm 0.7$  mmHg. In these rats (ICP  $>10$  mmHg), the ICP values at 30 minutes were  $56.6 \pm$

5.7% of baseline (n=4) in the exendin-4 treatment group compared to  $106.7 \pm 8.6\%$  of baseline (n=4) in the saline treatment group (**Fig. 7C**). In the rodents with elevated ICP, the ICP waveform was very unstable, with the appearance of B-waves characteristic of pathologically elevated ICP and a reduction in brain compliance (27). These were abolished in rats receiving exendin-4 but not saline (**Fig. 7D**).

## Discussion

The aim of the present study was to establish whether GLP-1 had a role in modulating CSF secretion and ICP. We were able to demonstrate that the GLP-1R agonist exendin-4 was able to reduce ICP in conscious healthy female rats and in a rat model of raised ICP. In addition, our results suggest that the ICP-lowering properties of exendin-4 may occur through reduced CSF secretion at the choroid plexus, implied by the reduction in  $\text{Na}^+ \text{K}^+$  ATPase activity in CPe cells. Furthermore, our data suggest that exendin-4 modulates CSF production in vitro through the GLP-1R/cAMP/PKA signaling pathway.

Alvarez et al. (15) first described the presence of the GLP-1R in the rat ependyma and choroid plexus by *in situ* hybridisation, but did not characterize the cellular localization of this receptor. Our studies corroborate these findings and demonstrate further that *GLP-1R* mRNA and protein are present in both rat and human choroid plexus. We localized the GLP-1R protein in tissue sections of the human choroid plexus to the CPe cells using a monoclonal antibody, and showed the presence of the receptor in the rat choroid plexus using fluorescently tagged exendin-4. We note that no specific antibody exists for mouse/rat tissue so rodent tissue was not examined for GLP-1R protein expression. In any case, our studies are in keeping with others showing localization of the GLP-1R in monkey kidney and human GLP-1R transfected cells (20, 21). G-protein coupled receptors undergo internalization, trafficking and recycling/degradation following agonist stimulation (28). We speculate that such dynamics

may allow the GLP-1R to be stimulated from both sides of the choroid plexus (**Fig. S2A**). Although GLP-1R mRNA and protein expression were in general low, it has recently been shown that activation of the receptor requires femto- to picomolar concentrations of GLP-1R, so even faced with low abundance, signaling would be expected in the presence of exendin-4 (29).

We successfully cultured monolayers of rat CPe cells, which we used as an in vitro cell culture model of the rat choroid plexus to assess CSF secretion. The Na<sup>+</sup> K<sup>+</sup> ATPase is localized to the apical surface and is the driving force for transporting Na<sup>+</sup> ions from the choroid plexus into the CSF against its concentration gradient. Many studies have demonstrated that modulation of Na<sup>+</sup> K<sup>+</sup> ATPase expression or activity directly correlates with CSF secretion (6, 30-33). We were able to show that exendin-4 reduces Na<sup>+</sup> K<sup>+</sup> ATPase activity, suggesting reduced CSF secretion at the choroid plexus. Previous studies have shown similar effects of exendin-4 on Na<sup>+</sup> K<sup>+</sup> ATPase activity in the renal system (34). In kidney proximal tubule epithelial cells and pancreatic beta cells, GLP-1 modulates Na<sup>+</sup> concentration through increased cAMP and PKA activation (18, 35). Using two different techniques, exendin-4 was seen to induce a concentration-dependent rise in cAMP in the choroid plexus, which was inhibited by the GLP-1R antagonist, exendin 9-39. Furthermore, a PKA inhibitor blocked the effects of exendin-4 on Na<sup>+</sup> K<sup>+</sup> ATPase activity, although we acknowledge that such approaches can be non-specific and further studies using specific knockout animals are required. Altogether, these data indicate that the cAMP/PKA-dependent pathway may be involved in the GLP-1R-mediated reduction in CSF secretion at the choroid plexus. In the kidney, GLP-1R agonist treatment increases diuresis through phosphorylation of the Na<sup>+</sup> H<sup>+</sup> exchanger (18, 36). There are PKA phosphorylation sites present on both the Na<sup>+</sup> H<sup>+</sup> exchanger and the Na<sup>+</sup> K<sup>+</sup> ATPase (37), therefore, in the choroid plexus, phosphorylation of either the Na<sup>+</sup> H<sup>+</sup> exchanger or the Na<sup>+</sup> K<sup>+</sup> ATPase may result in inhibition of Na<sup>+</sup> transport across the cells

and thus CSF production (**Fig. S2B-C**). In the choroid plexus, the Na<sup>+</sup> K<sup>+</sup> ATPase can also be phosphorylated by PKC (37). Interestingly, GLP-1R is able to signal through the PKC pathway in pancreatic beta cells (29, 38, 39). Therefore, the GLP-1R/PKC signaling pathway may also have a role in reducing CSF secretion and warrants further investigation.

The key finding of this study is that subcutaneous exendin-4 treatment is able to reduce ICP in vivo in normal rats and rats with raised ICP. In addition, the effect on ICP of a single administration of exendin-4 lasted for 24 hours and cumulative dosing reduced the pre-dose ICP. This suggests that exendin-4 may be able to maintain low ICP over a long period. This is an important advance, as there are very limited specific therapeutic options to clinically reduce and maintain low ICP under conditions of raised ICP. The main therapeutic agent for managing chronic raised ICP is acetazolamide, a carbonic anhydrase inhibitor. However, in idiopathic intracranial hypertension, acetazolamide is associated with limited efficacy and poor tolerability (48% withdrawal) (2), and is contraindicated for use in premature infants with post-haemorrhagic hydrocephalus (40). On the other hand, treatment with incretin mimetics is generally well tolerated, with the main side effects being transient nausea, constipation and diarrhea, and these drugs do not induce hypoglycemia (41). In patients taking the GLP-1R agonist liraglutide, drug withdrawal due to side effects was only 5.4% in the cohort receiving the highest dose (3mg; 12).

There are, however, a number of limitations to the present study. To determine the central actions of exendin-4 on ICP, we had to deliver exendin-4 directly into the brain's ventricular system. The injection itself may have a direct effect on ICP and could mask any changes in ICP relating to the treatment. To try to minimize these effects, we implanted an ICV cannula 2 days prior to the injection. Nonetheless, as it was not possible to completely seal the system, ICP showed a slight decrease in saline-treated rats. However, we were still able to establish a significant reduction in ICP with exendin-4 treatment. The study design was also

limited by the lack of blinding during the intervention, although the data were analyzed by different individuals with the same outcome. ICP was monitored continuously via automated software thus removing measurement bias. It will be of interest to study in the future, prolonged dosing in a rat model of hydrocephalus. However, this will require considerable technical optimization, given that ICP is notoriously difficult to measure in such models where recordings are typically only accurate immediately before euthanasia (42, 43).

GLP-1R agonists also have peripheral actions that have the potential to indirectly affect ICP. Whilst incretin mimetics have been shown to acutely increase heart rate and blood pressure (44), hypertension would be expected to cause the opposite effect to that seen here due to increased choroid plexus permeability and fluid secretion (45, 46). Indeed, our data imply that the effect of exendin-4 on ICP dynamics is through central mechanisms, since ICV infusion of exendin 9-39 partially inhibited the action of SC exendin-4. Exendin 9-39 may not have fully inhibited the actions of exendin-4, since the inhibitor was infused into the ventricle rather than being given as a bolus injection. However, it is also possible that the effects of exendin-4 are not fully mediated by GLP-1R and this requires further investigation. Previous studies have also demonstrated only moderate effects on attenuating exendin-4 induced food intake suppression at early time points following ICV bolus of exendin 9-39 (47). Nevertheless, the central actions of exendin-4 are further supported by the fact that exendin-4 lowered CSF pH whereas blood pH remained unchanged, which is supported by other studies showing that GLP-1 does not affect blood pH (19). It remains unclear how the subcutaneous administration of the GLP-1R agonist exendin-4 exerts its central effects on the choroid plexus. Following subcutaneous administration, circulating exendin-4 may cross the fenestrated capillaries in the choroid plexus and stimulate the GLP-1R on the basolateral side of the CPe cells. Otherwise, it is possible that exendin-4 crossed the blood brain barrier (48, 49) or entered the CSF via the circumventricular organs, where it is able to stimulate the receptors on the apical surface of the

CPe cells. Indeed, liraglutide readily crosses into the hypothalamic arcuate nucleus (50), and in vivo imaging studies in rodents using fluorescently-tagged ghrelin show passage of the gut peptide to the same region *via* fenestrated capillaries of the median eminence (51). Lastly, exendin-4 may stimulate vagal afferents that project to the nucleus tractus solitarius (11). This may lead to secretion of GLP-1 through a widespread network of fibres projecting to the third ventricle allowing GLP-1 to enter the CSF (Fig. S2A).

In summary, exendin-4 reduces Na<sup>+</sup> K<sup>+</sup> ATPase activity at the choroid plexus, implying a reduction in CSF secretion, and lowers ICP in conscious rats with and without elevated ICP. This work demonstrates that GLP-1R agonists may provide an alternative treatment for raised ICP in conditions such as idiopathic intracranial hypertension and hydrocephalus, and warrants further clinical investigation in humans.



## Materials and Methods

### Study design

The main aim of this study was to explore the potential of exendin-4, a GLP-1R agonist, to modulate CSF secretion and subsequently reduce ICP. Three experimental studies were performed: (i) in vitro analysis of the GLP-1R and downstream signaling pathway in human and rat choroid plexus, GLP-1R expression was determined through mRNA analysis, immunostaining of human choroid plexus tissue sections and fluorescently tagged exendin-4 binding to rat choroid plexus explants. The downstream signaling pathway was assessed in rat CPe cell culture by measuring cAMP generation and Na<sup>+</sup> K<sup>+</sup> ATPase activity. In vivo studies to determine the efficacy of exendin-4 to reduce ICP were conducted in (ii) healthy rats and (iii) in a pathological model of raised ICP, a rat model of hydrocephalus. The sample size (n=4-9 per experimental group) for the in vivo studies was based on the resources equation as the effects size was unknown. Exact numbers for each experiment are included below and in the figure legends. The investigators were not blinded when conducting or evaluating the experiments and the rats were randomly assigned to the treatment and control groups.

### Human tissue

Human choroid plexus samples were obtained from the Parkinson's UK Brain Bank at Imperial College London, under the ethical approval of the Wales Research Ethics Committee (Ref. No.

08/MRE09/31+5). Informed consent was obtained for the use of post mortem tissue for research. Samples were stored in RNALater at -80°C before being processed for qPCR following the protocol stated in the supplementary methods. Pooled human pancreas (540023), heart (540011) and ovary (540071) RNA was purchased from Agilent Technologies. Fresh choroid plexus samples were fixed in 4% formaldehyde before embedding in paraffin wax.

## **Experimental animals**

For the in vitro work, 150-200g female Sprague-Dawley rats (Charles River) were used at the University of Birmingham in accordance with the Animals and Scientific Procedures Act 1986, licensed by the UK Home Office and approved by the University of Birmingham Ethics Committee. For the in vivo studies, which were conducted in Rigshospitalet-Glostrup, 150-250g female Sprague-Dawley rats (Taconic) were housed in groups of 4, kept under a 12 hour light/dark cycle with free access to food and water. All experimental procedures were approved by the Danish Animal Experiments Inspectorate (license number 2014-15-0201-00256 and 2012-15-2934-00283). After treatments and surgical procedures, the rats were monitored daily for any adverse effects. Female rats were used to ensure the results were relevant to conditions such as idiopathic intracranial hypertension.

**Daily subcutaneous injection of exendin-4 in normal conscious rats.** On day 0, the epidural ICP probe was implanted and the animal allowed to recover. On day 2, 4 and 6, for the ICP recordings the rats were sedated with midazolam (2.5 mg/kg subcutaneous injection) in an infusion cage (Instech Laboratories), which had a swirl lever arm to ensure unhindered movement. A stable baseline ICP reading was recorded for around 30 minutes before the rats received a SC injection of either saline (n=9) or 20µg/kg exendin-4 (n=9). ICP was recorded

for a further 60 minutes after which the rat was returned to its normal cage. The daily SC injections of saline or exendin-4 were performed at similar times of the day for each rat.

**ICV injection of exendin-4 in anesthetized rats.** To determine whether the effects of exendin-4 on ICP were due to central activity the rats were fitted with an ICV cannula at the same time as the epidural ICP probe implantation and the rat allowed to recover. Subsequent ICP recordings during exendin-4 treatment were done under anaesthesia. A stable baseline ICP reading was recorded for around 30 minutes before the following treatments were then administered ICV in a counterbalance design: (1) 1µl saline (n=8) and (2) 0.3µg/1µl exendin-4 (n=6). ICP was recorded for a further 60 minutes after which the rat was allowed to recover. Injection treatments were separated by 2-3 days.

**Continuous ICV infusion of exendin 9-39 with SC injection of exendin-4 in conscious rats.** To determine whether the effects of exendin-4 on ICP are through the GLP-1R, rats were fitted with an osmotic pump attached to an ICV cannula containing either saline or exendin 9-39 at the same time as the epidural ICP probe implantation. On day 2 the rats were sedated, a stable baseline recorded before a SC injection of either saline or 20µg/kg exendin-4. ICP was then recorded for a further 60 minutes. The rats were therefore assigned to 3 treatment groups: (1) Saline filled osmotic pump with SC injection of exendin-4 (ICV saline + SC exendin-4; n=6); (2) exendin 9-39 filled osmotic pump with SC injection of saline (ICV exendin 9-39 + SC saline; n=5); and (3), exendin 9-39 filled osmotic pump with SC injection of exendin-4 (ICV exendin 9-39 + SC exendin-4; n=6).

**Exendin-4 dose response and time course experiment.** Rats underwent the same procedure as outlined in experiment 1. For the dosing experiment the rats were given either 1 (n=6), 3

(n=6) or 5µg/kg exendin-4 (n=6) and ICP recorded for 3 hours. For the time course experiment rats were given either saline (n=18 for ICP data but only 4 were used for choroid plexus analysis) and ICP recorded for 24 hours, or 5µg/kg exendin-4 and the ICP recorded for 6 (n=6), 12 (n=6) and 24 hours (n=12 for ICP data but only 6 were used for choroid plexus analysis). After each time point the rats were killed with an overdose of euthatol and transcardially perfused with ice cold PBS. The choroid plexus was then dissected, frozen immediately and stored at -80°C for qPCR and western blot analysis (described in detail in the supplementary methods).

**SC injection of exendin-4 in conscious hydrocephalic rats.** We used the kaolin model of hydrocephalus as our model of raised ICP. On day 0 the rats received an injection of kaolin to induce hydrocephalus and the rat allowed to recover. On day 6-8 the rats were fitted with an epidural ICP probe and the rat was then allowed to recover in the infusion cages still connected to the transducer so that the ICP could be continuously measured overnight to establish raised ICP. The following morning, after establishing the baseline ICP reading was stable, the rats received a SC injection of either saline (n=6, n=4 >10mmHg) or 20µg/kg exendin-4 (n=6; n=4 >10mmHg). ICP was then recorded for a further 60 minutes.

### **Statistical analysis**

Values are represented as mean and standard error of the mean (SEM). The majority of the data was analyzed using GraphPad Prism software, however, the time course experiment with 5µg/kg exendin-4 was analyzed using SPSS due to missing data points. For the ELISA cAMP analysis, the non-parametric Kruskal-Wallis test was used, and was followed by Mann-Whitney test (two-tailed) with the appropriate adjustment for multiple comparisons (Bonferroni). T-test or One-way ANOVA (followed by a post hoc Tukey test) was used for the

comparison of qPCR, western blot and Na<sup>+</sup> K<sup>+</sup> ATPase activity. Two-way ANOVA with Sidak's multiple comparison test was used for the comparison of ICP between two groups over a period of time. Values were considered statistically significant when P values were \*P<0.05, \*\*P<0.01, \*\*\*P<0.001, \*\*\*\*P<0.0001. Individual level data are included in table S1.

## Supplementary Materials

### Supplementary Materials and Methods

Fig. S1. Characterisation of primary rat choroid plexus epithelial cells *in vivo* and *in vitro*.

Fig. S2. Suggested route for GLP-1 action at the choroid plexus

Table S1. Individual level data corresponding to the different figures.

## References

1. L. Sakka, G. Coll, J. Chazal, Anatomy and physiology of cerebrospinal fluid. *European Annals of Otorhinolaryngology, Head and Neck Diseases* **128**, 309-316 (2011).
2. A. K. Ball, A. Howman, K. Wheatley, M. A. Burdon, T. Matthews, A. S. Jacks, M. Lawden, A. Sivaguru, A. Furmston, S. Howell, B. Sharrack, M. B. Davies, A. J. Sinclair, C. E. Clarke, A randomised controlled trial of treatment for idiopathic intracranial hypertension. *Journal of neurology* **258**, 874-881 (2011).
3. M. Wall, M. P. McDermott, K. D. Kiebert, J. J. Corbett, S. E. Feldon, D. I. Friedman, D. M. Katz, J. L. Keltner, E. B. Schron, M. J. Kupersmith, Effect of acetazolamide on visual function in patients with idiopathic intracranial hypertension and mild visual loss: the idiopathic intracranial hypertension treatment trial. *Jama* **311**, 1641-1651 (2014).
4. P. D. Brown, S. L. Davies, T. Speake, I. D. Millar, Molecular mechanisms of cerebrospinal fluid production. *Neuroscience* **129**, 957-970 (2004).
5. T. Speake, C. Whitwell, H. Kajita, A. Majid, P. D. Brown, Mechanisms of CSF secretion by the choroid plexus. *Microscopy research and technique* **52**, 49-59 (2001).
6. M. Pollay, B. Hisey, E. Reynolds, P. Tomkins, F. A. Stevens, R. Smith, Choroid plexus Na<sup>+</sup>/K<sup>+</sup>-activated adenosine triphosphatase and cerebrospinal fluid formation. *Neurosurgery* **17**, 768-772 (1985).
7. M. D. Parker, E. J. Myers, J. R. Schelling, Na<sup>+</sup>-H<sup>+</sup> exchanger-1 (NHE1) regulation in kidney proximal tubule. *Cellular and molecular life sciences : CMLS* **72**, 2061-2074 (2015).
8. H. H. Damkier, P. D. Brown, J. Praetorius, Cerebrospinal fluid secretion by the choroid plexus. *Physiological reviews* **93**, 1847-1892 (2013).
9. L. L. Baggio, D. J. Drucker, Biology of incretins: GLP-1 and GIP. *Gastroenterology* **132**, 2131-2157 (2007).

- 507 10. J. E. Campbell, D. J. Drucker, Pharmacology, physiology, and mechanisms of incretin hormone  
508 action. *Cell metabolism* **17**, 819-837 (2013).
- 509 11. P. J. Larsen, M. Tang-Christensen, J. J. Holst, C. Orskov, Distribution of glucagon-like peptide-1  
510 and other preproglucagon-derived peptides in the rat hypothalamus and brainstem.  
511 *Neuroscience* **77**, 257-270 (1997).
- 512 12. A. Astrup, S. Rossner, L. Van Gaal, A. Rissanen, L. Niskanen, M. Al Hakim, J. Madsen, M. F.  
513 Rasmussen, M. E. Lean, Effects of liraglutide in the treatment of obesity: a randomised,  
514 double-blind, placebo-controlled study. *Lancet* **374**, 1606-1616 (2009).
- 515 13. A. L. Alhadeff, H. J. Grill, Hindbrain nucleus tractus solitarius glucagon-like peptide-1 receptor  
516 signaling reduces appetitive and motivational aspects of feeding. *American journal of*  
517 *physiology. Regulatory, integrative and comparative physiology* **307**, R465-470 (2014).
- 518 14. A. Flint, A. Raben, A. Astrup, J. J. Holst, Glucagon-like peptide 1 promotes satiety and  
519 suppresses energy intake in humans. *The Journal of clinical investigation* **101**, 515-520 (1998).
- 520 15. E. Alvarez, I. Roncero, J. A. Chowen, B. Thorens, E. Blazquez, Expression of the glucagon-like  
521 peptide-1 receptor gene in rat brain. *Journal of neurochemistry* **66**, 920-927 (1996).
- 522 16. S. C. Cork, J. E. Richards, M. K. Holt, F. M. Gribble, F. Reimann, S. Trapp, Distribution and  
523 characterisation of Glucagon-like peptide-1 receptor expressing cells in the mouse brain. *Mol*  
524 *Metab* **4**, 718-731 (2015).
- 525 17. F. Marques, J. C. Sousa, G. Coppola, F. Gao, R. Puga, H. Brentani, D. H. Geschwind, N. Sousa, M.  
526 Correia-Neves, J. A. Palha, Transcriptome signature of the adult mouse choroid plexus. *Fluids*  
527 *Barriers CNS* **8**, 10 (2011).
- 528 18. L. R. Carraro-Lacroix, G. Malnic, A. C. Girardi, Regulation of Na<sup>+</sup>/H<sup>+</sup> exchanger NHE3 by  
529 glucagon-like peptide 1 receptor agonist exendin-4 in renal proximal tubule cells. *American*  
530 *journal of physiology. Renal physiology* **297**, F1647-1655 (2009).
- 531 19. R. O. Crajoinas, F. T. Oricchio, T. D. Pessoa, B. P. Pacheco, L. M. Lessa, G. Malnic, A. C. Girardi,  
532 Mechanisms mediating the diuretic and natriuretic actions of the incretin hormone glucagon-  
533 like peptide-1. *American journal of physiology. Renal physiology* **301**, F355-363 (2011).
- 534 20. C. Pyke, R. S. Heller, R. K. Kirk, C. Orskov, S. Reedtz-Runge, P. Kastrup, A. Hvelplund, L.  
535 Bardram, D. Calatayud, L. B. Knudsen, GLP-1 receptor localization in monkey and human  
536 tissue: novel distribution revealed with extensively validated monoclonal antibody.  
537 *Endocrinology* **155**, 1280-1290 (2014).
- 538 21. C. Pyke, L. B. Knudsen, The glucagon-like peptide-1 receptor--or not? *Endocrinology* **154**, 4-8  
539 (2013).
- 540 22. C. Widmann, W. Dolci, B. Thorens, Agonist-induced internalization and recycling of the  
541 glucagon-like peptide-1 receptor in transfected fibroblasts and in insulinomas. *The*  
542 *Biochemical journal* **310 ( Pt 1)**, 203-214 (1995).
- 543 23. A. Sinclair, M. Burdon, A. Ball, N. Nightingale, P. Good, T. Matthews, A. Jacks, M. Lawden, C.  
544 Clarke, E. Walker, J. Tomlinson, P. Stewart, S. Rauz, Low energy diet and intracranial pressure  
545 in women with idiopathic intracranial hypertension: prospective cohort study. *BMJ* **7**, 341  
546 (2010).
- 547 24. C. M. Herak-Kramberger, D. Breljak, M. Ljubojevic, M. Matokanovic, M. Lovric, D. Rogic, H.  
548 Brzica, I. Vrhovac, D. Karaica, V. Micek, J. I. Dupor, D. Brown, I. Sabolic, Sex-dependent  
549 expression of water channel AQP1 along the rat nephron. *American journal of physiology.*  
550 *Renal physiology* **308**, F809-821 (2015).
- 551 25. J. Hatta, T. Hatta, K. Moritake, H. Otani, Heavy water inhibiting the expression of transforming  
552 growth factor-beta1 and the development of kaolin-induced hydrocephalus in mice. *J*  
553 *Neurosurg* **104**, 251-258 (2006).
- 554 26. Y. Nakagawa, J. Cervos-Navarro, J. Artigas, Tracer study on a paracellular route in experimental  
555 hydrocephalus. *Acta Neuropathol* **65**, 247-254 (1985).
- 556 27. A. Spiegelberg, M. Preuss, V. Kurtcuoglu, B-waves revisited. *Interdiscip Neurosur* **6**, 13-17  
557 (2016).

- 558 28. S. S. Ferguson, Evolving concepts in G protein-coupled receptor endocytosis: the role in  
559 receptor desensitization and signaling. *Pharmacological reviews* **53**, 1-24 (2001).
- 560 29. M. Shigeto, R. Ramracheya, A. I. Tarasov, C. Y. Cha, M. V. Chibalina, B. Hastoy, K. Philippaert, T.  
561 Reinbothe, N. Rorsman, A. Salehi, W. R. Sones, E. Vergari, C. Weston, J. Gorelik, M. Katsura, V.  
562 O. Nikolaev, R. Vennekens, M. Zaccolo, A. Galione, P. R. Johnson, K. Kaku, G. Ladds, P.  
563 Rorsman, GLP-1 stimulates insulin secretion by PKC-dependent TRPM4 and TRPM5 activation.  
564 *The Journal of clinical investigation* **125**, 4714-4728 (2015).
- 565 30. M. Lindvall-Axelsson, P. Hedner, C. Owman, Corticosteroid action on choroid plexus: reduction  
566 in Na<sup>+</sup>-K<sup>+</sup>-ATPase activity, choline transport capacity, and rate of CSF formation. *Experimental*  
567 *brain research* **77**, 605-610 (1989).
- 568 31. M. Lindvall-Axelsson, C. Nilsson, C. Owman, B. Winblad, Inhibition of cerebrospinal fluid  
569 formation by omeprazole. *Experimental neurology* **115**, 394-399 (1992).
- 570 32. G. Fisone, G. L. Snyder, J. Fryckstedt, M. J. Caplan, A. Aperia, P. Greengard, Na<sup>+</sup>,K<sup>(+)</sup>-ATPase in  
571 the choroid plexus. Regulation by serotonin/protein kinase C pathway. *The Journal of*  
572 *biological chemistry* **270**, 2427-2430 (1995).
- 573 33. M. E. Han, H. J. Kim, Y. S. Lee, D. H. Kim, J. T. Choi, C. S. Pan, S. Yoon, S. Y. Baek, B. S. Kim, J. B.  
574 Kim, S. O. Oh, Regulation of cerebrospinal fluid production by caffeine consumption. *BMC*  
575 *neuroscience* **10**, 110 (2009).
- 576 34. S. Sancar-Bas, S. Gezinci-Oktayoglu, S. Bolkent, Exendin-4 attenuates renal tubular injury by  
577 decreasing oxidative stress and inflammation in streptozotocin-induced diabetic mice. *Growth*  
578 *Factors* **33**, 419-429 (2015).
- 579 35. Y. Miura, H. Matsui, Glucagon-like peptide-1 induces a cAMP-dependent increase of [Na<sup>+</sup>]<sub>i</sub>  
580 associated with insulin secretion in pancreatic beta-cells. *American journal of physiology.*  
581 *Endocrinology and metabolism* **285**, E1001-1009 (2003).
- 582 36. T. Rieg, M. Gerasimova, F. Murray, T. Masuda, T. Tang, M. Rose, D. J. Drucker, V. Vallon,  
583 Natriuretic effect by exendin-4, but not the DPP-4 inhibitor alogliptin, is mediated via the GLP-  
584 1 receptor and preserved in obese type 2 diabetic mice. *American journal of physiology. Renal*  
585 *physiology* **303**, F963-971 (2012).
- 586 37. A. G. Therien, R. Blostein, Mechanisms of sodium pump regulation. *American journal of*  
587 *physiology. Cell physiology* **279**, C541-566 (2000).
- 588 38. Y. Suzuki, H. Zhang, N. Saito, I. Kojima, T. Urano, H. Mogami, Glucagon-like peptide 1 activates  
589 protein kinase C through Ca<sup>2+</sup>-dependent activation of phospholipase C in insulin-secreting  
590 cells. *The Journal of biological chemistry* **281**, 28499-28507 (2006).
- 591 39. M. Shigeto, K. Kaku, Are both protein kinase A- and protein kinase C-dependent pathways  
592 involved in glucagon-like peptide-1 action on pancreatic insulin secretion? *Journal of diabetes*  
593 *investigation* **5**, 347-348 (2014).
- 594 40. C. A. Mazzola, A. F. Choudhri, K. I. Auguste, D. D. Limbrick, Jr., M. Rogido, L. Mitchell, A. M.  
595 Flannery, R. Pediatric Hydrocephalus Systematic, F. Evidence-Based Guidelines Task, Pediatric  
596 hydrocephalus: systematic literature review and evidence-based guidelines. Part 2:  
597 Management of posthemorrhagic hydrocephalus in premature infants. *J Neurosurg Pediatr* **14**  
598 **Suppl 1**, 8-23 (2014).
- 599 41. T. A. Wadden, P. Hollander, S. Klein, K. Niswender, V. Woo, P. M. Hale, L. Aronne, Weight  
600 maintenance and additional weight loss with liraglutide after low-calorie-diet-induced weight  
601 loss: the SCALE Maintenance randomized study. *International journal of obesity (2005)* **37**,  
602 1443-1451 (2013).
- 603 42. W. Luedemann, D. Kondziella, K. Tienken, P. Klinge, T. Brinker, D. B. von Rautenfeld, in  
604 *Intracranial Pressure and Brain Biochemical Monitoring*, M. Czosnyka, J. Pickard, P. Kirkpatrick,  
605 P. Smielewski, P. Hutchinson, Eds. (Springer Vienna, 2002), vol. 81, chap. 70, pp. 271-273.
- 606 43. O. Bloch, K. I. Auguste, G. T. Manley, A. S. Verkman, Accelerated progression of kaolin-induced  
607 hydrocephalus in aquaporin-4-deficient mice. *J Cereb Blood Flow Metab* **26**, 1527-1537 (2006).

44. J. M. Barragan, R. E. Rodriguez, J. Eng, E. Blazquez, Interactions of exendin-(9-39) with the effects of glucagon-like peptide-1-(7-36) amide and of exendin-4 on arterial blood pressure and heart rate in rats. *Regul Pept* **67**, 63-68 (1996).
45. L. A. Murtha, Q. Yang, M. W. Parsons, C. R. Levi, D. J. Beard, N. J. Spratt, D. D. McLeod, Cerebrospinal fluid is drained primarily via the spinal canal and olfactory route in young and aged spontaneously hypertensive rats. *Fluids Barriers CNS* **11**, 12 (2014).
46. C. E. Johanson, J. A. Duncan, 3rd, P. M. Klinge, T. Brinker, E. G. Stopa, G. D. Silverberg, Multiplicity of cerebrospinal fluid functions: New challenges in health and disease. *Cerebrospinal fluid research* **5**, 10 (2008).
47. S. E. Kanoski, S. M. Fortin, M. Arnold, H. J. Grill, M. R. Hayes, Peripheral and central GLP-1 receptor populations mediate the anorectic effects of peripherally administered GLP-1 receptor agonists, liraglutide and exendin-4. *Endocrinology* **152**, 3103-3112 (2011).
48. A. J. Kastin, V. Akerstrom, Entry of exendin-4 into brain is rapid but may be limited at high doses. *International journal of obesity and related metabolic disorders : journal of the International Association for the Study of Obesity* **27**, 313-318 (2003).
49. A. J. Kastin, V. Akerstrom, W. Pan, Interactions of glucagon-like peptide-1 (GLP-1) with the blood-brain barrier. *J Mol Neurosci* **18**, 7-14 (2002).
50. D. J. Hodson, R. K. Mitchell, E. A. Bellomo, G. Sun, L. Vinet, P. Meda, D. Li, W. H. Li, M. Bugliani, P. Marchetti, D. Bosco, L. Piemonti, P. Johnson, S. J. Hughes, G. A. Rutter, Lipotoxicity disrupts incretin-regulated human beta cell connectivity. *The Journal of clinical investigation* **123**, 4182-4194 (2013).
51. M. Schaeffer, F. Langlet, C. Lafont, F. Molino, D. J. Hodson, T. Roux, L. Lamarque, P. Verdie, E. Bourrier, B. Dehouck, J. L. Baneres, J. Martinez, P. F. Mery, J. Marie, E. Trinquet, J. A. Fehrentz, V. Prevot, P. Mollard, Rapid sensing of circulating ghrelin by hypothalamic appetite-modifying neurons. *Proc Natl Acad Sci U S A* **110**, 1512-1517 (2013).

## Acknowledgements

## Funding



A.S. is funded by an NIHR Clinician Scientist Fellowship (NIHR-CS-011-028) and by the Medical Research Council, UK (MR/K015184/1). D.J.H. was supported by Diabetes UK R.D. Lawrence (12/0004431), EFSD/Novo Nordisk Rising Star and Birmingham Fellowships, a Wellcome Trust Institutional Support Award, an MRC Project Grant (MR/N00275X/1) and an ERC Starting Grant (OptoBETA; 715884). This work was supported by a MRC confidence in concept grant, the West Midlands Neuroscience Teaching and Research Fund and the University of Birmingham Research Development Fund.

#### **Author contributions**

A.S. was responsible for the study concept. H.B., A.G., D.J.H. and A.S. conceived and designed the experiments; H.B. conducted the following in vitro experiments: immunohistochemistry, Na<sup>+</sup> K<sup>+</sup> ATPase activity assay, cAMP assay, rat qPCR and western blot, and FLEX analysis); C.W. performed human qPCR and cAMP assays; A.G. M.U. and J.M contributed to the immunohistochemistry data; H.B., M.U. J.M. and S.H. performed the ICP recordings; H.B. and M.U. analyzed the data; H.B., M.U., A.G., D.J.H., R.J and A.S co-wrote the manuscript and all authors reviewed the final version.

#### **Competing interests**

A.S. holds patent # PCT/GB2015/052453 related to this work entitled “elevated intracranial pressure treatment”. R.J. has given lectures for Pfizer, Berlin-Chemie, Norspan, Merck and Autonomic Technologies and has been a member of the advisory boards of Autonomic Technologies, Medotech and ElectroCore.

## Figure legends

**Fig. 1. GLP-1R expression in post-mortem human choroid plexus tissue in vitro.** (A) Representative image of haematoxylin and eosin staining of human choroid plexus tissue section demonstrating classic choroid plexus morphology. (B) The histogram shows *GLP-1R* mRNA expression in human pancreas (n=1), heart (n=1), ovary (n=1) and choroid plexus (n=5). (C-D) Representative images of GLP-1R staining of paraffin-embedded human choroid plexus counterstained with haematoxylin. Sections were incubated without primary antibody (C) and with the human GLP-1R antibody MAb 3F52 (D). (E-F) High magnification of the boxed regions shown in C and D respectively. Scale bars, 100µm, BV – blood vessel and CPe – choroid plexus epithelial cell.

**Fig. 2. Expression of GLP-1R after treatment with exendin-4 in rat choroid plexus in vitro.** (A) Representative images of rat choroid plexus after treatment with artificial CSF (aCSF) as control or fluorescently labelled exendin-4 (FLEX) in the presence or absence of the GLP-1R antagonist exendin 9-39. DAPI (blue) was used as a nuclear marker; scale bar, 50µm (insert, 25µm). (B-E) The histograms represent the fold change in mRNA expression of *Glp-1r* (B), *Na<sup>+</sup> K<sup>+</sup> atpase* (C), *Aqp1* (D) and *Nhe1* (E) (aCSF n=6; 3hr n=7, 6hr n=7) \*P<0.05, \*\*P<0.01; ANOVA with Tukey's multiple comparisons test.

**Fig. 3. Effect of exendin-4 treatment on cAMP and Na<sup>+</sup> K<sup>+</sup> ATPase activity in CPe cells.** (A-B) The histograms represent the amount of cAMP generated after incubation with control, exendin-4 with and without 1µM exendin 9-39 and forskolin (positive control) using two different methods of cAMP detection (A - control n=8, exendin-4 n=8, Forskolin n=5, B -

control n=5, 1nM n=5, 10nM n=6 and 100nM exendin-4 n=5; with 1μM exendin 9-39 n=6, n=5 and n=5 respectively). (C)  $\text{Na}^+ \text{K}^+$  ATPase activity was measured by determining the concentration of inorganic phosphate generated by the hydrolysis of ATP that was sensitive to ouabain ( $\text{Na}^+ \text{K}^+$  ATPase inhibitor) (control n=13, exendin-4 n=7; PKI n=8; exendin-4 + PKI n=8). \*P<0.05, \*\*P<0.01, \*\*\*P<0.001, \*\*\*\*P<0.0001, NS no significance. (A) Kruskal-Wallis followed by Mann-Whitney tests (Bonferroni correction); (B-C) ANOVA with Tukey's multiple comparisons test. Protein kinase A inhibitor, PKI.

**Fig. 4. Effect of exendin-4 on ICP in healthy conscious rats.** (A) Overview of the experimental design in normal rats. Rats were fitted with an epidural ICP probe and allowed to recover. Treatment was given daily for 5 days and ICP was recorded on days 2, 4 and 6, before and after the rats received a subcutaneous (SC) injection of either saline (n=9) or 20μg/kg exendin-4 (n=9). (B) Example ICP traces of saline (*blue*) and exendin-4 (*red*) treatment. Spikes in the trace represent when the animal was moving (\*) and accurate recording of ICP was confirmed by the response to jugular vein compression. (C-E) Line graphs showing the percentage of baseline ICP after SC injection of either saline or exendin-4 on day 2 (C), day 4 (D) and day 6 (E). (F-G) Histograms showing the pre-dose and 60 minutes post treatment ICP values (% of baseline on day 2) on days 2, 4 and 6 for exendin-4 (F) and saline (G). (H) Line graph of the % change in weight from day 2 (start of treatment) showing that both saline and exendin-4 treated rats lost weight but there was no significant difference between the groups on day 4 or 6. (I) Scatter plot of weight change (g) vs ICP change (mmHg) in the saline (*blue* n=4) and exendin-4 (*red* n=5) groups. (J-N) Histograms showing blood pH (J) and CSF pH (K), and the concentration of  $\text{Na}^+$  (L)  $\text{Cl}^-$  (M) and  $\text{Ca}^{2+}$  (N) in the CSF, 60 minutes after a SC injection of either saline or 20μg/kg exendin-4. (O) ICP was measured before and after a 1μl intracerebroventricular (ICV) injection of either saline (n=8) or 0.3μg exendin-4 (n=6). (P)

Exendin 9-39 was continually infused (4µg/µl/hr) into the lateral ventricle (ICV) and ICP was measured before and after a SC injection of either 20µg/kg exendin-4 (ICV exendin 9-39 + SC exendin-4, n=6) or saline (ICV exendin 9-39 + SC saline, n=5) and compared to continuous saline infusion (ICV Saline + SC exendin-4, n=6). \*P<0.05, \*\*P<0.01, \*\*\*P<0.001; (C-H, O-P) Two way ANOVA with Sidak's multiple comparison test; (J-N) T-test (two-tailed).

**Fig. 5. Effects of different doses of exendin-4 on ICP, mRNA and protein expression in healthy conscious rats.** (A-B) Dose-response of exendin-4's effects on ICP following SC administration of 1 (n=6), 3 (n=6), 5 (n=23) and 20 µg/kg (n=9) exendin-4 compared to saline (n=18) at 30 and 60 minutes. (C) Line graph showing the percentage of baseline ICP after treatment with 1, 3 or 5µg/kg exendin-4 measured over 3 hours. (D-G) The histograms show *Glp-1R* (D), *Na<sup>+</sup> K<sup>+</sup> atpase* (E), *Aqp1* (F) and *Nhe1* (G) mRNA expression in the rat choroid plexus after saline treatment (n=4) or treatment with 1 (n=5), 3 (n=6), 5 µg/kg (n=6) exendin-4. (H) Representative western blots and (I-K) semi-quantitative protein analysis for (I) *Na<sup>+</sup> K<sup>+</sup> ATPase* (112kDa) and (J) total AQP1, either non-glycosylated (NG, 29kDa) or glycosylated (G, 35kDa);  $\beta$ -actin (42kDa) loading control. (K) Histogram shows the ratio of G to NG AQP1. \*P<0.05, \*\*P<0.01, \*\*\*P<0.001, \*\*\*\*P<0.0001. (B-C) Two-way ANOVA with Sidak's multiple comparison test; (D-G and I-K) ANOVA with Tukey's multiple comparison test.

**Fig. 6. Effects of Exendin-4 time course on ICP, mRNA and protein expression in healthy conscious rats.** (A) Line graph showing the percentage of baseline ICP after a single SC injection of saline (n=18) or 5µg/kg exendin-4 (n=24) measured over 24 hours. (B-D) Histograms showing weight loss (B), water intake (C) and food intake (D) in rats treated with saline (n=4) or 5µg/kg exendin-4 at 3 (n=6), 6 (n=6) and 24 hours (n=6). (E-H) Histograms representing *Glp-1r* (E), *Na<sup>+</sup> K<sup>+</sup> atpase* (F), *Aqp1* (G) and *Nhe1* (H) mRNA expression in the

rat choroid plexus after treatment with saline (n=4) and 5µg/kg exendin-4 at 3 (n=6), 6 (n=5) and 24 hours (n=5). **(I)** Representative western blots and **(J-L)** semi-quantitative protein analysis for **(J)** Na<sup>+</sup> K<sup>+</sup> ATPase (112kDa) and **(K)** total AQP1 either nonglycosylated (NG, 29kDa) or glycosylated (G, 35kDa); β-actin (42kDa) loading control. **(L)** The histogram shows the ratio of G to NG AQP1. \*P<0.05, \*\*P<0.01, \*\*\*P<0.001, \*\*\*\*P<0.0001; (A-D) Two-way ANOVA with Sidak's multiple comparison test; (E-H and J-L) ANOVA with Tukey's multiple comparison test.

**Fig. 7. Effect of exendin-4 on ICP in a rat model of raised ICP (hydrocephalic).** **(A)** Overview of the experimental plan. Kaolin was injected into the cisterna magna to induce hydrocephalus. On Day 6 the ICP monitor was implanted under anaesthesia and ICP was recorded overnight to allow the ICP to normalize after implantation. On Day 7, the rats were given a SC injection of either saline (n=6) or 20µg/kg Exendin-4 (n=6), and ICP was recorded for a further 60 minutes. **(B)** Dot plot showing the individual baseline ICP values (mmHg) for the normal rats and rats injected with kaolin. The kaolin group had significantly higher baseline ICP values compared to the normal group, with 8/12 rats having an ICP value of >10mmHg. **(C)** Line graph showing the percentage of baseline ICP after treatment with either saline (dark blue, n=6) or exendin-4 (dark red, n=6). The groups could also be further divided into those with ICP >10mmHg in the saline group (light blue, n=4) and exendin-4 group (light red, n=4). **(D)** Example ICP trace in a hydrocephalic rat before and after treatment with exendin-4. Before treatment the rat exhibited pathological ICP B-waves (*b*), which were abolished following treatment with exendin-4. \*\*\*\* P<0.0001; (B) T-test (two tailed); (C) Two-way ANOVA with Sidak's multiple comparison test.