

Is carotid body physiological O₂ sensitivity determined by a unique mitochondrial phenotype?

Holmes, Andrew; Ray, Clare; Coney, Andrew; Kumar, Prem

DOI:

[10.3389/fphys.2018.00562](https://doi.org/10.3389/fphys.2018.00562)

License:

Creative Commons: Attribution (CC BY)

Document Version

Peer reviewed version

Citation for published version (Harvard):

Holmes, A, Ray, C, Coney, A & Kumar, P 2018, 'Is carotid body physiological O₂ sensitivity determined by a unique mitochondrial phenotype?', *Frontiers in Physiology*, vol. 9, 562. <https://doi.org/10.3389/fphys.2018.00562>

[Link to publication on Research at Birmingham portal](#)

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

Is carotid body physiological O₂ sensitivity determined by a unique mitochondrial phenotype?

Andrew P. Holmes¹, Clare J. Ray¹, Andrew M. Coney¹ & Prem Kumar^{1*}

¹Institute of Clinical Sciences, College of Medical and Dental Sciences, University of Birmingham, Birmingham, UK

Key words: carotid body, hypoxia, mitochondria, nitric oxide, arterial chemoreceptor, O₂ sensor, metabolism, mitochondrial inhibitors

Running title Carotid body mitochondria and O₂ sensing

Word count (excluding references and figure legends): 2995

* Corresponding author: P.Kumar

Email: p.kumar@bham.ac.uk

Table of contents category: Respiratory

Abstract

The mammalian carotid body (CB) is the primary arterial chemoreceptor that responds to acute hypoxia, initiating systemic protective reflex responses that act to maintain O₂ delivery to the brain and vital organs. The CB is unique in that it is stimulated at O₂ levels above those that begin to impact on the metabolism of most other cell types. Whilst a large proportion of the CB chemotransduction cascade is well defined, the identity of the O₂ sensor remains highly controversial. This short review evaluates whether the mitochondria can adequately function as acute O₂ sensors in the CB. We consider the similarities between mitochondrial poisons and hypoxic stimuli in their ability to activate the CB chemotransduction cascade and initiate rapid cardiorespiratory reflexes. We evaluate whether the mitochondria are required for the CB to respond to hypoxia. We also discuss if the CB mitochondria are different to those located in other non-O₂ sensitive cells, and what might cause them to have an unusually low O₂ binding affinity. In particular we look at the potential roles of competitive inhibitors of mitochondrial complex IV such as nitric oxide in establishing mitochondrial and CB O₂-sensitivity. Finally, we discuss novel signaling mechanisms proposed to take place within and downstream of mitochondria that link mitochondrial metabolism with cellular depolarization.

1 Introduction-the carotid body and hypoxia

The mammalian carotid body (CB) is a highly specialized sensory organ derived from the neural crest. The sensory units of the CB are the 'glomus' or 'type I' cells that respond to a variety of stimuli including hypoxia, hypercapnia, acidosis and hormones thereby allowing the CB to function as a polymodal receptor (Kumar and Bin-Jaliah, 2007; Ribeiro et al., 2013; Thompson et al., 2016). Type I cell activation leads to stimulation of adjacent chemoafferent fibers that relay sensory information into the central nervous system. The physiological consequence of CB stimulation is therefore the initiation of series of systemic protective reflexes characterized by increased ventilation, tachycardia, systemic vasoconstriction and adrenaline release from the adrenal medulla (Kumar, 2009).

There is now a general consensus that a series of key processes contribute to the CB hypoxic chemotransduction cascade. These include attenuation of outward K^+ current, type I cell depolarization, Ca^{2+} influx through L-type Ca^{2+} channels, neurosecretion and chemoafferent excitation (Kumar and Prabhakar, 2012; Lopez-Barneo et al., 2016). Single type I cells are exquisitely sensitive to O_2 and rapid (within ms) activation of the hypoxic chemotransduction cascade initiates at PO_2 s of between 20-40mmHg (Biscoe and Duchon, 1990; Buckler and Vaughan-Jones, 1994; Buckler and Turner, 2013); a level considerably greater than that at which cell metabolism is affected in O_2 -insensitive cells.

What remains highly controversial is the molecular identity of the specific O_2 sensor within the type I cell. We would argue that the physiological O_2 sensor should exhibit certain key features: 1. expression in the type I cell permitting intrinsic O_2 sensitivity; 2. the ability to bind O_2 ; 3. its binding of O_2 occurs over the physiological range at which the type I cell is stimulated; 4. it is required for the CB to be stimulated by hypoxia; and 5. it is able to activate the CB transduction cascade within milliseconds. Many proposed sensors fit one or two of these criteria but few have been shown to adequately comply with all five.

In mammalian cells, O_2 is the terminal electron acceptor in the mitochondrial respiratory chain. Continuous binding and reduction of O_2 in the CuB/haem a_3 (cytochrome a_3) binuclear center of complex IV drives mitochondrial electron transport and promotes activation of the mitochondrial ATP synthase. The long-established, mitochondrial hypothesis for chemoreception proposes that CB excitation induced by hypoxia is initiated by a reduction in O_2 dependent mitochondrial energy respiration. This review will briefly critique the current evidence as to whether the mitochondria can be considered the functionally relevant O_2 sensors within the CB.

2 Mitochondrial inhibitors mimic all aspects of the carotid body hypoxic chemotransduction cascade

All mitochondrial poisons induce chemoafferent excitation (Krylov and Anichkov, 1968; Mulligan et al., 1981; Obeso et al., 1989; Donnelly et al., 2014; Holmes et al., 2016; Holmes et al., 2017) leading to rapid increases in ventilation (Owen and Gesell, 1931), heart rate and arterial blood glucose (Alvarez-Buylla and de Alvarez-Buylla, 1988). Chemoafferent responses are rapid, dose dependent and reversible (Donnelly et al., 2014; Holmes et al., 2016) and the magnitude of the rise in chemoafferent frequency caused by saturating concentrations is consistent with those evoked by severe hypoxia or anoxia (Krylov and Anichkov, 1968; Mulligan et al., 1981; Obeso et al., 1989). Furthermore, mitochondrial inhibitors and uncouplers augment neurotransmitter secretion, confirming an action through the type I cell rather than the afferent nerve endings (Obeso et al., 1989; Rocher et al., 1991). Despite the strong consistency between all of the different types of mitochondrial poisons (both in the older and more recent

studies), it should be noted that some of the pharmacological agents used may not have acted selectively on the mitochondria and so the conclusions should be viewed with a certain degree of caution.

Mitochondrial poisons cause fast (within ms) type I cell depolarization and increases in $[Ca^{2+}]_i$. The size and timing of the $[Ca^{2+}]_i$ rise observed using many different mitochondrial inhibitors or uncouplers closely resembles that seen in hypoxia (Biscoe et al., 1989; Biscoe and Duchen, 1990; Wyatt and Buckler, 2004; Buckler, 2011). As with hypoxia (Buckler and Vaughan-Jones, 1994; Urena et al., 1994), the increases in $[Ca^{2+}]_i$ are dependent on cellular depolarization and extracellular Ca^{2+} influx through voltage gated Ca^{2+} channels (Wyatt and Buckler, 2004). TASK1/3, TREK-1, BK_{Ca} , $K_v4.1$ and $K_v4.3$, have all been shown to be expressed in the CB and inhibited by hypoxia (Buckler et al., 2000; Sanchez et al., 2002; Williams et al., 2004; Yamamoto and Taniguchi, 2006; Kim et al., 2009; Kreneisz et al., 2009; Turner and Buckler, 2013; Wang et al., 2017b). Of these, TASK1/3 and TASK-like K^+ currents are diminished by a multitude of mitochondrial inhibitors leading to membrane depolarization (Barbe et al., 2002; Wyatt and Buckler, 2004; Buckler, 2011; Turner and Buckler, 2013; Kim et al., 2015a).

Recent evidence has revealed the presence of TRP and other non-selective Ca^{2+} -activated cation currents in type I cells that are activated by hypoxia (Kumar et al., 2006; Kang et al., 2014; Kim et al., 2015b; Wang et al., 2017a). Although intriguing, the full functional relevance of these currents in type I cell O_2 -sensing remains to be further characterized, and in particular whether these currents can be upregulated to preserve O_2 sensing in the absence of TASK channels (Turner and Buckler, 2013). Current evidence suggests that mitochondrial inhibitors are also capable of increasing these inward depolarizing currents (Kim et al., 2015b; Wang et al., 2017a).

3 Mitochondria are necessary for the carotid body to respond to hypoxia

The presence of functional mitochondria does appear necessary for the CB to respond fully to hypoxia. For instance, cyanide, rotenone and FCCP all attenuate TASK channel currents in such a way that prevents any further reduction by hypoxia (Wyatt and Buckler, 2004). In intact CB preparations oligomycin, cyanide and azide all reduce or abolish subsequent chemoafferent responses to hypoxia (Mulligan et al., 1981; Donnelly et al., 2014). Some of the attenuation observed in these experiments may have been due to impairment of oxidative phosphorylation in the chemoafferent fibers, limiting their excitability. Any such impairment is not apparent in response to physiological levels of hypoxia, since the PO_2 that activates the type I cells occurs at a much higher level than those which would decrease mitochondrial function in the chemoafferent fibers. As such, chemoafferent responses to sustained hypoxia are better maintained than those in response to sustained high doses of mitochondrial inhibitors (Mulligan et al., 1981).

In a recent study the importance of mitochondrial complex I was tested by developing mice deficient in *Ndufs2* (a gene coding for NADH dehydrogenase [ubiquinone] iron-sulfur protein 2- a component of complex I that participates in ubiquinone binding) in tyrosine hydroxylase positive cells (Fernandez-Aguera et al., 2015). Type I cells isolated from these mice were insensitive to hypoxia; they lacked any hypoxia-induced K^+ current attenuation, $[Ca^{2+}]_i$ elevation or neurotransmitter release. Furthermore, these mice failed to increase respiratory frequency when breathing 10% O_2 . This work supported a previous study in which type I cell hypoxic chemosensitivity was abolished in the presence of rotenone (Ortega-Saenz et al., 2003). The authors propose a mechanism whereby exposure to hypoxia promotes reverse electron

transport and ROS/NADH generation via complex I which is driven by a high rate of succinate oxidation at complex II. Accordingly, they have recently shown that genetic and pharmacological deactivation of complex II completely blocks type I cell hypoxic sensitivity (Gao et al., 2017).

This intriguing and elegant hypothesis does however show some discrepancies with evidence from earlier reports. For instance, similar experiments performed on CBs with heterozygous *Sdh* knock out displayed an augmented, rather than depressed basal activity and had a completely preserved hypoxic response (Piruat et al., 2004). Furthermore, when rat type I cells were exposed to tetramethyl-p-phenylenediamine (TMPD) and ascorbate in the presence of rotenone, there was still a robust elevation in Ca^{2+} upon hypoxic stimulation (Wyatt and Buckler, 2004). This would suggest that feeding electrons into cytochrome c is sufficient to sustain type I cell hypoxic sensitivity even when complex I activity (and ROS generation) is inhibited. Complex IV activity rather than complex I and II may therefore be necessary for hypoxic chemotransduction. The same report also showed that application of H_2O_2 was unable to excite the type I cell directly. This observation is consistent with the lack of effect of multiple anti-oxidants used in other CB preparations and animal species (Sanz-Alfayate et al., 2001; Agapito et al., 2009; Gomez-Nino et al., 2009). Interestingly, using novel *ex vivo* CB culture techniques combined with FRET based ROS sensors, Bernardini and co-workers deduced that type I cell ROS actually decreases in hypoxia due to reduction in NADPH oxidase activity (an alternative ROS source) (Bernardini et al., 2015). Clearly there is a need for reconciliation between these findings.

4 The carotid body mitochondria are unique and have a low threshold for O_2

The evidence that mitochondria are required for CB O_2 chemotransduction and that mitochondrial inhibitors can cause chemoexcitation, is not enough to define them as the O_2 -sensors in the CB. Clearly, mitochondria are able to bind O_2 . However, the K_m of the cytochrome a_3 for O_2 is reported to be $<1\text{mmHg}$ in isolated mitochondria and between 1-5mmHg in dissociated cells and tissue preparations, with little variation existing between different cell types (Wilson et al., 1988; Tamura et al., 1989). This is far lower than the PO_2 at which the CB type I cells begin to be activated and, for this reason, is a common argument against the mitochondrial hypothesis.

However, there is now a substantial body of evidence indicating that the CB type I cell mitochondria are unique. Experiments performed by Mills & Jöbsis, were the first to identify an unusually low affinity cytochrome a_3 within the CB (Mills and Jobsis, 1970; 1972). Using absorbance spectra, they estimated that 43-67.5% of total cytochrome a_3 within the intact CB preparation had a remarkably low O_2 affinity. This fraction was reported to be almost 100% reduced at PO_2 s between 7-9mmHg and 50% reduced at a PO_2 as high as 90mmHg. In contrast, the remaining fraction was only 50% reduced at a PO_2 of approximately 0.8mmHg, comparable to cytochrome a_3 found in other tissues (Gnaiger, 2001). Thus, the CB appeared to express both low and high affinity subtypes of cytochrome a_3 . At that time, the specific cellular location(s) of each was unclear. Later experiments utilized the photolabile binding of CO, to deduce that saturation of cytochrome a_3 with CO prevented any additional chemoafferent excitation during hypoxia, implying that not only was the cytochrome a_3 in the CB unusual, it was also required for O_2 -sensing (Wilson et al., 1994; Lahiri et al., 1999). It should be pointed out that the concentrations of CO used in these studies could have directly modified the activity of the BK_{Ca} channel (Williams et al., 2004; Williams et al., 2008) and the generation of H_2S (Yuan et al., 2015) and as such some of the observations could be related to mechanisms independent of the mitochondria.

1
2 In dissociated rabbit type I cell clusters, mitochondrial electron transport begins to be inhibited
3 at a high PO₂ value of approximately 40mmHg (Duchen and Biscoe, 1992a). PO₂-NADH
4 response curves demonstrate a significant ‘right shift’ in type I cells compared to sensory
5 neurons, indicative of a heightened and distinctive O₂ sensitivity. In addition, mitochondrial
6 depolarization occurs at higher PO₂s compared to O₂-insensitive cells (Duchen and Biscoe,
7 1992b). More recent work has verified the “right shifts” in both PO₂-NADH and PO₂-rh-123
8 response curves in rat type I cells, confirming that the unusually low mitochondrial O₂ affinity
9 is conserved in multiple species (Turner and Buckler, 2013). By isolating complex IV activity
10 with a cocktail of mitochondrial inhibitors plus TMPD and ascorbate, the authors were able to
11 reveal that complex IV activity is a component of the mitochondria with the exceptionally low
12 O₂ affinity. Importantly, type I cell hypoxic response curves for electron transport inhibition,
13 mitochondrial depolarization and complex IV run-down display considerable overlap with the
14 rise in Ca²⁺, indicating that these processes are intimately linked. Therefore, it does appear that
15 type I cell mitochondria have a highly specialized *low affinity* for O₂ due to an altered
16 function/subtype of cytochrome a₃ in complex IV that predisposes CB energy metabolism to
17 being impaired at high O₂ tensions. It is likely that the high affinity cytochrome a₃ in the CB
18 described by Mills & Jöbsis is located in the non-O₂ sensing tissue such as the, nerve endings,
19 blood vessels and type II cells.

20
21 Understanding the mechanism linking a fall in mitochondrial O₂ consumption with K⁺ channel
22 inhibition (or cation channel activation) is contentious. As previously mentioned, there could be
23 a role for elevated mitochondrial ROS generation but this is still to be validated (Fernandez-
24 Aguera et al., 2015). Another possibility is an alteration in cytosolic nucleotides. Switching
25 from a cell attached to inside-out patch configuration diminishes background K⁺ channel
26 activity, suggesting that a basal level of an intracellular factor(s) activates TASK channels in
27 normoxia (Varas et al., 2007). Addition of 5mM MgATP in the inside-out configuration can
28 restore about 50% of this background K⁺ channel activity. Both mitochondrial inhibition and
29 hypoxia also significantly elevate free Mg²⁺, consistent with a decrease in MgATP. Thus, the
30 fall in MgATP during hypoxia is likely to attenuate a significant proportion of TASK-current
31 leading to depolarization. However, the remaining modulators that account for the other 50% of
32 TASK current are still to be identified.

33
34 Another proposed mediator of TASK channel activity that is sensitive to changes in cytosolic
35 nucleotide concentrations is AMPK (Wyatt et al., 2007). However, initial favorable studies
36 based on pharmacological evidence have since been challenged by the finding that the AMPK-
37 α₁α₂ deficient CB retains complete O₂-sensitivity (Mahmoud et al., 2016). Other groups have
38 also shown that pharmacological targeting of AMPK does not impact on the hypoxia-induced
39 K⁺ channel inhibition (Kim et al., 2014). Discrepancies may arise from the non-selectivity of
40 the drugs used to evaluate AMPK function and potential redundancy mechanisms known to
41 develop in genetic animal models (Nowak et al., 1997). A final hypothesis is that a build-up in
42 lactate upon mitochondrial inhibition in hypoxia activates the olfactory receptor Olfr78 (Chang
43 et al., 2015). However, the concentration of lactate necessary to elevate Ca²⁺ in an intact CB
44 preparation appears to be quite high (30mM) and whether local levels reach this threshold
45 during hypoxia is uncertain. We await a mechanism demonstrating how activation of Olfr78 (a
46 G-protein-coupled receptor) modulates TASK or cation channel activity. A summary of the
47 proposed O₂-sensitive mitochondrial signaling pathways in the CB is presented in Figure 1.

48 49 **5 What determines the low O₂ affinity of the carotid body mitochondria?**

1 We propose that there are 2 potential means to account for the extraordinary O₂ sensitivity of
2 type I cell mitochondria. First, there could be a high level of production of a cytosolic factor
3 that is able to freely diffuse into the mitochondria and then compete with O₂ binding in complex
4 IV (Figure 2). We predict that this competition would render mitochondrial electron transport
5 more susceptible to subsequent falls in O₂. We recently tested this by applying exogenous nitrite
6 to CBs and subsequently measuring hypoxic sensitivity (Holmes et al., 2016). Nitrite is reduced
7 within the mitochondria to generate local NO, a recognized competitive inhibitor of complex IV
8 (Brown and Cooper, 1994; Cleeter et al., 1994; Castello et al., 2006; Basu et al., 2008; Castello
9 et al., 2008). Moderate basal inhibition of the CB mitochondria by nitrite exaggerated the
10 subsequent chemoafferent excitation during hypoxia signifying an increase in CB O₂ sensitivity.
11 Therefore, we validated the idea that CB hypoxic sensitivity could be adjusted by a factor
12 capable of competing with O₂ in the mitochondria and suggested a physiological role for
13 endogenous NO in establishing type I cell mitochondrial O₂-sensitivity. Measurable amounts of
14 NO have been detected in mitochondrial of type I cells (Yamamoto et al., 2006). A possible
15 source is nitric oxide synthase 3 (NOS-3) given its location within the type I cell (Yamamoto et
16 al., 2006). Interestingly, mice with reduced NOS-3 have a dampened hypoxic ventilatory
17 response and a depressed CB function (Kline et al., 2000). One explanation for this is an
18 adaptation to chronic hypoxia brought about by reduced CB blood flow. However, this is
19 unlikely as there is no significant type I cell hyperplasia/hypertrophy (McGregor et al., 1984;
20 Tatsumi et al., 1991). Instead, the blunted CB activity could be due to the lack of NO acting on
21 the CB mitochondria. Consideration of the precise compartmentalization of NO should also be
22 taken into account. Whilst NO in the mitochondria induces chemostimulation, its action in other
23 regions is likely to have opposing effects via modulation of soluble guanylate cyclase and L-
24 type Ca²⁺/BK_{Ca} channels (Summers et al., 1999; Iturriaga et al., 2000; Silva and Lewis, 2002;
25 Valdes et al., 2003). In addition, other diffusible cytosolic factors have been implicated in CB
26 O₂ sensing including H₂S and CO (Peng et al., 2010; Yuan et al., 2015). Both of these gases are
27 capable of inhibiting type I cell mitochondria (Wilson et al., 1994; Lahiri et al., 1999; Buckler,
28 2011). Future experiments are required to evaluate if these substances act by setting type I cell
29 mitochondrial O₂ sensitivity.

30
31 A second explanation for the low O₂-affinity of the type I cell mitochondria is that it has a
32 unique gene expression profile (Figure 2). Exploring gene expression in the CB is challenging
33 due to its relatively small size and heterogeneity. However, advances in molecular biology
34 techniques now make it possible to perform whole genome analysis using just micrograms of
35 tissue or even single cells. RNA-sequencing analysis has now revealed a number of
36 mitochondrial related genes that have a particularly high expression in the type I cell (Zhou et
37 al., 2016). Of these *Ndufa4l2* and *Cox4i2* have been shown to be more highly expressed in the
38 CB compared with tissue from the superior cervical ganglion (Gao et al., 2017). Whether these
39 two genes contribute to the low mitochondrial O₂ affinity remains to be determined but these
40 findings do support the idea that CB mitochondria have a unique genetic signature encoding
41 their mitochondrial complexes. We would expect many more genetic studies to probe this
42 further until the type I cell mitochondria can be accurately modelled to pinpoint the structural
43 conformation underlying its low O₂ affinity. An interesting comparator may be the
44 mitochondria isolated from guinea pig CB which does not appear to have any inherent O₂
45 sensitivity (Gonzalez-Obeso et al., 2017).

46 47 **7 Conclusion**

48 On current evidence, it is very hard to disprove the mitochondrial hypothesis of CB O₂ sensing.
49 The mitochondria seem to fulfil all five criteria that we have proposed for adequate O₂-sensors.
50 What is less clear is a mechanistic understanding of how the low O₂ sensitivity of the CB

1 mitochondria is achieved and if mitochondria are involved in establishing pathological changes
2 in CB function.

3

4

References

- Agapito, M.T., Sanz-Alfayate, G., Gomez-Nino, A., Gonzalez, C., and Obeso, A. (2009). General redox environment and carotid body chemoreceptor function. *Am J Physiol Cell Physiol* 296(3), C620-631. doi: 00542.2008 [pii] 10.1152/ajpcell.00542.2008.
- Alvarez-Buylla, R., and de Alvarez-Buylla, E.R. (1988). Carotid sinus receptors participate in glucose homeostasis. *Respir Physiol* 72(3), 347-359.
- Barbe, C., Al-Hashem, F., Conway, A.F., Dubuis, E., Vandier, C., and Kumar, P. (2002). A possible dual site of action for carbon monoxide-mediated chemoexcitation in the rat carotid body. *J Physiol* 543(Pt 3), 933-945. doi: PHY_015750 [pii].
- Basu, S., Azarova, N.A., Font, M.D., King, S.B., Hogg, N., Gladwin, M.T., et al. (2008). Nitrite reductase activity of cytochrome c. *J Biol Chem* 283(47), 32590-32597. doi: M806934200 [pii] 10.1074/jbc.M806934200.
- Bernardini, A., Brockmeier, U., Metzen, E., Berchner-Pfannschmidt, U., Harde, E., Acker-Palmer, A., et al. (2015). Type I cell ROS kinetics under hypoxia in the intact mouse carotid body ex vivo: a FRET-based study. *Am J Physiol Cell Physiol* 308(1), C61-67. doi: 10.1152/ajpcell.00370.2013.
- Biscoe, T.J., and Duchen, M.R. (1990). Responses of type I cells dissociated from the rabbit carotid body to hypoxia. *J Physiol* 428, 39-59.
- Biscoe, T.J., Duchen, M.R., Eisner, D.A., O'Neill, S.C., and Valdeolmillos, M. (1989). Measurements of intracellular Ca²⁺ in dissociated type I cells of the rabbit carotid body. *J Physiol* 416, 421-434.
- Brown, G.C., and Cooper, C.E. (1994). Nanomolar concentrations of nitric oxide reversibly inhibit synaptosomal respiration by competing with oxygen at cytochrome oxidase. *FEBS Lett* 356(2-3), 295-298. doi: 0014-5793(94)01290-3 [pii].
- Buckler, K.J. (2011). Effects of exogenous hydrogen sulphide on calcium signalling, background (TASK) K channel activity and mitochondrial function in chemoreceptor cells. *Pflugers Arch* 463(5), 743-754. doi: 10.1007/s00424-012-1089-8.
- Buckler, K.J., and Turner, P.J. (2013). Oxygen sensitivity of mitochondrial function in rat arterial chemoreceptor cells. *J Physiol* 591(Pt 14), 3549-3563. doi: jphysiol.2013.257741 [pii] 10.1113/jphysiol.2013.257741.
- Buckler, K.J., and Vaughan-Jones, R.D. (1994). Effects of hypoxia on membrane potential and intracellular calcium in rat neonatal carotid body type I cells. *J Physiol* 476(3), 423-428.
- Buckler, K.J., Williams, B.A., and Honore, E. (2000). An oxygen-, acid- and anaesthetic-sensitive TASK-like background potassium channel in rat arterial chemoreceptor cells. *J Physiol* 525 Pt 1, 135-142. doi: PHY_0588 [pii].
- Castello, P.R., David, P.S., McClure, T., Crook, Z., and Poyton, R.O. (2006). Mitochondrial cytochrome oxidase produces nitric oxide under hypoxic conditions: implications for oxygen sensing and hypoxic signaling in eukaryotes. *Cell Metab* 3(4), 277-287. doi: S1550-4131(06)00071-4 [pii] 10.1016/j.cmet.2006.02.011.
- Castello, P.R., Woo, D.K., Ball, K., Wojcik, J., Liu, L., and Poyton, R.O. (2008). Oxygen-regulated isoforms of cytochrome c oxidase have differential effects on its nitric oxide production and on hypoxic signaling. *Proc Natl Acad Sci U S A* 105(24), 8203-8208. doi: 0709461105 [pii] 10.1073/pnas.0709461105.

- 1 Chang, A.J., Ortega, F.E., Riegler, J., Adison, D.V.M., and Krasnow, M.A. (2015). Oxygen
2 regulation of breathing through an olfactory receptor activated by lactate. *Nature*
3 527(7577), 240-+. doi: 10.1038/nature15721.
- 4 Cleeter, M.W., Cooper, J.M., Darley-USmar, V.M., Moncada, S., and Schapira, A.H. (1994).
5 Reversible inhibition of cytochrome c oxidase, the terminal enzyme of the mitochondrial
6 respiratory chain, by nitric oxide. Implications for neurodegenerative diseases. *FEBS*
7 *Lett* 345(1), 50-54. doi: 0014-5793(94)00424-2 [pii].
- 8 Donnelly, D.F., Kim, I., Mulligan, E.M., and Carroll, J.L. (2014). Non-additive interactions
9 between mitochondrial complex IV blockers and hypoxia in rat carotid body responses.
10 *Respir Physiol Neurobiol* 190, 62-69. doi: 10.1016/j.resp.2013.09.009.
- 11 Duchen, M.R., and Biscoe, T.J. (1992a). Mitochondrial function in type I cells isolated from
12 rabbit arterial chemoreceptors. *J Physiol* 450, 13-31.
- 13 Duchen, M.R., and Biscoe, T.J. (1992b). Relative mitochondrial membrane potential and
14 $[Ca^{2+}]_i$ in type I cells isolated from the rabbit carotid body. *J Physiol* 450, 33-61.
- 15 Fernandez-Aguera, M.C., Gao, L., Gonzalez-Rodriguez, P., Pintado, C.O., Arias-Mayenco, I.,
16 Garcia-Flores, P., et al. (2015). Oxygen Sensing by Arterial Chemoreceptors Depends
17 on Mitochondrial Complex I Signaling. *Cell Metab* 22(5), 825-837. doi:
18 10.1016/j.cmet.2015.09.004.
- 19 Gao, L., Bonilla-Henao, V., Garcia-Flores, P., Arias-Mayenco, I., Ortega-Saenz, P., and Lopez-
20 Barneo, J. (2017). Gene expression analyses reveal metabolic specifications in acute O-
21 2-sensing chemoreceptor cells. *Journal of Physiology-London* 595(18), 6091-6120. doi:
22 10.1113/jp274684.
- 23 Gnaiger, E. (2001). Bioenergetics at low oxygen: dependence of respiration and
24 phosphorylation on oxygen and adenosine diphosphate supply. *Respiration Physiology*
25 128(3), 277-297. doi: 10.1016/s0034-5687(01)00307-3.
- 26 Gomez-Nino, A., Agapito, M.T., Obeso, A., and Gonzalez, C. (2009). Effects of mitochondrial
27 poisons on glutathione redox potential and carotid body chemoreceptor activity. *Respir*
28 *Physiol Neurobiol* 165(1), 104-111. doi: S1569-9048(08)00298-X [pii]
29 10.1016/j.resp.2008.10.020.
- 30 Gonzalez-Obeso, E., Docio, I., Olea, E., Cogolludo, A., Obeso, A., Rocher, A., et al. (2017).
31 Guinea Pig Oxygen-Sensing and Carotid Body Functional Properties. *Frontiers in*
32 *Physiology* 8, 12. doi: 10.3389/fphys.2017.00285.
- 33 Holmes, A.P., Ray, C.J., Pearson, S.A., Coney, A.M., and Kumar, P. (2017). Ecto-5'-
34 nucleotidase (CD73) regulates peripheral chemoreceptor activity and cardiorespiratory
35 responses to hypoxia. *J Physiol*. doi: 10.1113/JP274498.
- 36 Holmes, A.P., Turner, P.J., Buckler, K.J., and Kumar, P. (2016). Moderate inhibition of
37 mitochondrial function augments carotid body hypoxic sensitivity. *Pflügers Archiv-*
38 *European Journal of Physiology* 468(1), 143-155. doi: 10.1007/s00424-015-1745-x.
- 39 Iturriaga, R., Villanueva, S., and Mosqueira, M. (2000). Dual effects of nitric oxide on cat
40 carotid body chemoreception. *J Appl Physiol* 89(3), 1005-1012.
- 41 Kang, D., Wang, J., Hogan, J.O., Vennekens, R., Freichel, M., White, C., et al. (2014). Increase
42 in cytosolic Ca^{2+} produced by hypoxia and other depolarizing stimuli activates a non-
43 selective cation channel in chemoreceptor cells of rat carotid body. *J Physiol* 592(Pt 9),
44 1975-1992. doi: 10.1113/jphysiol.2013.266957.
- 45 Kim, D., Cavanaugh, E.J., Kim, I., and Carroll, J.L. (2009). Heteromeric TASK-1/TASK-3 is
46 the major oxygen-sensitive background K^+ channel in rat carotid body glomus cells. *J*
47 *Physiol* 587(Pt 12), 2963-2975. doi: jphysiol.2009.171181 [pii]
48 10.1113/jphysiol.2009.171181.
- 49 Kim, D., Kang, D., Martin, E.A., Kim, I., and Carroll, J.L. (2014). Effects of modulators of
50 AMP-activated protein kinase on TASK-1/3 and intracellular Ca^{2+} concentration in

- rat carotid body glomus cells. *Respir Physiol Neurobiol* 195, 19-26. doi: 10.1016/j.resp.2014.01.020.
- Kim, D., Kim, I., Wang, J., White, C., and Carroll, J.L. (2015a). Hydrogen sulfide and hypoxia-induced changes in TASK (K2P3/9) activity and intracellular Ca(2+) concentration in rat carotid body glomus cells. *Respir Physiol Neurobiol* 215, 30-38. doi: 10.1016/j.resp.2015.04.012.
- Kim, I., Fite, L., Donnelly, D.F., Kim, J.H., and Carroll, J.L. (2015b). "Possible Role of TRP Channels in Rat Glomus Cells," in *Arterial Chemoreceptors in Physiology and Pathophysiology*, eds. C. Peers, P. Kumar, C.N. Wyatt, E. Gauda, C.A. Nurse & N. Prabhakar. (Berlin: Springer-Verlag Berlin), 227-232.
- Kline, D.D., Yang, T., Premkumar, D.R., Thomas, A.J., and Prabhakar, N.R. (2000). Blunted respiratory responses to hypoxia in mutant mice deficient in nitric oxide synthase-3. *J Appl Physiol* 88(4), 1496-1508.
- Kreneisz, O., Benoit, J.P., Bayliss, D.A., and Mulkey, D.K. (2009). AMP-activated protein kinase inhibits TREK channels. *J Physiol* 587(Pt 24), 5819-5830. doi: jphysiol.2009.180372 [pii] 10.1113/jphysiol.2009.180372.
- Krylov, S.S., and Anichkov, S.V. (1968). The effect of metabolic inhibitors on carotid chemoreceptors. *Arterial Chemoreceptors*, 103-109.
- Kumar, P. (2009). Systemic effects resulting from carotid body stimulation-invited article. *Adv Exp Med Biol* 648, 223-233. doi: 10.1007/978-90-481-2259-2_26.
- Kumar, P., and Bin-Jaliah, I. (2007). Adequate stimuli of the carotid body: more than an oxygen sensor? *Respir Physiol Neurobiol* 157(1), 12-21. doi: S1569-9048(07)00011-0 [pii] 10.1016/j.resp.2007.01.007.
- Kumar, P., Pearson, S., and Gu, Y.C. (2006). A role for TRP channels in carotid body chemotransduction? *Faseb Journal* 20(5), A1229-A1229.
- Kumar, P., and Prabhakar, N.R. (2012). Peripheral Chemoreceptors: Function and Plasticity of the Carotid Body. *Comprehensive Physiology*, 141-219.
- Lahiri, S., Ehleben, W., and Acker, H. (1999). Chemoreceptor discharges and cytochrome redox changes of the rat carotid body: role of heme ligands. *Proc Natl Acad Sci U S A* 96(16), 9427-9432.
- Lopez-Barneo, J., Gonzalez-Rodriguez, P., Gao, L., Fernandez-Aguera, M.C., Pardal, R., and Ortega-Saenz, P. (2016). Oxygen sensing by the carotid body: mechanisms and role in adaptation to hypoxia. *American Journal of Physiology-Cell Physiology* 310(8), C629-C642. doi: 10.1152/ajpcell.00265.2015.
- Mahmoud, A.D., Lewis, S., Juricic, L., Udoh, U.A., Hartmann, S., Jansen, M.A., et al. (2016). AMP-activated Protein Kinase Deficiency Blocks the Hypoxic Ventilatory Response and Thus Precipitates Hypoventilation and Apnea. *American Journal of Respiratory and Critical Care Medicine* 193(9), 1032-1043. doi: 10.1164/rccm.201508-1667OC.
- McGregor, K.H., Gil, J., and Lahiri, S. (1984). A morphometric study of the carotid body in chronically hypoxic rats. *J Appl Physiol* 57(5), 1430-1438.
- Mills, E., and Jobsis, F.F. (1970). Simultaneous measurement of cytochrome a3 reduction and chemoreceptor afferent activity in the carotid body. *Nature* 225(5238), 1147-1149.
- Mills, E., and Jobsis, F.F. (1972). Mitochondrial respiratory chain of carotid body and chemoreceptor response to changes in oxygen tension. *J Neurophysiol* 35(4), 405-428.
- Mulligan, E., Lahiri, S., and Storey, B.T. (1981). Carotid body O2 chemoreception and mitochondrial oxidative phosphorylation. *J Appl Physiol* 51(2), 438-446.
- Nowak, M.A., Boerlijst, M.C., Cooke, J., and Smith, J.M. (1997). Evolution of genetic redundancy. *Nature* 388(6638), 167-171. doi: 10.1038/40618.

- 1 Obeso, A., Almaraz, L., and Gonzalez, C. (1989). Effects of cyanide and uncouplers on
2 chemoreceptor activity and ATP content of the cat carotid body. *Brain Res* 481(2), 250-
3 257.
- 4 Ortega-Saenz, P., Pardal, R., Garcia-Fernandez, M., and Lopez-Barneo, J. (2003). Rotenone
5 selectively occludes sensitivity to hypoxia in rat carotid body glomus cells. *J Physiol*
6 548(Pt 3), 789-800. doi: 10.1113/jphysiol.2003.039693
7 2003.039693 [pii].
- 8 Owen, H., and Gesell, R. (1931). Peripheral and central chemical control of pulmonary
9 ventilation. *Proceedings of the Society for Experimental Biology and Medicine* 28(7),
10 0765-0766.
- 11 Peng, Y.J., Nanduri, J., Raghuraman, G., Souvannakitti, D., Gadalla, M.M., Kumar, G.K., et al.
12 (2010). H2S mediates O2 sensing in the carotid body. *Proc Natl Acad Sci U S A*
13 107(23), 10719-10724.
- 14 Piruat, J.I., Pintado, C.O., Ortega-Saenz, P., Roche, M., and Lopez-Barneo, J. (2004). The
15 mitochondrial SDHD gene is required for early embryogenesis, and its partial deficiency
16 results in persistent carotid body glomus cell activation with full responsiveness to
17 hypoxia. *Mol Cell Biol* 24(24), 10933-10940. doi: 24/24/10933 [pii]
18 10.1128/MCB.24.24.10933-10940.2004.
- 19 Ribeiro, M.J., Sacramento, J.F., Gonzalez, C., Guarino, M.P., Monteiro, E.C., and Conde, S.V.
20 (2013). Carotid body denervation prevents the development of insulin resistance and
21 hypertension induced by hypercaloric diets. *Diabetes* 62(8), 2905-2916.
- 22 Rocher, A., Obeso, A., Gonzalez, C., and Herreros, B. (1991). Ionic mechanisms for the
23 transduction of acidic stimuli in rabbit carotid body glomus cells. *J Physiol* 433, 533-
24 548.
- 25 Sanchez, D., Lopez-Lopez, J.R., Perez-Garcia, M.T., Sanz-Alfayate, G., Obeso, A., Ganfornina,
26 M.D., et al. (2002). Molecular identification of K α subunits that contribute to the
27 oxygen-sensitive K $^{+}$ current of chemoreceptor cells of the rabbit carotid body. *J Physiol*
28 542(Pt 2), 369-382. doi: PHY_18382 [pii].
- 29 Sanz-Alfayate, G., Obeso, A., Agapito, M.T., and Gonzalez, C. (2001). Reduced to oxidized
30 glutathione ratios and oxygen sensing in calf and rabbit carotid body chemoreceptor
31 cells. *J Physiol* 537(Pt 1), 209-220. doi: PHY_12621 [pii].
- 32 Silva, J.M., and Lewis, D.L. (2002). Nitric oxide enhances Ca(2+)-dependent K(+) channel
33 activity in rat carotid body cells. *Pflugers Arch* 443(5-6), 671-675. doi: 10.1007/s00424-
34 001-0745-1.
- 35 Summers, B.A., Overholt, J.L., and Prabhakar, N.R. (1999). Nitric oxide inhibits L-type Ca $^{2+}$
36 current in glomus cells of the rabbit carotid body via a cGMP-independent mechanism. *J*
37 *Neurophysiol* 81(4), 1449-1457.
- 38 Tamura, M., Hazeki, O., Nioka, S., and Chance, B. (1989). In vivo study of tissue oxygen
39 metabolism using optical and nuclear magnetic resonance spectroscopies. *Annu Rev*
40 *Physiol* 51, 813-834. doi: 10.1146/annurev.ph.51.030189.004121.
- 41 Tatsumi, K., Pickett, C.K., and Weil, J.V. (1991). Attenuated carotid body hypoxic sensitivity
42 after prolonged hypoxic exposure. *J Appl Physiol* 70(2), 748-755.
- 43 Thompson, E.L., Ray, C.J., Holmes, A.P., Pye, R.L., Wyatt, C.N., Coney, A.M., et al. (2016).
44 Adrenaline release evokes hyperpnoea and an increase in ventilatory CO $_2$ sensitivity
45 during hypoglycaemia: a role for the carotid body. *The Journal of physiology* 594(15),
46 4439-4452. doi: 10.1113/jp272191.
- 47 Turner, P.J., and Buckler, K.J. (2013). Oxygen and mitochondrial inhibitors modulate both
48 monomeric and heteromeric TASK-1 and TASK-3 channels in mouse carotid body type-
49 1 cells. *J Physiol* 591(Pt 23), 5977-5998. doi: 10.1113/jphysiol.2013.262022.

- 1 Urena, J., Fernandez-Chacon, R., Benot, A.R., Alvarez de Toledo, G.A., and Lopez-Barneo, J.
2 (1994). Hypoxia induces voltage-dependent Ca²⁺ entry and quantal dopamine secretion
3 in carotid body glomus cells. *Proc Natl Acad Sci U S A* 91(21), 10208-10211.
- 4 Valdes, V., Mosqueira, M., Rey, S., Del Rio, R., and Iturriaga, R. (2003). Inhibitory effects of
5 NO on carotid body: contribution of neural and endothelial nitric oxide synthase
6 isoforms. *Am J Physiol Lung Cell Mol Physiol* 284(1), L57-68. doi:
7 10.1152/ajplung.00494.2001
8 00494.2001 [pii].
- 9 Varas, R., Wyatt, C.N., and Buckler, K.J. (2007). Modulation of TASK-like background
10 potassium channels in rat arterial chemoreceptor cells by intracellular ATP and other
11 nucleotides. *J Physiol* 583(Pt 2), 521-536. doi: jphysiol.2007.135657 [pii]
12 10.1113/jphysiol.2007.135657.
- 13 Wang, J.J., Hogan, J.O., and Kim, D. (2017a). Voltage- and receptor-mediated activation of a
14 non-selective cation channel in rat carotid body glomus cells. *Respiratory Physiology &
15 Neurobiology* 237, 13-21. doi: 10.1016/j.resp.2016.12.005.
- 16 Wang, J.J., Hogan, J.O., Wang, R., White, C., and Kim, D. (2017b). Role of cystathionine-
17 gamma-lyase in hypoxia-induced changes in TASK activity, intracellular Ca²⁺ and
18 ventilation in mice. *Respiratory Physiology & Neurobiology* 246, 98-106. doi:
19 10.1016/j.resp.2017.08.009.
- 20 Williams, S.E., Brazier, S.P., Baban, N., Telezhkin, V., Muller, C.T., Riccardi, D., et al. (2008).
21 A structural motif in the C-terminal tail of slo1 confers carbon monoxide sensitivity to
22 human BK(Ca) channels. *Pflugers Archiv-European Journal of Physiology* 456(3), 561-
23 572. doi: 10.1007/s00424-007-0439-4.
- 24 Williams, S.E., Wootton, P., Mason, H.S., Bould, J., Iles, D.E., Riccardi, D., et al. (2004).
25 Hemoxxygenase-2 is an oxygen sensor for a calcium-sensitive potassium channel.
26 *Science* 306(5704), 2093-2097. doi: 1105010 [pii]
27 10.1126/science.1105010.
- 28 Wilson, D.F., Mokashi, A., Chugh, D., Vinogradov, S., Osanai, S., and Lahiri, S. (1994). The
29 primary oxygen sensor of the cat carotid body is cytochrome a3 of the mitochondrial
30 respiratory chain. *FEBS Lett* 351(3), 370-374. doi: 0014-5793(94)00887-6 [pii].
- 31 Wilson, D.F., Rumsey, W.L., Green, T.J., and Vanderkooi, J.M. (1988). The oxygen
32 dependence of mitochondrial oxidative phosphorylation measured by a new optical
33 method for measuring oxygen concentration. *J Biol Chem* 263(6), 2712-2718.
- 34 Wyatt, C.N., and Buckler, K.J. (2004). The effect of mitochondrial inhibitors on membrane
35 currents in isolated neonatal rat carotid body type I cells. *J Physiol* 556(Pt 1), 175-191.
36 doi: 10.1113/jphysiol.2003.058131
37 jphysiol.2003.058131 [pii].
- 38 Wyatt, C.N., Mustard, K.J., Pearson, S.A., Dallas, M.L., Atkinson, L., Kumar, P., et al. (2007).
39 AMP-activated protein kinase mediates carotid body excitation by hypoxia. *J Biol Chem*
40 282(11), 8092-8098. doi: M608742200 [pii]
41 10.1074/jbc.M608742200.
- 42 Yamamoto, Y., Konig, P., Henrich, M., Dedio, J., and Kummer, W. (2006). Hypoxia induces
43 production of nitric oxide and reactive oxygen species in glomus cells of rat carotid
44 body. *Cell Tissue Res* 325(1), 3-11. doi: 10.1007/s00441-006-0178-4.
- 45 Yamamoto, Y., and Taniguchi, K. (2006). Expression of tandem P domain K⁺ channel, TREK-
46 1, in the rat carotid body. *J Histochem Cytochem* 54(4), 467-472. doi: jhc.5A6755.2005
47 [pii]
48 10.1369/jhc.5A6755.2005.

1 Yuan, G., Vasavda, C., Peng, Y.J., Makarenko, V.V., Raghuraman, G., Nanduri, J., et al.
2 (2015). Protein kinase G-regulated production of H₂S governs oxygen sensing. *Sci*
3 *Signal* 8(373), ra37. doi: 10.1126/scisignal.2005846.
4 Zhou, T., Chien, M.-S., Kaleem, S., and Matsunami, H. (2016). Single cell transcriptome
5 analysis of mouse carotid body glomus cells. *The Journal of physiology* 594(15), 4225-
6 4251. doi: 10.1113/jp271936.
7
8

Figure Legends

Figure 1: Carotid body mitochondrial signaling mechanisms activated during hypoxia.

Hypoxia-induced mitochondrial inhibition is proposed to increase lactate generation (Chang et al., 2015), augment mitochondrial complex I reactive oxygen species (ROS) production (Fernandez-Aguera et al., 2015) or reduce mitochondrial ATP synthesis (Buckler and Turner, 2013). These changes are proposed to directly or indirectly (e.g. via Olf78 receptor activation, reduced MgATP concentration or stimulation of AMP-activated protein kinase; AMPK) modify ion channel function leading to resting membrane potential (RMP) depolarization. This causes opening of L-type Ca^{2+} channels, neurosecretion and an increase in discharge frequency of the adjacent chemoafferent fibers. TRP- transient receptor potential channel; NSC- non-selective cation channel; BK_{Ca} - large conductance Ca^{2+} -activated K^{+} channel; TASK-Twisk-related acid-sensitive K^{+} channel; TREK- TWIK-related K^{+} channel; AP-action potential.

Figure 2: Potential mechanisms underlying the unique low O_2 affinity mitochondria in the carotid body.

Inhibition of mitochondrial function in the carotid body type I cell occurs at PO_2 values well above those that inhibit metabolism in other O_2 insensitive cells. This activates a number of proposed signaling pathways (shown in blue). The unique low O_2 affinity of the carotid body mitochondria could be due to 1. a unique mitochondrial gene expression profile or 2. the presence of a competitive inhibitor such as NO (Holmes et al., 2016). The impact of lower mitochondrial O_2 affinity is to cause a right-shift in the PO_2 -chemoafferent response curve, thereby enhancing the functional O_2 sensitivity and allowing the carotid body to respond over a more physiological range of arterial PO_2 s. RET-reverse electron transport.


Acknowledgements

NA

Author contributions

APH, CJR, AMC and PK all contributed to the writing and editing of the manuscript.


Carotid body type I cell



1. Unique 'low O₂ affinity' mitochondrial gene expression profile

2. Presence of a competitive inhibitor e.g. NO

Type I cell mitochondrial electron transport chain



Whole carotid body chemoafferent O₂ sensitivity

