

Complete Closed Genome Sequence of Nontoxigenic Invasive *Corynebacterium diphtheriae* bv. *mitis* Strain ISS 3319

Azevedo Antunes, Camila; Richardson, Emily J.; Quick, Joshua; Fuentes-utrilla, Pablo; Isom, Georgia L.; Goodall, Emily C.; Möller, Jens; Hoskisson, Paul A.; Mattos-guaraldi, Ana Luiza; Cunningham, Adam F.; Loman, Nicholas J.; Sangal, Vartul; Burkovski, Andreas; Henderson, Ian R.

DOI:

[10.1128/genomeA.01566-17](https://doi.org/10.1128/genomeA.01566-17)

License:

Creative Commons: Attribution (CC BY)

Document Version

Publisher's PDF, also known as Version of record

Citation for published version (Harvard):

Azevedo Antunes, C, Richardson, EJ, Quick, J, Fuentes-utrilla, P, Isom, GL, Goodall, EC, Möller, J, Hoskisson, PA, Mattos-guaraldi, AL, Cunningham, AF, Loman, NJ, Sangal, V, Burkovski, A & Henderson, IR 2018, 'Complete Closed Genome Sequence of Nontoxigenic Invasive *Corynebacterium diphtheriae* bv. *mitis* Strain ISS 3319', *Genome Announcements*, vol. 6, no. 5, 01566-17, pp. e01566-17.
<https://doi.org/10.1128/genomeA.01566-17>

[Link to publication on Research at Birmingham portal](#)

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.



Complete Closed Genome Sequence of Nontoxigenic Invasive *Corynebacterium diphtheriae* bv. mitis Strain ISS 3319

Camila Azevedo Antunes,^{a,b} Emily J. Richardson,^c Joshua Quick,^c Pablo Fuentes-Utrilla,^c Georgia L. Isom,^c Emily C. Goodall,^c Jens Möller,^a  Paul A. Hoskisson,^d Ana Luiza Mattos-Guaraldi,^b  Adam F. Cunningham,^c Nicholas J. Loman,^c Vartul Sangal,^e  Andreas Burkovski,^a Ian R. Henderson^c

^aMicrobiology Division, Friedrich-Alexander University Erlangen, Nuremberg, Erlangen, Germany

^bLaboratory of Diphtheria and Corynebacteria of Clinical Relevance–LDCIC, Faculty of Medical Sciences, Rio de Janeiro State University, Rio de Janeiro, Brazil

^cInstitute of Microbiology and Infection, School of Biosciences, University of Birmingham, Birmingham, United Kingdom

^dStrathclyde Institute of Pharmacy and Biomedical Science, University of Strathclyde, Glasgow, United Kingdom

^eFaculty of Health and Life Sciences, Northumbria University, Newcastle upon Tyne, United Kingdom

ABSTRACT The genome sequence of the human pathogen *Corynebacterium diphtheriae* bv. mitis strain ISS 3319 was determined and closed in this study. The genome is estimated to have 2,404,936 bp encoding 2,257 proteins. This strain also possesses a plasmid of 1,960 bp.

Corynebacterium diphtheriae is the etiological agent of diphtheria, a toxigenic infection of the upper respiratory tract. Besides diphtheria, systemic infections caused by nontoxigenic strains have been reported (1–5). The molecular basis of toxin-independent pathogenicity of *C. diphtheriae* is poorly understood (6). Therefore, we sequenced the whole genome of *C. diphtheriae* bv. mitis ISS 3319, isolated from the throat of 9-year-old patient (7, 8). The strain lacks the *tox* gene but is able to adhere to and invade human epithelial cells and effectively colonize spleen and kidneys of mice (9–11).

Sequencing of libraries was carried out with the Oxford Nanopore MinION instrument and the Illumina HiSeq platform using a 250-bp paired-end protocol. The Illumina and MinION reads were assembled together using Unicycler with Pilon polishing, with a final hybrid assembly of two contigs of 1,604,547 and 800,125 bp. Two gaps of 124 bp and 140 bp were identified in the genome using BLAST searches of >500 bp from the ends of both the contigs against the published assembly of the same strain (GenBank accession no. JAQN000000000). The contigs were merged using the missing sequence after aligning them to the genome sequence of strain C7 (beta) (GenBank accession no. NC_016801), which belongs to the same sequence type, ST26, as ISS 3319 (7). Automatic annotation and identification of rRNAs and tRNAs were accomplished using the Prokka version 1.11 software tool (12).

The new sequence was compared to the previously published assembly of ISS 3319, which was also reannotated using Prokka to obtain an equivalence of annotation for comparative analysis. The annotated genomes were compared using Roary (13, 14); 2,195 genes are conserved between the assemblies, and only four single nucleotide polymorphisms (SNPs) were observed in the core genomic alignment. Most of the 26 genes reported to be specific to the previous assembly (GenBank accession no. JAQN000000000) showed significant similarities to several genes in the new assembly and may not be specific. The proteins encoded by the majority of these genes are short and fall within or near the identified genomic islands that were identified using IslandViewer 4 (15).

The closed genome of 2,404,936 bp contained 9 rRNAs, 53 tRNAs, 1 transfer-messenger RNA, 1 repeat region, 2,257 protein-coding sequences, and 53.46% GC

Received 15 December 2017 **Accepted** 19 December 2017 **Published** 1 February 2018

Citation Azevedo Antunes C, Richardson EJ, Quick J, Fuentes-Utrilla P, Isom GL, Goodall EC, Möller J, Hoskisson PA, Mattos-Guaraldi AL, Cunningham AF, Loman NJ, Sangal V, Burkovski A, Henderson IR. 2018. Complete closed genome sequence of nontoxigenic invasive *Corynebacterium diphtheriae* bv. mitis strain ISS 3319. Genome Announc 6:e01566-17. <https://doi.org/10.1128/genomeA.01566-17>.

Copyright © 2018 Azevedo Antunes et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Ian R. Henderson, i.r.henderson@bham.ac.uk.

content. Moreover, a circular plasmid of 1,960 bp with four coding sequences was also identified. A discontinuous mega-BLAST search of this plasmid sequence in GenBank revealed partial similarities (67% identity in 612-bp alignment) with plasmid pTA144 in *Moraxella* sp. (GenBank accession no. NC_001316) and the metamobilome of an uncultured strain (65% identity in 833-bp alignment; GenBank accession no. LN853682). A nonredundant translation alignment of the plasmid sequence showed a conserved domain with homology to the phage replication protein Cri, described as essential for DNA replication in *Vibrio cholerae* serotype O1 (16), and a sequence with 88% identity to a gene encoding an $\alpha\beta$ -hydrolase from a *Streptococcus* sp. (GenBank accession no. WP_002883881). Coinfections of *C. diphtheriae* with other pathogens, such as *Staphylococcus* spp., *Streptococcus* spp., *Arcanobacterium* spp., and *Pseudomonas* spp., were reported previously (8, 17, 18), and this close contact might support horizontal DNA transfer.

Accession number(s). The *C. diphtheriae* bv. mitis ISS 3319 whole-genome shotgun project has been deposited at DDBJ/EMBL/GenBank under the accession no. [CP025209](#) and [CP025210](#). The versions described in this paper are the second versions.

ACKNOWLEDGMENTS

Genome sequencing and bioinformatics were performed by MicrobesNG (<http://microbesng.uk>), which is supported by the BBSRC (grant BB/L024209/1). A.L.M.-G. and C.A.A. were supported by the Coordination for the Improvement of Higher Education Personnel (CAPES).

C.A.A. gratefully acknowledges the Förderung von Frauen in Forschung und Lehre (FFL) of the Friedrich-Alexander Universität Erlangen-Nürnberg.

REFERENCES

- Burkovski A. 2014. Diphtheria and its etiological agents, p 1–14. In Burkovski A (ed), *Corynebacterium diphtheriae* and related toxigenic species. Springer, Dordrecht, The Netherlands.
- Belchior E, Henry S, Badell E, Collet L, Benoit-Cattin T, de Montera AM, Guiso N, Patey O, Levy-Bruhl D, Filleul L, Chieze F, Olivier S. 2017. Diphtheria in Mayotte, 2007–2015. *Emerg Infect Dis* 23:1218–1220. <https://doi.org/10.3201/eid2307.170262>.
- Kolios AGA, Cozzio A, Zinkernagel AS, French LE, Kündig TM. 2017. Cutaneous *Corynebacterium* infection presenting with disseminated skin nodules and ulceration. *Case Rep Dermatol* 9:8–12. <https://doi.org/10.1159/000476054>.
- Santos LS, Sant'Anna LO, Ramos JN, Ladeira EM, Stavracakis-Peixoto R, Borges LL, Santos CS, Napoleão F, Camello TCF, Pereira GA, Hirata R, Vieira VV, Cosme LMSS, Sabbadini PS, Mattos-Guaraldi AL. 2015. Diphtheria outbreak in Maranhão, Brazil: microbiological, clinical and epidemiological aspects. *Epidemiol Infect* 143:791–798. <https://doi.org/10.1017/S0950268814001241>.
- Gomes DL, Martins CA, Faria LM, Santos LS, Santos CS, Sabbadini PS, Souza MC, Alves GB, Rosa AC, Nagao PE, Pereira GA, Hirata R, Mattos-Guaraldi AL. 2009. *Corynebacterium diphtheriae* as an emerging pathogen in nephrostomy catheter-related infection: evaluation of traits associated with bacterial virulence. *J Med Microbiol* 58:1419–1427. <https://doi.org/10.1099/jmm.0.012161-0>.
- Sangal V, Hoskisson PA. 2016. Evolution, epidemiology and diversity of *Corynebacterium diphtheriae*: new perspectives on an old foe. *Infect Genet Evol* 43:364–370. <https://doi.org/10.1016/j.meegid.2016.06.024>.
- Sangal V, Blom J, Sutcliffe IC, von Hunolstein C, Burkovski A, Hoskisson PA. 2015. Adherence and invasive properties of *Corynebacterium diphtheriae* strains correlates with the predicted membrane-associated and secreted proteome. *BMC Genomics* 16:765. <https://doi.org/10.1186/s12864-015-1980-8>.
- von Hunolstein C, Alfaroni G, Scopetti F, Pataracchia M, La Valle R, Franchi F, Pacciani L, Manera A, Giammanco A, Farinelli S, Engler K, De Zoysa A, Efstratiou A. 2003. Molecular epidemiology and characteristics of *Corynebacterium diphtheriae* and *Corynebacterium ulcerans* strains isolated in Italy during the 1990s. *J Med Microbiol* 52:181–188. <https://doi.org/10.1099/jmm.0.04864-0>.
- Ott L, Höller M, Gerlach RG, Hensel M, Rheinlaender J, Schäffer TE, Burkovski A. 2010. *Corynebacterium diphtheriae* invasion-associated protein (DIP1281) is involved in cell surface organization, adhesion and internalization in epithelial cells. *BMC Microbiol* 10:2. <https://doi.org/10.1186/1471-2180-10-2>.
- Ott L, Scholz B, Höller M, Hasselt K, Ensser A, Burkovski A. 2013. Induction of the NF- κ B signal transduction pathway in response to *Corynebacterium diphtheriae* infection. *Microbiology* 159:126–135. <https://doi.org/10.1099/mic.0.061879-0>.
- Puliti M, Von Hunolstein C, Marangi M, Bistoni F, Tissi L. 2006. Experimental model of infection with non-toxicogenic strains of *Corynebacterium diphtheriae* and development of septic arthritis. *J Med Microbiol* 55:229–235. <https://doi.org/10.1099/jmm.0.46135-0>.
- Seemann T. 2014. Prokka: rapid prokaryotic genome annotation. *Bioinformatics* 30:2068–2069. <https://doi.org/10.1093/bioinformatics/btu153>.
- Tange O. 2011. GNU parallel—the command-line power tool. *Linux: the USENIX magazine* 36:42–47.
- Page AJ, Cummins CA, Hunt M, Wong VK, Reuter S, Holden MT, Fookes M, Falush D, Keane JA, Parkhill J. 2015. Roary: rapid large-scale prokaryote pan genome analysis. *Bioinformatics* 31:3691–3693. <https://doi.org/10.1093/bioinformatics/btv421>.
- Bertelli C, Laird MR, Williams KP, Simon Fraser University Research, Lau BY, Hoar G, Winsor GL, Brinkman FS. 2017. IslandViewer 4: expanded prediction of genomic islands for larger-scale datasets. *Nucleic Acids Res* 45:W30–W35. <https://doi.org/10.1093/nar/gkx343>.
- Heidelberg JF, Eisen JA, Nelson WC, Clayton RA, Gwinn ML, Dodson RJ, Haft DH, Hickey EK, Peterson JD, Umayam L, Gill SR, Nelson KE, Read TD, Tettelin H, Richardson D, Ermolaeva MD, Vamathevan J, Bass S, Qin H, Dragoi I, Sellers P, McDonald L, Utterback T, Fleischmann RD, Nierman WC, White O, Salzberg SL, Smith HO, Colwell RR, Mekalanos JJ, Venter JC, Fraser CM. 2000. DNA sequence of both chromosomes of the cholera pathogen *Vibrio cholerae*. *Nature* 406:477–483. <https://doi.org/10.1038/35020000>.
- Gordon CL, Fagan P, Hennessy J, Baird R. 2011. Characterization of *Corynebacterium diphtheriae* isolates from infected skin lesions in the Northern Territory of Australia. *J Clin Microbiol* 49:3960–3962. <https://doi.org/10.1128/JCM.05038-11>.
- May ML, McDougall RJ, Robson JM. 2014. *Corynebacterium diphtheriae* and the returned tropical traveler. *J Travel Med* 21:39–44. <https://doi.org/10.1111/jtm.12074>.