

# The development of late-Holocene farmed landscapes

Smith, David; Hill, Geoff; Kenward, Harry

DOI:

[10.1177/0959683618804645](https://doi.org/10.1177/0959683618804645)

License:

None: All rights reserved

*Document Version*

Peer reviewed version

*Citation for published version (Harvard):*

Smith, D, Hill, G & Kenward, H 2018, 'The development of late-Holocene farmed landscapes: analysis of insect assemblages using a multi-period dataset', *The Holocene*, vol. 29, no. 1, pp. 45-63.  
<https://doi.org/10.1177/0959683618804645>

[Link to publication on Research at Birmingham portal](#)

**Publisher Rights Statement:**

(c) Sage 2018

Publication forthcoming in *Holocene*

**General rights**

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

**Take down policy**

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact [UBIRA@lists.bham.ac.uk](mailto:UBIRA@lists.bham.ac.uk) providing details and we will remove access to the work immediately and investigate.



## The Development of Late Holocene Farmed Landscapes: Analysis of Insect Assemblages using a Multi-period Dataset

Journal:	<i>The Holocene</i>
Manuscript ID	HOL-18-0023.R1
Manuscript Type:	Paper
Date Submitted by the Author:	n/a
Complete List of Authors:	Smith, David; University of Birmingham, Institute of Archaeology and Antiquity Kenward, Harry; University of York, Archaeology HILL, Geoff; University of Birmingham, Classics, Ancient History and Archaeology
Keywords:	Coleoptera, prehistoric farming, landscape development, field systems, econiche replacement, archaeoentomology
Abstract:	<p>Global agricultural intensification and expansion has led to the spread of a fairly cosmopolitan insect fauna associated with arable land and pasture. Studies of modern expansion and intensification of agriculture have shown profound effects in terms of declines in biodiversity, with implications for current nature conservation. However, modern entomological studies of farmland faunas do not consider if such effects occurred over a longer period of time or are merely a modern phenomenon. We examine the substantial British archaeoentomological dataset for the development of beetle (Coleoptera) faunas in a range of intensively farmed archaeological landscapes dating from the Late Neolithic through to the early Medieval period (c. 24,000 cal. BC – AD 900). The archaeological beetle fauna typically consisted of generalist species which still dominate modern farmland. Our analysis indicates that there is an essentially stable 'core group' of taxa that repeatedly occur regardless of period, location or the specific nature of the archaeological feature involved. On the basis of this result, we argue that the effects of the expansion of intensive farming on insect faunas seen in the modern world is a continuation of a longer pattern. We suggest that this is an example of human econiche replacement and ecosystem engineering. The approach taken here is applicable elsewhere in the world and we offer suggestions for future British and international research strategies.</p>

SCHOLARONE™  
Manuscripts

# The Development of Late Holocene Farmed Landscapes: Analysis of Insect Assemblages using a Multi-period Dataset

David Smith<sup>1</sup>, Geoff Hill<sup>1</sup> and Harry Kenward<sup>2</sup>

<sup>1</sup>Department of Classics, Ancient History and Archaeology, University of Birmingham, Edgbaston, Birmingham B15 2TT

<sup>2</sup> Department of Archaeology, The King's Manor, York, YO1 7EP

## Abstract

Global agricultural intensification and expansion has led to the spread of a fairly cosmopolitan insect fauna associated with arable land and pasture. Studies of modern expansion and intensification of agriculture have shown profound effects in terms of declines in biodiversity, with implications for current nature conservation. However, modern entomological studies of farmland faunas do not consider if such effects occurred over a longer period of time or are merely a modern phenomenon. We examine the substantial British archaeoentomological dataset for the development of beetle (Coleoptera) faunas in a range of intensively farmed archaeological landscapes dating from the Late Neolithic through to the early Medieval period (c. 24,000 cal. BC – AD 900). The archaeological beetle fauna typically consisted of generalist species which still dominate modern farmland. Our analysis indicates that there is an essentially stable 'core group' of taxa that repeatedly occur regardless of period, location or the specific nature of the archaeological feature involved. On the basis of this result, we argue that the effects of the expansion of intensive farming on insect faunas seen in the modern world is a continuation of a longer pattern. We suggest that this is an example of human econiche replacement and ecosystem engineering. The approach taken here is applicable elsewhere in the world and we offer suggestions for future British and international research strategies.

**Keywords:** Coleoptera, prehistoric farming, field systems, landscape development, econiche replacement, archaeoentomology

## Introduction

We live in a world where agricultural production, and consequently farmed landscapes, are expanding rapidly at the expense of natural 'wild' landscapes resulting in largescale loss of global biodiversity (e.g. Tilman, 1999, Foley et al. 2005; Tscharntke et al., 2005; Deininger and Byerlee, 2011; Zabel et al., 2014). The effects of modern agricultural intensification and expansion seems to be particularly dramatic and are presenting major issues for both agricultural production itself and also for nature conservation (e.g. Touhasca 1993, Holland and Reynolds 2003, Hutton and Giller 2003, Purtauf et al. 2005,). This decline in insect faunas as a result of farming is often portrayed as an essentially modern phenomenon rapidly approaching an 'ecological cliff edge' (Tilman 1999, Foley et al. 2005,

Denninger and Byerlee 2011;). However, there is little apparent understanding of how these effects may have operated over a longer timescale than that normally considered by ecologists (Ellis et al., 2013).

Many regions of the world currently do not have sufficient archaeoentomological data to address this issue; however, in Britain several decades of archaeoentomological research is available. Early work in southern Britain by Osborne (1978) and Robinson (1978, 1981a) described a number of Iron Age and Roman agricultural faunas which were dominated by a range of dung beetles, predatory ground beetles, click beetles and chafers. Both Robinson and Osborne noted the obvious contrast between these later prehistoric and Roman beetle faunas associated with farmed landscapes as compared with the woodland beetle faunas that dominate sites from the preceding, largely pre-agrarian, Mesolithic and Neolithic periods. Osborne (1978) suggested that post-Iron Age faunas indicated the presence of open grasslands, dominated by grazing animals and arable fields, and described this as 'culture-steppe'. In a later series of review papers, Robinson (e.g. 1983, 2000, 2003, 2013a, 2013b) outlined temporal changes in the insect faunas from a number of sites in the upper and lower Thames Valley. In particular, he observed a pattern such that many faunas from the Early Iron Age (c. 800 BC) onwards increasingly include taxa indicative of progressive woodland clearance and the development of pastureland, grassland and arable fields. Kenward (2009) also has produced similar results for the north of Britain.

These early studies often do not explicitly explore a number of issues surrounding the ecology, nature, origins and development of 'farmland beetle faunas'. More recently, as a result of developer-funded archaeology, many other archaeological sites associated with ditched and/or farmed landscapes have become available for study. The archaeoentomological data from these sites cover a wider range of archaeological periods, regions and deposit types, and most have a strong chronological dating programme.

This paper presents an analysis of the current dataset of archaeological Coleoptera (beetle) faunas from farmed landscapes (both arable and pastoral) dating between the Late Neolithic to Medieval (c. 2,4000 cal. BC – 900 AD) period in Britain. This analysis was undertaken to explore the following research questions:

- 1) What is the archaeological fauna recovered from farmed landscape? How does this compare to modern insect faunas from this habitat?
- 2) What is the ecology of the taxa that belong to this fauna?
- 3) To what extent do these faunas vary across periods, locations and by type of archaeological context? What other factors may account for any variation?
- 4) When did this fauna first appear in the archaeological record? What are the potential origins of this fauna and how quickly does it come to dominate in this changing landscape? What are the implications of this?

This paper also outlines a number of potential research directions for the study of insect faunas from archaeological farmland which are applicable worldwide.

## Archaeological background

The insect faunas considered come from 30 rural archaeological sites encompassing the British Neolithic (c. 4000–2200BC), Bronze Age (c. 2200-800 BC), Iron Age (800 BC – AD 47), Roman (AD 47 – 410) and early medieval periods (c. AD 900). Where sites are multiperiod, they have been divided into separate archaeological periods giving a total of 39 separate ‘chronological entities’. The location, dating, site type, details of publication and the number of samples analysed from the individual sites are presented in Table 1 and Figure 1. Where available, radiocarbon determinations for the sites mentioned are given in Table 1 and have been standardised and calibrated using the OxCAL programme (Reimer et al., 2013 OxCAL V.4.3, 2017; Bronk Ramsey 2009). Other chronology is based on archaeological finds or stratigraphic association.

Over the last 40 years our understanding of the development of the rural archaeological landscape of southern and central Britain has greatly changed. ‘Co-axial ditched field systems’ developed from around 1600 BC in upland areas; such as Dartmoor, Cumbria and the Pennines, and also in lower-lying areas such as Wessex, the North and South Downs, the Midlands and substantial areas of the Thames Valley (e.g. Fleming, 1988, 2007; Yates, 1999, 2007; Johnston, 2005; Chadwick, 2013; English, 2013). Earlier field systems have been primarily linked to stock rearing (Fleming, 1987; Yates, 1999; 2007; Fyfe et al., 2008, 2015). Ditched landscapes rapidly expanded in the Iron Age (c. 800 – 0 cal. BC) and the mid–Romano-British period (c. 100 – 200 cal. BC). These ditched field systems typically took the form of ‘strip fields’, ‘brick fields’ or ‘ladder enclosures’ and represent mixed farming landscapes incorporating arable fields, drove roads, waterholes and wells, the latter associated with livestock rearing (Riley, 1980; Taylor, 2007; Yates, 2007; Fyfe et al., 2008, 2015; Chadwick, 2013).

The development of such ditched agricultural landscapes at this period has often been attributed to an increase in agricultural production by a ‘top down’ autocratic authority responding to population rise and climatic downturn (i.e. Bradley, 1980; Fleming, 1987). However, alteration in landscape often is not synchronous with periods of dramatic climate change (Fyfe et al, 2015) and may relate to changing ideas of ownership and land tenure in response to social intensification and the production, exchange and consumption of elite materials (i.e. Brück, 2000; Johnston, 2005; Yates, 2007; Chadwick, 2013). The development of ditched farmlands appears to have coincided with a series of major changes to the British landscape (Huntley and Birks, 1983; Bell and Walker, 2005). Recent large-scale surveys of environmental archaeology by Dark (2006) and Fyfe and colleagues (2015) indicate that widespread clearance of woodland occurred in central and Southern England from around 1400–1200 BC, with a peak in clearance at around 500–200 BC. This loosely coincided with the onset of clay alluviation in most river systems, possibly in response to agricultural intensification (i.e. Macklin and Lewin, 2003; Macklin et al., 2005).

## Data collection and Analysis

### Sampling

1  
2  
3 Bulk samples of sediment were recovered from the sites listed in Table 1 during excavation from a  
4 range of archaeological features (mainly ditch, pit and well fills). Typically, the sample weight and  
5 volume ranged between 5–10 kg and 3–10 litres of sediment.  
6  
7

### 8 9 **Basic identification and quantification**

10 The bulk samples were prepared using the standard method of paraffin floatation outlined by Kenward  
11 and colleagues (1980). Waterlogged insect remains were sorted and identified under a low-power  
12 binocular microscope at magnifications between x15 and x45 by a number of different specialists  
13 (indicated in Table 1). Where achievable, the insect remains were identified to species level, often by  
14 direct comparison to specimens in reference collections. The nomenclature used in this paper follows  
15 Duff (2012) for the beetles and Stace (2010) for plants. The number of samples available per site  
16 varied from 1 to 22. The scores for the samples from each site have been amalgamated to generate a  
17 combined Minimum Number of Individuals (MNI) for each taxon. The full dataset consisted of 34,678  
18 individuals, representing 932 taxa.  
19  
20  
21  
22  
23  
24

### 25 **Analysis**

26 The degree of variation in the faunas over time, location and archaeological feature was analysed in  
27 the following ways:  
28  
29

30 **Ranked occurrence tables:** These were constructed for each of the main archaeological periods and  
31 were based on the MNI for each taxon. This analysis allowed the comparison between the most  
32 common taxa in each archaeological period and against modern faunas.  
33  
34

35 **Functional group analysis:** Taxa have been allocated to functional groups where possible (see Hill  
36 2015) and the relative proportions of these groupings were calculated. This analysis was undertaken  
37 to assess the extent to which the ecology of the faunas might differ between sites and periods.  
38

39 This approach is a revision of both Robinson's (1981a, 1983) 'landscape' and Kenward and Hall's  
40 (1995) synanthropic (associated with human activity) groupings. The functional group codes are  
41 displayed in Table 2. Ecological information comes from the Bugs CEP database (Buckland and  
42 Buckland 2006), Koch (1989, 1992) and the authors' field experience.  
43  
44

45 The aquatic and waterside ecological groupings are calculated as a percentage of the overall MNI for  
46 each 'chronological entity'. The percentages of terrestrial ecological groupings are calculated once the  
47 aquatic taxa have been removed from the MNI. In order to examine the extent of synanthropy  
48 (association with human settlement) shown by the faunas, two other summary statistics have been  
49 calculated for the terrestrial fauna:  
50  
51

- 52 1) The proportion of 'house fauna' recovered (a suite of beetles that have a particular  
53 affinity for human habitation and settlement – see Hall and Kenward, 1990; Smith,  
54 2012a)  
55
- 56 2) The Synanthropic Value (SV) has been calculated for the terrestrial fauna. In this  
57 statistic 'strongly' synanthropic species (*S*) carry more value (or weight) than 'typical'  
58  
59

(*T*) or weakly 'facultative' (*F*) species (Hall & Kenward 1990, Kenward, 1997; Hill, 2015) using this formula:  $SV = F + (2 \times T) + (3 \times S)$

**Similarity coefficients and Statistical ordination:** The degree of similarity of the faunas was assessed using Sørensen's similarity coefficient (Sørensen, 1948). This coefficient was chosen because it gives more weight to multiple occurrences and less weight to outliers, and its reciprocal is equivalent to Bray-Curtis distance (McCune and Grace, 2002; Borcard et al., 2011). The coefficient was calculated based on presence / absence of taxa.

A statistical ordination was carried out using non-metric multidimensional scaling (nMDS) within the 'R' statistical package (R Core Team, 2013). In terms of exploring community ecology data nMDS is generally considered to be an effective and passive ordination method. The ordination here is based on the proportions of the terrestrial functional group with aquatics and 'uncoded' taxa removed for all 39 'chronological entities'. We chose to restrict the nMDS analysis to the functional groups to determine if the faunas recovered are distinct or clustered according to their ecology and to prevent any 'superabundant' or rare species unduly influencing the ordination (e.g. Smith 1991, 2012a, 2013a, Hill 2015).

## Results

### **Defining the beetle fauna of archaeological farmed landscapes**

The ranked occurrence of taxa in each archaeological period is presented in Table 3. The ecology of the main species recovered is summarised in table Table 4. The relative proportions of functional groups recovered by archaeological site are presented in Figures 2 and 3 and by archaeological period in Figure 4.

The insect faunas from archaeological farmed landscapes appear to be dominated by a relatively narrow and repetitive range of terrestrial beetle taxa. Dung beetles, account for between 15%–40% of the terrestrial taxa, with 5%–10% belonging to the genus *Aphodius* (Figures 3 and 4). *Aphodius contaminatus*, *A. granarius*, *A. sphaelatus*, *A. prodromus*, *A. rufipes* and *A. fimetarius* are particularly common. *Geotrupes* spp., *Onthophagus fracticornis* and *O. joannae* also occur in some numbers, particularly in the Neolithic and Bronze Age material.

A large proportion of species (20%–40%) are associated with open and disturbed ground (Figure 4). Often this includes a range of predatory ground and rove beetles (e.g. *Bembidion* spp., *Calathus melanocephalus*, *C. fuscipes*, *Trechus quadristriatus*, *T. obtusus*, *Stenus* spp., *Amara* spp. *Xantholinus linearis*, *Rugilus* spp., *Philonthus* spp., *Harpalus rufipes* are particularly common). Phytophages associated with plants of disturbed ground are important (e.g. the weevil *Nedyus (Cidnorhinus) quadrimaculatus*, associated almost exclusively with stinging nettle (*Urtica dioica* L), and 'flea beetles' such as *Chaetocnema concinna*, *Phyllotreta* spp. and *Longitarsus* spp.). Other phytophage taxa, such as, *Mecinus pyraster*, *Mecinus (Gymnetron) pascuorum*, *Sitona* spp., *Rhinoncus* spp. *Gastrophysa* spp. and *Apion* spp., are associated with plants common in pasture. The 'click beetles' *Agrypnus murinus*, *Agriotes* spp., *Athous haemorrhoidalis* and the 'chafers'

1  
2  
3 *Phyllopertha horticola*, *Melolontha melolontha* and *Hoplia philanthus* also occur commonly in  
4 grassland. The larvae of click beetles and chafers can be pests of arable crops; but as Robinson  
5 (1983) pointed out, they are only a serious problem in newly ploughed fields from former grassland.  
6

7 Beyond this core group of recurrent species, there is a long tail of individual taxa that occur in small  
8 numbers, often only at one or two sites. These beetles probably represent small populations of  
9 individuals associated with isolated patches of 'wild' or weedy ground which will vary with local  
10 conditions. In addition, taphonomic effects can bring insects into archaeological deposits from some  
11 distance (Kenward, 1975, 1978).  
12  
13

14 Water beetles usually account for between 10%–40% of the individual assemblages but achieved  
15 60%–70% of all taxa at several locations (Figure 2). The aquatic fauna is dominated by three taxa  
16 (*Ochthebius minimus*, *Hydraena testacea* and small *Helophorus* spp.), which are indicative of slow-  
17 flowing, vegetated and often temporary waterbodies. *Colymbetes fuscus*, *Agabus bipustulatus*,  
18 various *Hydroporus* species and a range of Hydrophilidae are associated with this environment as  
19 well. Given that many of these faunas come from archaeological ditches or waterholes, the presence  
20 of significant numbers of water beetles is not surprising.  
21  
22  
23

24 Several phytophage species also are associated with aquatic environments and watersides.  
25 *Tanysphyrus lemnae* is associated with duckweeds (*Lemna* spp.), *Notaris acridulus* with reed sweet  
26 grass (*Glyceria maxima* (Hartm.) Holmb.), *Plateumaris braccata* with water reed (*Phragmites australis*  
27 (Cav.) Trin. ex Streud.) and *Donacia crassipes* with water lilies (*Nymphaea alba* L. and *Nuphar* spp.).  
28 Surprisingly many of the ditches from this survey did not yield large numbers of these waterside  
29 phytophages (functional groups MA and MFC – Figure 3), suggesting they were periodically  
30 maintained and cleared of vegetation, perhaps indeed by grazing. The Late Roman sites of Balby  
31 Carr and East Carr in the northern Trent Valley do, however, contain high proportions of these taxa,  
32 but these ditch systems appear to have been affected by rising water tables which lead to the  
33 development of carr (Knight et al., 2004).  
34  
35  
36  
37  
38  
39

#### 40 ***Comparing modern and past beetle faunas from farmland***

41  
42 There is extensive modern literature on several families of Coleoptera from farmland, since beetles  
43 have potential economic importance as pests or predators. These studies often concentrate on issues  
44 concerning resource partitioning, predator-prey relationships and response to different cultivation  
45 practices. They also tend to be restricted to a limited number of beetle families (Carabidae  
46 Staphylinidae, Scarabaeidae and some phytophage leaf beetles and weevils), rather than considering  
47 the fauna in its entirety (one exception is Robinson's (1983) survey of a variety of agricultural  
48 landscapes near Oxford undertaken for comparison with archaeological faunas). A ranked order list  
49 of taxa from a number of these modern studies is presented in Table 5. One limit to direct comparison  
50 between archaeological and modern coleopterous faunas is that they are sampled in very different  
51 ways (e.g. pitfall trapping vs. sediment sampling) and can represent very different time intervals (often  
52 one season for modern work as opposed to an unknown number of years or decades for the  
53 accumulation of archaeological deposits – Smith, 2017; Smith et al., 2010, 2014).  
54  
55  
56  
57  
58  
59  
60



1  
2  
3 Despite these issues, there is a remarkable similarity between insect faunas from farmed landscapes  
4 regardless of whether they are modern or from archaeological sites (compare Tables 3 and 5). Both  
5 modern and ancient faunas include the same range of xerophilous predatory ground and rove beetles  
6 from arable land and a similar range of dung beetles and phytophage weevils in grasslands. This  
7 implies that the 'farmland fauna' has been essentially stable in Britain for two and a half millennia.  
8  
9

### 10 11 12 ***Variation in the faunas between sites by period***

13 The similarity of the ranked occurrence of taxa (Table 3), the proportions of functional groups  
14 recovered when plotted chronologically (Figures 3 and 4) and the close clustering of the nMDS when  
15 displayed by period (Figure 5) suggest remarkable consistency for the majority of fauna associated  
16 with farmed landscapes over time.  
17  
18

19 However, two components of the 'farmland fauna', to an extent, change over time:  
20

- 21 1) There is a relative decline in species associated with animal dung and grassland  
22 between the Late Iron Age and the Romano-British period (see Figures 3 and 4).  
23 During the Bronze and Early Iron Age half of sites (8 out of 16) have dung beetle  
24 values exceeding 20%. This declines to a third of all sites (6 out of 19 sites) in the  
25 Late Iron Age and Romano-British periods.  
26
- 27 2) The values for 'house fauna', and the Synanthropic Values remain consistently low at  
28 these sites throughout all periods (see Figure 3 and Figure 4), with the exception of  
29 the Romano-British ditch deposits from Northfleet, Kent, Perry Oaks, Berkshire and  
30 St. Loyes, Oxfordshire, where synanthropes and grain pests are quite common.  
31

### 32 ***Variation in the faunas between sites by deposit and site type***

33  
34 In order to explore whether deposit type was an important factor in shaping this data the nMDS  
35 ordination was replotted with deposit types labelled (see Figure 6). No obvious clustering or  
36 separation of deposits is seen, suggesting that the insect fauna is ubiquitous, regardless of the type of  
37 archaeological feature it comes from.  
38  
39

40 The only clear distinction identified by the nMDS ordination is the presence of a strong gradient for the  
41 wet-loving taxa (MA and MFC fauna) towards the lower left-hand side of the plot. This suggests that  
42 the greatest variation between faunas was largely determined by moisture. The two sites that most  
43 strongly indicate the influence of wet-loving taxa are Balby Carr and East Carr which were mentioned  
44 above.  
45  
46

47 It could be argued that this survey of faunas results in such similarity because it concentrates on  
48 ditched farmland sites and ignores other landscapes. As a result, it may overemphasise the  
49 dominance of this suite of beetles and, indeed, this type of farmland landscape which exists in Britain  
50 from the Late Bronze Age onwards. However, a number of archaeological insect faunas from other  
51 landscapes have been recovered for this period, for example, the woodlands and heathlands of the  
52 Humberhead levels (Buckland, 1979; Whitehouse, 2000, 2004), the swamps and carrs of the  
53 Somerset Levels (e.g. Girling, 1977, 1979, 1982, 1985) and the floodplain of the Trent river (Smith  
54 and Howard, 2004; Greenwood and Smith, 2005). Crucially, all of these sites have insect faunas  
55 distinctly different in character from the farmland faunas discussed here and that they come from a  
56 range of landscapes that would not favour agricultural production in the first place.  
57  
58  
59  
60

### ***Another form of variation?***

The construction of matrices based on similarity coefficients using only presence-absence of taxa between faunas (see Figure 7) was undertaken to understand any variations by period and site type. In addition, this explored whether any presence of 'metacommunity level' regional variations (*sensu* Gilpin and Hanski, 1991; Hanski, 1998; Leibold et al., 2004) could be established. These plots indicate that there is little similarity between faunas (most of the plot has light shading, which indicates little or no similarity). This seems contradictory compared to the results outlined above. However, the use of only presence-absence in the analysis is probably significant. The presence of the very long 'tail' of less abundant, 'rare' and local species at the sites probably has a greater impact on the ordination than frequently occurring taxa which are limited to a comparatively narrow range. This is a problem already identified for this kind of ordination when used on occupation-site faunas by Carrott and Kenward (2001) and overcome with considerable success by excluding such rare taxa (or 'outliers').

Once the sites are organised by statistical similarity rather than by chronology (the right-hand plot in Figure 7), there does appear to be a pattern. Most of the sites from the Midlands and the Lower Thames Valley plot towards the top half of the diagram, those from the Upper Thames Valley plot in the middle and lower right, and a small group of other Midland sites plots at the bottom right. This could indicate weak regional variations (*sensu* Leibold et al., 2004; Brooks et al., 2008) or the effects of local 'mass effects' such as field size, heterogeneity of the landscape and the presence or absence of wild areas (e.g. Brooks et al. 2008).

However, this pattern also could be caused by a less edifying effect. The faunas appear to group by which archaeoentomologist studied them (the specialists have been named on Figure 7). In this case the analysis may be picking up on quite minor differences in the way in which the various archaeoentomologists have identified insect fragments and/or how precisely these are recorded. For example, the Helodidae (Scirtidae) has been classified in a number of different ways by specialists. This ranges from the conservative 'Helodidae Gen. et sp. indet.', through '?*Cyphon* sp.' to the possibly overoptimistic '*Cyphon pubescens* (F.)'. It is worth noting that one advantage of basing the nMDA analysis on ecological groupings, rather than the individual taxa, is it helps to negate this exact problem (e.g. Kenward 1978, Kenward and Hall 1995).

## **Discussion**

### ***Stability and change in the 'farmland' beetle fauna***

The results presented above clearly indicate that once agricultural landscapes developed, the majority of the insect fauna associated with them remained very stable, consistent and resilient over a considerable period of time reaching up to late medieval/pre-modern times.

The decline in the relative proportion of 'dung beetles' in the later Iron Age and the Roman periods merits further discussion. Robinson (2013a) also has suggested that there is a shift from *Onthophagus*-dominated faunas to *Aphodius*-dominated dung communities at the same time. Using

1  
2  
3 specific proportions of dung beetles to estimate proportions of landcover can be problematic (Smith et  
4 al., 2010, 2014) but this decline in dung beetles probably indicates that grazing and pasture were less  
5 dominant in the landscape from c. 400 - 200 BC onwards. Some (Fyfe et al., 2008, 2013; Trondman  
6 et al., 2015) have argued that there is evidence in the British pollen records for a change from  
7 predominantly pastoral economy towards mixed and arable farming at the same time. Van der Veen  
8 and O'Connor (1998) also suggest that the British landscape becomes predominantly arable during  
9 the warming effect of the 'Roman climatic optimum'.  
10  
11

12 It could be argued that in terms of deposit types and depositional environments, we are comparing  
13 apples with oranges. The majority of the Bronze and Early Iron Age samples are from relatively  
14 shallow features such as ditches and 'waterholes'; whereas the Later Iron Age and Romano-British  
15 samples derive from deeper features, such as pits or wells. Wells and pits often are thought to act as  
16 'pitfall' traps and collect insect faunas from a wider geographical area than shallower features (e.g.  
17 Hall et al. 1980; Kenward et al., 1986; Smith and Kenward, 2013). However, the nMDA analysis in  
18 Figure 6, where the sites are plotted together independently of feature type, suggests the relative  
19 depth of context is not actually a factor; therefore, a change in land use remains the most sensible  
20 explanation.  
21  
22  
23  
24

25 The increase in the proportions of synanthropic faunas at some Roman sites can be explained more  
26 easily. These deposits may have formed particularly close to settlement or, more likely, incorporated  
27 dumped settlement waste that contained numerous synanthropes and grain pests.  
28  
29  
30

### 31 ***The ecology of the 'farmland' fauna recovered***

32 This analysis has demonstrated that there is a very distinct 'farmland fauna' associated with  
33 archaeological agricultural landscapes. Many of the common species in this 'farmland fauna' such as  
34 the carabids and staphylinids, are fast moving, mobile ground-based predators which feed on small  
35 insects, snails and slugs and effectively exploit disturbed and temporary habitats. Modern ecological  
36 studies of these taxa describe them as being eurytopic and essentially opportunistic (e.g. Turin, 2000,  
37 Luff, 2002; Brooks et al., 2008; Betrand et al., 2016). Other species, such as the dung beetles and  
38 many of the phytophage taxa, also are adapted to exploit suitable temporary patches of habitat as  
39 they occur. Many of the species, especially the ground beetles, are to some degree xerophilous  
40 (Lindroth, 1974; Luff, 2007) and favour open environments with some disturbance. The species  
41 concerned, whether they are water beetles, ground beetles, phytophages or dung beetles, are all  
42 typically associated with the early stages of ecological successions (Brooks et al., 2008, Tonhasca,  
43 1993, Turin, 2000). What is apparent from our analysis is that through continuous human agency  
44 (uninterrupted farming in this case) what essentially is a short-lived, opportunistic fauna has had its  
45 longevity extended by millennia due to the stable and continuous generation of this farmland  
46 environment.  
47  
48  
49  
50  
51  
52  
53

54 Many of the beetle species encountered in farmland, both in the archaeological record and today, also  
55 appear to have high dispersal and breeding rates (the so called 'R' reproductive strategy) which  
56 enables them to rapidly distribute themselves across the landscape and quickly build up large  
57 populations. This R-strategy is seen as particularly important for species that take advantage of the  
58  
59  
60

1  
2  
3 early stages of successions in fragmented landscapes, where resources often are short-lived and  
4 separated by long distances (De Vries et al., 1996; Ribera et al., 2001; Bonsall and Hastings, 2004;  
5 Woodcock et al., 2010, 2014; Brooks et al., 2008; Bertrand, 2016).  
6  
7

### 9 ***Possible origins of the 'farmland' in Britain***

10  
11 The nature and ecology of the Later Holocene 'farmland fauna' are quite distinct from earlier insect  
12 communities dating to the Early and Middle Holocene (c. 9500 – 4000 cal. BC) which are dominated  
13 by numerous species associated with long-lived *urwald* woodland (e.g. Smith and Whitehouse, 2005;  
14 Whitehouse and Smith, 2010). These woodland faunas are ecologically complex, have many trophic  
15 levels and contain a wide range of highly specialised species which probably follow the so-called 'K'  
16 reproductive strategy (low dispersal and breeding rates). Conversely, the 'farmland fauna' identified  
17 here is relatively simple with shallow trophic levels, is eurytopic-dominated, and follows a clear 'R'  
18 strategy. This replacement of a 'natural' fauna by a 'farmland' fauna, from the Neolithic onwards,  
19 seems to have been very rapid where it happened (i.e. there are no faunas that could be regarded as  
20 transitional) and this process appears to have occurred repeatedly in different regions of Britain at  
21 several points in time. The earliest occurrence of the 'farmland fauna' seems to be at the late Neolithic  
22 sites of Silbury Hill, Wessex (2550–2535 cal BC) and Etton, Cambridgeshire (3725–3210 cal. BC).  
23 Despite their relatively early date these sites already contain most aspects of the 'farmland fauna'.  
24 However, both sites are associated with landscapes that were cleared of forest very early (c. 2600  
25 cal. BC) as compared to other parts of Britain, where clearance generally occurred from 1600 cal. BC  
26 onwards (Allen, 1997, 2000; Bell and Walker, 2005; Fyfe et al., 2015). Whether such early clearance  
27 of the Wessex Downs or the Cambridge Fen resulted from factors such as thin soils preventing  
28 reforestation, the development of a predominantly pastoral economy, or was related to cultural factors  
29 surrounding monument building and ritual use is not clear (Thomas, 1999, 2013; French et al., 2007,  
30 2012; Wilkinson and Straker, 2008). However, these sites do demonstrate that the 'farmland fauna'  
31 was present in lowland Britain from the late Neolithic onwards and, presumably, had the potential to  
32 develop wherever woodland was cleared.  
33  
34  
35  
36  
37  
38  
39  
40

41 Most of the taxa which later become associated with archaeological farmland probably existed  
42 naturally within a range of small, localised ecological niches (niche in this instance being the place,  
43 function and ecological requirements of an individual insect) as part of the Early Holocene woodland  
44 habitat (habitat being the wide environment in which this insect can occur). Andersen (2000) and  
45 Buckland (2005) suggest that species typical of dry anthropogenic habitats, such as farmland or  
46 heathland, were originally steppe species which recolonised open landscapes at the end of last  
47 glaciation. As woodland returned during the Early Holocene, they became associated with temporary  
48 glades and open areas. These glades could have resulted from storm damage, forest fires, river bank  
49 erosion, beavers and/or actions of grazing animals (e.g. Coles and Orme, 1988; Vera, 2000; Coles,  
50 2006). Large herbivores may have a particular significance in maintaining openings in the canopy  
51 (Vera 2000). There has been considerable argument recently over the physical extent of such  
52 woodland clearings in the Early and Mid-Holocene woodland (e.g. Vera, 2000; Mitchel, 2005).  
53 Additionally, a number of recent studies using estimates of landscape cover based on regional pollen  
54 proportions has indicated that open areas may have occurred more often than previously thought in  
55  
56  
57  
58  
59  
60

1  
2  
3 Britain during the Early and Mid-Holocene (Fyfe et al., 2013, 2015, Trondman et al., 2015,  
4 Woodbridge et al., 2014). Ecologically, any Coleoptera associated with these glades would have to be  
5 able to exploit these spatially and temporally limited open environments before woodland  
6 regeneration occurred. They would exhibit many of the same forms of ecological adaptation we have  
7 described above for the 'farmland fauna' itself. A substantial part of the beetle 'farmland fauna' has  
8 actually been recovered, albeit in small numbers, from a range of Early and Mid-Holocene woodland  
9 faunas (Buckland and Buckland, 2006).

12 It was only with large scale woodland clearance and the development of extensive farmed landscapes  
13 from the Middle Iron Age onwards that these small limited niches in woodland begin to coalesce into  
14 the larger anthropogenic habitat of open farmland and pasture. For the 'farmland fauna' beetles, it is  
15 their ability to exploit open, dry and disturbed landscapes that allows them to become dominant. This  
16 is a 'ragbag' of a fauna, which probably never existed together in nature prior to this point. They have  
17 not changed the niches they use in essence but have come together as a fauna to exploit a new  
18 habitat made through human agency. The ecological history and origins of the synanthropic fauna  
19 from archaeological settlement is also similar, with taxa that previously did not occur together in  
20 nature coexisting in the new habitat provided by human habitation (Kenward and Allison 1995).

25 It is possible; however, that some of the species associated with open farmland may have dispersed  
26 alongside Neolithic farming practices as they spread across Europe. This is probably the case for  
27 some field pests, such as the bruchid 'pea weevils' and some leaf beetles. A similar origin also has  
28 been suggested for most of the pests of stored grain (see Smith and Kenward 2011).

## 33 **Conclusions and implications**

### 35 **Implications for modern agricultural expansion and the development of insect faunas and** 36 **biodiversity**

38 The archaeoentomological story described here – a decline in insect biodiversity, the growing  
39 dominance of eurytopic species, and a shallowing of trophic levels as agricultural landscapes  
40 developed – has clear echoes in the modern world. A number of studies have examined the effect of  
41 modern agricultural intensification on insect faunas (e.g. Firbank et al., 2008; Brooks et al., 2008;  
42 Bertrand et al., 2016). These suggest that the loss of wild areas, increased cultivation, development of  
43 monocultures, decline of fallow crops, and rapid crop rotation all reduce the number and diversity of  
44 beetles present in the modern landscape (Brooks et al., 2008, Fournier and Loreau, 1999; Pettit and  
45 Usher, 1998; Bertrand et al., 2016, Purtauf et al., 2005, Woodcock et al., 2014). Woodcock and  
46 colleagues (2008, 2010) have indicated that overgrazing, loss of surrounding 'wild' areas, decline of  
47 traditional grazing techniques and hay cutting times all can have a devastating effect on beetle  
48 diversity in grasslands. Crucially, this leads to a decline in plant diversity, a resulting drop in  
49 phytophage species richness and then a drop in the quantity and diversity of predatory beetles  
50 (Brooks et al., 2008; Dombos 2001; Lagerlöf et al., 2002; Purtauf et al., 2005). Recently, Hutton and  
51 Giller (2003) have shown that intensification and 'improvement' of pasture can lead to a rapid decline  
52 in the number and variety of dung beetles.

1  
2  
3 One of the major implications of the archaeoentomological study presented here is that the modern  
4 decline in insect biodiversity and the shallowing of trophic levels resulting from modern intensive  
5 agriculture is not a new phenomenon; it is part of a continuum of effects with deep temporal roots that  
6 stretch back to at least 3,500 years in Britain. The main differences between the past and present are  
7 merely a matter of the scale and perhaps the speed of the effect. Presumably this process may span  
8 a much longer period in other parts of the world, such as the Fertile Crescent of the Near East or the  
9 'rice bowl' of China, where there is longer agrarian history leading back to at least 9,500 BC (Ellis et  
10 al., 2013).

### 16 ***Is this an example of Human niche creation?***

17 'Niche creation' is a relatively new concept in evolutionary biology; it suggests that organisms have  
18 the capacity to modify their environments so they can influence their own and other species evolution,  
19 ecology and distribution (Jones et al., 1994, 1997; Laland et al., 1996, 1999, 2000, 2001; Kendal et al.  
20 2011). Humans are undoubtedly efficient 'niche creators', for example the shaping of Pleistocene  
21 (Sterelny, 2011; Boivin et al., 2016), agricultural (Rowley-Conwy and Layton, 2011; Gerbault et al.,  
22 2011; Smith, DB 2011), and urban landscapes (Boivin et al., 2016). Intentional human 'niche  
23 creation', or in this case 'habitat creation', can pronouncedly affect the distribution of other animals  
24 and plants. For the insects concerned, there were obvious winners and losers. The creation of farmed  
25 landscapes is profoundly destructive for 'urwaldrelikt' specialists such as *Prosomis mandibularis*, but  
26 beneficial for open ground generalists, such as *Pterostichus melanarius*.

### 33 ***Implications for further study***

34  
35 As is the case in every large survey of data, many issues raised here are in need of further  
36 investigation. One obvious flaw is the shortage of insect faunas related to farmland from the Neolithic,  
37 Saxon and Medieval periods in Britain. The lack of suitable Neolithic sites, in part, relates to an  
38 absence of evidence for large-scale agriculture or grazing outside a limited number of locations  
39 (Moffett et al., 1989; Thomas, 1999, 2013; Richmond, 1999; Bell and Walker, 2005; Stevens and  
40 Fuller, 2012). The shortage of rural archaeoentomological faunas from the Saxon and Medieval  
41 periods is partially explained by present archaeological research priorities, which tend only to prioritise  
42 the excavation of prehistoric field systems. Gathering more archaeoentomological data associated  
43 with agriculture from historic periods should be a national priority in Britain. The present dataset is  
44 also largely biased towards the south and midlands of England and again identifies the urgent need  
45 for archaeoentomological sampling of farmland environments to occur in other regions of Britain as a  
46 national priority.

47  
48 We also, with some reason, have assumed that the adaptation of Coleoptera from the 'urwald'  
49 woodland biome to culture-steppe was rapid, but this can only be tested on sites which span the  
50 transition itself.

51  
52 The rarity of comparable archaeological studies beyond Britain has restricted this analysis to a limited  
53 geographical area. It is likely that preservation of insect faunas will be rare in many areas but similar  
54 work should be undertaken in other parts of the world where possible. It would be fascinating to try to  
55  
56  
57  
58  
59  
60

1  
2  
3 establish how insect faunas are affected by the development and spread of farming from the 8th  
4 millennium BC onwards in the Near East and continental Europe. Examining how the insect faunas of  
5 North America developed in response to farming, whether first nation maize-based agriculture, or  
6 European introduced agrarian practices, also would be fascinating.  
7  
8  
9

## 10 **ACKNOWLEDGEMENTS**

11  
12 The authors would like to thank the individual archaeological field units who have supported this work  
13 and frequently have funded the analyses. Any unpublished insect faunas that have been used in this  
14 analysis have been included with the permission of the individual field units and we are grateful for  
15 these permissions. The statistical analysis included in this paper was supported by a University of  
16 Birmingham College of Arts and Law Research Assistantship grant which provided funding at a vital  
17 moment to Geoff Hill to participate on this project. The paper was read and edited at several points by  
18 Wendy Smith who made many helpful corrections and suggestions. Figure 1 was prepared by Helen  
19 Moulden. The authors would like to thank the referees of this paper for many useful suggestions and  
20 criticism. A referee's comments were particularly helpful in terms of how to restructure the paper.  
21  
22

## 23 **REFERENCES**

- 24 Allen, MJ (1997) Environment and Land-use: the economic development of the communities who built  
25 Stonehenge. *Proceedings of the British Academy* 92: 115–144.
- 26 Allen, MJ (2000). High resolution mapping of Neolithic and Bronze Age chalkland landscapes and  
27 landuse. In: Fairbairn, A.S. (ed.) *Plants in Neolithic Britain and Beyond*. Oxford: Oxbow Books, pp. 9–  
28 26.
- 29 Andersen, J (2000). What is the origin of the carabid beetle fauna of dry anthropogenic habitats in  
30 western Europe? *Journal of Biogeography* 27, 795–806.
- 31 Bell, M and Walker, MJC (2005). *Late Quaternary Environmental Change* (Second edition). Harlow:  
32 Pearson.
- 33  
34 Bertrand, C, Burel, F and Baudry, J (2016) Spatial and temporal heterogeneity of the crop mosaic  
35 influences carabid beetles in agricultural landscapes. *Landscape Ecology* 31:451–466
- 36 Blake, S, McCracken, DI, Eyre, MD, Garside, A and Foster, GN (2003). The relationship between the  
37 classification of Scottish ground beetle assemblages (Coleoptera, Carabidae) and the National  
38 Vegetation Classification of British plant communities. *Ecography* 26: 602–616.
- 39 Boivin, NL, Zeder, MA, Fuller, DQ, Crowther, A, Larson, G, Erlandson, JM, Denham, T and Petraglia,  
40 MD (2016) Ecological consequences of human niche construction: Examining long-term  
41 anthropogenic shaping of global species distributions. *Proceedings of the National Academy of  
42 Sciences* 133: 6388–6396.
- 43 Bonsall, MB and Hastings, A (2004) Demographic and environmental stochasticity in predator–prey  
44 metapopulation dynamics. *Journal of Animal Ecology* 73: 1043–1055.
- 45  
46 Borcard, D, Gillet, F, Legendre, P (2011) *Numerical Ecology with R*. Dordrecht: Springer.
- 47 Bradley, R, (1980) Subsistence, exchange and technology: a social framework for the Bronze Age in  
48 southern England c. 1400-700 BC. In: Barrett, JC, Bradley, R (eds.), *Settlement and Society in the  
49 British Later Bronze Age*. (British Archaeological Reports, British Series 83). Oxford: British  
50 Archaeological Reports, pp. 57-76.
- 51 Bronk Ramsey, C (2009) Bayesian analysis of radiocarbon dates. *Radiocarbon* 51(1): 337–360.
- 52 Brooks, DR, Perry, JN, Clark, SJ, Heard, MS, Firbank, LG, Holdgate, R, Mason, NS, Shortall,  
53 CR, Skellern, MP and Woiwod, IP (2008) National-scale metacommunity dynamics of carabid beetles  
54 in UK farmland. *Journal of Animal Ecology* 77: 265–274.
- 55  
56 Brück, J, (2000) Settlement, landscape and social identity: the Early/Middle Bronze Age transition in  
57 Wessex, Sussex and the Thames Valley. *Oxford Journal of Archaeology* 19: 273–300.  
58  
59  
60

- 1  
2  
3 Buckland, PC (1979) *Thorne Moors: A Palaeoecological Study of a Bronze Age Site: A Contribution to*  
4 *the History of the British Insect Fauna* (Department of Geography, University of Birmingham,  
5 Occasional Publication Number 8). Birmingham: University of Birmingham.
- 6 Buckland, PC (2005) Lowland Heaths – a paleoentomological view. In: Prendergast, HVG (ed.)  
7 *Heathlands, Past, Present and Future* (The Proceedings of the 8th National Heathland Conference).  
8 Lewes: East Sussex County Council.
- 9 Buckland, PI and Buckland, PC (2006) *BugsCEP Coleopteran Ecology Package* (IGBP PAGES/World  
10 Data Centre for Paleoclimatology Data Contribution Series # 2006-116. NOAA/NCDC  
11 Paleoclimatology Program, Boulder CO, USA). URL: <http://www.ncdc.noaa.gov/paleo/insect.html> or  
12 <http://www.bugscep.com>.
- 13 Carrott, J and Kenward, H (2001) Species associations among insect remains from urban  
14 archaeological deposits and their significance in reconstructing the past human environment. *Journal*  
15 *of Archaeological Science* 28: 887–905.
- 16 Chadwick, A (2013) Some fishy things about scales: macro- and microapproaches to Later Prehistoric  
17 and Romano-British field systems. *Landscapes*, 14: 13–32.
- 18 Chowne, P, Girling, M and Greig, JRA (1986). Excavations at an Iron Age Defended Enclosure at  
19 Tattershall Thorpe, Lincolnshire. *Proceedings of the Prehistoric Society* 52: 159–188.
- 20 Coles, B . (2006) *Beavers in Britain's Past* (Wetland Archaeology Research Project Occasional Paper  
21 19). Oxford: Oxbow Books.
- 22 Coles, J and Orme, B (1988) *Homo sapiens* or *Castor fiber*? *Antiquity* 57: 95–102.
- 23 Dark, P (2006) Climate deterioration and land-use change in the first millennium BC: perspectives  
24 from the British palynological record. *Journal of Archaeological Science* 33: 1381–1395.
- 25 Deininger, K. and Byerlee, D. (2011). *Rising Global Interest in Farmland: Can It Yield Sustainable and*  
26 *Equitable Benefits?*. Washington DC: World Bank.  
27 (https://siteresources.worldbank.org/DEC/Resources/Rising-Global-Interest-in-Farmland.pdf  
28 downloaded 13/07/2017).
- 29 De Vries, HH, den Boer, PJ and van Dijk, TS (1996) Ground beetle species in heathland fragments  
30 in relation to survival, dispersal, and habitat preference. *Oecologia* 107: 332–342
- 31 Dombos M (2001) Collembola of loess grassland: effects of grazing and landscape on community  
32 composition. *Soil Biology and Biochemistry* 33: 2037–2045.
- 33 Duff, A.G (2012) *Checklist of Beetles of the British Isles* (second edition). Iver: Pemberley Books.
- 34 Ellis, EC, O'Kaplan, JE, Fuller, DQ, Vavrus, S, Goldewijk, KK and Verburg, PH (2013) Used planet: A  
35 global history. *Proceedings of the National Academy of Science* 110: 7978–7985.
- 36 English, J (2013) *Pattern and Progress: Field Systems of the Second and Early First Millennia BC in*  
37 *Southern Britain* (British Archaeological Reports, British Series 587). Oxford: Archaeopress.
- 38 Eyre, MD, Luff, ML, Staley, JR and Telfer, MG (2003) The relationship between British ground  
39 beetles (Coleoptera, Carabidae) and land cover. *Journal of Biogeography* 30: 719–730.
- 40 Firbank, LG, Petit, S, Smart, S, Blain, A and Fuller, RJ (2008) Assessing the impacts of agricultural  
41 intensification on biodiversity: A British perspective. *Philosophical Transactions of the Royal Society:*  
42 *Biological Sciences* 363: 777–787
- 43 Fleming, A (1987) Co-axial field systems: some questions of time and space. *Antiquity* 61: 188–202.
- 44 Fleming, A (1988) *The Dartmoor Reaves: Investigating Prehistoric Land Divisions* (first edition).  
45 London: Batsford.
- 46 Fleming, A (2007). *The Dartmoor Reaves: Investigating Prehistoric Land Divisions* (second edition).  
47 Bollington: Windgather Press..
- 48 Foley JA, DeFries, R, Asner, G., Barford, C, Bonan, G, Carpenter, SR, Chapin, FS, Coe, MT, Daily,  
49 GC, Gibbs, HK, Helkowski, JH, Holloway, T, Howard, EA, Kucharik, CJ, Monfreda, C, Patz, JA,  
50 Prentice, IC, Ramankutty, N and Snyder, PK (2005) Global consequences of land use. *Science* 309  
51 (5734): 570–574.
- 52 Fournier, E and Loreau, M (1999) Effects of newly planted hedges on ground-beetle diversity  
53 (Coleoptera. Carabidae) in an agricultural landscape. *Ecography* 22: 87–97.



- 1  
2  
3 French, C, Lewis, H, Allen, M, Green, M, Scaife, R and Gardiner, J (2007). *Prehistoric Landscape*  
4 *Development and Human Impact in the Upper Allen Valley, Cranborne Chase, Dorset*. Cambridge:  
5 McDonald Institute for Archaeological Research.
- 6 French, C, Scaife, R, Allen, MJ, Parker Pearson, N, Pollard, J, Richards, C, Thomas, J, Welham, K  
7 (2012) Durrington Walls to West Amesbury by way of Stonehenge: a major transformation of the  
8 Holocene landscape. *The Antiquities Journal* 92: 1–36.
- 9 Fyfe, RM, Brück, J, Johnston, R, Lewis, H, Roland, TP, and Wickstead, H (2008) Historical context  
10 and chronology of Bronze Age land enclosure on Dartmoor, UK. *Journal of Archaeological Science*  
11 35: 2250–2261.
- 12 Fyfe, RM, Twiddle, C, Sugita, S, Gaillard, MJ, Barratt, P, Caseldine, CJ, Dodson, J, Edwards, KJ,  
13 Farrell, M & Froyd, C (2013) The Holocene vegetation cover of Britain and Ireland: overcoming  
14 problems of scale and discerning patterns of openness. *Quaternary Science Reviews* 73: 132-148
- 15 Fyfe, RM, Woodbridge, J and Roberts, N (2015) From forest to farmland: pollen-inferred land cover  
16 change across Europe using the pseudobiomization approach. *Global Change Biology* 21: 1197–  
17 1212
- 18 Gerbault, P, Liebert, A, Iltan, Y, Powell, A, Currat, M, Burger, J, Swallow, DM and Thomas, GM (2011)  
19 Evolution of lactase persistence: an example of human niche construction. *Philosophical Transactions*  
20 *of the Royal Society (B)* 366: 863–877.
- 21 Gilpin, ME and Hanski, I (1991) *Metapopulation Dynamics: Empirical and Theoretical Investigations*.  
22 London: Academic Press.
- 23  
24 Girling, MA (1977) Fossil insect assemblages from Rowland's track. *Somerset Levels Papers* 3: 51–  
25 60.
- 26 Girling, MA (1979) Fossil insects from the Sweet track. *Somerset Levels Papers* 5: 84–93.
- 27  
28 Girling, MA (1982) Fossil insect faunas from forest sites. In: Bell, M and Limbrey, S (eds.)  
29 *Archaeological Aspects of Woodland Ecology* (British Archaeological Reports, International Series  
30 146). Oxford: British Archaeological Reports, pp.129–146
- 31 Girling, MA (1985) An 'old forest' beetle fauna from a Neolithic and Bronze Age peat deposit at  
32 Stileway. *Somerset Levels Papers* 11: 80–85.
- 33 Greenwood, M and Smith, DN (2005) A survey of Coleoptera from sedimentary deposits from the  
34 Trent Valley. In: Smith, DN, Brickley, MB and Smith, W (eds.) *Fertile Ground: Papers in Honour of*  
35 *Professor Susan Limbrey* (AEA symposia no. 22). Oxford: Oxbow Books, pp. 53–67.
- 36 Greenwood, MT, Bickerton, MA, Castella, E, Large, ARG and Petts, GE (1991) The use of  
37 Coleoptera (Arthropoda: Insecta) for flood plain characterization on the River Trent. *U.K. Regulated*  
38 *Rivers: Research and Management* 6: 321–332.
- 39 Hall AR and Kenward HK (1990) *Environmental Evidence from the Colonia* (The Archaeology of  
40 York 14/6). London: Council for British Archaeology.
- 41  
42 Hall, AR, Kenward, HK and Williams, D (1980) *Environmental Evidence from Roman Deposits in*  
43 *Skeldergate* (Archaeology of York 14/3). London: Council for British Archaeology.
- 44 Hanski, I (1998). *Metapopulation Ecology*. Oxford: Oxford University Press.
- 45 Hill, GE (2015) *Reconstructing Holocene landscape cover using sub-fossil Coleoptera*. PhD Thesis,  
46 Queens University Belfast, UK.
- 47  
48 Hill, GE (2016) *The Insect Remains from Late Iron Age Deposits from Enderby, Leicestershire, UK*.  
49 (University of Birmingham Environmental Archaeology Services Report 249). Unpublished report to  
50 University of Leicester Archaeology Service.
- 51 Holland, JM and Reynolds, CM (2003) The impact of soil cultivation on arthropod (Coleoptera and  
52 Araneae) emergence on arable land. *Pedobiologia* 47: 181–191.
- 53 Huntley, B and Birks, HJB (1983) *An Atlas of Past and Present Pollen Maps for Europe: 0–13000*  
54 *years ago*. Cambridge: Cambridge University Press.
- 55 Hutton, SA and Giller, PS (2003) The effects of the intensification of agriculture on northern temperate  
56 dung beetle communities. *Journal of Applied Ecology* 40: 994–1007.
- 57  
58 Jones, CG, Lawton, JH and Shachak, M (1994) Organisms as ecosystem engineers. *Oikos* 69: 373–  
59 386.

- 1  
2  
3 Jones, CG, Lawton, JH and Shachak, M (1997) Positive and negative effects of organisms as  
4 physical ecosystem engineers. *Ecology* 78: 1946–1957.
- 5 Johnston, R (2005) Pattern without a plan: rethinking the Bronze Age co-axial field systems on  
6 Dartmoor, south-west England. *Oxford Journal of Archaeology* 24: 1–21.
- 7 Kendal, J, Tehrani, JJ and Odling-Smee, J (2011) Human niche construction in interdisciplinary  
8 focus. *Philosophical Transactions of the Royal Society (B)* 366: 785–792.
- 9 Kenward HK (1975) Pitfalls in the environmental interpretation of insect death assemblages. *Journal*  
10 *of Archaeological Science* 2: 85–94.
- 11 Kenward HK (1978) *The Analysis of Archaeological Insect Assemblages: A New Approach*.  
12 (Archaeology of York, 19/1). London: Council for British Archaeology for York Archaeological Trust.
- 13 Kenward HK (1997) Synanthropic insects and the size, remoteness and longevity of archaeological  
14 occupation sites: applying concepts from biogeography to past “islands” of human occupation.  
15 *Quaternary Proceedings* 5: 135–152.
- 16 Kenward, HK (2009) Northern Regional Review of Environmental Archaeology: Invertebrates in  
17 archaeology in the North of England. Swindon: English Heritage. ([http://services.english-heritage.org.uk/ResearchReportsPdfs/012\\_2009WEB.pdf](http://services.english-heritage.org.uk/ResearchReportsPdfs/012_2009WEB.pdf))
- 18 Kenward, HK and Allison, EP (1995) Rural origins of the urban insect fauna. In: Hall, AR and  
19 Kenward, H. (eds.) *Urban–Rural Connections: Perspectives from Environmental Archaeology*  
20 (Symposia of the Association for Environmental Archaeology 12). Oxford: Oxbow books, pp. 55–78.
- 21 Kenward, HK and Hall AR (1995) *Biological Evidence from Anglo-Scandinavian Deposits at 16–22*  
22 *Coppergate* (The Archaeology of York 14/7). London: Council for British Archaeology.
- 23 Kenward, HK and Hall, AR (1997). Enhancing bio-archaeological interpretation using Indicator  
24 groups: stable manure as a paradigm. *Journal of Archaeological Science* 24: 663–673.
- 25 Kenward, HK, Hall, AR and Jones, AKG (1980) A tested set of techniques for the extraction of plant  
26 and animal macrofossils from waterlogged archaeological deposits. *Science and Archaeology* 22: 3–  
27 15.
- 28 Kenward, HK, Hall, AR and Jones, AKG (1986) *Environmental Evidence from a Roman Well and*  
29 *Anglian Pits in the Legionary Fortress* (Archaeology of York 14/2). London: Council for British  
30 Archaeology.
- 31 Kenward, HK, Hall, AR and McComish, JM (2004). Archaeological implications of plant and  
32 invertebrate remains from fills of a massive Post-Medieval cut at Low Fishergate, Doncaster, UK.  
33 *Environmental Archaeology* 9: 61–74
- 34 Knight, D, Howard, AJ and Leary, R (2004) The Romano-British Landscape. In: Knight, D. and  
35 Howard, AJ (eds.) *Trent Valley Landscapes*. King’s Lynn: Heritage, pp. 115–152.
- 36 Koch, K (1989) *Die Kafer Mitteleuropas* (Ökologie Band 2). Krefeld: Goecke and Evers.
- 37 Koch, K (1992) *Die Kafer Mitteleuropas* (Ökologie Band 3). Krefeld: Goecke and Evers.
- 38 Kroos, S and Schaefer, S (1998) The effect of different farming systems on epigeic arthropods: a  
39 five-year study on the rove beetle fauna (Coleoptera: Staphylinidae) of winter wheat. *Agriculture,*  
40 *Ecosystems and Environment* 69: 121–133.
- 41 Lagerlöf, J, Goffre, B and Vincent, D (2002) The importance of field boundaries for earthworms  
42 (Lumbricidae) in the Swedish agricultural landscape. *Agriculture, Ecosystems and Environment* 89:  
43 91–103.
- 44 Laland, KN, Odling-Smee, FJ and Feldman, MW (1996) On the evolutionary consequences of Niche  
45 construction. *Journal of Evolutionary Biology* 9: 293–316.
- 46 Laland, KN, Odling-Smee, FJ and Feldman, MW (1999) Evolutionary consequences of Niche  
47 construction and their implications for ecology. *Proceedings of the National Academy of Science USA*  
48 96: 10242–10247.
- 49 Laland, KN, Odling-Smee, FJ and Feldman, MW (2000) Niche construction, biological evolution and  
50 cultural change. *Behavioural Brain Sciences* 23: 131–175.
- 51 Laland, KN, Odling-Smee, J and Feldman, MW (2001) Cultural niche construction and human  
52 evolution. *Journal of Evolutionary Biology* 34: 22–33.
- 53  
54  
55  
56  
57  
58  
59  
60

- 1  
2  
3 Leibold, MA, Holyoak, M, Mouquet, N, Amarasekare, P, Chase, JM, Hoopes, MF Holt, RD, Shurin, JB,  
4 Law, R, Tilman, D, Loreau, M and Gonzalez, A (2004) The metacommunity concept: a framework for  
5 multi-scale community ecology. *Ecology Letters* 7: 601–613.
- 6 Lindroth CH (1974) *Coleoptera: Carabidae*. (Handbooks for the Identification of British Insects. IV.  
7 part 2). London: Royal Entomological Society.
- 8 Luff, ML (1989) Use of Carabids as environmental indicators in grassland and cereals. *Annales*  
9 *Zoologici Fennici* 33: 185–195.
- 10 Luff, ML (2002) Carabid assemblage organisation and species composition. In: J.M. Holland (ed.)  
11 *The Agroecology of Carabid Beetles* Andover: Intercept, pp. 41–80.
- 12 Luff, M (2007) *Coleoptera: Carabidae* (Handbooks for the Identification of British Insects 4/2 -  
13 second edition). London: Royal Entomological Society.
- 14 Macklin, MG, Johnstone, E and Lewin, J (2005) Pervasive and long-term forcing of Holocene river  
15 instability and flooding in Great Britain by centennial-scale climate change, *The Holocene* 15: 937–  
16 943.
- 17 Macklin, MG and Lewin, J (2003) River sediments, great floods and centennial scale Holocene  
18 climate change, *Journal of Quaternary Science* 18: 101–105.
- 19 McCune, B and Grace, JB (2002) *Analysis of Ecological Communities*. Glenden Beach, Oregon,  
20 USA: MjM Software.
- 21 Mitchell, FJG (2005) How open were European primeval forests? Hypothesis testing using  
22 palaeoecological data. *Journal of Ecology* 93: 168–177.
- 23 Moffett, L, Robinson, MA and Straker, V (1989) Cereals, fruits and nuts: charred plant remains from  
24 Neolithic sites in England and Wales and the Neolithic economy. In: Milles, A, Williams, D and  
25 Gardiner, N (eds.) *The Beginnings of Agriculture* (British Archaeological Reports, International Series  
26 496), Oxford: British Archaeological Reports, pp 243–261.
- 27 Okansen (2015) *Multivariate Analysis of Ecological Communities in R: vegan tutorial* Retrieved from  
28 the University of Oulu : <http://cc.oulu.fi/~jarioksa/opetus/metodi/vegantutor.pdf>
- 29 Osborne, PJ (1969) An insect fauna of Late Bronze Age date from Wilsford, Wiltshire. *Journal of*  
30 *Animal Ecology* 38: 555–566.
- 31 Osborne, PJ (1979) Insect remains. In: Smith, C (ed.) *Fisherwick: The Reconstruction of an Iron*  
32 *Age Landscape* (British Archaeological Reports, British Series 61). Oxford: British Archaeological  
33 Reports, pp 85–87 and 189–193.
- 34 Osborne, PJ (1978) Insect evidence for the effect of man on the lowland landscape. In: Limbrey, S,  
35 and Evans, JG (eds.) *The Effects of Man on the Landscape: The Lowland Zone* (Council for British  
36 Archaeology Research Report 21). London: Council for British Archaeology, pp. 32–34.
- 37 OxCAL V.4.3 (2017). *OxCAL on-line*. Oxford: Oxford Radiocarbon Accelerator Unit. (URL  
38 <https://c14.arch.ox.ac.uk/oxcal.html>)
- 39 Pettit, S and Usher, MB (1998) Biodiversity in agricultural landscapes: the ground beetle  
40 communities of woody uncultivated habitats. *Biodiversity and Conservation* 7: 1549–1561.
- 41 Purtauf, T, Dauber, J and Wolters, V (2005) The response of carabids to landscape simplification  
42 differs between trophic groups. *Oecologia* 142: 458–464.
- 43 R Core Team (2013) *R: A Language and Environment for Statistical Computing*. Vienna, Austria: R  
44 Foundation for Statistical Computing. (URL <http://www.R-project.org/>).
- 45 Reimer, PJ, Bard, E, Bayliss, A, Beck, JW, Blackwell, PG, Bronk Ramsey, C, Grootes, PM,  
46 Guilderson, TP, Haffidason, H, Hajdas, I, Hatt, ŽC, Heaton, TJ, Hoffmann, DL, Hogg, AG., Hughen,  
47 KA, Kaiser, KF, Kromer, B, Manning, SW, Niu, M, Reimer, RW, Richards, DA, Scott, EM, Southon,  
48 JR, Staff, RA, Turney, CSM and van der Plicht, J (2013) IntCal13 and Marine13 Radiocarbon Age  
49 Calibration Curves 0-50,000 Years cal BP. *Radiocarbon* 55(4).
- 50 Ribera, I, Doleded, S, Downie, IS and Forster, GN (2001) Effects of Land Disturbance and stress on  
51 species traits of ground beetle assemblages. *Ecology* 82: 1112–1129.
- 52 Richmond, A (1999) *Preferred Economies* (British Archaeological Reports, British Series 290).  
53 Oxford: British Archaeological Reports.
- 54 Riley, DN (1980) *Early Landscape from the Air. Studies of Crop Marks in South Yorkshire and North*  
55 *Nottinghamshire*. Sheffield: Department of Archaeology and Prehistory.  
56  
57  
58  
59  
60

- 1  
2  
3 Robinson, MA (1978). A comparison between the effects of man on the environment of the first  
4 gravel terrace and flood-plain of the Upper Thames Valley during the Iron Age and Roman periods. In:  
5 Limbrey, S and Evans, JG (eds.) *The Effect of Man on the Landscape: the Lowland Zone* (Council for  
6 British Archaeology Research Report, British Series 21). London: Council for British Archaeology, pp.  
7 35–43.
- 8 Robinson, MA (1979) The biological evidence. In: Lambrick, G. and Robinson, MA (eds.) *Iron Age*  
9 *and Roman Riverside Settlements at Farmoor, Oxfordshire* (Council for British Archaeology  
10 Research Report, British Series 32). London: Council for British Archaeology, pp 77–133.
- 11 Robinson, MA (1981a) The use of ecological groupings of Coleoptera for comparing sites. In: Jones,  
12 M. and Dimbleby, G (eds.) *The Environment of Man: The Iron Age to the Anglo-Saxon Period* (British  
13 Archaeological Reports, British Series 87). Oxford: British Archaeological Reports, pp. 251–286.
- 14 Robinson, MA (1981b) Roman waterlogged plant and invertebrate evidence. In: Hinchliffe, J and  
15 Thomas, R (eds.) Archaeological investigations at Appleford. *Oxoniensia* 45: 90–106.
- 16 Robinson, MA (1983) Arable/pastoral ratios from insects? In: M. Jones (ed.) *Integrating the*  
17 *Subsistence Economy* (British Archaeological Reports, International Series 181). Oxford: British  
18 Archaeological Reports, pp 19–47.
- 19 Robinson, M (1992) Insect assemblages. In: Butterworth, CA and Lobb, SJ (eds) *Excavations in the*  
20 *Burghfield Area, Berkshire: Developments in the Bronze Age and Saxon Landscapes* (Wessex  
21 Archaeological Report 1). Salisbury: Wessex Archaeology, pp 143–148.
- 22 Robinson, M (1993) The scientific evidence. In: Allen, TG and Robinson, MA (eds.) *The Prehistoric*  
23 *Landscape and Iron Age Enclosed Settlement at Mingies Ditch, Hardwich-with-Yelford Oxon* (Oxford  
24 Archaeological Unit, Thames Valley Landscapes. The Windrush Valley 2). London: Council for British  
25 Archaeology, pp 101–141.
- 26 Robinson, M. (1997) The insects. In: Whittle, A. (ed.) *Sacred Mound Holy Rings: Silbury Hill and the*  
27 *West Kennet Palisade Enclosures: A Later Neolithic Complex in North Wiltshire* Oxford: Oxbow  
28 Books, pp 36–46.
- 29 Robinson, M (1998). Insect assemblages. In: Pryor, F (ed.) *Etton: Excavations at a Neolithic*  
30 *Causewayed Enclosure near Maxey Cambridgeshire, 1982–1987*. London: English Heritage, pp.  
31 337–348.
- 32 Robinson, MA (2000) Coleopteran evidence for the elm decline, Neolithic activity in woodland,  
33 clearance and the use of the landscape. In: Fairbairn, AS (ed.) *Plants in Neolithic Britain and*  
34 *Beyond* (Neolithic Studies Group Seminar Papers 5). Oxford: Oxbow Books, pp. 27–36.
- 35 Robinson, M (2003) *English Heritage Reviews of Environmental Archaeology: Midlands Region*  
36 *Insects*. (URL: <http://services.english-heritage.org.uk/ResearchReportsPdfs/009-2003.pdf>).
- 37 Robinson, M (2006) Analysis of insects from Middle Bronze Age to Roman deposits from Perry  
38 Oaks. In: Brown, L, Lewis, J and Smith, AD (eds.) *Landscape Evolution in the Middle Thames Valley:*  
39 *Perry Oaks Volume 1: Heathrow Terminal 5 Excavations*. London: Framework Archaeology.
- 40 Robinson, M (2013a) The relative abundance of *Onthophagus* species in British assemblages of  
41 dung beetles as evidence for Holocene climate change. *Environmental Archaeology* 18: 132–142.
- 42 Robinson, M (2013b) Holocene archaeological evidence of extinct and very rare British  
43 Scarabaeoidea. *Environmental Archaeology* 18: 143–53.
- 44 Robinson, M, Dickson, JH and Greig, JRA (1984) Waterlogged plant and invertebrate evidence. In:  
45 Miles, D (ed.) *Archaeology at Barton Court Farm, Abingdon, Oxon*. (Microfiche chapters VIII & IX).  
46 Council for British Archaeology Research Report, British Series 50). London: Council for British  
47 Archaeology, pp. 9C1–9D8.
- 48 Rowley-Conwy, P and Layton, R (2011) Foraging and farming as niche construction: stable and  
49 unstable adaptations. *Philosophical Transactions of the Royal Society (B)* 366: 849–862.
- 50 Smith, DB (2011) General patterns of niche construction and the management of 'wild' plant and  
51 animal resources by small-scale pre-industrial societies. *Philosophical Transactions of the Royal*  
52 *Society (B)* 366: 836–848.
- 53 Smith, D.N. (1991). *An Investigation of the Potential of Modern Analogue Faunas to Act as*  
54 *Comparisons to Palaeontological Samples from Archaeological Farm Sites*. PhD dissertation,  
55 University of Sheffield.
- 56  
57  
58  
59  
60

- 1  
2  
3 Smith, DN (1997) The insects from Blaco Hill (now called East Carr, Mattersey). (URL:  
4 [http://archaeologydataservice.ac.uk/archiveDS/archDownload?t=arch-1352-](http://archaeologydataservice.ac.uk/archiveDS/archDownload?t=arch-1352-1/dissemination/pdf/Nottinghamshire/GL26021 B.pdf)  
5 [1/dissemination/pdf/Nottinghamshire/GL26021 B.pdf](http://archaeologydataservice.ac.uk/archiveDS/archDownload?t=arch-1352-1/dissemination/pdf/Nottinghamshire/GL26021 B.pdf))
- 6 Smith, DN (2002) The insect remains. In: Coates, GA (ed.) *Prehistoric and Romano-British*  
7 *Landscape: Excavations at Whitemoor Hays Quarry, Staffordshire, 1997–1999*. (British  
8 Archaeological Reports, British Series 340). Oxford: British Archaeological Reports, pp 67–72.
- 9 Smith, DN (2006). Insect remains. In: Jones, L, Woodward, A and Buteux, S (eds.) *Iron Age,*  
10 *Roman and Saxon Occupation at Grange Park: Excavations at Courteenhall, Northamptonshire 1999*  
11 (Birmingham Archaeology Monograph Series 1 / British Archaeological Reports; British Series 425).  
12 Oxford: Archaeopress, pp. 143–146.
- 13 Smith, DN (2009) *Insect remains from Kingsmead Quarry, Horton, Berkshire*. (University of  
14 Birmingham Environmental Archaeology Services Report 175). Unpublished report to Wessex  
15 Archaeology.
- 16 Smith, DN (2011a) [Romano-British] insect remains In: Jones, A (ed) *Excavations at Little Paxton*  
17 *Quarry, Cambridgeshire, 1992–1998* (Birmingham Archaeology Monograph Series 10 / British  
18 Archaeological Reports, British Series 545). Oxford: British Archaeological Reports, pp. 265–277.
- 19 Smith, DN (2011b) Insects from Northfleet In: Barnett, C, McKinley, JI, Stafford, E, Grimm, JM and  
20 Stevens, C.J. (eds.) *Settling the Ebbsfleet Valley: High Speed 1 Excavations at Springhead and*  
21 *Northfleet, Kent, The Late Iron Age, Roman, Saxon and Medieval Landscape (Volume 3: Late Iron*  
22 *Age to Roman Human remains and Environmental Reports)*. Oxford/ Salisbury: Oxford/ Wessex  
23 Archaeology, pp. 88–90.
- 24 Smith, DN (2011c) The insects from Loyes, Exeter. (University of Birmingham Environmental  
25 Archaeology Services Report 196). Unpublished report to Exeter Archaeology.
- 26 Smith, DN (2012a) *Insects in the City: An Archaeoentomological Perspective on London's Past*  
27 (British Archaeological Reports, British Series 561). Oxford: Archaeopress.
- 28 Smith, D.N. (2012b) The insect remains. In: Grant, MJ, Norcott, D and Stevens, CJ (eds). *Olympic*  
29 *Park Palaeo-environmental Appendix*. Salisbury: Wessex Archaeology. (URL:  
30 [http://archaeologydataservice.ac.uk/archiveDS/archiveDownload?t=arch-1175-](http://archaeologydataservice.ac.uk/archiveDS/archiveDownload?t=arch-1175-1/dissemination/pdf/PDFA-SpecialistReports/1_Olympic_Palaeo-environmental-PDFA.pdf)  
31 [1/dissemination/pdf/PDFA-SpecialistReports/1\\_Olympic\\_Palaeo-environmental-PDFA.pdf](http://archaeologydataservice.ac.uk/archiveDS/archiveDownload?t=arch-1175-1/dissemination/pdf/PDFA-SpecialistReports/1_Olympic_Palaeo-environmental-PDFA.pdf))
- 32 Smith, DN (2012c) The insects from the waterhole at Lockington XA 58.2006. (University of  
33 Birmingham Environmental Archaeology Services Report 203). Unpublished report to University of  
34 Leicester Archaeology Service).
- 35 Smith, D.N. 2013a. Defining an 'indicator package' to allow identification of 'cess pits' in the  
36 archaeological record. *Journal of Archaeological Science* 40, 526–43.
- 37 Smith, DN (2013b) The insects from Brickets Wood, M25 (University of Birmingham Environmental  
38 Archaeology Services Report 219). Unpublished report to Oxford Archaeology South.
- 39 Smith, DN (2014a) *The Insect Remains from Kingsmead Quarry Horton (71808)* (University of  
40 Birmingham Environmental Archaeology Services Report 226). Unpublished report to Wessex  
41 Archaeology.
- 42 Smith, DN (2014b) *The Insect Remains from Cambridge North West* (University of Birmingham  
43 Environmental Archaeology Services Report 224 and 232). Unpublished report to Cambridge  
44 Archaeology Unit.
- 45 Smith, DN (2015a) Insect Remains. In: Powell, A, Barclay, A, Mephram, L and Stevens, C (eds)  
46 *Imperial College Sports Ground and RMC Land, Harlington*. (Wessex Archaeology Report 33).  
47 Salisbury: Wessex Archaeology, pp 276–283.
- 48 Smith, DN (2015b) Insect remains from Romano British pit (P68). In: G. Hughes and A. Woodward  
49 (eds). *The Iron Age and Romano-British Settlement at Crick Covert Farm: Excavations 1997–1998*.  
50 Oxford: Archaeopress, pp. 284–289.
- 51 Smith, DN (2017) Locating where archaeological sites occur in intertidal sequences: The use of  
52 archaeoentomological data as a proxy for tidal regime. *Journal of Archaeological Science* 82: 1–16.
- 53 Smith, DN and Howard, AJ (2004) Identifying changing fluvial conditions in low gradient alluvial  
54 archaeological landscapes: Can Coleoptera provide insights into changing discharge rates and  
55 floodplain evolution? *Journal of Archaeological Science* 31: 109–20.

- 1  
2  
3 Smith, DN and Kenward, HK (2011). Roman grain pests in Britain: Implications for grain supply and  
4 agricultural production. *Britannia* 42: 243–62.
- 5 Smith, DN and Kenward HK (2013) 'Well, Sextus, what can we do with this?' The disposal and use  
6 of insect-infested grain in Roman Britain. *Environmental Archaeology* 17: 141–50.
- 7 Smith, DN and Langham, E (2000) Waterlogged insect remains. In: Palmer, SC (ed.) Archaeological  
8 Excavations in the Arrow Valley, Warwickshire. *Transactions of the Birmingham and Warwickshire*  
9 *Archaeological Society* 103: pp. 179–81.
- 10 Smith, DN and Tetlow, EP (1997) The insect remains. In: Jones, L (ed.) Archaeological Excavation  
11 of a Brickwork Plan Field System at Catesby Business Park, Balby Carr, Doncaster, South Yorkshire  
12 2002. *Yorkshire Archaeology Journal* 79, pp 48–50.
- 13 Smith, DN and Tetlow, EP (2007). The insect remains. In T. Gent and S. Reed, S. A Bronze Age field  
14 and figure at Hillfarrence. *Somerset Archaeology and Natural History*, 150: pp. 120–122.
- 15 Smith, DN and Whitehouse, N (2005) Not seeing the Trees for the Woods; a palaeontomological  
16 perspective on Holocene woodland composition. In: Smith, DN, Brickley, MB and Smith, W (eds.)  
17 *Fertile Ground: Papers in Honour of Professor Susan Limbrey* (Association for Environmental  
18 Archaeology Symposia No. 22). Oxford: Oxbow Books, pp. 136–161.
- 19 Smith, DN, Nayyar, K, Schreve, D, Thomas, R and Whitehouse, N (2014) Can dung beetles from the  
20 palaeoecological and archaeological record indicate herd concentration and the identity of  
21 herbivores? *Quaternary International* 341: 119–130.
- 22 Smith, DN, Whitehouse, N, Bunting, MJ and Chapman, H (2010) Can we characterise 'openness' in  
23 the Holocene palaeoenvironmental record? Analogue studies from Dunham Massey deer park and  
24 Epping Forest, England. *The Holocene* 20: 215–29.
- 25 Sørensen, T (1948) A method of establishing groups of equal amplitude in plant sociology based on  
26 similarity of species and its application to analyses of the vegetation on Danish commons. *Kongelige*  
27 *Danske Videnskabernes Selskab* 5: 1–34.
- 28 Stace, C (2010) *New Flora of the British Isles*. (third edition). Cambridge: Cambridge University  
29 Press.
- 30 Sterelny, K (2011) From hominins to humans: how *sapiens* became behaviourally modern.  
31 *Philosophical Transactions of the Royal Society (B)* 366: 809–822.
- 32 Stevens, CJ and Fuller, DQ (2012) Did Neolithic farming fail? The case for a Bronze Age agricultural  
33 revolution in the British Isles. *Antiquity* 86: 707–722.
- 34 Taylor, J (2007) *An Atlas of Roman Rural Settlement in England* (Council for British Archaeology  
35 Research Report 151). London: Council for British Archaeology.
- 36 Tetlow, E. (2010) Insect remains (Specialist data: Section 17). In: Lewis, J (ed) *Landscape Evolution*  
37 *in the Lower Thames Valley: Heathrow Terminal 5 excavations Volume 2*. London: Framework  
38 Archaeology. (URL: <http://www.framearch.co.uk/t5/evidence/>).
- 39 Thomas, J (1999) *Understanding the Neolithic*. London: Routledge.
- 40 Thomas, J (2013) *The birth of Neolithic Britain: An Interpretive Account*. Oxford: Oxford University  
41 Press.
- 42 Tilman, D (1999) Global environmental impacts of agricultural expansion: The need for sustainable  
43 and efficient practices. *Proceedings of the National Academy of Science USA* 96: 5995–6000.
- 44 Tonhasca, A (1993) Carabid beetle assemblages under diversified agroecosystems. *Entomologia*  
45 *Experimentalis et Applicata* 68: 279-285.
- 46 Trondman, A-K, Gaillard, M-J, Mazier, F, Sugita, S, Fyfe, R, Nielsen, AB, Twiddle, C, Barratt, P, Birks,  
47 HJB and Bjune, AE (2015). Pollen-based quantitative reconstructions of Holocene regional vegetation  
48 cover (plant-functional types and land-cover types) in Europe suitable for climate modelling *Global*  
49 *Change Biology* 2: 676-69.
- 50 Tscharrntke, T, Klein, AM, Kruess, A, Steffan-Dewenter, I and Thies., C (2005) Landscape  
51 perspectives on agricultural intensification and biodiversity – ecosystem service management.  
52 *Ecology Letters* 8: 857–874.
- 53 Turin, H (2000) *The Netherlands Ground Beetles: Distribution and Ecology*. Utrecht: KNNV Uitgeverij.
- 54  
55  
56  
57  
58  
59  
60

- 1  
2  
3 van der Veen, M and O'Connor, T (1998) The expansion of agricultural production in late Iron Age  
4 and Roman Britain. In: Bailey, J (ed.) *Science in Archaeology: An Agenda for the Future*. London:  
5 English Heritage, pp. 127–43.
- 6 Vera, FWM (2000) *Grazing Ecology and Forest History*. Oxford: CABI Publishing.
- 7 Whitehouse, NJ (2000) Forest fires and insects: palaeoentomological research from a sub-fossil  
8 burnt forest. *Palaeogeography, Palaeoclimatology, Palaeoecology* 164: 231–246.
- 9  
10 Whitehouse NJ (2004) Mire ontogeny, environmental and climate change inferred from fossil beetle  
11 successions from Hatfield Moors, eastern England. *The Holocene* 14: 79–93.
- 12 Whitehouse, N and Smith D (2010). How fragmented was the British Holocene wildwood?  
13 Perspectives on the 'Vera' grazing debate from the fossil beetle record. *Quaternary Science Reviews*  
14 29: 539–53.
- 15 Wilkinson, KN and Straker, V (2008). Neolithic and Early Bronze Age environments in South-west  
16 England. In: Webster, CJ (ed.) *The Archaeology of South-west England. South-west Archaeological*  
17 *Research Frameworks: Resource Assessment and Research Agenda*. Taunton: Somerset County  
18 Council, pp 63–74.
- 19 Woodbridge, J, Fyfe, RM, Roberts, N, Downey, S, Edinborough, K & Shennan, S (2014) The impact  
20 of the Neolithic agricultural transition in Britain: a comparison of pollen-based land-cover and  
21 archaeological C-14 date-inferred population change. *Journal of Archaeological Science* 51: 216-224
- 22 Woodcock, BA, Edwards, AR, Lawson, CS, Westbury, DB, Brook, JH, Harris, SJ, Brown, VK and  
23 Mortimer, SR (2008) Contrasting success in the restoration of plant and phytophagous beetle  
24 assemblages of species-rich mesotrophic grasslands. *Oecologia* 154: 773–783.
- 25 Woodcock, Vogiatzakis, BA, Duncan, IA, Westbury, B, Lawson, CS, Edwards, AR, Brook, AJ, Harris,  
26 SJ, Lock, KA, Maczey, N, Masters, G, Valerie, K., Brown, VK, and Mortimer, SR (2010) The role of  
27 management and landscape context in the restoration of grassland phytophagous beetles. *Journal of*  
28 *Applied Ecology* 47: 366–376.
- 29 Woodcock, BA, Harrower, C, Redhead, J, Edwards, M, Vanbergen, AJ, Heard, MS, Roy, DB and  
30 Pywell, RF (2014) National patterns of functional diversity and redundancy in predatory ground  
31 beetles and bees associated with key UK arable crops. *Journal of Applied Ecology* 51: 142–151
- 32 Yates, DT (1999) Bronze Age field systems in the Thames Valley. *Oxford Journal of Archaeology* 18:  
33 157–170.
- 34 Yates, DT (2007). *Land, Power and Prestige: Bronze Age Field Systems in Southern England*.  
35 Oxford: Oxbow Books.
- 36  
37 Zabel, F, Putzenlechner, B and Mauser, W (2014) Global agricultural land resources – a high  
38 resolution suitability evaluation and its perspectives until 2100 under climate change conditions. *PLoS*  
39 *ONE* 9: e107522. doi:10.1371/journal.pone.0107522.
- 40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

## Figures and Tables

Figure 1: Location map for the archaeological sites in this survey

Figure 2. The relative proportions of the broader aquatic, wetland and terrestrial functional groups

Figure 3: The relative proportions of the terrestrial functional groups for the archaeological sites

Figure 4. The relative proportions of functional groups (dung, foul material, open and woodland) and synanthropic groups by archaeological period

Figure 5: nMDS analysis for of the taxa from each individual 'chronological entity' by archaeological period (see key for details of each archaeological period)

Figure 6: nMDS analysis for of the taxa from each individual 'chronological entity' by archaeological feature (see key for details of each archaeological feature)

Figure 7: Dissimilarity matrices. On a simple score of presence/ absence, sites' faunas are largely dissimilar to each other (i.e. large tail of less common species heavily influential). However, those sites that are similar cluster to a large extent according to the person carrying out the analysis.

Table 1. Location, dating, feature type and reference for the archaeological sites included in this survey

Table 2. Functional groups used in this analysis (based on Hill 2015)

Table 3: Rank order and % MNI for the 29 most frequently recovered Coleoptera by period for the sites in this survey. Note that some taxa (*Chaetocnema concinna*, *Aphodius sphaelatus*, *A. fimetarius*, *Megasternum 'boletophagum'*, here rendered as '*concinnum*', *Hydrobius fuscipes*, *Anotylus sculpturatus* and *Carpelimus bilineatus*) have been split since these sites were recorded

Table 4. Summary table of the insects thought to be particularly important in archaeological farms by ecological groupings and then by taxonomic order

Table 5. Beetles typically recorded in modern ecological surveys from arable, grassland and pasture (taxa ordered by frequency)



**Table 1. Location, dating, feature type and reference for the archaeological sites included in this survey**

Site	Number of samples	Type of feature	Radiocarbon or estimated calendar dates	Period	Reference
Etton, Cambridgeshire	11	base of enclosure ditch of Causeway Camp	3725-3670 to 3310-3210 cal. BC	Early Neolithic	Robinson, 1998
Silbury Hill, Wiltshire	7	old ground surface and peat stack below mound	2550 – 2535 cal BC	Mid to late Neolithic	Robinson, 1997
Wilsford Shaft, Wiltshire	1	base of shaft / well below Bronze Age barrow	1470– 1290 cal BC	Early Bronze Age	Osborne, 1969
Terminal 5 Heathrow, Greater London (Bronze Age),	22	ditches and waterholes	ca. 1650 – 900 BC	Mid to late Bronze Age	Tetlow, 2010
Perry Oaks (terminal 5) Heathrow, Greater London (Bronze Age)	6	Bronze Age pits and waterholes	1450 – 1210 cal BC and 1380 – 1040 Cal BC	Bronze Age	Robinson, 2006
Hillfarence, Somerset	5	field ditch and pit	1450-1130 cal BC	Mid to late Bronze Age	Smith and Tetlow, 2007
Horton Quarry 71808 Berkshire	2	waterholes	ca. 1200 – 700 BC	Mid to late Bronze Age	Smith, 2014a
Kingsmead Horton Quarry (BA), Berkshire	13	waterholes	ca. 1200 – 700 BC	Mid to late Bronze Age	Smith, 2009
Imperial Sports Ground), Greater London (Bronze Age)	3	waterhole	1210-910 cal. BC	Mid to late Bronze Age	Smith, 2015a
Anslow's Cottages, Burghfield, Berkshire	5	fluvial succession	Single date of 91-431 cal. BC	Mid to late Bronze Age	Robinson, 1992
Terminal 5 Heathrow (Late Bronze Age - Iron Age)	26	ditches and waterholes	One sample 1690-1500 cal BC. 25 dated 1500-50 Cal BC	Iron Age	Tetlow, 2010
Whitemoor Haye, Staffordshire	1	series of recuts of terminal end of ring ditch	ca. 500 – 200 BC	Mid Iron Age	Smith, 2002
Kingsmead Horton Quarry, Berkshire (Late Bronze Age – Iron Age)	5	ditches and waterholes	ca. 1000 – 0 BC	Late Bronze Age - Iron Age	Smith, 2009
Enderby, Leicestershire	8	waterholes and pit features	200 – 00 BC? Estimated dates from archaeology	Late Iron Age	Hill, 2016
Olympic Park, Greater London.	10	ditches	750-260 cal. BC	Mid to late Iron Age	Smith, 2012b
Tattershall Thorpe, Lincolnshire	15	Iron Age enclosure ditch	ca. 770-200 BC based on archaeology	Mid to late Iron Age	Chowne et al., 1986
Fisherwick, Staffordshire	3	enclosure ditch	380 ca. BC – 70 cal AD	Late Iron Age	Osborne, 1979
Minges Ditches, Oxfordshire	12	enclosure ditches	390-40 cal. BC	Late Iron Age	Robinson, 1993
Farmoor, Oxfordshire (Iron Age)	6	Iron Age sumps and gullies	382 cal. BC-76 cal. AD	Late Iron Age	Robinson, 1979
Little Paxton, Cambridgeshire	10	waterholes, brickfield ditches	80 – 250 AD dated by pottery	Late First century to mid third century	Smith, 2011a
Lockington, Derbyshire, waterholes	2	waterholes	Estimated period dates from archaeology	'Roman'	Smith, 2012c

Site	Number of samples	Type of feature	Radiocarbon or estimated calendar dates	Period	Reference
Imperial Sports Ground, Greater London (Roman)	2	wells	240-510 cal. AD	Roman	Smith, 2015a
Brickets Wood Bund, Hertfordshire	3	waterholes	<i>Estimated period dates from archaeology</i>	'Roman'	Smith, 2013b
Perry Oaks (Terminal 5) Heathrow, Greater London (Roman)	4	waterholes	<i>c100-250 AD and c 200 -300 AD based on archaeology</i>	Roman	Robinson, 2006
Kingsmead Horton Quarry (Roman)	5	waterholes or pits		Roman	Smith, 2009
Terminal 5 Heathrow (Roman)	5	ditches and waterholes	One date of 170cal BC to 220 AD. Other features estimated from archaeology	Early to late Roman	Tetlow, 2010
St. Loyes College, Exeter	4	military wells (dumped stabling material)	<i>60 – 100 AD estimated dates from archaeology</i>	Early Roman	Smith, 2011c
Northfleet Roman Villa, Kent	8	well and cistern	<i>80 – 120 AD estimated dates from archaeology</i>	Early Roman	Smith, 2011b
Balby Carr, Catesby Buisness Park, South Yorkshire	4	brickfield Roman ditch system	<i>100 – 130 AD estimated dates from archaeology</i>	Early Roman	Smith and Tetlow 1997
Grange Park, Well. Courteenhall, Northamptonshire	1	Roman timber lined well	<i>50 -450 AD estimated dates from pottery in fill deposit</i>	2nd century	Smith, 2006
Daventry Rail Frieght Terminal (Covert Farm)	3	waterhole	<i>c. 150 – 225 AD based on pottery assemblage</i>	Mid 2 <sup>nd</sup> to mid third century	Smith, 2015b
Cambridge north West 13, Cambridgeshire	4	three wells	<i>100 BC – 250 AD? Estimated dates from archaeology</i>	Late Iron Age to mid Roman?	Smith, 2014b
Farmoor, Oxfordshire (Roman)	7	Roman wells	<i>ca. 150 – 350 AD</i>	Mid to late Roman (dated by pottery)	Robinson, 1979
Appleford, Oxfordshire	4	Roman wells and waterholes	<i>200 – 400 AD Estimated dates from archaeology and pottery</i>	Mid to late Roman	Robinson, 1981b
East Carr, Mattersey, Nottinghamshire	10	sub rectangular Roman field systems	<i>361 cal. BC to 252 cal. AD but 200 – 300 AD Estimated dates from archaeology</i>	Mid late Roman	Smith 1997
Cambridge North West 12, Cambridgeshire	4	three timber lined wells	<i>200 – 450 AD estimated dates from archaeology and pottery</i>	2nd, 3rd and 4th century	Smith, 2014b
Salford Priors, Warwickshire	4	ditch associated with villa	<i>300 – 450 AD estimated dates from archaeology</i>	3rd or 4th century	Smith and Langham, 2000
Barton Court Farm, Oxfordshire	6	ditches and other features	<i>c. 350 – 500 AD estimated dates from archaeology</i>	Late Roman to early Saxon	Robinson et al., 1984
Terminal 5 Heathrow (Medieval)	1	ditch	<i>1200 – 1600 AD estimated from archaeology</i>	Medieval	Tetlow, 2010

Radiocarbon dates calibrated using OXCAL and are to 95% probability (2 sigma). Where several dates exist from different features at the site these have been combined to give an extended age range.

Table 2. Functional groups used in this analysis (based on Hill 2015)

	Functional Group	Code	Definition
True Aquatics	Aquatic	<b>A</b>	Beetles which spend the majority of their adult life in water. <i>Not included in terrestrial sum.</i>
	Riparian	<b>R</b>	Hygrophilous taxa, littoral, usually in the bare waterlogged soils besides water. Also associated with emergent vegetation. <i>Included in terrestrial sum.</i>
Wetland & Waterside taxa	Marsh and Aquatic Plants	<b>MA</b>	Chrysomelidae and Curculionidae species which feed exclusively on marsh and aquatic plants. <i>Included in terrestrial sum.</i>
	Marsh, Fen and Carr	<b>MFC</b>	Hygrophilous, and often eurytopic taxa, found across a variety of semi-aquatic environments, such as marsh, swamp, fen, and floodplains. <i>Included in terrestrial sum.</i>
Generalists	Foul Material	<b>FM</b>	Species living on various types of foul (decaying) organic material. Sometimes, but in most cases not exclusively synanthropic. Foul material includes dung, but these taxa are not dung specialists. <i>Included in terrestrial sum.</i>
Open landscapes	Dung	<b>DUNG</b>	Taxa strongly associated with the faeces of herbivores. <i>Included in terrestrial sum.</i>
	Open and Disturbed	<b>OD</b>	Taxa found in open and vegetated, or disturbed and relatively bare conditions, wet or dry (but not strictly 'wetlands'). <i>Included in terrestrial sum.</i>
Woodland associates	Edge of, or Light, Woodland	<b>ELW</b>	Species which show strong preference to forest margins, forest-steppe, copses/felled trees within woodlands, open or pasture woods, pine heaths, hedgerows, single or sun-exposed trees (e.g. certain Elateridae), or whose larval and adult stage alternate between open spaces and forest (e.g. certain Cerambycidae). <i>Included in terrestrial sum.</i>
	Woodlands and Trees	<b>WT</b>	Includes the Coleoptera which feed on wood in varying stages of decay, leaves, fruit, and bark and live wood, fungal feeders and predators strictly associated with woodland. Except where a taxon can be defined within <b>ELW</b> . <i>Included in terrestrial sum.</i>
<b>Uncoded or Ubiquitous</b>		<b>u</b>	Taxa to which none of the other FGs can be applied owing to either lack of taxonomic resolution or ubiquity. <i>Not included in terrestrial sum.</i>

**Table 3: Rank order and % MNI for the 29 most frequently recovered Coleoptera by period for the sites in this survey. Note that some taxa (*Chaetocnema concinna*, *Aphodius sphaelatus*, *A. fimetarius*, *Megasternum 'boletophagum'*, here rendered as '*concinnum*', *Hydrobius fuscipes*, *Anotylus sculpturatus* and *Carpelimus bilineatus*) have been split since these sites were recorded. The second column lists the rank occurrence scores for the entire archaeological dataset, regardless of archaeological period, followed by the percentage of individuals as part of the whole fauna for each taxon.**

Rank order	Total for all periods	%MNI	Neolithic faunas	%MNI	Bronze age	%MNI	Iron age	%MNI	Romano-British	%MNI	Post Roman	
	39 sites in total		2 sites in period		8 sites in period		9 sites in period		18 sites in period		2 sites in period	
1	<i>Aphodius</i> spp.	7.4	<i>Megasternum concinnum</i> (Marsh.)	10.1	<i>Aphodius</i> spp.	8.9	<i>Aphodius</i> spp.	10.6	<i>Aphodius</i> spp.	5.1	<i>Helophorus brevipalpis</i> Bedel	8.4
2	<i>Ochthebius</i> spp.	3.2	<i>Ochthebius</i> spp.	7.1	<i>Geotrupes</i> spp.	6.7	<i>Helophorus brevipalpis</i> Bedel	4.7	<i>Ochthebius</i> spp.	3.7	<i>Platystethus</i> spp.	8.2
3	<i>Helophorus</i> spp.	2.7	<i>Aphodius</i> spp.	6.5	<i>Onthophagus fracticornis</i> (Preysl.)	4.7	<i>Cercyon</i> spp.	4.0	<i>Helophorus</i> spp.	3.7	<i>Aphodius</i> spp.	4.5
4	<i>Apion</i> spp.	2.4	<i>Stenus</i> spp.	4.3	<i>Calathus melanocephalus</i> (L.)	4.7	<i>Ochthebius minimus</i> (F.)	2.8	<i>Latridius minutus</i> (group)	2.6	<i>Meligethes</i> spp.	3.6
5	<i>Megasternum concinnum</i> (Marsh.)	2.2	<i>Apion</i> spp.	4.3	<i>Calathus fuscipes</i> (Goeze)	2.8	<i>Apion</i> spp.	2.8	<i>Apion</i> spp.	2.4	<i>Apion</i> spp.	3.2
6	<i>Cercyon</i> spp.	2.1	<i>Helophorus</i> spp.	3.5	<i>Onthophagus joannae</i> (Goljan)	2.6	<i>Phyllopertha horticola</i> (L.)	2.6	<i>Megasternum concinnum</i> (Marsh.)	1.9	<i>Latridius minutus</i> (group)	2.8
7	<i>Latridius minutus</i> (group)	1.9	<i>Xantholinus linearis</i> (Ol.)/ <i>longiventris</i> Heer	2.0	<i>Ochthebius</i> spp.	1.9	<i>Ochthebius</i> spp.	2.5	<i>Anotylus sculpturatus</i> (Grav.)	1.8	Ceutorhynchinae indet.	2.6
8	<i>Helophorus brevipalpis</i> Bedel	1.8	<i>Calathus melanocephalus</i> (L.)	1.9	Aleocharinae indet.	1.8	<i>Platystethus cornutus</i> (Grav.)	2.0	Aleocharinae indet.	1.8	<i>Longitarsus</i> spp.	2.4
9	<i>Anotylus sculpturatus</i> Grav.	1.4	<i>Ochthebius minimus</i> (F.)	1.9	<i>Helophorus</i> spp.	1.7	<i>Megasternum concinnum</i> (Marsh.)	1.9	<i>Cercyon</i> spp.	1.7	<i>Helophorus grandis</i> (Ill.)	2.4
10	<i>Geotrupes</i> spp.	1.4	Aleocharinae indet.	1.7	<i>Phyllotreta</i> spp.	1.5	<i>Helophorus</i> spp.	1.8	<i>Philonthus</i> spp.	1.5	<i>Phyllotreta vittula</i> (Redt.)	2.2
11	<i>Ochthebius minimus</i> (F.)	1.4	<i>Aphodius granarius</i> (L.)	1.4	<i>Hydraena testacea</i> Curt.	1.3	<i>Latridius minutus</i> (group)	1.8	<i>Platystethus cornutus</i> (Grav.)	1.3	<i>Oxyomus sylvestris</i> (Scop.)	1.9
12	Aleocharinae indet.	1.4	<i>Calathus fuscipes</i> (Goeze)	1.2	<i>Anobium punctatum</i> (Geer)	1.3	<i>Chaetocnema concinna</i> (Marsh.)	1.6	<i>Platystethus arenarius</i> (Fourcr.)	1.3	<i>Phyllotreta nigripes</i> (F.)	1.9
13	<i>Platystethus cornutus</i> (Grav.)	1.3	<i>Tachyporus</i> spp.	1.2	<i>Megasternum concinnum</i> (Marsh.)	1.2	<i>Aphodius contaminatus</i> (Hbst.)	1.4	<i>Lesteva longelytrata</i> (Goeze)	1.3	<i>Megasternum</i> (Marsh.)	1.5
14	<i>Philonthus</i> spp.	1.2	<i>Phyllopertha horticola</i> (L.)	1.2	<i>Apion</i> spp.	1.2	<i>Anotylus sculpturatus</i> (Grav.)	1.4	<i>Amara</i> spp.	1.2	Corticariinae indet.	1.5
15	<i>Calathus melanocephalus</i> (L.)	1.2	<i>Barynotus obscurus</i> (F.)	1.2	<i>Phyllopertha horticola</i> (L.)	1.2	<i>Tanysphyrus lemnae</i> (Payk.)	1.3	<i>Anobium punctatum</i> (Geer)	1.2	<i>Cercyon</i> spp.	1.3

Rank order	Total for all periods	%MNI	Neolithic faunas	%MNI	Bronze age	%MNI	Iron age	%MNI	Romano-British	%MNI	Post Roman	
16	<i>Chaetocnema concinna</i> (Marsh.)	1.2	<i>Rugilus</i> spp.	1.2	<i>Ptinus fur</i> (L.)	1.0	<i>Carpelimus bilineatus</i> (Steph.)	1.1	<i>Oxyomus sylvestris</i> (Scop.)	1.1	<i>Chaetocnema concinna</i> (Marsh.)	1.3
17	<i>Phyllopertha horticola</i> (L.)	1.1	<i>Philonthus</i> spp.	1.2	<i>Anotylus sculpturatus</i> (Grav.)	1.0	<i>Trechus obtusus</i> Er./ <i>quadristriatus</i> (Schr.)	1.1	<i>Phyllotreta</i> spp.	1.1	<i>Hydrobius fuscipes</i> (L.)	1.3
18	<i>Stenus</i> spp.	1.1	<i>Agrypnus murinus</i> (L.)	1.2	<i>Stenus</i> spp.	1.0	<i>Hydrobius fuscipes</i> (L.)	1.0	<i>Anotylus rugosus</i> (F.)	1.1	<i>Tachyporus</i> spp.	1.3
19	<i>Anobium punctatum</i> (Geer)	1	<i>Longitarsus</i> spp.	1.2	<i>Chaetocnema concinna</i> (Marsh.)	0.9	<i>Aphodius sphaelatus</i> (Panz.)	1.0	<i>Aphodius contaminatus</i> (Hbst.)	1.1	Cryptophagidae indet.	1.3
20	<i>Calathus fuscipes</i> (Goeze)	1	<i>Mecinus pyraister</i> (Hbst.)	1.2	<i>Aphodius granarius</i> (L.)	0.9	<i>Bembidion</i> spp.	1.0	<i>Stenus</i> spp.	1.1	<i>Tachinus</i> spp.	1.1
21	<i>Aphodius contaminatus</i> (Hbst.)	0.9	<i>Geotrupes</i> spp.	1.1	<i>Longitarsus</i> spp.	0.8	<i>Philonthus</i> spp.	0.9	<i>Chaetocnema concinna</i> (Marsham)	1.0	<i>Geotrupes</i> spp.	0.9
22	<i>Phyllotreta</i> spp.	0.9	<i>Gabrius</i> spp.	1.0	<i>Limnebius</i> spp.	0.8	<i>A. prodromus</i> Brahm/ <i>sphaelatus</i> (Panzer)	0.9	<i>Xantholinus</i> spp.	1.0	<i>Anotylus rugosus</i> (F.)	0.9
23	<i>Longitarsus</i> spp.	0.9	<i>Hydrothassa glabra</i> (Hbst.)	1.0	<i>Xantholinus linearis</i> (Ol.)/ <i>longiventris</i> Heer	0.8	<i>Anobium punctatum</i> (Geer)	0.9	<i>Anotylus nitidulus</i> (Grav.)	0.9	<i>Trechus obtusus</i> Er./ <i>quadristriatus</i> (Schr.)	0.9
24	<i>Oxyomus sylvestris</i> (Scop.)	0.9	<i>Tachinus</i> spp.	0.9	<i>Mecinus pyraister</i> (Hbst.)	0.8	<i>Anotylus rugosus</i> (F.)	0.8	Corticariinae indet.	0.9	<i>Atomaria</i> spp.	0.9
25	<i>Anotylus rugosus</i> (F.)	0.9	<i>Quedius</i> spp.	0.9	<i>Philonthus</i> spp.	0.8	<i>Longitarsus</i> spp.	0.7	<i>Brachypterus urticae</i> (F.)	0.9	<i>Aphodius rufipes</i> (Geer)	0.9
26	<i>Amara</i> spp.	0.8	<i>Anotylus rugosus</i> (F.)	0.8	<i>Oxyomus sylvestris</i> (Scop.)	0.8	<i>Aphodius fimetarius</i> (L.)	0.7	<i>Longitarsus</i> spp.	0.9	<i>Aspidapion aeneum</i> (F.)	0.9
27	<i>Lesteva longoelytrata</i> (Goeze)	0.8	<i>Atomaria</i> spp.	0.8	<i>Brachypterus urticae</i> (F.)	0.7	Scirtidae indet.	0.7	<i>Bembidion</i> spp.	0.9	<i>Phyllotreta atra</i> (F.)	0.9
28	<i>Platystethus arenarius</i> (Fourcr.)	0.8	<i>Onthophagus joannae</i> (Goljan)	0.8	<i>Chaetocnema</i> spp.	0.7	<i>Aphodius prodromus</i> (Brahm)	0.7	<i>Ochthebius minimus</i> (Group)	0.9	<i>Ochthebius</i> spp.	0.7
29	<i>Onthophagus fracticornis</i> (Preyssl.)	0.8	<i>Phyllobius</i> spp.	0.8	<i>Latridius minutus</i> (group)	0.7	<i>Limnebius</i> spp.	0.6	<i>Ptinus fur</i> (L.)	0.8	<i>Philonthus</i> spp.	0.7

Table 4 Summary table of the insects thought to be particularly important in archaeological farms by ecological groupings and then by taxonomic order based on Table 3 and the authors' previous experience.

PART OF FAUNA	SPECIES	ECOLOGICAL NOTES
TYPICAL GROUND BEETLES	<i>Notiophilus biguttatus</i> , <i>Nebria brevicollis</i> , <i>Loricera pilicornis</i> , <i>Clivina fossor</i> , <i>Trechus quadristriatus</i> , <i>T. obtusus</i> , <i>Pterostichus madidus</i> , <i>P. melanarius</i> , <i>Anchomenus (Agonum) dorsalis</i> , <i>Amara spp.</i> , <i>Harpalus affinis</i> , <i>Harpalus rufipes</i> , <i>Calathus fuscipes</i> , <i>Calathus melanocephalus</i> , <i>Syntomus truncatellus</i> , <i>Dromius linearis</i>	A range of 'generalist' predators favoured by open and often disturbed ground. This may be ploughed fields or grassland. <i>C. fossor</i> seems to be typical of soft damp ground, <i>S. truncatellus</i> and <i>D. linearis</i> of open sandy ground.
TYPICAL OF GRASSLANDS	Range of Elateridae 'click beetles' Typically various <i>Agriotes</i> spp. and <i>Athous haemorrhoidalis</i> . The 'Garden Chafer' <i>Phyllopertha horticola</i> , the 'cockchafer' <i>Melolontha melolontha</i> and 'the Welsh Chafer' <i>Hoplia philanthis</i> .	These species tend to be typical of old grassland, where they feed on the roots of grasses.
PLANT FEEDERS	<b><u>Grassland and meadow</u></b> <i>Gastrophysa polygoni</i> , <i>G. viridula</i> , <i>Apion frumentarium</i> , <i>Perapion violaceum</i> , <i>P. hydrolapathi</i> (on docks) <i>Sitona lepidus</i> , <i>S. suturalis</i> , <i>S. humeralis</i> , <i>Hypera</i> spp. (Clover) <i>Coelositona cambricus</i> , <i>S. waterhousei</i> (birds foot trefoil) <i>Oxystoma cracca</i> , <i>O. cerdo</i> , <i>O. pomonae</i> (Vetches) <i>Cleonis piger</i> (thistles) <b><u>Disturbed ground and wasteland</u></b> <i>Aspidapion aeneum</i> (common mallow) <i>Rhinoncus pericarpus</i> , <i>Rhinoncus castor</i> (Docks) <i>Ceutorhynchus contractus</i> (poppies and mignonettes) <i>Ceutorhynchus erysimi</i> (Shepard's purse) <i>Brachypterus urticae</i> , <i>Parethelcus (Ceutorhynchus) pollinarius</i> , <i>Nedyus quadrimaculatus</i> (stinging nettle) <i>Mecinus pyraster</i> , <i>M. (Gymnetron) labile</i> , <i>M. (Gymnetron) pascuorum</i> (lanceolate plantain)	There is substantial overlap between the species which occur in grassland and meadow and those in disturbed ground.  All these species feed on plants typical of grassland. Often they are carried into settlement in cut hay (Kenward and Hall, 1997).
NEAR TO DAMP AREAS AND MUDDY GROUND	<i>Lesteva longoelytrata</i> , <i>Carpelimus 'bilineatus'</i> , <i>Anotylus nitidulus</i> , <i>Platystethus cornutus</i>	These species are typical of open, wet and disturbed ground both in agricultural land and within settlements. They may often be associated with areas of trampled or poached ground around waterholes and channels, but this may be overly simplistic.
DUNG BEETLES	<i>Sphaeridium scarabaeoides</i> , <i>S. lunatum</i> , <i>Cercyon impressus</i> , <i>Cercyon haemorrhoidalis</i> , <i>Cercyon melanocephalus</i> , <i>Cercyon unipunctatus</i> , <i>Platystethus arenarius</i> , <i>Geotrupes spp.</i> , <i>Onthophagus spp.</i> , <i>Aphodius fossor</i> , <i>Aphodius rufipes</i> , <i>Aphodius luridus</i> , <i>A. contaminatus</i> , <i>A. sphacelatus</i> , <i>A. prodromus</i> , <i>A. porcus</i> , <i>A. fimetarius</i> , <i>A. ater</i> , <i>A. granarius</i> etc.	This is a selection of the most common of a wide range of species associated with herbivore dung. Several other species are recovered, some of which from the Bronze and Iron Ages are rare at the present day (Robinson, 2013b). Several are also thought to be capable of breeding in wet waste materials around human settlement (Kenward <i>et al.</i> , 2004). These dung associates can be very numerous in some sites where they can account for up to 40% – 60% of the terrestrial fauna, suggesting nearby grazing or pasture.
TREES	Various woodborers - <i>Anobium punctatum</i> , <i>Grynobius planus</i> , <i>Lyctus linearis</i> , <i>Phloeotribus rhododactylus</i> , <i>Hylesinus (Lepersinus) varius</i> , <i>Hylesinus toranio (oleiperda)</i> , (ash). Phytophages - <i>Andrion (Sitona) regensteiniense</i> (broom and gorse), <i>Rhynchites</i> spp. s. lat. (rosaceous shrubs such as blackthorn, hawthorn and bramble), <i>Rhynchaenus</i> spp. (leaf miners), <i>Curculio</i> spp. (nut weevils)	Most archaeological sites from agricultural ditches and rural landscapes contain very few of these taxa.

Table 5. Beetles typically recorded in modern ecological surveys from arable, grassland and pasture (taxa ordered by frequency).


Habitat type	Reference	Sampling method	Families	Species in order of occurrence
<b>Arable land</b>				
Arable lands in Scotland	Blake et al. , 2003	Pitfall trapping	Carabidae	<i>Nebria brevicollis</i> , <i>Loricera pilicornis</i> , <i>Pterostichus strenuus</i> , <i>Amara familiaris</i> , <i>Anchomenus dorsalis</i> , <i>Synuchus</i> spp. <i>Amara apricaria</i>
Middle of open crop fields in Southern England	Fournier and Loreau, 1999	Pitfall trapping	Carabidae	<i>Pterostichus melanarius</i> and <i>Trechus obtusus</i> favoured, <i>Bembidion obtusum</i> , <i>Notiophilus biguttatus</i> , <i>Nebria brevicollis</i> <i>Badister sodalis</i> also present
Mixed arable land on floodplain of river Trent	Greenwood et al., 1991	Pitfall trapping	Carabidae and Staphylinidae	<i>N. brevicollis</i> , <i>Pterostichus madidus</i> , <i>Anchomenus dorsalis</i> , <i>Amara familiaris</i> , <i>Bembidion obtusum</i> , <i>Drusilla caniculata</i> .
Ploughed land in Sussex, England	Holland and Reynolds, 2003	Pitfall trapping	Carabidae	<i>Nebria brevicollis</i> , <i>Anotylus</i> spp., <i>Harpalus rufipes</i> , <i>Pterostichus madidus</i> , <i>Harpalus affinis</i> , <i>Trechus quadristriatus</i> , <i>P. melanarius</i> , <i>Calathus fuscipes</i> , <i>Amara</i> spp., <i>Bembidion obtusum</i> , <i>Xantholinus</i> spp, <i>Notiophilus biguttatus</i> and <i>Poecilus cupreus</i> .
Ploughed fields in northern Germany	Kroos and Schaefer, 1998	Pitfall trapping	Staphylinidae	<i>Tachyporus hypnorum</i> , <i>Anotylus inustus</i> , <i>Lesteva longolytrata</i> , <i>Philonthus fuscipennis</i> , <i>Ocyopus nitens</i> , <i>Omalium caesum</i> , <i>Philonthus rotundicollis</i> and <i>Coprophilus striatulus</i>
Arable land, Binsley Lane and South Hinksey, Oxford	Robinson, 1983	Pitfalls and sweep-netting	Whole fauna	<b>Pitfall trapping</b> <i>Harpalus rufipes</i> , <i>Tachinus rufipes</i> , <i>Pterostichus melanarius</i> , <i>Amara similata</i> , Aleocharinae gen. & spp. indet., <i>Anchomenus dorsalis</i> , <i>Ceutorhynchus typhae</i> , <i>Enicmus transversus</i> , <i>Tachyporus solutus</i> . <b>Sweep-netting</b> <i>Ceutorhynchus floralis</i> , <i>Phyllotreta</i> spp., <i>Meligethes</i> spp.
<b>Grassland, meadow and pasture</b>				
Large number of undisturbed grassland sites across UK	Eyre et al., 2003	Pitfall trapping	Carabidae	Dominated by <i>Abax parallelepipedus</i> , <i>Pterostichus niger</i> and <i>Platynus assimilis</i> ; dry grassland also with <i>Calathus fuscipes</i> , <i>Amara aenea</i> , <i>Harpalus rufipes</i> . Dampier grassland with <i>Bembidion obtusum</i> , <i>Harpalus affinis</i> , <i>Dromius melanocephalus</i> .
Series of coastal grasslands	Luff, 1989	Pitfall trapping	Carabidae	<i>Badister bipustulatus</i> , <i>Amara aenea</i> , <i>A. communis</i> . <i>A. tibialis</i> , <i>Syntomus foveatus</i>
Mesotrophic Grasslands in Scotland	Blake et al., 2003	Pitfall trapping	Carabidae	<i>Notiophilus biguttatus</i> , <i>Bembidion lampros</i> , <i>Anchomenus dorsalis</i> , <i>B. guttula</i> , <i>B. lampros</i> , <i>B. aeneum</i>
Grassland on floodplain of River Trent	Greenwood et al., 1991	Pitfall trapping	Carabidae and Staphylinidae	<i>Nebria brevicollis</i> , <i>Bembidion aeneum</i> , <i>B. guttula</i> , <i>Philonthus cognatus</i> <i>Aleochara bipustula</i> .
Mesotrophic meadowland Sussex	Woodcock et al., 2008	Suction collection	Chrysomelidae and Curculionidae	<i>Longitarsus pratensis</i> , <i>L. luridus</i> , <i>Sitona lepidus</i> , <i>S. lineatus</i> , <i>L. melanocephalus</i> , <i>Trichosirocalus troglodytes</i> , <i>Protapion fulvipes</i> , <i>Sphaeroderma rubidum</i> , <i>Ischnopteraion loti</i> , <i>Protapion assimile</i> .

Habitat type	Reference	Sampling method	Families	Species in order of occurrence
Calcareous and five mesotrophic grassland sites in southern England	Woodcock et al., 2010	Suction collection	Chrysomelidae and Curculionidae	<b>Calcareous grassland</b> <i>Longitarsus pratensis</i> , <i>Sitona lineatus</i> , <i>L. luridus</i> , <i>Trachyphloeus alternans</i> , <i>L. parvulus</i> , <i>Catapion pubesens</i> , <i>S. hispidulus</i> , <i>L. atricillus</i> , <i>Protapion dichroum</i> , <i>C. seniculus</i> . <b>Mesotrophic grassland</b> <i>L. pratensis</i> , <i>P. trifolii</i> , <i>P. fulvipes</i> , <i>L. melanocephalus</i> , <i>Sitona lepidus</i> , <i>S. lineatus</i> , <i>P. assimile</i> , <i>L. atricillus</i> , <i>S. hispidulus</i> , <i>P. apricans</i>
Hay meadow, Pixey mead, Oxford	Robinson, 1983	Pitfalling and sweep netting	Whole fauna	<b>Pit fall trapping</b> <i>Pterostichus versicolor</i> , <i>Peocilus cupreus</i> , <i>Amara communis</i> , <i>Oedostethus quadripustulatus</i> , <i>Pterostichus melanarius</i> , <i>Longitarsus luridus</i> , <i>Bembidion gilvipes</i> , <i>Clivina fossor</i> , <i>Harpalus rufipes</i> , <i>Agriotes obscurus</i> . <b>Sweep netting</b> <i>Protapion trifolii</i> , <i>Cantharis nigra</i> , <i>Cantharis rufa</i> .
Grazed Pasture, Port Meadow, Oxford	Robinson, 1983	Pitfalls and sweep netting	Whole fauna	<b>Pitfall trapping</b> <i>Philonthus cognatus</i> , <i>Loricera pilicornis</i> , <i>Bembidion guttula</i> , <i>Tachinus signatus</i> , <i>Aleocharinae</i> gen and spp, indet. <i>Notaris acridulus</i> , <i>Helophorus brevipalpis</i> , <i>Bembidion properans</i> , <i>B. lunulatum</i> , <i>Philonthus laminatus</i> , <i>Aphodius fossor</i>
Grazed pasture in Southern Ireland	Hutton and Giller, 2003	Baited pitfall traps	Scarabaeidae, Hyrophilidae and Histeridae 'dung beetles'	<i>Aphodius prodromus</i> , <i>A. sphacelatus</i> , <i>A. ater</i> , <i>A. rufipes</i> , <i>A. depressus</i> , <i>Sphaeridium lunatum</i> , <i>S. scarabaeoides</i> and <i>Margarinotus ventralis</i> accounted for 95% of variation



1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47

Figure 1: Location of the archaeological sites used in this survey



Review

Figure 2: The relative proportions of the broader aquatic, wetland and terrestrial functional group for each site in chronological order (N – Neolithic, BA – Bronze Age, IA – Iron age, RB – Romano British, S – Saxon, M – Medieval).




Figure 3: The relative proportions of the terrestrial functional groups by site in chronological order (N – Neolithic, BA – Bronze Age, IA – Iron age, RB – Romano British, S – Saxon, M – Medieval).




Figure 4: The relative proportions of functional groups (dung, foul material, open and woodland) and synthropic groups by archaeological period using box and whisker plots to indicate the range and mean for each functional group. Outliers have the same numbers as are given for the sites in Figure 1. The number of sites in each archaeological period is indicated below the middle row of the diagrams.




Figure 5: nMDS analysis for of the taxa from each individual 'chronological entity' by archaeological period (see key for details of each archaeological period)






Figure 6: nMDS analysis for of the taxa from each individual 'chronological entity' by archaeological feature (see key for details of each archaeological feature)




1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47

Figure 7: Dissimilarity matrices for the insect faunas from the sites examined based on simple score of presence/absence. The left-hand diagram presents the sites ordered alphabetically by period. The right-hand diagram re-orders the sites by the degree to which their archaeoentomological faunas are similar, regardless of period; assemblages with similar faunas plot closer together. Results are presented as a heat map of the similarity matrix with blocks indicating the highest degree of similarity.

Dissimilarity Matrix



Ordered Dissimilarity Matrix



## Figures and Tables

Figure 1: Location map for the archaeological sites in this survey

Figure 2. The relative proportions of the broader aquatic, wetland and terrestrial functional groups

Figure 3: The relative proportions of the terrestrial functional groups for the archaeological sites

Figure 4. The relative proportions of functional groups (dung, foul material, open and woodland) and synanthropic groups by archaeological period

Figure 5: nMDS analysis for of the taxa from each individual 'chronological entity' by archaeological period (see key for details of each archaeological period)

Figure 6: nMDS analysis for of the taxa from each individual 'chronological entity' by archaeological feature (see key for details of each archaeological feature)

Figure 7: Dissimilarity matrices. On a simple score of presence/ absence, sites' faunas are largely dissimilar to each other (i.e. large tail of less common species heavily influential). However, those sites that are similar cluster to a large extent according to the person carrying out the analysis.

Table 1. Location, dating, feature type and reference for the archaeological sites included in this survey

Table 2. Functional groups used in this analysis (based on Hill 2015)

Table 3: Rank order and % MNI for the 29 most frequently recovered Coleoptera by period for the sites in this survey. Note that some taxa (*Chaetocnema concinna*, *Aphodius sphaelatus*, *A. fimetarius*, *Megasternum 'boletophagum'*, here rendered as '*concinnum*', *Hydrobius fuscipes*, *Anotylus sculpturatus* and *Carpelimus bilineatus*) have been split since these sites were recorded

Table 4. Summary table of the insects thought to be particularly important in archaeological farms by ecological groupings and then by taxonomic order

Table 5. Beetles typically recorded in modern ecological surveys from arable, grassland and pasture (taxa ordered by frequency)



**Table 1. Location, dating, feature type and reference for the archaeological sites included in this survey**

Site	Number of samples	Type of feature	Radiocarbon or estimated calendar dates	Period	Reference
Etton, Cambridgeshire	11	base of enclosure ditch of Causeway Camp	3725-3670 to 3310-3210 cal. BC	Early Neolithic	Robinson, 1998
Silbury Hill, Wiltshire	7	old ground surface and peat stack below mound	2550 – 2535 cal BC	Mid to late Neolithic	Robinson, 1997
Wilsford Shaft, Wiltshire	1	base of shaft / well below Bronze Age barrow	1470– 1290 cal BC	Early Bronze Age	Osborne, 1969
Terminal 5 Heathrow, Greater London (Bronze Age),	22	ditches and waterholes	ca. 1650 – 900 BC	Mid to late Bronze Age	Tetlow, 2010
Perry Oaks (terminal 5) Heathrow, Greater London (Bronze Age)	6	Bronze Age pits and waterholes	1450 – 1210 cal BC and 1380 – 1040 Cal BC	Bronze Age	Robinson, 2006
Hillfarence, Somerset	5	field ditch and pit	1450-1130 cal BC	Mid to late Bronze Age	Smith and Tetlow, 2007
Horton Quarry 71808 Berkshire	2	waterholes	ca. 1200 – 700 BC	Mid to late Bronze Age	Smith, 2014a
Kingsmead Horton Quarry (BA), Berkshire	13	waterholes	ca. 1200 – 700 BC	Mid to late Bronze Age	Smith, 2009
Imperial Sports Ground), Greater London (Bronze Age)	3	waterhole	1210-910 cal. BC	Mid to late Bronze Age	Smith, 2015a
Anslow's Cottages, Burghfield, Berkshire	5	fluvial succession	Single date of 91-431 cal. BC	Mid to late Bronze Age	Robinson, 1992
Terminal 5 Heathrow (Late Bronze Age - Iron Age)	26	ditches and waterholes	One sample 1690-1500 cal BC. 25 dated 1500-50 Cal BC	Iron Age	Tetlow, 2010
Whitemoor Haye, Staffordshire	1	series of recuts of terminal end of ring ditch	ca. 500 – 200 BC	Mid Iron Age	Smith, 2002
Kingsmead Horton Quarry, Berkshire (Late Bronze Age – Iron Age)	5	ditches and waterholes	ca. 1000 – 0 BC	Late Bronze Age - Iron Age	Smith, 2009
Enderby, Leicestershire	8	waterholes and pit features	200 – 00 BC? Estimated dates from archaeology	Late Iron Age	Hill, 2016
Olympic Park, Greater London.	10	ditches	750-260 cal. BC	Mid to late Iron Age	Smith, 2012b
Tattershall Thorpe, Lincolnshire	15	Iron Age enclosure ditch	ca. 770-200 BC based on archaeology	Mid to late Iron Age	Chowne et al., 1986
Fisherwick, Staffordshire	3	enclosure ditch	380 ca. BC – 70 cal AD	Late Iron Age	Osborne, 1979
Minges Ditches, Oxfordshire	12	enclosure ditches	390-40 cal. BC	Late Iron Age	Robinson, 1993
Farmoor, Oxfordshire (Iron Age)	6	Iron Age sumps and gullies	382 cal. BC-76 cal. AD	Late Iron Age	Robinson, 1979
Little Paxton, Cambridgeshire	10	waterholes, brickfield ditches	80 – 250 AD dated by pottery	Late First century to mid third century	Smith, 2011a
Lockington, Derbyshire, waterholes	2	waterholes	Estimated period dates from archaeology	'Roman'	Smith, 2012c

Site	Number of samples	Type of feature	Radiocarbon or estimated calendar dates	Period	Reference
Imperial Sports Ground, Greater London (Roman)	2	wells	240-510 cal. AD	Roman	Smith, 2015a
Brickets Wood Bund, Hertfordshire	3	waterholes	<i>Estimated period dates from archaeology</i>	'Roman'	Smith, 2013b
Perry Oaks (Terminal 5) Heathrow, Greater London (Roman)	4	waterholes	<i>c100-250 AD and c 200 -300 AD based on archaeology</i>	Roman	Robinson, 2006
Kingsmead Horton Quarry (Roman)	5	waterholes or pits		Roman	Smith, 2009
Terminal 5 Heathrow (Roman)	5	ditches and waterholes	One date of 170cal BC to 220 AD. Other features estimated from archaeology	Early to late Roman	Tetlow, 2010
St. Loyes College, Exeter	4	military wells (dumped stabling material)	<i>60 – 100 AD estimated dates from archaeology</i>	Early Roman	Smith, 2011c
Northfleet Roman Villa, Kent	8	well and cistern	<i>80 – 120 AD estimated dates from archaeology</i>	Early Roman	Smith, 2011b
Balby Carr, Catesby Buisness Park, South Yorkshire	4	brickfield Roman ditch system	<i>100 – 130 AD estimated dates from archaeology</i>	Early Roman	Smith and Tetlow 1997
Grange Park, Well. Courteenhall, Northamptonshire	1	Roman timber lined well	<i>50 -450 AD estimated dates from pottery in fill deposit</i>	2nd century	Smith, 2006
Daventry Rail Frieght Terminal (Covert Farm)	3	waterhole	<i>c. 150 – 225 AD based on pottery assemblage</i>	Mid 2 <sup>nd</sup> to mid third century	Smith, 2015b
Cambridge north West 13, Cambridgeshire	4	three wells	<i>100 BC – 250 AD? Estimated dates from archaeology</i>	Late Iron Age to mid Roman?	Smith, 2014b
Farmoor, Oxfordshire (Roman)	7	Roman wells	<i>ca. 150 – 350 AD</i>	Mid to late Roman (dated by pottery)	Robinson, 1979
Appleford, Oxfordshire	4	Roman wells and waterholes	<i>200 – 400 AD Estimated dates from archaeology and pottery</i>	Mid to late Roman	Robinson, 1981b
East Carr, Mattersey, Nottinghamshire	10	sub rectangular Roman field systems	<i>361 cal. BC to 252 cal. AD but 200 – 300 AD Estimated dates from archaeology</i>	Mid late Roman	Smith 1997
Cambridge North West 12, Cambridgeshire	4	three timber lined wells	<i>200 – 450 AD estimated dates from archaeology and pottery</i>	2nd, 3rd and 4th century	Smith, 2014b
Salford Priors, Warwickshire	4	ditch associated with villa	<i>300 – 450 AD estimated dates from archaeology</i>	3rd or 4th century	Smith and Langham, 2000
Barton Court Farm, Oxfordshire	6	ditches and other features	<i>c. 350 – 500 AD estimated dates from archaeology</i>	Late Roman to early Saxon	Robinson et al., 1984
Terminal 5 Heathrow (Medieval)	1	ditch	<i>1200 – 1600 AD estimated from archaeology</i>	Medieval	Tetlow, 2010

Radiocarbon dates calibrated using OXCAL and are to 95% probability (2 sigma). Where several dates exist from different features at the site these have been combined to give an extended age range.

Table 2. Functional groups used in this analysis (based on Hill 2015)

	Functional Group	Code	Definition
True Aquatics	Aquatic	<b>A</b>	Beetles which spend the majority of their adult life in water. <i>Not included in terrestrial sum.</i>
	Riparian	<b>R</b>	Hygrophilous taxa, littoral, usually in the bare waterlogged soils besides water. Also associated with emergent vegetation. <i>Included in terrestrial sum.</i>
Wetland & Waterside taxa	Marsh and Aquatic Plants	<b>MA</b>	Chrysomelidae and Curculionidae species which feed exclusively on marsh and aquatic plants. <i>Included in terrestrial sum.</i>
	Marsh, Fen and Carr	<b>MFC</b>	Hygrophilous, and often eurytopic taxa, found across a variety of semi-aquatic environments, such as marsh, swamp, fen, and floodplains. <i>Included in terrestrial sum.</i>
Generalists	Foul Material	<b>FM</b>	Species living on various types of foul (decaying) organic material. Sometimes, but in most cases not exclusively synanthropic. Foul material includes dung, but these taxa are not dung specialists. <i>Included in terrestrial sum.</i>
Open landscapes	Dung	<b>DUNG</b>	Taxa strongly associated with the faeces of herbivores. <i>Included in terrestrial sum.</i>
	Open and Disturbed	<b>OD</b>	Taxa found in open and vegetated, or disturbed and relatively bare conditions, wet or dry (but not strictly 'wetlands'). <i>Included in terrestrial sum.</i>
Woodland associates	Edge of, or Light, Woodland	<b>ELW</b>	Species which show strong preference to forest margins, forest-steppe, copses/felled trees within woodlands, open or pasture woods, pine heaths, hedgerows, single or sun-exposed trees (e.g. certain Elateridae), or whose larval and adult stage alternate between open spaces and forest (e.g. certain Cerambycidae). <i>Included in terrestrial sum.</i>
	Woodlands and Trees	<b>WT</b>	Includes the Coleoptera which feed on wood in varying stages of decay, leaves, fruit, and bark and live wood, fungal feeders and predators strictly associated with woodland. Except where a taxon can be defined within <b>ELW</b> . <i>Included in terrestrial sum.</i>
<b>Uncoded or Ubiquitous</b>		<b>u</b>	Taxa to which none of the other FGs can be applied owing to either lack of taxonomic resolution or ubiquity. <i>Not included in terrestrial sum.</i>

**Table 3: Rank order and % MNI for the 29 most frequently recovered Coleoptera by period for the sites in this survey. Note that some taxa (*Chaetocnema concinna*, *Aphodius sphaelatus*, *A. fimetarius*, *Megasternum 'boletophagum'*, here rendered as '*concinnum*', *Hydrobius fuscipes*, *Anotylus sculpturatus* and *Carpelimus bilineatus*) have been split since these sites were recorded. The second column lists the rank occurrence scores for the entire archaeological dataset, regardless of archaeological period, followed by the percentage of individuals as part of the whole fauna for each taxon.**

Rank order	Total for all periods	%MNI	Neolithic faunas	%MNI	Bronze age	%MNI	Iron age	%MNI	Romano-British	%MNI	Post Roman	
	39 sites in total		2 sites in period		8 sites in period		9 sites in period		18 sites in period		2 sites in period	
1	<i>Aphodius</i> spp.	7.4	<i>Megasternum concinnum</i> (Marsh.)	10.1	<i>Aphodius</i> spp.	8.9	<i>Aphodius</i> spp.	10.6	<i>Aphodius</i> spp.	5.1	<i>Helophorus brevipalpis</i> Bedel	8.4
2	<i>Ochthebius</i> spp.	3.2	<i>Ochthebius</i> spp.	7.1	<i>Geotrupes</i> spp.	6.7	<i>Helophorus brevipalpis</i> Bedel	4.7	<i>Ochthebius</i> spp.	3.7	<i>Platystethus</i> spp.	8.2
3	<i>Helophorus</i> spp.	2.7	<i>Aphodius</i> spp.	6.5	<i>Onthophagus fracticornis</i> (Preysl.)	4.7	<i>Cercyon</i> spp.	4.0	<i>Helophorus</i> spp.	3.7	<i>Aphodius</i> spp.	4.5
4	<i>Apion</i> spp.	2.4	<i>Stenus</i> spp.	4.3	<i>Calathus melanocephalus</i> (L.)	4.7	<i>Ochthebius minimus</i> (F.)	2.8	<i>Latridius minutus</i> (group)	2.6	<i>Meligethes</i> spp.	3.6
5	<i>Megasternum concinnum</i> (Marsh.)	2.2	<i>Apion</i> spp.	4.3	<i>Calathus fuscipes</i> (Goeze)	2.8	<i>Apion</i> spp.	2.8	<i>Apion</i> spp.	2.4	<i>Apion</i> spp.	3.2
6	<i>Cercyon</i> spp.	2.1	<i>Helophorus</i> spp.	3.5	<i>Onthophagus joannae</i> (Goljan)	2.6	<i>Phyllopertha horticola</i> (L.)	2.6	<i>Megasternum concinnum</i> (Marsh.)	1.9	<i>Latridius minutus</i> (group)	2.8
7	<i>Latridius minutus</i> (group)	1.9	<i>Xantholinus linearis</i> (Ol.)/ <i>longiventris</i> Heer	2.0	<i>Ochthebius</i> spp.	1.9	<i>Ochthebius</i> spp.	2.5	<i>Anotylus sculpturatus</i> (Grav.)	1.8	Ceutorhynchinae indet.	2.6
8	<i>Helophorus brevipalpis</i> Bedel	1.8	<i>Calathus melanocephalus</i> (L.)	1.9	Aleocharinae indet.	1.8	<i>Platystethus cornutus</i> (Grav.)	2.0	Aleocharinae indet.	1.8	<i>Longitarsus</i> spp.	2.4
9	<i>Anotylus sculpturatus</i> Grav.	1.4	<i>Ochthebius minimus</i> (F.)	1.9	<i>Helophorus</i> spp.	1.7	<i>Megasternum concinnum</i> (Marsh.)	1.9	<i>Cercyon</i> spp.	1.7	<i>Helophorus grandis</i> (Ill.)	2.4
10	<i>Geotrupes</i> spp.	1.4	Aleocharinae indet.	1.7	<i>Phyllotreta</i> spp.	1.5	<i>Helophorus</i> spp.	1.8	<i>Philonthus</i> spp.	1.5	<i>Phyllotreta vittula</i> (Redt.)	2.2
11	<i>Ochthebius minimus</i> (F.)	1.4	<i>Aphodius granarius</i> (L.)	1.4	<i>Hydraena testacea</i> Curt.	1.3	<i>Latridius minutus</i> (group)	1.8	<i>Platystethus cornutus</i> (Grav.)	1.3	<i>Oxyomus sylvestris</i> (Scop.)	1.9
12	Aleocharinae indet.	1.4	<i>Calathus fuscipes</i> (Goeze)	1.2	<i>Anobium punctatum</i> (Geer)	1.3	<i>Chaetocnema concinna</i> (Marsh.)	1.6	<i>Platystethus arenarius</i> (Fourcr.)	1.3	<i>Phyllotreta nigripes</i> (F.)	1.9
13	<i>Platystethus cornutus</i> (Grav.)	1.3	<i>Tachyporus</i> spp.	1.2	<i>Megasternum concinnum</i> (Marsh.)	1.2	<i>Aphodius contaminatus</i> (Hbst.)	1.4	<i>Lesteva longelytrata</i> (Goeze)	1.3	<i>Megasternum</i> (Marsh.)	1.5
14	<i>Philonthus</i> spp.	1.2	<i>Phyllopertha horticola</i> (L.)	1.2	<i>Apion</i> spp.	1.2	<i>Anotylus sculpturatus</i> (Grav.)	1.4	<i>Amara</i> spp.	1.2	Corticariinae indet.	1.5
15	<i>Calathus melanocephalus</i> (L.)	1.2	<i>Barynotus obscurus</i> (F.)	1.2	<i>Phyllopertha horticola</i> (L.)	1.2	<i>Tanysphyrus lemnae</i> (Payk.)	1.3	<i>Anobium punctatum</i> (Geer)	1.2	<i>Cercyon</i> spp.	1.3

Rank order	Total for all periods	%MNI	Neolithic faunas	%MNI	Bronze age	%MNI	Iron age	%MNI	Romano-British	%MNI	Post Roman	
16	<i>Chaetocnema concinna</i> (Marsh.)	1.2	<i>Rugilus</i> spp.	1.2	<i>Ptinus fur</i> (L.)	1.0	<i>Carpelimus bilineatus</i> (Steph.)	1.1	<i>Oxyomus sylvestris</i> (Scop.)	1.1	<i>Chaetocnema concinna</i> (Marsh.)	1.3
17	<i>Phyllopertha horticola</i> (L.)	1.1	<i>Philonthus</i> spp.	1.2	<i>Anotylus sculpturatus</i> (Grav.)	1.0	<i>Trechus obtusus</i> Er./ <i>quadristriatus</i> (Schr.)	1.1	<i>Phyllotreta</i> spp.	1.1	<i>Hydrobius fuscipes</i> (L.)	1.3
18	<i>Stenus</i> spp.	1.1	<i>Agrypnus murinus</i> (L.)	1.2	<i>Stenus</i> spp.	1.0	<i>Hydrobius fuscipes</i> (L.)	1.0	<i>Anotylus rugosus</i> (F.)	1.1	<i>Tachyporus</i> spp.	1.3
19	<i>Anobium punctatum</i> (Geer)	1	<i>Longitarsus</i> spp.	1.2	<i>Chaetocnema concinna</i> (Marsh.)	0.9	<i>Aphodius sphaelatus</i> (Panz.)	1.0	<i>Aphodius contaminatus</i> (Hbst.)	1.1	Cryptophagidae indet.	1.3
20	<i>Calathus fuscipes</i> (Goeze)	1	<i>Mecinus pyraaster</i> (Hbst.)	1.2	<i>Aphodius granarius</i> (L.)	0.9	<i>Bembidion</i> spp.	1.0	<i>Stenus</i> spp.	1.1	<i>Tachinus</i> spp.	1.1
21	<i>Aphodius contaminatus</i> (Hbst.)	0.9	<i>Geotrupes</i> spp.	1.1	<i>Longitarsus</i> spp.	0.8	<i>Philonthus</i> spp.	0.9	<i>Chaetocnema concinna</i> (Marsham)	1.0	<i>Geotrupes</i> spp.	0.9
22	<i>Phyllotreta</i> spp.	0.9	<i>Gabrius</i> spp.	1.0	<i>Limnebius</i> spp.	0.8	<i>A. prodromus</i> Brahm/ <i>sphaelatus</i> (Panzer)	0.9	<i>Xantholinus</i> spp.	1.0	<i>Anotylus rugosus</i> (F.)	0.9
23	<i>Longitarsus</i> spp.	0.9	<i>Hydrothassa glabra</i> (Hbst.)	1.0	<i>Xantholinus linearis</i> (Ol.)/ <i>longiventris</i> Heer	0.8	<i>Anobium punctatum</i> (Geer)	0.9	<i>Anotylus nitidulus</i> (Grav.)	0.9	<i>Trechus obtusus</i> Er./ <i>quadristriatus</i> (Schr.)	0.9
24	<i>Oxyomus sylvestris</i> (Scop.)	0.9	<i>Tachinus</i> spp.	0.9	<i>Mecinus pyraaster</i> (Hbst.)	0.8	<i>Anotylus rugosus</i> (F.)	0.8	Corticariinae indet.	0.9	<i>Atomaria</i> spp.	0.9
25	<i>Anotylus rugosus</i> (F.)	0.9	<i>Quedius</i> spp.	0.9	<i>Philonthus</i> spp.	0.8	<i>Longitarsus</i> spp.	0.7	<i>Brachypterus urticae</i> (F.)	0.9	<i>Aphodius rufipes</i> (Geer)	0.9
26	<i>Amara</i> spp.	0.8	<i>Anotylus rugosus</i> (F.)	0.8	<i>Oxyomus sylvestris</i> (Scop.)	0.8	<i>Aphodius fimetarius</i> (L.)	0.7	<i>Longitarsus</i> spp.	0.9	<i>Aspidapion aeneum</i> (F.)	0.9
27	<i>Lesteva longoelytrata</i> (Goeze)	0.8	<i>Atomaria</i> spp.	0.8	<i>Brachypterus urticae</i> (F.)	0.7	Scirtidae indet.	0.7	<i>Bembidion</i> spp.	0.9	<i>Phyllotreta atra</i> (F.)	0.9
28	<i>Platystethus arenarius</i> (Fourcr.)	0.8	<i>Onthophagus joannae</i> (Goljan)	0.8	<i>Chaetocnema</i> spp.	0.7	<i>Aphodius prodromus</i> (Brahm)	0.7	<i>Ochthebius minimus</i> (Group)	0.9	<i>Ochthebius</i> spp.	0.7
29	<i>Onthophagus fracticornis</i> (Preyssl.)	0.8	<i>Phyllobius</i> spp.	0.8	<i>Latridius minutus</i> (group)	0.7	<i>Limnebius</i> spp.	0.6	<i>Ptinus fur</i> (L.)	0.8	<i>Philonthus</i> spp.	0.7

Table 4 Summary table of the insects thought to be particularly important in archaeological farms by ecological groupings and then by taxonomic order based on Table 3 and the authors' previous experience.

PART OF FAUNA	SPECIES	ECOLOGICAL NOTES
TYPICAL GROUND BEETLES	<i>Notiophilus biguttatus</i> , <i>Nebria brevicollis</i> , <i>Loricera pilicornis</i> , <i>Clivina fossor</i> , <i>Trechus quadristriatus</i> , <i>T. obtusus</i> , <i>Pterostichus madidus</i> , <i>P. melanarius</i> , <i>Anchomenus (Agonum) dorsalis</i> , <i>Amara spp.</i> , <i>Harpalus affinis</i> , <i>Harpalus rufipes</i> , <i>Calathus fuscipes</i> , <i>Calathus melanocephalus</i> , <i>Syntomus truncatellus</i> , <i>Dromius linearis</i>	A range of 'generalist' predators favoured by open and often disturbed ground. This may be ploughed fields or grassland. <i>C. fossor</i> seems to be typical of soft damp ground, <i>S. truncatellus</i> and <i>D. linearis</i> of open sandy ground.
TYPICAL OF GRASSLANDS	Range of Elateridae 'click beetles' Typically various <i>Agriotes</i> spp. and <i>Athous haemorrhoidalis</i> . The 'Garden Chafer' <i>Phyllopertha horticola</i> , the 'cockchafer' <i>Melolontha melolontha</i> and 'the Welsh Chafer' <i>Hoplia philanthis</i> .	These species tend to be typical of old grassland, where they feed on the roots of grasses.
PLANT FEEDERS	<b><u>Grassland and meadow</u></b> <i>Gastrophysa polygoni</i> , <i>G. viridula</i> , <i>Apion frumentarium</i> , <i>Perapion violaceum</i> , <i>P. hydrolapathi</i> (on docks) <i>Sitona lepidus</i> , <i>S. suturalis</i> , <i>S. humeralis</i> , <i>Hypera</i> spp. (Clover) <i>Coelositona cambricus</i> , <i>S. waterhousei</i> (birds foot trefoil) <i>Oxystoma cracca</i> , <i>O. cerdo</i> , <i>O. pomonae</i> (Vetches) <i>Cleonis piger</i> (thistles) <b><u>Disturbed ground and wasteland</u></b> <i>Aspidapion aeneum</i> (common mallow) <i>Rhinoncus pericarpus</i> , <i>Rhinoncus castor</i> (Docks) <i>Ceutorhynchus contractus</i> (poppies and mignonettes) <i>Ceutorhynchus erysimi</i> (Shepard's purse) <i>Brachypterus urticae</i> , <i>Parethelcus (Ceutorhynchus) pollinarius</i> , <i>Nedyus quadrimaculatus</i> (stinging nettle) <i>Mecinus pyraster</i> , <i>M. (Gymnetron) labile</i> , <i>M. (Gymnetron) pascuorum</i> (lanceolate plantain)	There is substantial overlap between the species which occur in grassland and meadow and those in disturbed ground.  All these species feed on plants typical of grassland. Often they are carried into settlement in cut hay (Kenward and Hall, 1997).
NEAR TO DAMP AREAS AND MUDDY GROUND	<i>Lesteva longoelytrata</i> , <i>Carpelimus 'bilineatus'</i> , <i>Anotylus nitidulus</i> , <i>Platystethus cornutus</i>	These species are typical of open, wet and disturbed ground both in agricultural land and within settlements. They may often be associated with areas of trampled or poached ground around waterholes and channels, but this may be overly simplistic.
DUNG BEETLES	<i>Sphaeridium scarabaeoides</i> , <i>S. lunatum</i> , <i>Cercyon impressus</i> , <i>Cercyon haemorrhoidalis</i> , <i>Cercyon melanocephalus</i> , <i>Cercyon unipunctatus</i> , <i>Platystethus arenarius</i> , <i>Geotrupes spp.</i> , <i>Onthophagus spp.</i> , <i>Aphodius fossor</i> , <i>Aphodius rufipes</i> , <i>Aphodius luridus</i> , <i>A. contaminatus</i> , <i>A. sphacelatus</i> , <i>A. prodromus</i> , <i>A. porcus</i> , <i>A. fimetarius</i> , <i>A. ater</i> , <i>A. granarius</i> etc.	This is a selection of the most common of a wide range of species associated with herbivore dung. Several other species are recovered, some of which from the Bronze and Iron Ages are rare at the present day (Robinson, 2013b). Several are also thought to be capable of breeding in wet waste materials around human settlement (Kenward <i>et al.</i> , 2004). These dung associates can be very numerous in some sites where they can account for up to 40% – 60% of the terrestrial fauna, suggesting nearby grazing or pasture.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47

TREES	<p>Various woodborers - <i>Anobium punctatum</i>, <i>Grynobius planus</i>, <i>Lyctus linearis</i>, <i>Phloeotribus rhododactylus</i>, <i>Hylesinus (Leperisinus) varius</i>, <i>Hylesinus toranio (oleiperda)</i>, (ash).  Phytophages - <i>Andrion (Sitona) regensteinense</i> (broom and gorse), <i>Rhynchites</i> spp. s. lat. (rosaceous shrubs such as blackthorn, hawthorn and bramble), <i>Rhynchaenus</i> spp. (leaf miners), <i>Curculio</i> spp. (nut weevils)</p>	<p>Most archaeological sites from agricultural ditches and rural landscapes contain very few of these taxa.</p>
-------	--	---

For Peer Review

Table 5. Beetles typically recorded in modern ecological surveys from arable, grassland and pasture (taxa ordered by frequency).


Habitat type	Reference	Sampling method	Families	Species in order of occurrence
<b>Arable land</b>				
Arable lands in Scotland	Blake et al. , 2003	Pitfall trapping	Carabidae	<i>Nebria brevicollis</i> , <i>Loricera pilicornis</i> , <i>Pterostichus strenuus</i> , <i>Amara familiaris</i> , <i>Anchomenus dorsalis</i> , <i>Synuchus</i> spp. <i>Amara apricaria</i>
Middle of open crop fields in Southern England	Fournier and Loreau, 1999	Pitfall trapping	Carabidae	<i>Pterostichus melanarius</i> and <i>Trechus obtusus</i> favoured, <i>Bembidion obtusum</i> , <i>Notiophilus biguttatus</i> , <i>Nebria brevicollis</i> <i>Badister sodalis</i> also present
Mixed arable land on floodplain of river Trent	Greenwood et al., 1991	Pitfall trapping	Carabidae and Staphylinidae	<i>N. brevicollis</i> , <i>Pterostichus madidus</i> , <i>Anchomenus dorsalis</i> , <i>Amara familiaris</i> , <i>Bembidion obtusum</i> , <i>Drusilla caniculata</i> .
Ploughed land in Sussex, England	Holland and Reynolds, 2003	Pitfall trapping	Carabidae	<i>Nebria brevicollis</i> , <i>Anotylus</i> spp., <i>Harpalus rufipes</i> , <i>Pterostichus madidus</i> , <i>Harpalus affinus</i> , <i>Trechus quadristriatus</i> , <i>P. melanarius</i> , <i>Calathus fuscipes</i> , <i>Amara</i> spp., <i>Bembidion obtusum</i> , <i>Xantholinus</i> spp, <i>Notiophilus biguttatus</i> and <i>Poecilus cupreus</i> .
Ploughed fields in northern Germany	Kroos and Schaefer, 1998	Pitfall trapping	Staphylinidae	<i>Tachyporus hypnorum</i> , <i>Anotylus inustus</i> , <i>Lesteva longoelytrata</i> , <i>Philonthus fuscipennis</i> , <i>Ocypus nitens</i> , <i>Omalium caesum</i> , <i>Philonthus rotundicollis</i> and <i>Coprophilus striatulus</i>
Arable land, Binsley Lane and South Hinksey, Oxford	Robinson, 1983	Pitfalls and sweep-netting	Whole fauna	<b>Pitfall trapping</b> <i>Harpalus rufipes</i> , <i>Tachinus rufipes</i> , <i>Pterostichus melanarius</i> , <i>Amara similata</i> , Aleocharinae gen. & spp. indet., <i>Anchomenus dorsalis</i> , <i>Ceutorhynchus typhae</i> , <i>Enicmus transversus</i> , <i>Tachyporus solutus</i> . <b>Sweep-netting</b> <i>Ceutorhynchus floralis</i> , <i>Phyllotreta</i> spp., <i>Meligethes</i> spp.
<b>Grassland, meadow and pasture</b>				
Large number of undisturbed grassland sites across UK	Eyre et al., 2003	Pitfall trapping	Carabidae	Dominated by <i>Abax parallelepipedus</i> , <i>Pterostichus niger</i> and <i>Platynus assimilis</i> ; dry grassland also with <i>Calathus fuscipes</i> , <i>Amara aenea</i> , <i>Harpalus rufipes</i> . Dampier grassland with <i>Bembidion obtusum</i> , <i>Harpalus affinus</i> , <i>Dromius melanocephalus</i> .
Series of coastal grasslands	Luff, 1989	Pitfall trapping	Carabidae	<i>Badister bipustulatus</i> , <i>Amara aenea</i> , <i>A. communis</i> . <i>A. tibialis</i> , <i>Syntomus foveatus</i>
Mesotrophic Grasslands in Scotland	Blake et al., 2003	Pitfall trapping	Carabidae	<i>Notiophilus biguttatus</i> , <i>Bembidion lampros</i> , <i>Anchomenus dorsalis</i> , <i>B. guttula</i> , <i>B. lampros</i> , <i>B. aeneum</i>
Grassland on floodplain of River Trent	Greenwood et al., 1991	Pitfall trapping	Carabidae and Staphylinidae	<i>Nebria brevicollis</i> , <i>Bembidion aeneum</i> , <i>B. guttula</i> , <i>Philonthus cognatus</i> <i>Aleochara bipustula</i> .
Mesotrophic meadowland Sussex	Woodcock et al., 2008	Suction collection	Chrysomelidae and Curculionidae	<i>Longitarsus pratensis</i> , <i>L. luridus</i> , <i>Sitona lepidus</i> , <i>S. lineatus</i> , <i>L. melanocephalus</i> , <i>Trichosirocalus troglodytes</i> , <i>Protapion fulvipes</i> , <i>Sphaeroderma rubidum</i> , <i>Ischnopteration loti</i> , <i>Protapion assimile</i> .



1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47

Habitat type	Reference	Sampling method	Families	Species in order of occurrence
Calcareous and five mesotrophic grassland sites in southern England	Woodcock et al., 2010	Suction collection	Chrysomelidae and Curculionidae	<b>Calcareous grassland</b> <i>Longitarsus pratensis</i> , <i>Sitona lineatus</i> , <i>L. luridus</i> , <i>Trachyploeus alternans</i> , <i>L. parvulus</i> , <i>Catapion pubesens</i> , <i>S. hispidulus</i> , <i>L. atricillus</i> , <i>Protapion dichroum</i> , <i>C. seniculus</i> . <b>Mesotrophic grassland</b> <i>L. pratensis</i> , <i>P. trifolii</i> , <i>P. fulvipes</i> , <i>L. melanocephalus</i> , <i>Sitona lepidus</i> , <i>S. lineatus</i> , <i>P. assimile</i> , <i>L. atricillus</i> , <i>S. hispidulus</i> , <i>P. apricans</i>
Hay meadow, Pixey mead, Oxford	Robinson, 1983	Pitfalling and sweep netting	Whole fauna	<b>Pit fall trapping</b> <i>Pterostichus versicolor</i> , <i>Peocilus cupreus</i> , <i>Amara communis</i> , <i>Oedostethus quadripustulatus</i> , <i>Pterostichus melanarius</i> , <i>Longitarsus luridus</i> , <i>Bembidion gilvipes</i> , <i>Clivina fossor</i> , <i>Harpalus rufipes</i> , <i>Agriotes obscurus</i> . <b>Sweep netting</b> <i>Protapion trifolii</i> , <i>Cantharis nigra</i> , <i>Cantharis rufa</i> .
Grazed Pasture, Port Meadow, Oxford	Robinson, 1983	Pitfalls and sweep netting	Whole fauna	<b>Pitfall trapping</b> <i>Philonthus cognatus</i> , <i>Loricera pilicornis</i> , <i>Bembidion guttula</i> , <i>Tachinus signatus</i> , <i>Aleocharinae</i> gen and spp, indet. <i>Notaris acridulus</i> , <i>Helophorus brevipalpis</i> , <i>Bembidion properans</i> , <i>B. lunulatum</i> , <i>Philonthus laminatus</i> , <i>Aphodius fossor</i>
Grazed pasture in Southern Ireland	Hutton and Giller, 2003	Baited pitfall traps	Scarabaeidae, Hyrophilidae and Histeridae 'dung beetles'	<i>Aphodius prodromus</i> , <i>A. sphacelatus</i> , <i>A. ater</i> , <i>A. rufipes</i> , <i>A. depressus</i> , <i>Sphaeridium lunatum</i> , <i>S. scarabaeoides</i> and <i>Margarinotus ventralis</i> accounted for 95% of variation

Figure 1: Location of the archaeological sites used in this survey



view

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47

**Figure 2: The relative proportions of the broader aquatic, wetland and terrestrial functional group for each site in chronological order (N – Neolithic, BA – Bronze Age, IA – Iron age, RB – Romano British, S – Saxon, M – Medieval).**





Figure 3: The relative proportions of the terrestrial functional groups by site in chronological order (N – Neolithic, BA – Bronze Age, IA – Iron age, RB – Romano British, S – Saxon, M – Medieval).



1  
2  
3  
4  
5  
6  
7 **Figure 4: The relative proportions of functional groups (dung, foul material, open and woodland) and synthropic groups by archaeological period using**  
8 **box and whisker plots to indicate the range and mean for each functional group. Outliers have the same numbers as are given for the sites in Figure 1.**  
9 **The number of sites in each archaeological period is indicated below the middle row of the diagrammes.**  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47





Figure 5: nMDS analysis for of the taxa from each individual 'chronological entity' by archaeological period (see key for details of each archaeological period)



1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47

Figure 6: nMDS analysis for of the taxa from each individual 'chronological entity' by archaeological feature (see key for details of each archaeological feature)




Figure 7: Dissimilarity matrices for the insect faunas from the sites examined based on simple score of presence/absence. The left-hand diagram presents the sites ordered alphabetically by period. The right-hand diagram re-orders the sites by the degree to which their archaeoentomological faunas are similar, regardless of period; assemblages with similar faunas plot closer together. Results are presented as a heat map of the similarity matrix with blocks indicating the highest degree of similarity.

