# UNIVERSITY BIRMINGHAM University of Birmingham Research at Birmingham

# Delayed reactive distractor suppression in aging populations

Ashinoff, Brandon; Geng, Joy; Mevorach, Carmel

*DOI:* 10.1037/pag0000336

*License:* None: All rights reserved

Document Version Peer reviewed version

# Citation for published version (Harvard):

Ashinoff, B, Geng, J & Mevorach, C 2019, 'Delayed reactive distractor suppression in aging populations', *Psychology and Aging*, vol. 34, no. 3, pp. 418-430. https://doi.org/10.1037/pag0000336

Link to publication on Research at Birmingham portal

#### Publisher Rights Statement: Checked for eligibility 06/02/2019

©American Psychological Association, 2019. This paper is not the copy of record and may not exactly replicate the authoritative document published in the APA journal. Please do not copy or cite without author's permission. The final article is available, upon publication, at: http://dx.doi.org/10.1037/pag0000336

Copyright 2019 APA, all rights reserved

#### **General rights**

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

•Users may freely distribute the URL that is used to identify this publication.

•Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.

•User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?) •Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

#### Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

# Delayed Reactive Distractor Suppression in Aging Populations

Brandon K. Ashinoff

University of Birmingham and Columbia University

Joy Geng

University of California - Davis

Carmel Mevorach

University of Birmingham

Authors Note:

Brandon K. Ashinoff, School of Psychology and Centre for Human Brain Health, University of Birmingham, Birmingham, United Kingdom; Department of Psychiatry, Columbia University; Joy Geng, Department of Psychology, University of California – Davis; Carmel Mevorach, School of Psychology and Centre for Human Brain Health, University of Birmingham, Birmingham, United Kingdom. Correspondence regarding this manuscript should be addressed to Brandon K. Ashinoff: bka2115@cumc.columbia.edu.

The authors would like to thank Denise Clissett for her critical role in the recruitment of older participants, as well as Samantha Ellis, Charlie-Louise Milward, and Leah Supra for their contributions to this project as undergraduate research assistants.

This paper is based on chapter 3 of author B.K.A.'s dissertation (Ashinoff, 2017). All raw data and experimental code for this study is publicly available on the Open Science Framework (OSF): <a href="https://osf.io/3h7ux/">https://osf.io/3h7ux/</a>

Date: 05/02/2019 – This manuscript has not been published yet.

# Abstract

Previous studies have tended to infer that reactive control is intact in aging populations because of evidence that proactive control is impaired and that older participants appear to favor reactive control strategies. However, most of these studies did not compare reactive control in young and older participants directly. In our study, a young (18-21 years old) and older (60 + years old) cohort engaged in a task that assesses reactive distractor suppression where subjects had to discriminate between an upright and inverted t-shape in the presence of a salient or non-salient distractor. In previous studies using this paradigm (DiQuattro and Geng, 2010) young participants reactively used the salient distractor as an anti-cue and performed better (faster RT and higher accuracy) when it was present. It was found that older participants were not able to reactively suppress the salient distractor with a 200 msec display but were able to do so with a 600 msec display. It was concluded that the initiation of reactive suppression is delayed for older participants, but that effective suppression is possible given enough time.

Keywords: Cognitive Aging, Cognitive Control, Distractor Suppression, Reactive Control, Salience Suppression

#### Introduction

2 It is generally accepted that normal aging can lead to declines in cognitive performance 3 (Braver and Barch, 2002; Craik and Salthouse, 2011; Andrews-Hannah et al, 2007; Geerligs et al, 4 2014a; 2014b; Grady, 2012; Larson et al, 2016; Li et al, 2001; 2016; Persson et al, 2006; Zanto et al, 5 2010). More specifically, there is a plethora of evidence highlighting impairments in inhibition 6 mechanisms (Hasher and Zacks, 1988; Bauer et al, 2012; Mayas et al, 2012; Gazzaley et al, 2005; but 7 see Frings et al, 2015 for contrasting findings). Previous studies have highlighted the increased 8 interference experienced from irrelevant distractors in old age across different experimental 9 paradigms such as global/local tasks (Tsvetanov et al, 2013; Mevorach et al, 2016), reading with 10 distractor tasks (Darowski et al, 2008), response inhibition (Anguera and Gazzaley, 2012), as well as 11 inhibition in the context of WM tasks (Gazzaley et al, 2005). These findings and others fit with the 12 notion that aging is associated with a general impairment in a central inhibition mechanism (the 13 inhibitory deficit theory; Hasher and Zacks, 1988), which manifests in various inhibition related 14 scenarios. The focus on inhibitory processes in old age is especially relevant because there is 15 evidence that distractor inhibition is crucial in mediating cognitive control in general (Darowski et al., 16 2008). It should be noted that although it is often assumed that inhibition deficits result in impaired 17 cognition, there is some evidence to suggest that it may improve cognition under some 18 circumstances (Amer, Campbell, and Hasher, 2016).

19 In contrast, the notion that an all-encompassing inhibition impairment is associated with age 20 has been challenged by studies showing impairments only on subsets of inhibition tasks (Rey-21 Mermet et al, 2018a; 2018b). For instance, Kramer et al (1994) found age related inhibition deficits 22 in a stop-signal task, but not in a response competition or spatial pre-cueing task. Furthermore, even 23 when inhibition impairments occur across tasks, evidence suggests they may be independent. For 24 instance, Anguera and Gazzaley (2012; Sebastian et al, 2013) assessed motor inhibition in a stop 25 signal task and sensory filtering within the context of a delayed recognition task in young and old 26 participants. Critically, they showed that motor and sensory inhibition were independently impaired

as a function of aging. More recent studies have built upon this conclusion, further highlighting
distinct age effects on different cognitive inhibitory functions and their potential neural correlates
(Vadaga et al, 2015; Bloemendaal et al, 2016).

30 The dual mechanisms theory of proactive and reactive cognitive control (Braver, 2012) 31 suggests a potential explanation for the failure to identify a general inhibition impairment in old age. 32 Rather than a single inhibition mechanism, the DMC differentiates between two modes of control: 33 Proactive, an attentional biasing mechanism which mediates behavioral responses to a given 34 stimulus in advance; and reactive, which is a "late correction" mechanism that allows one to alter 35 behavioral plans "in the moment" when suddenly presented with new and relevant information. 36 Consequently, it is possible that only one of these inhibition mechanisms is affected by age, or that 37 they are affected to different degrees. Indeed, previous studies have identified a proactive pattern 38 of performance in young participants and a reactive one in old participants (Braver et al., 2005; 39 Paxton et al, 2008). Interestingly, even when performance across age groups was equivalent, brain 40 activity that was consistent with impairments to proactive control (Vadaga et al, 2015) and increased 41 reliance on reactive control (Paxton et al., 2008) was documented in the older cohorts. Similar brain 42 dynamics have also been recorded in the context of a task switching paradigm as older participants 43 (relative to younger participants) showed reduced sustained activation, a hallmark of proactive 44 control (Braver, 2012), and increased transient activation, a hallmark of reactive control (Braver, 45 2012), during switch trials in the anterior prefrontal cortex (Jimura and Braver 2010). These findings 46 are consistent with the idea that older participants show more reactive control related- and less 47 proactive control related activity than younger participants.

The above evidence points to a selective impairment in inhibition in old age – proactive processes appear to decline with age, while reactive processes may be intact. However, there are two major issues with this interpretation. First, most studies of the DMC and aging have highlighted the activation of reactive processes in older participants and proactive process in young participants 52 in the same task, but did not directly compare reactive processes across age groups. Second, in many 53 of these studies (typically using the AxCPT paradigm; Braver et al, 2005; Paxton et al, 2008; Braver et 54 al, 2009) utilising either proactive or reactive control yield different performance benefits. As such, it is possible that the increased use of reactive control in old age represents an unconscious strategic 55 56 bias rather than specific impairment in proactive control and intact reactive control. For example, 57 one of the early examples for inhibition deficits was reported by Hasher et al (1991) in a study measuring inhibitory function in young and older participants using a negative priming task 58 59 (assessing the persistence of inhibition of a distractor by switching its role to a target on subsequent 60 trials). Hasher et al (1991) found that young participants showed persistence of inhibition from one 61 trial to the next, but older participants showed no effects, suggesting impaired inhibitory function. 62 They argued that this reflected impairment in a central inhibition mechanism. However, an 63 alternative interpretation is that the younger participants were engaging proactive control and that 64 the persistence of inhibition was an artifact of their anticipating the state of the target and distractor 65 items, whereas older participants engaged reactive control and therefore did not show negative 66 priming because they didn't anticipate their state. Crucially, this arguably conferred a strategic 67 benefit to the older participants in this context since they were not biased away from the target on 68 switch trials.

69 A few notable studies have addressed the issue of comparing reactive control between age 70 groups. One set of studies have focused on "proportion congruence" manipulations in conflict 71 resolution tasks, such as Stroop tasks where congruent (no-conflict) and incongruent (conflict) 72 displays are contrasted (Bugg and Crump, 2012; also see Bugg, 2015 for examples using a flanker 73 task). Proportion congruence studies manipulate the ratio of congruent and incongruent trials within 74 a block (list-wise) or for a specific target type (item-specific; e.g., dogs within a list of animals) and 75 measure how such changes in frequency modulate performance differences between the congruent 76 and incongruent conditions. The typical finding in such studies with young adults is that the 77 congruency effects (difference in performance between congruent and incongruent displays) are

78 larger in the condition with more congruent trials, than in the condition with more incongruent trials 79 (Bugg and Chahani, 2011; Bugg, Jacobi, and Chahani, 2011; Bugg, Jacoby, and Toth, 2008). 80 Importantly, the two versions of this paradigm (list-wide and item-specific) arguably tap proactive 81 and reactive control mechanisms, separately. List-wide manipulations enable participants to adopt a 82 more stringent control in anticipation of trials throughout a block (proactive), while item-specific 83 versions do not. In the latter case, the proportion congruence manipulation is applied to different 84 items that are randomly intermixed and therefore, whether more or less stringent control is needed 85 cannot be predicted before and item is presented (reactive). In fact, Gonthier et al (2016) showed 86 distinct doubly-dissociated behavioral signatures for item-specific (reactive) vs list-wise (proactive) 87 proportion congruence manipulations, supporting the notion that item-specific proportion 88 congruence manipulations tap into reactive control mechanisms. Using such paradigms, Bugg et al 89 (2014a; 2014b) compared performance of young and old adults. Specifically, Bugg et al., (2014b) 90 found that both young and older participants exhibited the standard item-specific proportion 91 congruence effect, where the mostly congruent items had larger congruency effects than the mostly 92 incongruent items. Since reactive control was argued to be necessary to produce this effect, Bugg et 93 al (2014b) concluded that reactive control was spared in older participants.

94 Much like the "item-specific" proportion congruency effect, "sequential" congruency effects 95 have also been used as a measure of reactive control. "Sequential" congruency effects refer to the 96 phenomenon that congruency effects for trials immediately following an incongruent trial are 97 reduced compared to congruency effects for trials immediately following a congruent trial (In other 98 words, analyzing trials n+1 and grouping them based on trial n). Typically, both young and older 99 participants show these sequential congruency effects of equal magnitude, but older participants 100 show overall larger congruency effects and longer overall response times (even after accounting for 101 speed of processing deficits; Puccioni & Vallesi, 2012; West & Moore, 2005; but see Aschenbrenner 102 and Balota, 2016 for a study where older participants showed a larger sequential congruency effect). 103 In other words, although older participants seem to take longer to complete the task, these studies

have suggested that reactive control is effectively intact in older cohorts. In contrast, Xiang et al
(2016) found no "sequential" congruency effects for older participants, with respect to both
response time and accuracy, arguing that reactive control might be impaired in aging. However, they
did not account for generalized slowing effects or speed-accuracy trade-offs, which makes their
results difficult to interpret.

109 The purpose of this study is to further assess reactive distractor suppression in aging 110 populations. It should be noted here that reactive and proactive cognitive control are essentially 111 umbrella terms that can refer to two different mechanisms of engaging a wide range of similar 112 cognitive abilities, including distractor suppression, target selection, conflict resolution, and so on. Despite this, most studies tend to use the general term of proactive or reactive control, rather than 113 114 specify the cognitive mechanisms being investigated. This is relevant because it may be that 115 proactive and reactive control of some cognitive abilities may be intact, while others may be 116 impaired. Although the DMC generally argues for a general impairment within proactive control 117 mechanisms, it may be valuable to adopt a more precise perspective. Therefore, we refer to the process of reactive distractor suppression throughout our study, rather than reactive control. 118

119 In order to stringently assess age-related differences in reactive distractor suppression, 120 performance should be compared in a task that yields clear benefits when reactive distractor 121 suppression is engaged and eliminates or significantly reduces the role of proactive distractor 122 suppression in young participants too. Here, we describe such a task which specifically taps reactive 123 distractor suppression mechanisms. DiQuattro and Geng (2011) investigated the brain mechanisms 124 that are involved in processing contextually relevant, but not task relevant stimuli. While in an fMRI 125 scanner, they had participants (Mean age = 23.8; Age Range: 18 - 39 yrs) perform a visual search 126 task for a low contrast target in the presence of either a high or low contrast non-target (50% predictability), each of which would appear in one of two pre-defined locations; participants could 127 128 not predict the location of the distractor on a given trial. Despite being task irrelevant, the salient

non-target was contextually relevant as the presence of the high contrast non-target informed the
participant that the target was in the other location, effectively triggering a reactive distractor
suppression. They found that participants were both faster and more accurate on trials with a salient
distractor, compared to a similar (to the target) distractor.

133 Geng and DiQuattro (2010; experiment 1) used a variant of this paradigm with eye tracking 134 showing that, when the distractor was unpredictable, participants made a saccade to distractor first 135 on ~68% of trials, necessitating reactive rapid rejection instead of proactively inhibiting the 136 distractor (saccading towards the target first; for similar trials, it was ~50% rapid rejection and ~50% 137 inhibition). They argued that rapid rejection was only needed when inhibition failed. In other words, 138 the engagement of reactive control in this task pre-supposes a failure of proactive control (Geng, 139 2014). Crucially though, even when a saccade was first made towards the distractor, there was still a 140 performance benefit on salient trials. Even when proactive inhibition failed, young participants were 141 able to reactively rapidly reject the distractor in a beneficial manner. Therefore, the presence of a 142 performance benefit (i.e. better performance on salient trials) in this task must be attributed to both 143 a failure of proactive (inhibition) and the engagement of reactive (rapid rejection) processes.

144 The fMRI analysis in DiQuattro and Geng (2011) provided converging evidence for the notion 145 that this task engages reactive distractor suppression. It revealed that the left TPJ and left inferior 146 frontal gyrus (IFG) were significantly more active when there was a high salience non-target 147 compared to a low salience non-target. Dynamic causal modelling revealed a network in which left 148 TPJ projects to left IFG that, in turn projects to the frontal eye fields (FEF). The authors interpreted 149 this to mean that the ventral attention network (TPJ and IFG in this study) that is typically associated 150 with bottom up attention (Corbetta and Shulman, 2002), updates control signals to the dorsal 151 attentional network (FEF in this study). They refer to this network as an attentional circuit breaker that can reorient attention when top-down (i.e. proactive) attentional processes don't work or lead 152 to counterproductive outcomes. Importantly, they further suggest that the TPJ and IFG are 153 154 effectively generating a "reactive" control signal as a consequence of the stimulus presentation

(Braver et al, 2009; Braver, 2012). Additional converging evidence comes from studies of young
adults with high expression of psychosis proneness, who have been shown to favor reactive
distractor suppression relative to proactive distractor suppression (Abu-Akel et al, 2016a; 2016b;
2016c). In the t-task described above, performance benefits from the presence of the salient
distractor scale with the expression of psychosis proneness (Abu-Akel et al., 2018), highlighting that
this paradigm is sensitive enough to detect variations in the magnitude of engagement of reactive
distractor suppression.

162 Moreover, another advantage of this task is that it avoids engaging general non-perceptual 163 inhibition processes. The stimuli within the distractor is never a valid response option (a sideways t 164 shape instead of upright or inverted), so participants won't be primed to make a specific response if 165 they do process the distractor, which would then need to be inhibited. In addition (unlike stop-signal 166 tasks), the "correct" response never changes within the course of a trial so they never have to switch 167 responses. Indeed, older participants appear to be impaired in a stop-signal task (Kramer et al, 1994), which tests participants' ability to cancel a motor response while it is already being executed 168 169 (and therefore presumably measures reactive inhibition). However, the requirement to cancel an 170 already executable motor response may relate to other processes (primarily motor 171 cancelation/response inhibition) which may be independent or more complex than reactive 172 inhibition per se. (Swick, Ashley, and Turken, 2011; Kolodny, Mevorach, and Shalev, 2017). In fact, 173 Anguerra and Gazzaley (2012) have highlighted the independence of impairments in response 174 inhibition and perceptual inhibition (but not exclusively reactive inhibition) in old age. In our study, 175 by design, only differences in perceptual inhibition are likely to change behavior.

This paradigm is distinct from the proportion congruence studies in two important ways. First, proportion congruence effects rely on the implicit learning of the associations between different items and their likelihood of conflict (Blais et al, 2012). It is possible that proportion congruence effects may be less susceptible to age-related decline due to their reliance on implicit learning (Cohen-Shikora, Diede, & Bugg, 2018). However, in our task, there is an even probability of all trial types and making a correct response does not depend on any element of previous trials.
Thus, our task may tap into a more temporally bounded form of reactive control. Second, our
paradigm manipulated stimulus presentation duration to assess the timing of reactive control. As far
as we know, no previous studies have done this.

In the present study we compared old and young participants' performance on the T-search task to directly assess if reactive inhibition is impaired or intact in old age. If in older participants reactive distractor inhibition is indeed intact (irrespective of a possible proactive impairment), it is expected they will derive a benefit from the presence of the salient distractor compared to a similar distractor. Conversely, if older participants have impaired reactive distractor suppression, it is expected that they show minimal benefit from the presence of a salient distractor.

192Experiment 1a193Methods194IRB Approval

All experiments in this study were approved by the Science, Technology, Engineering and
 Mathematics Ethical Review Committee at the University of Birmingham.

197 Participants

198 25 young participants and 26 older participants participated in two successive behavioral 199 experiments. Three subjects (2 older, 1 younger) were excluded from the analysis due to poor 200 performance (Overall accuracy across conditions < 60%; A cut-off of 60% was chosen to balance 201 between exclude participants who were guessing or did not understand the task, and including 202 participants who simply had low accuracy), resulting in 24 young participants (Mean Age: 18.8, SEM 203 of Age: .19, Age Range: 18 -21; 23 Females) and 24 older participants' (Mean Age: 70.1, SEM of Age: 204 1.63, Age Range: 60 - 82; 13 Females) data being analysed. All participants had normal or corrected 205 to normal vision. The order of the tasks was counterbalanced to account for possible fatigue and

206 order effects. The other experiment is reported elsewhere and has no relevance to the current 207 investigation. Young participants were recruited from the undergraduate population in the school of 208 psychology at the University of Birmingham, UK. They were compensated for their participation with 209 course credits. The older participants were recruited from a volunteer pool maintained by the School 210 of Psychology at the University of Birmingham. They were compensated for 1.5 hours of their time 211 with a one-time payment of £7. All participants had to sign an informed consent form prior to the 212 study. Participants' were healthy with no history of head injury, mental health issues or neurological 213 disorders. The old participants were screened for decline in cognitive functions using the Montreal 214 Cognitive Assessment (MoCA). All of the older participants scored within the normal range (Mean 215 Score: 27.5, SEM of Score: .23).

216 Power Analysis

217 An a priori statistical power analysis was performed for sample size estimation, based on data from 218 DiQuattro and Geng (2011; N = 21). They found that the presence of a salient distractor significant 219 improved both response time and accuracy for healthy young participants, with an effect size of 220 cohen's d = 2.9 and d = 1.7, respectively. These are considered to be extremely large using Cohen's 221 (1988) criteria. With an alpha = .05 and power = 0.95, the projected sample size needed to detect an 222 effect size of 1.7 (GPower 3.1.9.2; Faul et al, 2007; 2009) is approximately N = 6 for a repeated 223 measures ANOVA with a within-between subjects interaction. However, one possibility is that the 224 effect size will be much smaller for the older participants. Therefore, as an additional check the 225 projected sample size needed to detect an effect size of .3 (with all the same other parameters) is 226 approximately N = 40 (20 per group). Thus, our sample size of 51 should be adequate to assess the 227 main objectives of this study, unless the effect size for older participants is lower than .3. This power 228 analysis also applies to experiment 2 which has a sample size of 39.

229 Stimuli and Procedure

230 Participants were presented with five blocks of 46 trials each. Color was defined using RGB 231 color coordinates. The background color of the display was grey [100 100 100]. On all trials, target 232 and non-target stimuli were displayed. Each stimulus was a square whose center was 6.5 degrees of 233 visual angle (horizontally 6.3 degrees; vertically 1 degree; all measures of visual angle were 234 calculated assuming a viewing distance of 50 cm) diagonally left or right and below the center of the 235 screen. Each square subtended 1.8 degrees of visual angle. The target square was dark grey [120 120 236 120]. In the target square, a vertical line with a width of .224 degrees of visual angle bisected the 237 square. A second horizontal line also appeared to create a 'T'-like shape (Figure 1). These lines were 238 a dark grey [80 80 80]. On half of trials, the horizontal line was .281 degrees of visual angle above 239 the center of the square, creating an 'Upright' T (Figure 1) and on the other half, the horizontal line 240 was .281 degrees of visual angle below the center of the square creating an 'Inverted' T. The color of 241 the non-target square depended on the trial type. On 'Similar' trials, the color was the same as the target square. On 'Salient' trials, the non-target square was white [255 255 255]. In the non-target 242 243 square, an horizontal line with a height of .224 degrees of visual angle bisected the square. A second 244 vertical line also appeared to create a sideways 'T'-like shape (Figure 1). On 'Similar' trials, the line 245 color was the same as inside the target square. On 'Salient' trials, the line color was black [0 0 0]. On 246 50% of trials, the vertical line was .281 degrees of visual angle right of the center of the square, 247 creating a clockwise rotated "T". On 50% of the trials, the vertical line was .281 degrees of visual 248 angle left of the center of the square creating a counter-clockwise rotated "T".

In each block there were 50% "Salient" trials and 50% "Similar" trials, randomly intermixed. On any given trial there was a 50% chance that the target would appear in the left position and 50% chance that it would appear in the right position. Participants had to identify if there was an upright or inverted "T" stimulus on each trial by pressing the "H" or "B" keys, respectively. These buttons were chosen because the "H" key is positioned above the "B" key on the keyboard, mimicking the spatial orientation of the target stimuli, where the upright "T" stimulus has a horizontal line above the center of the stimulus square and the inverted "T" stimulus has a horizontal line below thecenter of the stimulus square.

257 Every trial began with a white [255 255 255] fixation cross presented at the center of the 258 screen, which persisted throughout the trial (including during 'blank' screens). Each trial began with 259 blank screen. The "fixation" time was randomly selected based on a uniform distribution of times 260 between 1500 - 2000 msec (Figure 1). Next, the appropriate stimulus (depending on the trial) was 261 displayed for 200 msec. Participants could respond starting when the stimulus was presented. After 262 the stimulus was removed, the participant was presented with blank screen until they made a 263 response. Once a response was made, the next trial would begin. Participants were given the 264 chance to take short breaks in between blocks (< 5 min). Each session began with 20 practice trials. 265 During the practice, participants received visual feedback such that if they made an identification 266 error, the fixation cross changed to red for 250 msec before turning back to white for the rest of the 267 fixation time.



Figure 1. Diagram of the reactive distractor inhibition t-task. Participants were presented with either a salient or similar stimulus on any give trial. In the

salient example the correct response would be to press the H-key to indicate an upright target. In the similar example the correct response would be to
 press the B-key to indicate an inverted target.

### **Results & Discussion**

273 Response time in msec (RT) and accuracy rate (i.e. proportion of correct responses) were 274 measured as dependent variables. All values are presented as mean +/- standard error of the mean. 275 The data was cleaned to account for outliers. For each participant, response time data that was 276 greater than and less than 2 standard deviations from the participants' individual mean was 277 excluded from all analyses. The individual mean response time was calculated separately for each 278 salience condition (salient trials and similar trials). This resulted in the loss of an average of 4.37% 279 (SEM = .22%) of the response time data per older participant and 3.71% (SEM = .23%) per young 280 participant. An independent samples t-test revealed that these values were significantly different 281 (t(46) = -2.039, p = .047, d = .58). This is attributable to the notion that older participants tend to 282 exhibit greater variability in cognitive performance (Hultsch and MacDonald, Chapter 4 in Dixon et 283 al, 2004; Morse, 1993) than younger participants. As such, their response time distributions would 284 be wider and they would have more trials that would fall outside of 2 standard deviations from the 285 mean.

286 For the accuracy data, a rationalized arcsine transformation (Equation 1) was applied to each participants' overall accuracy in each salience condition. This was done to account for possible 287 288 violations of normality that can arise in binomially distributed data. The rationalized arcsine 289 transformation was designed to normalize the data in accordance with the arcsine transformation 290 while maintaining a more intuitive scale for interpretation of results (Studebaker, 1985). Since the 291 transformation don't reflect true probabilities, they will not be reported with a % symbol, but for 292 interpretation purposes the values do approximate the raw data, albeit skewed towards larger 293 values than the raw data.

$$RArcSine = \left(46.47324337 * \left(2 * arcsin(\sqrt{Accuracy})\right)\right) - 23$$

Equation 1. This rationalized arcsine transformation calculates the standard arcsine transformation,
 then adjusts the value such that a proportion of .5 will have a transform of 50, rather than 1.5708.

296 Transformed accuracy values (Figure 2a) were analysed using a repeated measures ANOVA 297 with Saliency (salient distractor vs similar distractor) as within subject factor and participants age 298 group (Young vs Older) as a between subject factor. The main effects of salience (F(1,46) = 2.368, p =.131,  $\eta_p^2$  = .049) and age group (F(1,46) = .446, p = .508,  $\eta_p^2$  = .010), were not significant, but there 299 was a significant interaction between salience and age group (F(1,46) = 5.132, p = .028,  $\eta_p^2$  = .100). 300 301 Planned comparisons with paired samples t-tests comparing performance in the salient and similar conditions within each age group revealed that for older participants there was no significant 302 303 difference in accuracy for salient (90.99 +/- 2.88) and similar trials (92.00 +/- 1.78, t(23) = -.505, p = -.505, p304 .618, d = .10). In contrast, young participants were more accurate on salient trials (96.28  $\pm$  2.52) 305 than on similar trials (90.98 +/- 2.56; t(23) = 2.739, p = .012; d = .56).

Moreover, an independent samples t-test revealed there was no significant difference in RArcSine transformed accuracy during the similar condition between the young (90.98+/- 2.56) and older (92.00 +/- 1.78), participants (t(46) = -.329, p = .744, d = .09), which highlights that the group differences we observed were not due to greater difficulty in identifying the stimuli by the older group. It also confirms that the benefit effect for the young participants was specifically due to enhanced performance in the presence of a salient distractor.

312 For the Response Time data, the raw response times were transformed into z-scores to 313 account for generalized speed of processing deficits reported in aging populations that can be confounded with inhibition deficits (Craik and Salthouse, 2011; Salthouse, 1994; 1996; 2000; 314 315 Salthouse and Meinz, 1995). We applied a z-score transformation to each individual subject's 316 response time data by subtracting their overall mean response time from their condition mean 317 (Salient or Similar), then dividing by the standard deviation of their condition means. To be clear, this 318 was done separately for their salient and similar condition data. This procedure has been used 319 previously to account for speed of processing deficits (Tsvetanov et al, 2013) in aging populations 320 and is described in more detail in Faust et al (1999).

321 The z-RT data were analysed using a repeated measures ANOVA with Saliency (salient 322 distractor vs similar distractor) as within subject factor and participants age group (Young vs Older) as a between subject factor (Figure 2b). The analysis revealed a main effect of saliency (F(1,46) = 323 15.173, p < .001,  $\eta_p^2$  = .248), where participants were quicker to respond to salient trials (z-RT = -.103 324 +/- .025) than to similar trials (z-RT = .071 + /- .020; Smaller values reflect faster response times). This 325 326 effect is a typical result for this paradigm and supports the notion that participants are engaged in 327 reactive cognitive control and utilise the salient distractor as an anti-cue (DiQuattro and Geng, 2011; Geng and DiQuattro, 2010). There was not a main effect of age group (F(1,46) = 2.629, p = .112,  $\eta_p^2$  = 328 .054, but there was an interaction between salience and age group (F(1,46) = 7.881, p = .007,  $\eta_p^2$  = 329 330 .146).

331 We conducted planned comparisons using a paired samples t-test to compare z-RTs for the 332 two salience conditions in each group. For the older participants, there was no significant difference 333 in z-RT across salience conditions (t(23) = -.782, p = .442, d = .16; Salient: z-RT = -.033 +/- .034; 334 Similar: z-RT = .015 + /-.029). However, younger participants responded significantly quicker (t(23) = -4.665, p < .001, d = .95) in the salient condition (z-RT = -.17 +/- .038) compared with the similar 335 336 condition (z-RT = .127 +/- .027). These results suggest that the main effect of saliency was primarily 337 driven by the younger participants and that the older participants showed no benefit in performance 338 when the salient distractor appeared.





- 340 Figure 2. (a) Graph reflecting mean of RArcSine transformed accuracy for salient and similar
- 341 conditions across age groups. (b) Graph reflecting z-scored response time data (msec) for salient and
- 342 similar conditions across age groups. PES stands for partial eta squared and d stands for cohen's d.

#### **Experiment 1b**

The identical performance we documented for older participants in the similar and salient 345 346 trials could potentially stem from reduced visual contrast sensitivity in this age group (Roberts and 347 Allen, 2016; Pardhan, 2004; Owsley et al, 1983; Sekuler et al, 1980). One possible complication in 348 any aging study of higher order cognitive abilities - such as reactive cognitive control - is that low 349 quality information due to age-related impairments to lower level perceptual abilities may cascade 350 through the information processing stream affecting performance (the information degradation 351 hypothesis; Monge and Madden, 2016; but see Houston et al, 2016 for a counter-perspective). 352 Notably, Porto et al (2016) found that controlling for visual acuity scaled and/or eliminated an age-353 related reduction in posterior P3b amplitude during a visual oddball task. The posterior P3b 354 amplitude is generally presumed to be indicative of higher-level decision making and executive 355 functions, although it should be noted that the specific role of the posterior P3b amplitude is still 356 under a great deal of debate (See Polich, 2007 for a more in-depth discussion). 357 Indeed, if our older participants were not sensitive to the contrast differences between 358 salient and similar non-targets then their performance in the two conditions would be equivalent. To 359 exclude this possibility, we invited a subset of the older participants who took part in experiment 1a 360 back and assessed whether or not they are able to distinguish between the two contrast conditions 361 (Salient vs Similar). In experiment 1b we presented the exact same stimuli to a set of older 362 participants, but instead of responding to the t-shapes, they had to indicate if the box colors of the 363 two elements of the display (target and non-target) were the same or different. If the participants 364 can successfully distinguish between the salient and non-salient stimuli, then we can be confident 365 that age-related impairments to visual contrast sensitivity are not influencing our results.

366

#### Methods

367 Participants
368 5 older participants (Mean Age: 70.8, SEM of Age: 2.35, Age Range: 65 - 78; 1 Female)
369 participated in the experiment. The older participants were recruited from the initial cohort who

370 participated in Experiment 1a. They were compensated for 1.5 hours of their time with a one-time 371 payment of £7. All participants had to sign an informed consent form prior to the study. Participants' 372 were healthy with no history of head injury, mental health issues or neurological disorders. The 373 older participants were screened for decline in cognitive functions using the Montreal Cognitive 374 Assessment (MoCA). All of the older participants scored within the normal range (>= 26 out of 30). 375 Stimuli and Procedure 376 The stimuli and procedure were exactly the same as in experiment 1a except that instead of 377 indicating if the target were upright or inverted, participants pressed the "h" key to indicate if the 378 two stimulus boxes were the same color (similar trials) and the "b" key if they were different (salient 379 trials).

380

# **Results and Discussion**

Overall raw accuracy across the 5 participants was very high (Mean = 98% +/- .68%). The data was transformed to a rationalized arcsine measure, consistent with experiment 1a. To assess the RArcSine transformed accuracy results, a one-way one-sample t-test was conducted to determine if a statistically significant difference existed between the average RArcSine and an RArcSine of 50 (equivalent to a proportion correct of .5). The t-test was significant, t(4) = 19.8, p < .001, d = 8.85, suggesting that older participants were able to successfully distinguish between the salient and non-salient stimulus boxes.

The response time was not z-transformed like in experiment 1a since there is only an older participants group and we do not need to account for group differences. However, the response time data was cleaned to account for outliers using the same procedure described in experiment 1a. This resulted in the loss of an average of 4.69% (SEM = .59%) of the response time data for salient trials and 3.82% (SEM = .89%) of the response time data for similar trials. A paired samples t-test revealed that these were not significantly different (t(4) = 1.12, p = .326, d = .5). The ability to distinguish successfully between these scenarios was evidenced by virtually identical response times for same (Mean = 592 +/- 37.4 msec) and different (593 +/-37.5 msec) trials (t(4) = -.029, p = .977, d
= .013), confirming that older participants were not simply increasing accuracy during different trials
by taking more time.

398 Overall, these results exclude the possibility that the lack of benefit for older participants, 399 observed in Experiment 1a was attributed to impairments to visual contrast sensitivity yielding the 400 salient and similar trials identical for our older participants. It should be noted that although there is 401 a small sample size in this study, the consistently high accuracy (≥ 96%) across all participants and 402 their age range allows us to be relatively confident that these results are a reasonable estimation of 403 our participants' visual contrast sensitivity.

404

# Experiment 2

405 Although we show that general age-related processing speed deficits can't fully account for 406 our data (as overall RTs were not associated with the benefit measure for older participants), it is 407 still possible that certain aspects of processing speed affect older participants ability to utilise 408 reactive distractor suppression this task. For instance, if older adults take longer to accumulate 409 evidence that will yield a disengagement decision from a non-target element, it might be the case 410 that even though they are sensitive to the contrast differences it takes them similar amount of time 411 to reach a decision to disengage. In experiment 1b, it took the older participants 592 msec to simply 412 discriminate between the salient and similar conditions, whereas in experiment 1a they were only given 200 msec to use the salient distractor. Thus, it is possible that in old age effective reactive 413 414 distractor suppression is possible given enough time.

This idea would fit with the argument that older participants favor reactive control in the first place because it takes longer to accumulate neural resources in old age and reactive control typically requires fewer resources over a shorter period of time compared to proactive control (Grady, 2012). Consequently, slow resource accumulation could theoretically impact reactive control as well. In particular, it would result in a delayed initiation of reactive inhibition. If the initiation of reactive distractor suppression is delayed in older cohorts, then we would expect them to be able to
effectively utilise reactive distractor suppression as long as they are given enough time to engage
reactive suppression mechanisms.

To address the possibility that there is an age-related delay in the initiation of reactive inhibition processes, Experiment 2 was conducted in which the stimulus presentation time was extended from 200 msec to 600 msec. If the initiation of reactive inhibition takes longer in old age, then a longer presentation time should allow older participants to take advantage of the salient distractor, and we would expect to see a performance benefit. Alternatively, if reactive inhibition is simply less effective in old age we expect to replicate the results from experiment 1a and find no performance benefit for the old participants.

430

431 Participants

# Methods

432 20 young participants (Mean Age: 19.2, SEM of Age: .20, Age Range: 18 -21; 18 Females) and 433 19 older participants (Mean Age: 69.47, SEM of Age: 1.12, Age Range: 62 - 78; 11 Females) 434 participated in the experiment. Like experiment 1a, we planned to exclude any participants with less 435 than 60% overall accuracy, but no participants met this criterion and none were excluded from 436 analysis. Young participants were recruited from the undergraduate population in the school of 437 psychology at the University of Birmingham, UK. They were compensated for their participation with 438 course credits. The older participants were recruited from a volunteer pool maintained by the School 439 of Psychology at the University of Birmingham. They were compensated for 1.5 hours of their time 440 with a one-time payment of £7. All participants had to sign an informed consent form prior to the 441 study. Participants' were healthy with no history of head injury, mental health issues or neurological disorders. The older participants were screened for decline in cognitive functions using the Montreal 442 443 Cognitive Assessment (MoCA). All of the older participants scored within the normal range (>= 26 444 out of 30).

445 Stimuli and Procedure

- 446 The stimuli and procedure were exactly the same as in experiment 1a, except that the 447 stimulus was presented for 600 msec instead of 200 msec.
- 448 **Results and Discussion** 449 Response time in msec (RT) and accuracy rate (i.e. proportion of correct responses) were 450 measured as dependent variables. The response time data was cleaned to account for outliers. For 451 each participant, response time data that was greater than and less than 2 standard deviations from 452 the mean was excluded from all analyses. The mean response time was calculated separately for 453 each salience condition (salient trials and similar trials). This resulted in the loss of an average of 454 3.64% (SEM = .22%) of the response time data, per older participant and 3.61% (SEM = .24%) per 455 young participant. An independent samples t-test revealed that these values were not significantly different (t(37) = -.093, p = .927). All values are presented as mean +/- standard error of the mean. 456
- Accuracy data was rationalized arcsine (RArcSine) transformed and response time data was z-transformed using the same procedures as in experiment 1a. RArcSine transformed accuracy and ztransformed response time were analysed using a repeated measures ANOVA with Saliency (salient distractor vs similar distractor) as within subject factor and participants age group (Young vs Older) as a between subject factor. Data is reported as mean +/- standard error of the mean.
- For RArcSine transformed accuracy (Figure 3a), the main effect of salience was significant (F(1,37) = 39.75, p < .001,  $\eta_p^2$  = 0.518), driven by more accurate responses during salient trials (109.99 +/- 1.68) relative to similar trials (102.07 +/- 1.56). However, the main effect of age group (F(1,37) = .05, p = .824,  $\eta_p^2$  = 0.001) and the interaction (F(1,37) = 1.3, p = .261,  $\eta_p^2$  = 0..034) were not significant. This data suggests that, in terms of RArcSine transformed accuracy, the young and older participants were equally effective at using the salient distractor as an anti-cue when the stimulus presentation time was extended.

469 For response time (Figure 3b), the main effect of salience was significant (F(1,37) = 84.654, p < .001,  $\eta_p^2$  = 0.696) driven by relatively faster responses during salient trials (zRT: -.451 +/- .053) 470 compared to similar trials (zRT: .271 +/- .026), suggesting that the salient distractor provided a 471 benefit to performance. However, the main effect of age group (F(1,37) = 1.211, p = .278,  $\eta_p^2$  = 472 0.032), and the interaction were not significant (F(1,37) = 3.696, p = .062,  $\eta_p^2$  = 0.091). Given the 473 marginal p-value for the interaction, we conducted an exploratory analysis to assess whether there 474 is potential evidence the interaction is driven by a lack of performance benefit (i.e. the difference 475 476 between salient and similar performance) for the older adults. Crucially, simple effects revealed that 477 the difference in performance between the salient and similar conditions was significant for both the young (p < .001,  $\eta_p^2$  = 0.632; Salient: -.542; Similar: .330) and older (p < .001,  $\eta_p^2$  = 0.411; Salient: -478 479 .359; Similar: .212) cohorts. Further simple effects revealed that there was no difference in performance between older and younger participants for the salient trials (p = .093,  $\eta_p^2$  = 0.075; 480 Older: -.359; Younger: -.542), but there was a difference in performance for the similar trials (p = 481 .031,  $\eta_p^2$  = 0.120; Older: .212; Younger: .330; Older participants were less slowed during similar 482 483 trials). This verifies that despite the trending interaction, both age groups showed a clear 484 performance benefit in the presence of a salient distractor, even if it is attenuated in the older 485 cohort.

These data suggest that given enough time older participants are able to use the salient distractor as an anti-cue as effectively as the younger participants. This suggests that older participants may take longer to initiate reactive inhibition, possibly due to the slower accumulation of neural resources, but that given enough time reactive inhibition can be effectively implemented.



491 Figure 3. (a) Graph reflecting mean RArcSine transformed accuracy for salient and similar conditions

492 across age groups. (b) Graph reflecting z-scored response time data (msec) for salient and similar

493 conditions across age groups.

# **General Discussion**

495 The purpose of this study was to assess the effect of aging on reactive suppression in older 496 populations by comparing performance in a task that relies on reactive suppression in young 497 participants too. In experiment 1a, with a 200 msec display, we found that young participants were 498 able to effectively use a salient distractor as an anti-cue to benefit performance in terms of both 499 accuracy and response time, demonstrating effective reactive distractor suppression. Older 500 participants on the other hand showed no change in performance when the salient distractor was 501 present in the display. Importantly, the lack of performance benefit for old participants could not be 502 attributed to reduced contrast sensitivity as a subset of the older participants were shown to be able 503 to distinguish between low and high contrast items in experiment 1b. However, experiment 2 504 showed that older participants could engage reactive inhibition if given enough time. When the 505 stimulus presentation time was extended to 600 msec, older participants showed better 506 performance during salient trials than similar trials (in both accuracy and response time) at an 507 equivalent magnitude to the young participants. These data suggest that older participants have a 508 delayed initiation of reactive inhibition processes that scales with age, but that given enough time 509 effective reactive inhibition is possible. Prior literature typically shows a) impairments to proactive control in aging and b) a shift from proactive to reactive mechanisms in aging. Despite the dearth of 510 511 studies directly investigating reactive control deficits, this has led to an implicit (and sometimes 512 explicit) assumption throughout prior literature that older participants shift to reactive control 513 because proactive control is impaired and reactive control is intact. Our study challenges this notion 514 as we did find an age-related deficit in a measure reflecting reactive distractor inhibition. That being 515 said, it is possible that reactive control is less impaired than proactive control, which induces a shift, 516 but that is a different issue than shifting due to no deficit.

517 One concern for this type of research in general is that inhibition deficits may in fact be 518 attributed to a generalized deficit in processing speed (Salthouse and Meintz, 1995; Salthouse, 2000; 519 Verhaeghen and De Meersman; 1998). Since most studies that identify inhibition deficits measure response time, it could appear as if there were impaired response times in a specific inhibition task for older participants, when in fact they are simply overall slower. However, even after accounting for this possibility, inhibition deficits still persist in many inhibition tasks (Verhaeghen and Cerella, 2002). In our study, we z-transformed the response time data to account for this possibility and still found no performance benefit in the 200 msec condition (Experiment 1a). As such, we would argue that the age-related lack of a benefit we report cannot be attributed to general speed of processing deficits and is more likely associated with a delay in reactive distractor suppression specifically.

527 A second concern with respect to this specific study is that the deficit observed may in fact 528 be completely or partially due to age-related impairments in attentional orienting rather than reactive inhibition. The nature of orienting attention is complex and there are many variables to 529 530 consider, particularly with respect to aging (see Erel and Levy, 2016 for a comprehensive review), 531 but the most relevant aspects in the context of our study are covert and overt orienting, and 532 exogenous and endogenous orienting. Participants in the current study were instructed to keep their 533 eyes focused on the fixation point throughout the trials ostensibly to encourage covert attention (no 534 eye movements) which is also likely given the short presentation times in experiment 1a, however 535 eye tracking was not employed so the use of overt attention cannot be ruled out. Regardless, 536 research shows that older participants typically do not have impairments in covert attentional 537 orienting (Jennings et al, 2007), and that while deficits in overt attention tasks have been reported 538 (Kingstone et al, 2002), it has been argued that they can be attributed to deficits in motor control 539 over eye movements (Chen and Machado, 2016; Dowiasch et al, 2015; Warren et al, 2013; Crawford 540 et al, 2013; Klein et al, 2000, Ross et al, 1999) rather than attentional control (Erel and Levy, 2016).

541 With regard to exogenous and endogenous orienting during trials with a salient distractor, it 542 is arguable that this study engages both. Geng and DiQuattro (2010) showed that salient distractors 543 could facilitate performance (using a similar t-task paradigm to our study) using a combination of 544 two attentional strategies: inhibition, where saccades toward the salient distractor are actively 545 inhibited, and rapid rejection, where a saccade toward the salient distractor is quickly disengaged 546 and redirected towards the target. Inhibition took place on target-first trials, when the first saccade 547 went towards the target, and rapid rejection followed by inhibition took place on distractor-first 548 trials, when the first saccade went towards the salient distractor. Within the t-task, the process of 549 rapid rejection essentially consists of three phases: Orienting attention towards the salient 550 distractor, disengaging attention from the salient distractor, and reorienting attention towards the 551 target/inhibiting the salient distractor. The initial orienting of eye movements towards the salient 552 distractor is a classic example of overt exogenous orienting. However, disengaging, reorienting, and 553 inhibiting only begins because the participants recognize the distractor as such, making the target no 554 longer in an unpredictable location. This suggests that endogenous orienting is an integral part of 555 reactive distractor suppression processes which are likely important for performance in our task.

556 Previous studies have typically reported intact (Waszak et al, 2010; larocci et al, 2009; 557 Jennings et al, 2007; Folk and Hoyer, 1992; Craik and Byrd, 1982) or even enhanced (Langley et al, 558 2011a; 2011b; Mahoney et al, 2010) exogenous orienting in aging. In contrast, endogenous attention 559 is sometimes reported to be impaired (Olk and Kingstone, 2009; Bojko et al, 2004; Brodeur and Enns, 560 1997; Greenwood et al, 1993; see also Erel and Levy, 2016). Furthermore, impaired (i.e. slow) 561 attentional disengagement (Owsley, 2016; Greenwood and Parasuraman, 1994) has also been 562 documented in older populations. Consequently, if spatial endogenous disengagement and orienting 563 is impaired in old age it may well be the case that these impairments also manifest in impaired 564 reactive distractor suppression.

565 Nevertheless, disengagement and endogenous orienting are likely engaged in both salient 566 and similar trials in our task as both trials may involve the initial selection of the non-target item 567 (Geng & DiQuattro, 2010). Thus, an impairment in these processes should have affected 568 performance in both trial types. However, the older participants in Experiment 1a did not show such 569 a general impaired performance. In fact, accuracy was the same for both young (86%) and older 570 (86.2%) participants during the similar condition in experiment 1a. As such, the performance 571 patterns we report point to a difference in the efficiency of processes that are specifically utilised 572 when a salient distractor is present. One reason behind this could simply be a slower processing 573 capacity for salient items in old age (that is overcome when the input is presented for longer). This 574 seems unlikely as Experiment 1b demonstrated that the older participants showed virtually no 575 difference in response time when discriminating between similar (two low contrast stimuli) and 576 salient (one low and one high contrast stimuli) conditions. If salient stimuli required more processing 577 time in old age, we would have expected to see longer response times on trials where a salient item 578 was present in the display. A second, and seemingly more likely explanation, is that old participants 579 are exhibiting impairments in the reactive suppression of salient information (rather than having 580 difficultly processing salient information in the first place). In fact, impaired suppression of salient 581 information in old age has previously been reported by Tsvetanov et al (2013) in the context of a 582 proactive inhibition task. It is therefore possible that older adults exhibit impairments in salience suppression in general, regardless of whether reactive or proactive inhibition is called upon. 583

584 A final possibility is that 600 msec is too long to still be considered "reactive." According to 585 Irlbacher et al (2014), reactive control can be parsed into an early and late mechanism and that each 586 mechanism provides a unique method of identifying and resolving conflict. In the context of working 587 memory inhibition, both mechanisms resolve interference that occurs when a familiar stimulus is 588 identified but must be ignored. Familiarity-inhibition models favor speed over accuracy and are 589 considered to be a quick and early acting reactive control mechanism, engaging around 300-450 ms 590 after stimulus presentation (Du et al, 2008). In these models, the interference is resolved through 591 the inhibition of the familiar stimulus (Mecklinger et al, 2003). On the other hand, context retrieval 592 models favor accuracy over speed and are considered to be a slower and late acting reactive control 593 mechanism, engaging around 550 ms after stimulus presentation (Zhang et al, 2010). In these 594 models, the interference is resolved by selecting for the relevant target features more strongly (by 595 retrieving the appropriate contextual information; Badre and Wagner, 2005; 2007). Based on this, it

is reasonable to believe that reactive processes were used by participants with a presentation timeof 600 msec.

598 However, despite indications that our data reflect an impairment in reactive distractor 599 suppression, a major limitation was that we did not use eye-tracking and therefore cannot tease 600 apart the relative contribution of proactive and reactive distractor suppression during this task. In 601 other words, there is no direct evidence that young and old participants are completing the task in a 602 similar manner (i.e. specifically via reactive control mechanisms). That being said, Geng and 603 DiQuattro (2010) showed that young participants had a failure of proactive control mechanisms (i.e. 604 a saccade was made to the salient distractor) on ~68% of trials, leading necessarily to the 605 engagement of reactive control mechanisms (i.e. rapid rejection). Importantly, performance on 606 these "reactive" trials was still better than on control trials on which the first saccade went to a non-607 salient distractor, suggesting that the reactive rejection of the salient distractor was facilitated even 608 when it initially captured attention. Based on this we can infer that a similar proportion of "proactive 609 failure" trials likely occurred in our study for the young participants. Further, in conjunction with 610 well-characterized prior literature that has shown age-related deficit across proactive control mechanisms, we also infer that "proactive failures" are probably more common in our older cohort. 611 612 This would suggest that our older cohort likely relied more on reactive control mechanisms to 613 complete the task than the younger cohort. However, this is ultimately speculative and must be 614 confirmed in future research with other methods.

Overall, this study suggests that older participants exhibit an age-related delay in the initiation of reactive inhibition. The nature of this impairment is hypothesized to be specific to reactive salience suppression, since the underlying rapid rejection and inhibition processes appear intact when there is no salient distractor present (as evidenced by equivalent performance across age groups in experiment 1a during similar trials). Future research will have to investigate the underlying impairment that leads to this delay in the initiation of reactive distractor suppression. For

621	example, reactive distractor suppression may be engaged late because of the slower accumulation
622	of neural resources (Grady, 2012), or because the attentional capture process is intact but the delay
623	is in the transition to inhibition/rapid rejection, or because both attentional capture and
624	inhibition/rapid rejection are independently delayed. Of course, these possibilities are not
625	necessarily mutually exclusive. An alternative explanation that could account for this data is that
626	there is a "fast" and a "slow" reactive control mechanism and that only the faster reactive
627	mechanism is impaired. Future research will have to distinguish between these possibilities.
628	
629	
630	
631	
632	
633	
634	
635	
636	
637	
638	
639	
640	
641	

# References

643	Abu-Akel, A., Apperly, I. A., Wood, S. J., Hansen, P. C., & Mevorach, C. (2016). Autism
644	Tendencies and Psychosis Proneness Interactively Modulate Saliency Cost.
645	Schizophrenia Bulletin, sbw066. https://doi.org/10.1093/schbul/sbw066
646	Abu-Akel, A., Apperly, I., Spaniol, M., Geng, J., & Mevorach, C. (2016). Context-given
647	benefits: Saliency-based selection as a function of autism and psychosis traits. Journal
648	of Vision, 16(12), 16. https://doi.org/10.1167/16.12.16
649	Abu-Akel, A., Apperly, I., Spaniol, M. M., Geng, J. J., & Mevorach, C. (2018). Diametric
650	effects of autism tendencies and psychosis proneness on attention control irrespective
651	of task demands. Scientific Reports, 8(1), 8478. https://doi.org/10.1038/s41598-018-
652	26821-7
653	Abu-Akel, A. M., Apperly, I. A., Wood, S. J., & Hansen, P. C. (2016). Autism and psychosis
654	expressions diametrically modulate the right temporoparietal junction. Social
655	Neuroscience, 0(0), 1-13. https://doi.org/10.1080/17470919.2016.1190786
656	Amer, T., Campbell, K. L., & Hasher, L. (2016). Cognitive Control As a Double-Edged
657	Sword. Trends in Cognitive Sciences, 20(12), 905–915.
658	https://doi.org/10.1016/j.tics.2016.10.002
659	Andrews-Hanna, J. R., Snyder, A. Z., Vincent, J. L., Lustig, C., Head, D., Raichle, M. E., &
660	Buckner, R. L. (2007). Disruption of Large-Scale Brain Systems in Advanced Aging.
661	Neuron, 56(5), 924–935. https://doi.org/10.1016/j.neuron.2007.10.038
662	Anguera, J. A., & Gazzaley, A. (2012). Dissociation of motor and sensory inhibition
663	processes in normal aging. Clinical Neurophysiology, 123(4), 730-740.
664	https://doi.org/10.1016/j.clinph.2011.08.024
665	Aschenbrenner, A. J., & Balota, D. A. (2017). Dynamic adjustments of attentional control in
666	healthy aging. Psychology and Aging, 32(1), 1. https://doi.org/10.1037/pag0000148

- 667 Ashinoff, B. K. (2017). The behavioral and neural correlates of cognitive control across the
- 668 *lifespan* (PhD). University of Birmingham. http://etheses.bham.ac.uk/7738/
- 669 Badre, D., & Wagner, A. D. (2005). Frontal lobe mechanisms that resolve proactive
- 670 interference. *Cerebral Cortex*, *15*(12), 2003–2012.
- 671 https://doi.org/10.1093/cercor/bhi075
- Badre, David, & Wagner, A. D. (2007). Left ventrolateral prefrontal cortex and the cognitive
- 673 control of memory. *Neuropsychologia*, 45(13), 2883–2901.
- https://doi.org/10.1016/j.neuropsychologia.2007.06.015
- Bauer, E., Gebhardt, H., Gruppe, H., Gallhofer, B., & Sammer, G. (2012). Altered negative
- priming in older subjects: first evidence from behavioral and neural level. *Frontiers in Human Neuroscience*, 6, 270. https://doi.org/10.3389/fnhum.2012.00270
- Blais, C., & Bunge, S. (2009). Behavioral and Neural Evidence for Item-specific
- 679 Performance Monitoring. *Journal of Cognitive Neuroscience*, 22(12), 2758–2767.
- 680 https://doi.org/10.1162/jocn.2009.21365
- Blais, C., Harris, M. B., Guerrero, J. V., & Bunge, S. A. (2012). Rethinking the role of
- automaticity in cognitive control. *The Quarterly Journal of Experimental Psychology*,

683 65(2), 268–276. https://doi.org/10.1080/17470211003775234

Bloemendaal, M., Zandbelt, B., Wegman, J., van de Rest, O., Cools, R., & Aarts, E. (2016).

685 Contrasting neural effects of aging on proactive and reactive response inhibition.

- 686 *Neurobiology of Aging*. https://doi.org/10.1016/j.neurobiolaging.2016.06.007
- Bojko, A., Kramer, A. F., & Peterson, M. S. (2004). Age equivalence in switch costs for
- 688 prosaccade and antisaccade tasks. *Psychology and Aging*, *19*(1), 226–234.
- 689 https://doi.org/10.1037/0882-7974.19.1.226
- Braver, T. S. (2012). The variable nature of cognitive control: a dual mechanisms framework.

691 *Trends in Cognitive Sciences*, *16*(2), 106–113.

- 692 https://doi.org/10.1016/j.tics.2011.12.010
- Braver, T. S., & Barch, D. M. (2002). A theory of cognitive control, aging cognition, and
  neuromodulation. *Neuroscience & Biobehavioral Reviews*, 26(7), 809–817.
- 695 https://doi.org/10.1016/S0149-7634(02)00067-2
- Braver, T. S., Paxton, J. L., Locke, H. S., & Barch, D. M. (2009). Flexible neural mechanisms
  of cognitive control within human prefrontal cortex. *Proceedings of the National*
- 698 *Academy of Sciences*, *106*(18), 7351–7356. https://doi.org/10.1073/pnas.0808187106
- 699 Braver, T. S., Satpute, A. B., Rush, B. K., Racine, C. A., & Barch, D. M. (2005). Context
- 700 Processing and Context Maintenance in Healthy Aging and Early Stage Dementia of
- the Alzheimer's Type. *Psychology and Aging*, 20(1), 33–46.
- 702 https://doi.org/10.1037/0882-7974.20.1.33
- Brodeur, D. A., & Enns, J. T. (1997). Covert visual orienting across the lifespan. *Canadian Journal of Experimental Psychology*, *51*(1), 20–35.
- Bugg, J. M., Jacoby, L. L., & Toth, J. P. (2008). Multiple levels of control in the Stroop task.
- 706 *Memory & Cognition*, 36(8), 1484–1494. https://doi.org/10.3758/MC.36.8.1484
- 707 Bugg, J. M. (2014a). Conflict-Triggered Top-Down Control: Default Mode, Last Resort, or
- No Such Thing? Journal of Experimental Psychology. Learning, Memory, and
- 709 *Cognition*, 40(2), 567–587. https://doi.org/10.1037/a0035032
- 710 Bugg, J. M. (2014b). Evidence for the sparing of reactive cognitive control with age.
- 711 *Psychology and Aging*, 29(1), 115–127. https://doi.org/10.1037/a0035270
- 712Bugg, J. M. (2015). The relative attractiveness of distractors and targets affects the coming
- and going of item-specific control: Evidence from flanker tasks. *Attention*,
- 714 *Perception, & Psychophysics*, 77(2), 373–389. https://doi.org/10.3758/s13414-014-
- 715 0752-x
- 716 Bugg, J. M., & Chanani, S. (2011). List-wide control is not entirely elusive: Evidence from

- 717 picture-word Stroop. Psychonomic Bulletin & Review, 18(5), 930-936.
- https://doi.org/10.3758/s13423-011-0112-y 718
- Bugg, J. M., & Crump, M. J. C. (2012). In Support of a Distinction between Voluntary and 719 720 Stimulus-Driven Control: A Review of the Literature on Proportion Congruent
- Effects. Frontiers in Psychology, 3. https://doi.org/10.3389/fpsyg.2012.00367 721
- Bugg, J. M., Jacoby, L. L., & Chanani, S. (2011). Why it is too early to lose control in 722
- 723 accounts of item-specific proportion congruency effects. Journal of Experimental *Psychology: Human Perception and Performance*, *37*(3), 844-859. 724
- 725 Bush, G., Whalen, P. J., Rosen, B. R., Jenike, M. A., McInerney, S. C., & Rauch, S. L.
- (1998). The counting stroop: An interference task specialized for functional 726
- neuroimaging-validation study with functional MRI. Human Brain Mapping, 6(4), 727
- 728 270-282. https://doi.org/10.1002/(SICI)1097-0193(1998)6:4<270::AID-
- 729 HBM6>3.0.CO;2-0
- Chen, P. L., & Machado, L. (2016). Age-related deficits in voluntary control over saccadic 730
- 731 eye movements: consideration of electrical brain stimulation as a therapeutic strategy.
- *Neurobiology of Aging*, 41, 53–63. 732
- https://doi.org/10.1016/j.neurobiolaging.2016.02.010 733
- Cohen-Shikora, E. R., Diede, N. T., & Bugg, J. M. (2018). The flexibility of cognitive 734
- 735 control: Age equivalence with experience guiding the way. *Psychology and Aging*,
- 736 33(6), 924–939. https://doi.org/10.1037/pag0000280
- Corbetta, M., & Shulman, G. L. (2002). Control of goal-directed and stimulus-driven 737 attention in the brain. Nature Reviews Neuroscience, 3(3), 201–215.
- 739 https://doi.org/10.1038/nrn755
- Craik, F. I., & Byrd, M. (1982). Aging and cognitive deficits. Aging and cognitive processes. 740
- *Springer*, 191–211. 741

- 742 Craik, F. I. M., & Salthouse, T. A. (2011). *The Handbook of Aging and Cognition: Third*743 *Edition*. Psychology Press.
- 744 Crawford, T. J., Higham, S., Mayes, J., Dale, M., Shaunak, S., & Lekwuwa, G. (2013). The
- role of working memory and attentional disengagement on inhibitory control: effects
- of aging and Alzheimer's disease. *AGE*, *35*(5), 1637–1650.
- 747 https://doi.org/10.1007/s11357-012-9466-y
- Darowski, E. S., Helder, E., Zacks, R. T., Hasher, L., & Hambrick, D. Z. (2008). Age-related
  differences in cognition: The role of distraction control. *Neuropsychology*, 22(5),
- 750 638–644. https://doi.org/10.1037/0894-4105.22.5.638
- 751 DiQuattro, N. E., & Geng, J. J. (2011). Contextual Knowledge Configures Attentional
- 752 Control Networks. *The Journal of Neuroscience*, *31*(49), 18026–18035.
- 753 https://doi.org/10.1523/JNEUROSCI.4040-11.2011
- Dixon, R. A., Bäckman, L., Park, D. C., Backman, P. of P. at U. of G. D. of P. S. D. of N. and
- G. M. L., & Nilsson, L.-G. (2004). *New Frontiers in Cognitive Aging*. Oxford
  University Press.
- Dowiasch, S., Marx, S., Einhäuser, W., & Bremmer, F. (2015). Effects of aging on eye
- movements in the real world. *Frontiers in Human Neuroscience*, 9.
- 759 https://doi.org/10.3389/fnhum.2015.00046
- 760 Du, Y., Xiao, Z., Song, Y., Fan, S., Wu, R., & Zhang, J. X. (2008). An electrophysiological
- signature for proactive interference resolution in working memory. *International*
- *Journal of Psychophysiology*, 69(2), 107–111.
- 763 https://doi.org/10.1016/j.ijpsycho.2008.03.007
- Erel, H., & Levy, D. A. (2016). Orienting of visual attention in aging. *Neuroscience* &
- 765 *Biobehavioral Reviews*, 69, 357–380. https://doi.org/10.1016/j.neubiorev.2016.08.010
- Faul, F., Erdfelder, E., Buchner, A., & Lang, A.-G. (2009). Statistical power analyses using

767	G*Power 3.1: Tests for correlation and regression analyses. Behavior Research
768	Methods, 41(4), 1149–1160. https://doi.org/10.3758/BRM.41.4.1149
769	Faul, F., Erdfelder, E., Lang, AG., & Buchner, A. (2007). G*Power 3: A flexible statistical
770	power analysis program for the social, behavioral, and biomedical sciences. Behavior
771	Research Methods, 39(2), 175–191. https://doi.org/10.3758/BF03193146
772	Faust, M. E., Balota, D. A., Spieler, D. H., Ferraro, F. R., Faust, M. E., Psychology, D. O.,
773	Department, P. (1999). Individual differences in information-processing rate and
774	amount: Implications for group differences in response latency. Psychological
775	Bulletin, 777–799.
776	Folk, C. L., & Hoyer, W. J. (1992). Aging and shifts of visual spatial attention. Psychology
777	and Aging, 7(3), 453-465. https://doi.org/10.1037//0882-7974.7.3.453
778	Frings, C., Schneider, K. K., & Fox, E. (2015). The negative priming paradigm: An update
779	and implications for selective attention. Psychonomic Bulletin & Review, 22(6),
780	1577-1597. https://doi.org/10.3758/s13423-015-0841-4
781	Gazzaley, A., Cooney, J. W., Rissman, J., & D'Esposito, M. (2005). Top-down suppression
782	deficit underlies working memory impairment in normal aging. Nature Neuroscience,
783	8(10), 1298–1300. https://doi.org/10.1038/nn1543
784	Geerligs, L., Maurits, N. M., Renken, R. J., & Lorist, M. M. (2014). Reduced specificity of
785	functional connectivity in the aging brain during task performance. Human Brain
786	Mapping, 35(1), 319-330. https://doi.org/10.1002/hbm.22175
787	Geerligs, L., Saliasi, E., Maurits, N. M., Renken, R. J., & Lorist, M. M. (2014). Brain
788	mechanisms underlying the effects of aging on different aspects of selective attention.
789	NeuroImage, 91, 52-62. https://doi.org/10.1016/j.neuroimage.2014.01.029
790	Geng, J. J., & DiQuattro, N. E. (2010). Attentional capture by a perceptually salient non-
791	target facilitates target processing through inhibition and rapid rejection. Journal of

- Glaser, W. R., & Glaser, M. O. (1989). Context effects in stroop-like word and picture
  processing. *Journal of Experimental Psychology. General*, *118*(1), 13–42.
- Gonthier, C., Braver, T. S., & Bugg, J. M. (2016). Dissociating proactive and reactive control
- in the Stroop task. *Memory & Cognition*, 44(5), 778–788.
- 797 https://doi.org/10.3758/s13421-016-0591-1
- 798 Grady, C. (2012). The cognitive neuroscience of ageing. *Nature Reviews Neuroscience*,
- 799 *13*(7), 491–505. https://doi.org/10.1038/nrn3256
- Greenwood, Pamela M., & Parasuraman, R. (1994). Attentional disengagement deficit in
- 801 nondemented elderly over 75 years of age. *Aging, Neuropsychology, and Cognition,*
- 802 *1*(3), 188–202. https://doi.org/10.1080/13825589408256576
- Greenwood, P.M., Parasuraman, R., & Haxby, J. V. (1993). Changes in visuospatial attention
  over the adult lifespan. *Neuropsychologia*, *31*(5), 471–485.
- 805 https://doi.org/10.1016/0028-3932(93)90061-4
- Hasher, L., Stoltzfus, E. R., Zacks, R. T., & Rypma, B. (1991). Age and inhibition. *Journal of Experimental Psychology: Learning, Memory, and Cognition*, 17(1), 163.
- 808 Hasher, L., & Zacks, R. T. (1988). Working Memory, Comprehension, and Aging: A Review
- and a New View. In G. H. Bower (Ed.), *Psychology of Learning and Motivation* (Vol.
- 810 22, pp. 193–225). Academic Press. Retrieved from
- 811 http://www.sciencedirect.com/science/article/pii/S0079742108600419
- Houston, J. R., Bennett, I. J., Allen, P. A., & Madden, D. J. (2016). Visual Acuity does not
- 813 Moderate Effect Sizes of Higher-Level Cognitive Tasks. *Experimental Aging*
- 814 *Research*, 42(3), 221–263. https://doi.org/10.1080/0361073X.2016.1156964
- 815 Iarocci, G., Enns, J. T., Randolph, B., & Burack, J. A. (2009). The modulation of visual
- 816 orienting reflexes across the lifespan. *Developmental Science*, *12*(5), 715–724.

817 https://doi.org/10.1111/j.1467-7687.2009.00810.x

- 818 Irlbacher, K., Kraft, A., Kehrer, S., & Brandt, S. A. (2014). Mechanisms and neuronal
  819 networks involved in reactive and proactive cognitive control of interference in
- working memory. *Neuroscience & Biobehavioral Reviews*, 46, Part 1, 58–70.
- 821 https://doi.org/10.1016/j.neubiorev.2014.06.014
- Jennings, J. M., Dagenbach, D., Engle, C. M., & Funke, L. J. (2007). Age-related changes
- and the attention network task: an examination of alerting, orienting, and executive
- function. *Neuropsychology, Development, and Cognition. Section B, Aging,*
- 825 *Neuropsychology and Cognition*, *14*(4), 353–369.
- 826 https://doi.org/10.1080/13825580600788837
- Jimura, K., & Braver, T. S. (2010). Age-Related Shifts in Brain Activity Dynamics during
  Task Switching. *Cerebral Cortex*, 20(6), 1420–1431.
- 829 https://doi.org/10.1093/cercor/bhp206
- Jimura, K., Locke, H. S., & Braver, T. S. (2010). Prefrontal cortex mediation of cognitive
- 831 enhancement in rewarding motivational contexts. *Proceedings of the National*
- 832 *Academy of Sciences*, *107*(19), 8871–8876. https://doi.org/10.1073/pnas.1002007107
- 833 Kingstone, A., Klein, R., Morein-Zamir, S., Hunt, A., Fisk, J., & Maxner, C. (2002).
- 834 Orienting attention in aging and Parkinson's disease: Distinguishing modes of control.
- *Journal of Clinical and Experimental Neuropsychology*, 24(7), 951–967.
- 836 https://doi.org/10.1076/jcen.24.7.951.8387
- 837 Klein, C., Fischer, B., Hartnegg, K., Heiss, W. H., & Roth, M. (2000). Optomotor and
- 838 neuropsychological performance in old age. *Experimental Brain Research*, 135(2),
- 839 141–154. https://doi.org/10.1007/s002210000506
- 840 Kolodny, T., Mevorach, C., & Shalev, L. (2017). Isolating response inhibition in the brain:
- Parietal versus frontal contribution. *Cortex*, 88, 173–185.

- 842 https://doi.org/10.1016/j.cortex.2016.12.012
- Kramer, A. F., Humphrey, D. G., Larish, J. F., & Logan, G. D. (1994). Aging and inhibition:
  Beyond a unitary view of inhibitory processing in attention. *Psychology and Aging*,
  9(4), 491–512. https://doi.org/10.1037/0882-7974.9.4.491
- Langley, L. K., Friesen, C. K., Saville, A. L., & Ciernia, A. T. (2011). Timing of reflexive
  visuospatial orienting in young, young-old, and old-old adults. *Attention, Perception,*
- *and Psychophysics*, *73*(5), 1546–1561. https://doi.org/10.3758/s13414-011-0108-8
- 849 Langley, L. K., Gayzur, N. D., Saville, A. L., Morlock, S. L., & Bagne, A. G. (2011). Spatial
- distribution of attentional inhibition is not altered in healthy aging. *Attention*,
- *Perception, and Psychophysics*, *73*(3), 766–783. https://doi.org/10.3758/s13414-0100059-5
- Larson, M. J., Clayson, P. E., Keith, C. M., Hunt, I. J., Hedges, D. W., Nielsen, B. L., & Call,
- V. R. A. (2016). Cognitive control adjustments in healthy older and younger adults:
- 855 Conflict adaptation, the error-related negativity (ERN), and evidence of generalized
  856 decline with age. *Biological Psychology*, *115*, 50–63.
- 857 https://doi.org/10.1016/j.biopsycho.2016.01.008
- Li, S.-C., Lindenberger, U., & Sikström, S. (2001). Aging cognition: from neuromodulation
  to representation. *Trends in Cognitive Sciences*, 5(11), 479–486.
- 860 https://doi.org/10.1016/S1364-6613(00)01769-1
- 861 Li, T., Yao, Y., Cheng, Y., Xu, B., Cao, X., Waxman, D., ... Feng, J. (2016). Cognitive
- training can reduce the rate of cognitive aging: a neuroimaging cohort study. *BMC*
- 863 *Geriatrics*, 16, 12. https://doi.org/10.1186/s12877-016-0194-5
- Lustig, C., Hasher, L., & Zacks, R. T. (2007). Inhibitory deficit theory: Recent developments
  in a "new view." *Inhibition in Cognition*, *17*, 145–162.
- Mahoney, J. R., Verghese, J., Goldin, Y., Lipton, R., & Holtzer, R. (2010). Alerting,

- 867 orienting, and executive attention in older adults. *Journal of the International*
- 868 *Neuropsychological Society: JINS*, *16*(5), 877–889.
- 869 https://doi.org/10.1017/S1355617710000767
- 870 Mayas, J., Fuentes, L. J., & Ballesteros, S. (2012). Stroop interference and negative priming
- 871 (NP) suppression in normal aging. *Archives of Gerontology and Geriatrics*, 54(2),
- 872 333–338. https://doi.org/10.1016/j.archger.2010.12.012
- 873 Mecklinger, A., Weber, K., Gunter, T. C., & Engle, R. W. (2003). Dissociable brain
- 874 mechanisms for inhibitory control: effects of interference content and working
- 875 memory capacity. *Cognitive Brain Research*, *18*(1), 26–38.
- 876 https://doi.org/10.1016/j.cogbrainres.2003.08.008
- Mevorach, C., Spaniol, M. M., Soden, M., & Galea, J. M. (2016). Age-dependent distractor
  suppression across the vision and motor domain. *Journal of Vision*, *16*(11), 27.
- 879 https://doi.org/10.1167/16.11.27
- 880 Monge, Z. A., & Madden, D. J. (2016). Linking Cognitive and Visual Perceptual Decline in
- 881 Healthy Aging: The Information Degradation Hypothesis. *Neuroscience and*
- 882 *Biobehavioral Reviews*, 69, 166–173. https://doi.org/10.1016/j.neubiorev.2016.07.031
- Morse, C. K. (1993). Does variability increase with age? An archival study of cognitive
  measures. *Psychology and Aging*, 8(2), 156–164.
- 885 Olk, B., & Kingstone, A. (2009). A new look at aging and performance in the antisaccade
  886 task: The impact of response selection. *European Journal of Cognitive Psychology*,
- 887 21(2–3), 406–427. https://doi.org/10.1080/09541440802333190
- 888 Owsley, C. (2016). Vision and Aging. *Annual Review of Vision Science*, 2(1), 255–271.
- 889 https://doi.org/10.1146/annurev-vision-111815-114550
- 890 Pardhan, S. (2004). Contrast sensitivity loss with aging: sampling efficiency and equivalent
- 891 noise at different spatial frequencies. *JOSA A*, *21*(2), 169–175.

- 893 Paxton, J. L., Barch, D. M., Racine, C. A., & Braver, T. S. (2008). Cognitive Control, Goal
- 894 Maintenance, and Prefrontal Function in Healthy Aging. *Cerebral Cortex*, 18(5),
- 895 1010–1028. https://doi.org/10.1093/cercor/bhm135
- Polich, J. (2007). Updating P300: An Integrative Theory of P3a and P3b. *Clinical*
- 897 *Neurophysiology : Official Journal of the International Federation of Clinical*
- 898 *Neurophysiology*, *118*(10), 2128–2148. https://doi.org/10.1016/j.clinph.2007.04.019
- 899 Porto, F. H. G., Tusch, E. S., Fox, A. M., Alperin, B., Holcomb, P. J., & Daffner, K. R.
- 900 (2016). One of the most well-established age-related changes in neural activity
- 901 disappears after controlling for visual acuity. *NeuroImage*, *130*, 115–122.
- 902 https://doi.org/10.1016/j.neuroimage.2016.01.035
- Puccioni, O., & Vallesi, A. (2012). Conflict resolution and adaptation in normal aging: The
  role of verbal intelligence and cognitive reserve. *Psychology and Aging*, 27(4), 1018.
- 905 https://doi.org/10.1037/a0029106
- 906 Rey-Mermet, A., & Gade, M. (2018). Inhibition in aging: What is preserved? What declines?
- 907 A meta-analysis. *Psychonomic Bulletin & Review*, 25(5), 1695–1716.
- 908 https://doi.org/10.3758/s13423-017-1384-7
- 909 Rey-Mermet, A., Gade, M., & Oberauer, K. (2018). Should we stop thinking about
- 910 inhibition? Searching for individual and age differences in inhibition ability. *Journal*
- 911 *of Experimental Psychology: Learning, Memory, and Cognition, 44*(4), 501–526.
- 912 https://doi.org/10.1037/xlm0000450
- 913 Roberts, K. L., & Allen, H. A. (2016). Perception and Cognition in the Ageing Brain: A Brief
- 914 Review of the Short- and Long-Term Links between Perceptual and Cognitive
- 915 Decline. *Frontiers in Aging Neuroscience*, 8.
- 916 https://doi.org/10.3389/fnagi.2016.00039

917	Ross, R. G., Olincy, A., Harris, J. G., Radant, A., Adler, L. E., Compagnon, N., & Freedman,
918	R. (1999). The effects of age on a smooth pursuit tracking task in adults with
919	schizophrenia and normal subjects. Biological Psychiatry, 46(3), 383-391.
920	https://doi.org/10.1016/S0006-3223(98)00369-2
921	Salthouse, T.A. (1994). The Aging of Working Memory. <i>Neuropsychology</i> , 8(4), 535–543.
922	Salthouse, Timothy A. (1996). The processing-speed theory of adult age differences in
923	cognition. Psychological Review, 103(3), 403-428. https://doi.org/10.1037/0033-
924	295X.103.3.403
925	Salthouse, Timothy A. (2000). Aging and measures of processing speed. Biological
926	Psychology, 54(1-3), 35-54. https://doi.org/10.1016/S0301-0511(00)00052-1
927	Salthouse, Timothy A., & Meinz, E. J. (1995). Aging, Inhibition, Working Memory, and
928	Speed. The Journals of Gerontology Series B: Psychological Sciences and Social
929	Sciences, 50B(6), P297–P306. https://doi.org/10.1093/geronb/50B.6.P297
930	Sebastian, A., Baldermann, C., Feige, B., Katzev, M., Scheller, E., Hellwig, B., Klöppel,
931	S. (2013). Differential effects of age on subcomponents of response inhibition.
932	Neurobiology of Aging, 34(9), 2183–2193.
933	https://doi.org/10.1016/j.neurobiolaging.2013.03.013
934	Sekuler, R., & Hutman, L. P. (1980). Spatial Vision and Aging. I: Contrast Sensitivity.
935	Journal of Gerontology, 35(5), 692-699. https://doi.org/10.1093/geronj/35.5.692
936	Strauss, G. P., Allen, D. N., Jorgensen, M. L., & Cramer, S. L. (2005). Test-Retest Reliability
937	of Standard and Emotional Stroop Tasks: An Investigation of Color-Word and
938	Picture-Word Versions. Assessment, 12(3), 330-337.
939	https://doi.org/10.1177/1073191105276375
940	Stroop, J. R. (1935). Studies of interference in serial verbal reactions. Journal of
941	Experimental Psychology, 18(6), 643-662. https://doi.org/10.1037/h0054651

- Studebaker, G. A. (1985). A Rationalized Arcsine Transform. *Journal of Speech, Language*, *and Hearing Research*, 28. https://doi.org/10.1044/jshr.2803.455
- Swick, D., Ashley, V., & Turken, U. (2011). Are the neural correlates of stopping and not
  going identical? Quantitative meta-analysis of two response inhibition tasks.
- 946 *NeuroImage*, *56*(3), 1655–1665. https://doi.org/10.1016/j.neuroimage.2011.02.070
- 947 Titz, C., Behrendt, J., Menge, U., & Hasselhorn, M. (2008). A Reassessment of Negative
- 948 Priming Within the Inhibition Framework of Cognitive Aging: There Is More in It
- 949 Than Previously Believed. *Experimental Aging Research*, *34*(4), 340–366.
- 950 https://doi.org/10.1080/03610730802273936
- 951 Tsvetanov, K. A., Mevorach, C., Allen, H., & Humphreys, G. W. (2013). Age-related
- differences in selection by visual saliency. *Attention, Perception, & Psychophysics*,
  75(7), 1382–1394. https://doi.org/10.3758/s13414-013-0499-9
- 954 Vadaga, K. K., Blair, M., & Li, K. Z. H. (2016). Are Age-Related Differences Uniform
- 955 Across Different Inhibitory Functions? *The Journals of Gerontology Series B:*
- 956 *Psychological Sciences and Social Sciences*, 71(4), 641–649.
- 957 https://doi.org/10.1093/geronb/gbv002
- Verhaeghen, P., & Cerella, J. (2002). Aging, executive control, and attention: a review of
  meta-analyses. *Neuroscience & Biobehavioral Reviews*, 26(7), 849–857.
- 960 https://doi.org/10.1016/S0149-7634(02)00071-4
- Verhaeghen, P., & De Meersman, L. (1998). Aging and the Stroop effect: A meta-analysis. *Psychology and Aging*, *13*(1), 120–126. https://doi.org/10.1037/0882-7974.13.1.120
- 963 Warren, D. E., Thurtell, M. J., Carroll, J. N., & Wall, M. (2013). Perimetric evaluation of
- saccadic latency, saccadic accuracy, and visual threshold for peripheral visual stimuli
- 965 in young compared with older adults. *Investigative Ophthalmology and Visual*
- 966 *Science*, 54(8), 5778–5787. https://doi.org/10.1167/iovs.13-12032

- 967 Waszak, F., Li, S.-C., & Hommel, B. (2010). The development of attentional networks:
- 968 Cross-sectional findings from a life span sample. *Developmental Psychology*, 46(2),
  969 337–349. https://doi.org/10.1037/a0018541
- 970 West, R., & Moore, K. (2005). Adjustments of Cognitive Control in Younger and Older
- 971 Adults. Cortex, 41(4), 570–581. https://doi.org/10.1016/S0010-9452(08)70197-7
- 972 Xiang, L., Zhang, B., Wang, B., Jiang, J., Zhang, F., & Hu, Z. (2016). The Effect of Aging on
- 973 the Dynamics of Reactive and Proactive Cognitive Control of Response Interference.

974 Frontiers in Psychology, 7. https://doi.org/10.3389/fpsyg.2016.01640

- 275 Zanto, T. P., Toy, B., & Gazzaley, A. (2010). Delays in neural processing during working
- 976 memory encoding in normal aging. *Neuropsychologia*, 48(1), 13–25.
- 977 https://doi.org/10.1016/j.neuropsychologia.2009.08.003
- 278 Zhang, J. X., Wu, R., Kong, L., Weng, X., & Du, Y. (2010). Electrophysiological correlates
  of proactive interference in the 'Recent Probes' verbal working memory task.

980 *Neuropsychologia*, 48(7), 2167–2173.

981 https://doi.org/10.1016/j.neuropsychologia.2010.04.008