

## Interest in orthodontic tooth alignment in adult patients affected by periodontitis

Hirschfeld, Josefine; Reichardt, Elisabeth; Sharma, Praveen; Hilber, Anna; Meyer-marcotty, Philipp; Stellzig-eisenhauer, Angelika; Schlagenhauf, Ulrich; Sickel, Franka E.

DOI:

[10.1002/JPER.18-0578](https://doi.org/10.1002/JPER.18-0578)

License:

Other (please specify with Rights Statement)

*Document Version*

Peer reviewed version

*Citation for published version (Harvard):*

Hirschfeld, J, Reichardt, E, Sharma, P, Hilber, A, Meyer-marcotty, P, Stellzig-eisenhauer, A, Schlagenhauf, U & Sickel, FE 2019, 'Interest in orthodontic tooth alignment in adult patients affected by periodontitis: a questionnaire-based cross-sectional pilot study', *Journal of Periodontology*, vol. 90, no. 9, pp. 957-965. <https://doi.org/10.1002/JPER.18-0578>

[Link to publication on Research at Birmingham portal](#)

### **Publisher Rights Statement:**

Checked for eligibility 18/02/2019

This is the peer reviewed version of the following article: Hirschfeld, J, Reichardt, E, Sharma, P, et al. Interest in orthodontic tooth alignment in adult patients affected by periodontitis: A questionnaire-based cross-sectional pilot study. *J Periodontol.* 2019; 90: 957– 965., which has been published in final form at: <https://doi.org/10.1002/JPER.18-0578>. This article may be used for non-commercial purposes in accordance with Wiley Terms and Conditions for Use of Self-Archived Versions.

### **General rights**

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

### **Take down policy**

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact [UBIRA@lists.bham.ac.uk](mailto:UBIRA@lists.bham.ac.uk) providing details and we will remove access to the work immediately and investigate.



**Interest in orthodontic tooth alignment in adult patients affected by periodontitis: a questionnaire-based cross-sectional pilot study.**

Journal:	<i>Journal of Periodontology</i>
Manuscript ID	JOP-18-0578.R2
Manuscript Type:	Original Article
Date Submitted by the Author:	30-Jan-2019
Complete List of Authors:	Hirschfeld, Josefine; University of Birmingham College of Medical and Dental Sciences, School of Dentistry Reichardt, Elisabeth; Charité Universitätsmedizin Berlin Centrum für Zahn-, Mund- und Kieferheilkunde, Orthodontics, Dentofacial Orthopedics and Pedodontics Sharma, Praveen; University of Birmingham College of Medical and Dental Sciences, School of Dentistry Hilber, Anna; Universitätsklinikum Würzburg Klinik und Poliklinik für Mund- Kiefer- und Plastische Gesichtschirurgie, Orthodontics Meyer-Marcotty, Philipp; Georg-August-Universität Göttingen Universitätsmedizin, Center for Dentistry, Oral Medicine and Craniomaxillofacial Surgery Stellzig-Eisenhauer, Angelika; Universitätsklinikum Würzburg Klinik und Poliklinik für Mund- Kiefer- und Plastische Gesichtschirurgie, Orthodontics Schlagenhauf, Ulrich; Universitätsklinikum Würzburg Klinik und Poliklinik für Mund- Kiefer- und Plastische Gesichtschirurgie, Periodontology Sickel, Franka; Universitätsklinikum Würzburg Klinik und Poliklinik für Mund- Kiefer- und Plastische Gesichtschirurgie, Orthodontics
Key Words:	Periodontitis, Orthodontics, Treatment planning

1  
2  
3 **Interest in Orthodontic Tooth Alignment in Adult Patients Affected by Periodontitis: A**  
4  
5 **Questionnaire-Based Cross-Sectional Pilot Study**  
6  
7  
8  
9

10 **Running Title: Interest in Orthodontic Treatment in Periodontitis Patients**  
11  
12  
13

14 **Dr.** Josefine Hirschfeld<sup>\*</sup>, **Dr.** Elisabeth Reichardt<sup>†</sup>, **Dr.** Praveen Sharma<sup>\*</sup>, **Dr.** Anna Hilber<sup>‡</sup>,  
15  
16 **Prof.** Philipp Meyer-Marcotty<sup>§</sup>, **Prof.** Angelika Stellzig-Eisenhauer<sup>‡</sup>, **Prof.** Ulrich  
17  
18 Schlagenhauf<sup>¶</sup>, **Dr.** Franka E. SICKEL<sup>‡</sup>  
19  
20  
21  
22  
23

24 <sup>\*</sup>Department of Restorative Dentistry, Birmingham Dental School and Hospital,  
25  
26 Birmingham, UK

27 <sup>†</sup> Department of Orthodontics, Dentofacial Orthopedics and Pedodontics, Charité, Berlin,  
28  
29 Germany

30 <sup>‡</sup>Department of Orthodontics, University Hospital Würzburg, Würzburg, Germany

31 <sup>§</sup> Department of Orthodontics, University Medical Center Göttingen, Göttingen, Germany

32 <sup>¶</sup> Department of Periodontology, University Hospital Würzburg, Würzburg, Germany  
33  
34  
35  
36  
37  
38

39 **Correspondence:**

40 Josefine Hirschfeld  
41 Birmingham Dental School and Hospital  
42 5 Mill Pool Way, B5 7EG Birmingham, UK  
43 Phone: +44 (0) 121 4665114  
44 Email: [j.hirschfeld@bham.ac.uk](mailto:j.hirschfeld@bham.ac.uk)  
45  
46  
47

48 **Summary Sentence:**

49 The findings of this study show that a considerable number of adult patients with moderate to  
50 severe periodontitis is interested in undergoing orthodontic treatment in order to improve oral  
51 and overall health, function and esthetics.  
52  
53  
54  
55

56 **3 Figures; 2 Tables; 50 References; 3475 Words**  
57  
58  
59  
60

## ABSTRACT

**Background:** Orthodontic treatment can successfully align pathologically migrated teeth and lead to improvement of periodontal stability in patients with periodontitis. Periodontic-orthodontic approaches have gained increasing attention in the past years. Here, we investigated the interest of adults affected by chronic periodontitis in undergoing orthodontic treatment as well as patient-related and tooth-related influence factors.

**Methods:** Periodontal and orthodontic measurements/indices were taken from 115 adult patients with moderate to severe periodontitis. The study participants answered a questionnaire investigating patient demographics, quality of life aspects and their interest in undergoing orthodontic treatment. Correlations between clinical data, questionnaire responses and this interest were analyzed by means of an age- and gender-adjusted multiple regression model.

**Results:** Two thirds of the participants were interested in orthodontic therapy and indicated long-term healthy and esthetically appealing teeth as their main motives. A significant correlation was found between subjectively felt impaired dental esthetics and an interest in orthodontic treatment. However, there were no correlations with the severity of periodontitis, tooth alignment or patient demographics, including gender. Older patients were significantly more often interested in orthodontic treatment. Most participants had never been provided with information about orthodontic treatment options for adults.

**Conclusion:** A considerable number of adult patients with periodontitis is interested in orthodontics to improve tooth alignment. However, severity of periodontitis and tooth misalignment or demographic factors may not be indicative thereof. Therefore, dental

1  
2  
3 practitioners need to be aware of patients wishing to align their teeth and to provide them  
4  
5 with the relevant information and, if appropriate, enable interdisciplinary treatment planning.  
6  
7  
8  
9  
10

11  
12 **Keywords:** periodontitis, orthodontics, adult, surveys and questionnaires, quality of life,  
13  
14 esthetics  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

For Peer Review

## INTRODUCTION

Periodontitis is a chronic inflammatory disease of the gingival tissues and alveolar bone triggered by oral biofilms <sup>1</sup>. If left untreated, in susceptible individuals, this inflammation results in periodontal attachment loss and eventually tooth loss <sup>2</sup>. Periodontitis can also lead to pathologic tooth migration (PTM), which often motivates patients to seek periodontal therapy <sup>3</sup>. PTM occurs in 30-56% of the patients, mainly caused by diminished periodontal tissue support, mechanical pressure of edematous gingiva, dysfunctions or malocclusion <sup>4,5</sup>. As a consequence, teeth become extruded, protruded and gaps can develop between the anterior teeth. This appearance can worsen after molar loss, often leading to mesial tipping of posterior teeth and resulting in a deeper bite with increased anterior occlusal loading. Dysbalanced perioral muscular forces and bruxism can further diminish oral function and phonetic problems may occur <sup>6,7</sup>. Periodontal therapy alone is unlikely to completely resolve the esthetic challenges of such patients. Due to soft tissue shrinkage after successful periodontal therapy, cervical areas often become exposed and black triangles develop. Thus, the oral/dental appearance of these patients is impaired which affects their dentofacial attractiveness <sup>8</sup>. These developments can drastically reduce the patient's self-confidence and quality of life (QOL) <sup>9,10</sup>. Orthodontic treatment is aimed at correcting malpositioned teeth and in the past decades, the number of orthodontic therapies in adults has continuously increased in the United States and Europe <sup>11,12</sup>.

Realignment of malpositioned teeth can substantially help maintaining the natural dentition, control periodontal breakdown and restore oral function <sup>13-16</sup>, however, it is crucial to achieve a stable and non-inflamed periodontal condition, as well as a high level of compliance regarding oral hygiene, prior to orthodontic intervention <sup>17</sup>. The correction of some orthodontic problems, such as excessively tipped molars, traumatic deep-bites and flared and

1  
2  
3 spaced incisors, may be particularly beneficial in periodontally compromised individuals<sup>18</sup>.  
4  
5 Despite possible, temporary, compromise in esthetics and comfort as well as financial effort  
6  
7 during orthodontic therapy, a considerable number of adult patients pursue orthodontic  
8  
9 therapy to rectify PTM and optimize their dental esthetics<sup>19</sup>. According to previous reports,  
10  
11 the key motives for patients seeking orthodontic treatment are improvements in esthetic  
12  
13 appearance, functional and phonetic rehabilitation, maintenance of teeth and an improvement  
14  
15 in QOL<sup>20-23</sup>. Oral health related QOL is a subjective parameter and is evaluated by assessing  
16  
17 functional restrictions of the orofacial system, orofacial pain, dentofacial esthetics as well as  
18  
19 psychosocial influence of oral health.  
20  
21  
22  
23  
24  
25

26 Given that periodontitis and periodontal therapy significantly impact on dental esthetics and  
27  
28 orthodontic therapy may reduce this impact, there is little information in the literature  
29  
30 regarding the interest of adult patients with periodontitis in undergoing orthodontic therapy.  
31  
32 Therefore, the first aim of this study was to characterize, in a cohort of adult patients in  
33  
34 Germany affected by moderate to severe periodontitis, the patient- and tooth-related factors  
35  
36 associated with seeking orthodontic treatment. The second aim of this study was to  
37  
38 investigate predictors of an interest in orthodontic treatment (IOT) within this cohort.  
39  
40  
41  
42  
43  
44

## 45 **METHODS**

### 46 **Patients**

47  
48  
49 A total of 115 patients were recruited for this study. Patients were recruited between 02/2013  
50  
51 and 06/2013 from the Periodontology department at the Dental Hospital of the University of  
52  
53 Würzburg, Germany. All participants were undergoing non-surgical periodontal therapy in  
54  
55 regular recall visits and were chosen for the study based on their basic periodontal  
56  
57 examination (BPE) scores. Inclusion criteria were  
58  
59  
60

- aged >18 years
- BPE of  $\geq 3$  in at least three sextants
- diagnosis of generalized chronic periodontitis, defined by: a predominantly horizontal pattern of radiographic bone loss, the teeth affected (non-molar-incisor bone loss pattern), patient age and the initial presence of plaque and/or calculus.

Exclusion criteria were

- extensive prosthetic restorations such as partial dentures
- gaps of more than one missing tooth, in order to exclude the possibility that potential oral discomfort assessed in the questionnaire was caused by large spaces alone
- dysgnathia such as cleft lip/palate.

All patients participated in the study after signing informed consent and the study was approved by the ethics committee of the University Hospital Würzburg, Germany (approval number: AZ 41/12).

### Measures of periodontal status

To assess the periodontal status of patients, periodontal probing was performed of all teeth at six sites for single-rooted teeth and eight or ten sites for upper or lower molars, respectively.

Bleeding on probing (BOP) and gingival recessions were also recorded for each site. All measurements were taken by one examiner (A.H.). The updated case definition by the Centers for Disease Control/American Academy of Periodontology (CDC/AAP) working group (2012)<sup>24</sup> was used to classify periodontal health into none/mild/moderate/severe periodontitis.

### Measures of orthodontic malocclusions

Bite registrations were taken in habitual occlusion and alginate impressions of both jaws were



1  
2  
3 taken. Orthodontic study models were made and trimmed three-dimensionally in order to  
4  
5 measure and evaluate malpositions using a sliding caliper (Munich Pattern). Little's  
6  
7 irregularity index and contact point deviations of upper incisors were surveyed by measuring  
8  
9 the distances from their anatomical interproximal contact area parallel to the occlusal plane.  
10  
11 Five values (in mm) in each jaw were summed up, from the mesial contact area of one canine  
12  
13 to the mesial contact area of the other canine in the same jaw. To estimate dental  
14  
15 attractiveness, the esthetic component (AC) of the Index of Orthodontic Treatment Need  
16  
17 (IOTN) was assessed (IOTN/AC) <sup>25</sup>.  
18  
19  
20  
21  
22  
23

### 24 **Questionnaire**

25  
26 All 115 study participants filled out a questionnaire in German. The questionnaire was  
27  
28 divided into seven subsections and contained a total of 34 questions. Section 1 collected  
29  
30 demographic data (age, gender, family status, rural or urban living environment, education,  
31  
32 profession, income, smoking status). Sections 2-6 were derived from the German Oral Health  
33  
34 Impact Profile (OHIP-G49), assessing oral health related QOL in adults <sup>26</sup> and referred to the  
35  
36 time period of the preceding month. For this survey, selected OHIP-G49 questions were  
37  
38 modified to focus on possible periodontitis-related and tooth alignment-related discomfort.  
39  
40 Section 7 investigated attitudes towards orthodontic intervention and previous orthodontic  
41  
42 experience. The questionnaire translated from German to English is shown in the  
43  
44  
45  
46

### 47 **Supplementary Table 1.**

### 51 **Statistical analysis**

52  
53 A priori sample size calculation was performed using the following parameters: one tailed,  
54  
55 effect size = 0.23; alpha level = 0.05 and power = 0.8, indicating that a total sample size of  
56  
57  
58  
59  
60

1  
2  
3 n=115 was required <sup>¶</sup>. We hypothesized a positive correlation between IOT and increased  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

orthodontic and periodontal clinical parameters. The effect size was calculated according to the standard deviations published by Masood et al., who assessed the impact of malocclusion on oral health-related QOL by means of an OHIP questionnaire <sup>27</sup>. To address the aims of the study, patient- and tooth-related factors as well as QOL-related influence factors associated with seeking orthodontic treatment were assessed, considering clinical data and answers obtained from the questionnaire. Data were analyzed using a multiple regression model with the help of a statistical software <sup>#</sup>.

## RESULTS

### Cohort description

Of the 115 patients enrolled in the study, 60% (n=69) were women. The participants had a mean age of  $56.8 \pm 9.3$  years. This cohort lived mostly (61%) in an urban environment and with a partner (81%). 63% were employees or self-employed with a regular monthly income, which was predominantly in the range of 1,500-3,000€ (58%) and lower (31%). 73% of the participants had a non-university degree (vocational training degree). Most patients had a statutory health insurance (77%) and 13% were privately insured (**Table 1**). Amongst the patients were 17 smokers.

### Periodontal assessment of the cohort

Given the inclusion criteria, all participants were classified as either “moderate” (n=36) or

---

<sup>¶</sup> G\*Power Statistical Power Analyses software version 3.1.1, University of Düsseldorf, Germany (Faul et al., 2007)

<sup>#</sup> STATA 15 software, StataCorp. 2017, College Station, TX, USA

1  
2  
3 “severe” (n=79) periodontitis according to the 2012 CDC/AAP classification<sup>24</sup>. The mean  
4 number of teeth, including wisdom teeth, was 25.7. Mean PPD, CAL and recessions were  
5 2.3mm, 2.8mm and 0.5mm, respectively (**Table 2**).  
6  
7  
8  
9

### 10 11 12 **Orthodontic assessment of the cohort**

13  
14 In our cohort, 23.5% showed an IOTN/AC score of >5, indicating a moderate to definite  
15 treatment need<sup>28</sup> (**Table 2**). The mean Little’s index (lower anterior teeth) was 7.5mm,  
16  
17 whereas the mean contact point deviation (upper anterior teeth) was 8.2mm.  
18  
19

### 20 21 22 **Characteristics of patients seeking orthodontic treatment**

23  
24 In this cohort, 78 patients (68%) expressed an IOT. This group had a mean age of 54.4 (SD  
25 7.9) years and 56% were female. Patients with severe periodontitis were more likely to  
26 express an interest in orthodontic treatment. 27% of patients who expressed an interest in  
27 orthodontics had an IOTN/AC >5, with a mean contact point deviation of 7.8mm (SD  
28 2.9mm) and a mean Little’s index of 7.7mm (SD 3.3mm) (**Table 2**). Of those patients with an  
29 IOT, half had been provided with information about treatment options. Although more than  
30 one third (36%) of all participants had undergone orthodontic treatment previously, no  
31 significant correlation between past treatment and current IOT was detectable.  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44

45 Of the patients who reported an interest in orthodontic treatment, half preferred treatment  
46 durations of six months or one year (**Figure 1a**). The majority of patients (72%) indicated a  
47 preference for transparent aligners as an orthodontic appliance (**Figure 1b**) and expressed  
48 that visibility of the orthodontic appliance would be a matter of moderate to high concern for  
49 them. 22% of patients interested in orthodontic treatment indicated that they would be willing  
50 to undergo treatment which included dental extractions. Patients were also asked to indicate  
51 which treatment costs would be acceptable for them. 44% were not willing to pay for this  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 service, whereas 36% would agree to invest 1,000€ or more (20%) (**Figure 1c**).

### 8 **Motives for orthodontic treatment**

9  
10 Those who indicated IOT, most frequently reported maintaining their own teeth,  
11 improvement of esthetic appearance, enhancement of overall well-being and improving  
12 functional aspects as their main motives to undergo this treatment (**Figure 2**). There was a  
13 correlation between self-observed (questionnaire) esthetic impairment due to malpositioned  
14 teeth and IOT. This correlation was seen after dichotomization of the 5-step Likert scale  
15 answers (never vs. ever) ( $p < 0.001$ ,  $OR = 2.169 - 12.171$ ,  $CI = 95\%$ ). Interestingly, there was no  
16 statistically significant correlation between the orthodontic clinical data collected and self-  
17 observed misaligned teeth. However, this result is inversely associated with IOT: 68% of  
18 those patients who felt esthetically impaired due to their tooth position were not interested in  
19 orthodontic treatment. Furthermore, there was no correlation between IOT and other QOL-  
20 related factors such as tooth-related self-consciousness, avoidance of smiling or worry and  
21 depression (data not shown). None of the OHIP-G49 derived questionnaire results related to  
22 QOL impairment due to tooth positions, the orofacial system and gingivae showed to have an  
23 influence on IOT in our regression model.  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44

### 45 **IOT is independent of clinical parameters but associated with age**

46 Two thirds (67.8%) of the participants (71% of the men, 63% of the women) indicated an  
47 IOT. Interestingly, a higher trend towards an interest in orthodontic treatment was observed  
48 with increasing age ( $p = 0.055$ ,  $OR = 0.99 - 1.1$ ,  $CI = 95\%$ ). Multiple regression analysis showed  
49 that neither the periodontal parameters (mean clinical attachment loss,  $p = 0.264$ , recession,  
50  $p = 0.515$ ) nor orthodontic misalignment measured by IOTN/AC (dichotomized in scores 1-5  
51 and 6-10), Little's irregularity index and contact point deviation of upper anterior teeth had  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 an influence on IOT ( $p=0.245$ ,  $0.268$  and  $0.075$  respectively). Adjustment for age and gender  
4  
5 in the regression model confirmed these results. Furthermore, patient demographics like  
6  
7 monthly income, education, type of health insurance, profession, living environment (rural  
8  
9 versus urban) and family status had no influence on IOT (**Table 1**).

### 14 **Information sources of orthodontic treatment possibilities for adults**

15  
16 All participants were asked to report sources from which they had obtained information about  
17  
18 orthodontic treatment for adults. 57% indicated that they had never been informed before,  
19  
20 whereas dentists and orthodontist were a major information source for many others. Internet  
21  
22 and media such as magazines or television only played a minor role (**Figure 3**). Of those  
23  
24 patients with an IOT, half had been provided with information about treatment options.  
25  
26 Although more than one third (36%) of all participants had undergone orthodontic treatment  
27  
28 previously, no significant correlation between past treatment and current IOT was detectable  
29  
30 (p=0.3).  
31  
32  
33  
34  
35  
36  
37

### 38 **DISCUSSION**

39  
40 The patient population with a history of periodontitis and pathologic tooth migrations is  
41  
42 likely to increase in orthodontic clinics over the coming years, as in ageing populations a  
43  
44 prolonged maintenance of teeth is observed <sup>29</sup>. This study characterizes for the first time a  
45  
46 cohort of periodontitis patients regarding their interest in orthodontic tooth alignment in  
47  
48 relation to their periodontal and orthodontic indices as well as patient demographics. In our  
49  
50 study cohort, the majority of participants were over 50 years old, reflecting the increasing  
51  
52 incidence of chronic periodontitis in aged individuals <sup>30</sup>. As an important finding of this  
53  
54 study, a large proportion (68%) of our cohort expressed an IOT. Those who indicated IOT  
55  
56 most frequently reported maintaining own teeth, improvement of esthetic appearance,  
57  
58  
59  
60

1  
2  
3 enhancement of overall well-being and improving functional aspects as their main motives.

4  
5 This indicates that many patients are aware of the importance of oral and overall health and  
6  
7 are seeking to improve both.  
8  
9

10  
11  
12 It was previously reported that orthodontic therapy may not only ameliorate periodontal  
13  
14 parameters but could also help patients to gain better self-assurance and psychological well-  
15  
16 being and to provide amended possibilities of oral hygiene<sup>31-33</sup>. Past surveys also revealed  
17  
18 that esthetic appearance and dentofacial harmony are key drivers for a patient's decision to  
19  
20 undergo orthodontic treatment and that patients are often aware of tooth deviation and  
21  
22 malocclusion<sup>20-22</sup>. Dental esthetics and function often decrease with the progression of  
23  
24 periodontitis, recession formation and general factors like tooth wear and discoloration, all of  
25  
26 which are typically increased with age<sup>34</sup>. Interestingly, however, our study showed that self-  
27  
28 reported impaired dental esthetics due to misaligned teeth were not positively correlated with  
29  
30 IOT. Moreover, this reported esthetic impairment was not corresponding to the measured  
31  
32 orthodontic indices. These results indicate that patients judge their own dental esthetics  
33  
34 individually and independently of orthodontic treatment need, and that orthodontic indices  
35  
36 and even self-reported dental esthetic impairment are not predictive factors of IOT. Similarly,  
37  
38 our analysis revealed that there were no differences between men and women in expressing  
39  
40 IOT, being in accordance with previous reports<sup>35, 36</sup>.  
41  
42  
43  
44  
45  
46  
47  
48

49 In our cohort, one third showed a moderate to definite orthodontic treatment need, however,  
50  
51 higher IOTN/AC scores or contact point deviations were not correlated with IOT. It is  
52  
53 possible that the presence of periodontitis, regular periodontal treatment visits and therefore  
54  
55 increased dental awareness may influence these patients' judgments and change their oral and  
56  
57 dental health priorities. Nearly half of those study participants having a history of orthodontic  
58  
59  
60

1  
2  
3 intervention was interested in retreatment, although there was no significant correlation  
4  
5 between previous treatment and current interest. The investigation of psychosocial aspects in  
6  
7 orthodontically treated and untreated adults in a study by Demasure-Trockels et al. reported  
8  
9 that there appears to be a higher degree of sensitivity and self-observation in already-treated  
10  
11 patients, leading to a higher motivation for orthodontic re-intervention <sup>37</sup>. In our cohort, the  
12  
13 frequent periodontal treatment visits may have a similar effect, supported by the fact that  
14  
15 most patients reported that their oral health awareness had increased due to the frequent recall  
16  
17 visits.  
18  
19  
20  
21  
22  
23

24 Our results further show that many patients (46% of those with IOT) would favor treatment  
25  
26 periods of one year or less, although 36% indicated that treatment duration would not play an  
27  
28 important role in the decision to undergo orthodontic treatment. More permanent treatment  
29  
30 modes like fixed orthodontic braces (36%) and dental extractions (22%) were indicated as  
31  
32 acceptable by a considerable proportion, which may reflect a more serious interest and higher  
33  
34 motivation in these patients. At the same time, although the majority of participants would be  
35  
36 prepared to fund their orthodontic treatment partially or in full, a significant proportion (44%)  
37  
38 indicated that treatment costs would not be acceptable for them. Thus, potential treatment  
39  
40 devices and costs need to be considered and discussed with periodontal patients wishing to  
41  
42 undergo orthodontic therapy.  
43  
44  
45  
46  
47  
48

49 More than half of the study participants had never been introduced to orthodontic treatment  
50  
51 possibilities for adults. The other half had received information from dental professionals and  
52  
53 through their own childrens' treatment, but rarely through mass media and internet. A  
54  
55 previous study showed that individuals who receive mass media messages containing health-  
56  
57 related information, can recall these messages at a high rate <sup>38</sup>. Therefore, to better inform  
58  
59  
60

1  
2  
3 adult periodontitis patients about orthodontic treatment options, audiovisual media, social  
4  
5 media and the internet could become an important information source in the future <sup>39</sup>.

6  
7  
8 Communication between patients and dentists should also be improved in this area, as it is an  
9  
10 indispensable way to advise patients individually and in detail as well as to ensure that  
11  
12 patients obtain reliable and high quality information from the media and internet <sup>40</sup>.

13  
14  
15  
16  
17 The wish to undergo orthodontic treatment is likely to be influenced by socio-cultural  
18  
19 background and the society's esthetic expectations and ideals. In Germany, as in other  
20  
21 industrialized nations, esthetic appearance and dentofacial harmony play an important role in  
22  
23 society. It was shown that established norms for dental and facial appearance do not vary  
24  
25 widely among industrialized nations <sup>41</sup>. These norms and ideals are broadcasted through the  
26  
27 media, having a strong impact on the behavior and decision-making of beauty-conscious  
28  
29 societies, and leading to an increased demand for esthetic improvement from the public <sup>42</sup>.  
30  
31 Moreover, harmonious dental appearances and oral health were shown to have an impact on  
32  
33 employment and professional success <sup>43, 44</sup>.

34  
35  
36  
37  
38  
39  
40 The option to undergo orthodontic treatment is also often influenced by financial aspects.  
41  
42 Orthodontic treatment of adult patients to correct malpositioned teeth is currently not funded  
43  
44 by the statutory health insurance in Germany <sup>45</sup>, whereas some private health insurances may  
45  
46 contribute to or fully cover the costs arising from such treatment. Therefore, for most patients  
47  
48 this treatment would be self-funded and may not always be affordable. The per capita net  
49  
50 (after taxes) income in Germany in the year 2011 was 2,174 € (2,756 US\$) and, for  
51  
52 comparison, in the USA 2,984 US\$ (using purchasing power parity exchange rates) <sup>46</sup>. This  
53  
54 average income is comparable between Germany and the USA when considering living  
55  
56 expenses: the consumer price index (CPI) for Germany in 2011 was 102.1 and for the USA  
57  
58  
59  
60



1  
2  
3 103.2 (2010 = 100) <sup>47</sup>. According to the Gebührenordnung für Zahnärzte (GOZ, dental fee  
4  
5 schedule), the costs of orthodontic treatment, on average and for treatment of all teeth, ranges  
6  
7 between 4,000 and 6,000 € ( $\approx$  4,500 – 6,800 US\$) <sup>48</sup>. However, treatment cost can  
8  
9 substantially deviate from this average, depending on the specific case complexity, treatment  
10  
11 duration, techniques and materials used.  
12  
13  
14  
15  
16

17 A limitation of this study is that it does not allow for comparison with IOT in periodontally  
18  
19 healthy individuals, which may reveal a potential influence of existing periodontitis or  
20  
21 periodontal therapy on IOT. Moreover, the questionnaire used in this pilot study requires to  
22  
23 be fully validated in the next step, which is also aimed at expanding the study cohort and to  
24  
25 include patients with a wider range of PTM severity. Nevertheless, as most questions in our  
26  
27 questionnaire were derived from the OHIP-G49, we argue that potential biases derived from  
28  
29 the questionnaire design, if any, may be minor and inconsequential for the outcomes of this  
30  
31 pilot survey. Additionally, the inclusion of further orthodontic indices such as the IOTN/DC  
32  
33 in a future study may reveal statistically significant correlations with self-observed PTM and  
34  
35 IOT. Expanding the study cohort should also be aimed at increasing the sample size for the  
36  
37 purpose of fully validating the questionnaire, and to increase the generalizability of our  
38  
39 findings to the wider population of patients affected by periodontitis.  
40  
41  
42  
43  
44  
45  
46

## 47 **CONCLUSION**

48  
49 As there do not appear to be factors that can reliably predict a patient's wish to undergo  
50  
51 orthodontic tooth alignment other than the explicit statement thereof, improved  
52  
53 communication between patients and dental practitioners as well as between periodontists  
54  
55 and orthodontists can help to identify patients interested in correcting the position of their  
56  
57 pathologically migrated teeth <sup>49, 50</sup>. However, this communication between the dental  
58  
59  
60

1  
2  
3 practitioner and the patient should also be aimed at carefully managing expectations  
4  
5 regarding treatment cost, duration, risks and possible outcomes, as these are key to attaining  
6  
7 high satisfaction with treatment outcomes in both parties <sup>17</sup>. Importantly, this group of  
8  
9 patients needs to be made aware of the necessity to stabilize their periodontal condition and  
10  
11 maintain high standards of oral hygiene prior to any orthodontic intervention.  
12  
13

14 The results of this study can facilitate addressing the treatment wishes of adult periodontitis  
15  
16 patients, as they demonstrate that adult patients with periodontitis are often interested in  
17  
18 orthodontic tooth alignment, however, this interest was not related to demographic factors,  
19  
20 subjective tooth-related physical or psychological discomfort and clinical indices in our  
21  
22 cohort.  
23  
24  
25  
26  
27  
28  
29

### 30 **ACKNOWLEDGMENTS**

31  
32 The authors wish to acknowledge the participation of all study participants. The Center for  
33  
34 Dental, Oral and Maxillofacial Medicine of the University of Würzburg ;provided the study  
35  
36 premises and technical equipment. Moreover, we thank Sabine Karl from the Faculty of  
37  
38 Mathematics and Informatics of the University of Würzburg for her valuable contribution to  
39  
40 statistical data analysis as well as Nicole Petersen from the Department of Periodontology  
41  
42 (Würzburg) for her help with recruiting patients.  
43  
44  
45  
46  
47  
48  
49

### 50 **CONFLICT OF INTEREST STATEMENT**

51  
52 The authors declare that they have no conflict of interest. This study was financially  
53  
54 supported by the German Society of Periodontology (DGParo; Regensburg, Germany) and  
55  
56 German Society of Dental Oral and Maxillofacial Medicine (DGZMK; Düsseldorf,  
57  
58 Germany).  
59  
60

## REFERENCES

1. Meyle J, Chapple I. Molecular aspects of the pathogenesis of periodontitis. *Periodontol 2000* 2015;69:7-17.
2. Needleman I, Garcia R, Gkraniias N, et al. Mean annual attachment, bone level, and tooth loss: A systematic review. *J Clin Periodontol* 2018;45 Suppl 20:S112-s129.
3. Brunsvold MA. Pathologic tooth migration. *J Clin Periodontol* 2005;76:859-866.
4. Martinez-Canut P, Carrasquer A, Magan R, Lorca A. A study on factors associated with pathologic tooth migration. *J Clin Periodontol* 1997;24:492-497.
5. Dadlani H, Ramachandra SS, Mehta DS. Spontaneous correction of pathologically migrated teeth with periodontal therapy alone. *J Indian Soc Periodontol* 2013;17:531-534.
6. Towfighi PP, Brunsvold MA, Storey AT, Arnold RM, Willman DE, McMahan CA. Pathologic migration of anterior teeth in patients with moderate to severe periodontitis. *J Periodontol* 1997;68:967-972.
7. Rathod SR, Kolte AP, Chintawar S. The dynamic relationship between pathological migrating teeth and periodontal disease. *J Indian Soc Periodontol* 2013;17:762-764.
8. Tugnait A, Clerehugh V. Gingival recession-its significance and management. *J Dent* 2001;29:381-394.
9. Zhang M, McGrath C, Hagg U. The impact of malocclusion and its treatment on quality of life: a literature review. *Int J Paediatr Dent* 2006;16:381-387.
10. Klages U, Bruckner A, Zentner A. Dental aesthetics, self-awareness, and oral health-related quality of life in young adults. *Eur J Orthod* 2004;26:507-514.
11. Khan RS, Horrocks EN. A study of adult orthodontic patients and their treatment. *Br J Orthod* 1991;18:183-194.

- 1  
2  
3 12. Proffit WR, Fields HWJ, Moray LJ. Prevalence of malocclusion and orthodontic  
4 treatment need in the United States: estimates from the NHANES III survey. *Int J*  
5  
6 *Adult Orthodon Orthognath Surg* 1998;13:97-106.  
7  
8
- 9  
10 13. Ong MMA, Wang HL. Periodontic and orthodontic treatment in adults. *Am J Orthod*  
11  
12 *Dentofacial Orthop* 2002;122:420–428.  
13
- 14 14. Re S, Corrente G, Abundo R, Cardaropoli D. Orthodontic treatment in periodontally  
15 compromised patients: 12-year report. *Int J Periodontics Restorative Dent*  
16  
17 2000;20:31-39.  
18
- 19 20 15. Corrente G, Abundo R, Re S, Cardaropoli D, Cardaropoli G. Orthodontic movement  
21 into infrabony defects in patients with advanced periodontal disease: a clinical and  
22 radiological study. *J Periodontol* 2003;74:1104-1109.  
23  
24  
25  
26  
27
- 28 29 16. Panwar M, Jayan B, Arora V, Singh S. Orthodontic management of dentition in  
30 patients with periodontally compromised dentition. *J Indian Soc Periodontol*  
31  
32 2014;18:200-204.  
33  
34
- 35 36 17. Gkantidis N, Christou P, Topouzelis N. The orthodontic-periodontic interrelationship  
37 in integrated treatment challenges: a systematic review. *J Oral Rehabil* 2010;37:377-  
38  
39 390.  
40  
41
- 42 43 18. Antoun JS, Mei L, Gibbs K, Farella M. Effect of orthodontic treatment on the  
44 periodontal tissues. *Periodontol 2000* 2017;74:140-157.  
45  
46
- 47 48 19. Gazit-Rappaport T, Haisraeli-Shalish M, Gazit E. Psychosocial reward of orthodontic  
49 treatment in adult patients. *Eur J Orthod* 2010;32:441-446.  
50
- 51 52 20. Espeland LV, Stenvik A. Perception of personal dental appearance in young adults:  
53 Relationship between occlusion, awareness, and satisfaction. *Am J Orthod*  
54  
55 *Dentofacial Orthop* 1991;100:234–241.  
56  
57  
58  
59  
60

- 1  
2  
3 21. McKiernan EX, McKiernan F, Jones ML. Psychological profiles and motives of  
4 adults seeking orthodontic treatment. *Int J Adult Orthodon Orthognath Surg*  
5  
6 1992;7:187-198.  
7  
8
- 9  
10 22. Meyer-Marcotty P, Alpers GW, Gerdes AB, Stellzig-Eisenhauer A. Impact of facial  
11 asymmetry in visual perception: a 3-dimensional data analysis. *Am J Orthod*  
12  
13 *Dentofacial Orthop* 2010;137:168.e161-168; discussion 168-169.  
14  
15
- 16 23. Pabari S, Moles DR, Cunningham SJ. Assessment of motivation and psychological  
17 characteristics of adult orthodontic patients. *Am J Orthod Dentofacial Orthop*  
18  
19 2011;140:e263-272.  
20  
21
- 22 24. Eke PI, Page RC, Wei L, Thornton-Evans G, Genco RJ. Update of the case definitions  
23 for population-based surveillance of periodontitis. *J Periodontol* 2012;83:1449-1454.  
24  
25
- 26 25. Brook PH, Shaw WC. The development of an index of orthodontic treatment priority.  
27  
28 *Eur J Orthod* 1989;11:309-320.  
29  
30
- 31 26. John MT, Patrick DL, Slade GD. The German version of the Oral Health Impact  
32 Profile--translation and psychometric properties. *Eur J Oral Sci* 2002;110:425-433.  
33  
34
- 35 27. Masood Y, Masood M, Zainul NNB, Araby NBAA, Hussain SF, Newton T. Impact of  
36 malocclusion on oral health related quality of life in young people. *Health Qual Life*  
37 *Outcomes* 2013;11:25-25.  
38  
39
- 40 28. Kapoor P, Singh H. Evaluation of esthetic component of the index of orthodontic  
41 treatment need: The orthodontists' perspective. *Indian J Dent* 2015;6:181-184.  
42  
43
- 44 29. Tonetti MS, Bottenberg P, Conrads G, et al. Dental caries and periodontal diseases in  
45 the ageing population: call to action to protect and enhance oral health and well-being  
46 as an essential component of healthy ageing - Consensus report of group 4 of the joint  
47 EFP/ORCA workshop on the boundaries between caries and periodontal diseases. *J*  
48 *Clin Periodontol* 2017;44 Suppl 18:S135-s144.  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

- 1  
2  
3 30. Eke PI, Wei L, Borgnakke WS, et al. Periodontitis prevalence in adults  $\geq$  65 years  
4 of age, in the USA. *Periodontol 2000* 2016;72:76-95.  
5  
6  
7
- 8 31. de Couto Nascimento V, de Castro Ferreira Conti AC, de Almeida Cardoso M,  
9 Valarelli DP, de Almeida-Pedrin RR. Impact of orthodontic treatment on self-esteem  
10 and quality of life of adult patients requiring oral rehabilitation. *Angle Orthod*  
11 2016;86:839-845.  
12  
13
- 14 32. Johal A, Alyaqoobi I, Patel R, Cox S. The impact of orthodontic treatment on quality  
15 of life and self-esteem in adult patients. *Eur J Orthod* 2015;37:233-237.  
16  
17
- 18 33. Hunt O, Hepper P, Johnston C, Stevenson M, Burden D. Professional perceptions of  
19 the benefits of orthodontic treatment. *Eur J Orthod* 2001;23:315-323.  
20  
21
- 22 34. Carvalho TS, Lussi A. Age-related morphological, histological and functional  
23 changes in teeth. *J Oral Rehabil* 2017;44:291-298.  
24  
25
- 26 35. Soderfeldt B, Palmqvist S, Arnbjerg D. Factors affecting attitudes toward dental  
27 appearance and dental function in a Swedish population aged 45-69 years. *Community*  
28 *Dent Health* 1993;10:123-130.  
29  
30
- 31 36. Al-Zarea BK. Satisfaction with appearance and the desired treatment to improve  
32 aesthetics. *Int J Dent* 2013;2013:912368.  
33  
34
- 35 37. Demasure-Trockels P, Katsaros C, Besold G, Berg R. [The psychosocial aspects in  
36 orthodontically treated and untreated adults with similar, clearly visible tooth  
37 malalignments]. *J Orofac Orthop* 1995;56:77-83.  
38  
39
- 40 38. Marcus BH, Owen N, Forsyth LH, Cavill NA, Fridinger F. Physical activity  
41 interventions using mass media, print media, and information technology. *Am J Prev*  
42 *Med* 1998;15:362-378.  
43  
44
- 45 39. Jorgensen G. Social media basics for orthodontists. *Am J Orthod Dentofacial Orthop*  
46 2012;141:510-515.  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

- 1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60
40. Cline RJ, Haynes KM. Consumer health information seeking on the Internet: the state of the art. *Health Educ Res* 2001;16:671-692.
41. Cons NC, Jenny J, Kohout FJ, Freer TJ, Eismann D. Perceptions of occlusal conditions in Australia, the German Democratic Republic and the United States of America. *Int Dent J* 1983;33:200-206.
42. Aschheim KW. *Esthetic dentistry: A clinical approach to techniques and materials, third edition*; 2014: 1-588.
43. Pithon MM, Nascimento CC, Barbosa GC, Coqueiro Rda S. Do dental esthetics have any influence on finding a job? *Am J Orthod Dentofacial Orthop* 2014;146:423-429.
44. Hyde S, Satariano WA, Weintraub JA. Welfare dental intervention improves employment and quality of life. *J Dent Res* 2006;85:79-84.
45. Bundesausschuss der Zahnärzte und Krankenkassen. Richtlinien des Bundesausschusses der Zahnärzte und Krankenkassen für die kieferorthopädische Behandlung. In: Bundesanzeiger, ed. 226:24-966. Cologne: Bundesanzeiger Verlag, 2004.
46. OECD. Taxing Wages - Comparative tables: Net income after taxes - US dollars PPP exchange rates. Available at: <https://stats.oecd.org/index.aspx?queryid=55145>. 2019.
47. World Bank. Consumer price index (2010 = 100) - International Monetary Fund, International Financial Statistics and data files. Available at: <https://data.worldbank.org/indicator/FP.CPI.TOTL?end=2011&start=2010>. 2019.
48. Bundeszahnärztekammer. Gebührenordnung für Zahnärzte / Dental Fee Schedule (GOZ). Ed. Berlin: Federal Government of Germany, 2012.
49. Bills D. Collaboration between the general dentist and orthodontist: an essential partnership to ensure orthodontic success. *Compend Contin Educ Dent* 2013;34:262-267.

- 1  
2  
3 50. Mavreas D, Athanasiou AE. Orthodontics and its interactions with other dental  
4 disciplines. *Prog Orthod* 2009;10:72-81.  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

For Peer Review



**FIGURE LEGENDS**

**Figure 1.** Patients' (n=78) acceptance of orthodontic therapy duration and appliances. **A)**

Willingness to tolerate orthodontic therapy length, **B)** acceptable orthodontic appliances

(multiple answers were possible), **C)** acceptable treatment costs payable. 1000 € ≈ 1100 US\$,

5000 € ≈ 5700 US\$, US\$ rounded to the nearest 100. Currency conversion as of January

2019 (source: <https://www.xe.com/currencyconverter/>).

**Figure 2.** Motives for orthodontic treatment in male and female participants (n=78) interested

in undergoing orthodontic tooth alignment (multiple answers were possible).

**Figure 3.** Previous sources of information regarding orthodontic treatment option for adult

patients employed by all study participants (multiple answers were possible).

## TABLES

**Table 1.** Description of the cohort demographics and association with IOT.




Demographic parameters	Whole cohort			p-value*
	(n=115)	IOT (n=78)	No IOT (n=37)	
Age (years)	56.8 (9.3)	54.4 (7.9)	57.9 (9.7)	0.055
Female (%)	60	56.4	67.6	0.310
Private Insurance (%)	13.0	16.7	5.4	0.138
Urban residence (%)	60.9	61.5	59.5	0.841
Monthly income** (after tax, in €) %				0.461
<1500 (<1700 US\$)	30.6	31.9	27.8	
1500-3000	58.3	59.7	55.6	
>3000 (>3400 US\$)	11.1	8.3	16.7	
Previous orthodontic treatment (% yes)	35.7	32.1	43.2	0.298

\* comparing those interested in orthodontics VS not. Values are mean and (SD), unless otherwise stated. \*\* Currency conversion as of January 2019, US\$ rounded to the nearest 100 (source: <https://www.xe.com/currencyconverter/>).

**Table 2.** Description of the cohort's clinical parameters and association with IOT.

Clinical parameters	Whole cohort (n=115)	IOT (n=78)	No IOT (n=37)	p- value*
Periodontal diagnoses (%)				0.291
Moderate periodontitis	31.3	24.3	34.6	
Severe periodontitis	68.7	75.7	65.4	
Number of teeth	25.7 (2.8)	25.8 (2.8)	25.5 (2.7)	0.564
Mean PPD (mm)	2.3 (0.4)	2.2 (0.4)	2.3 (0.4)	0.374
Mean CAL (mm)	2.8 (0.7)	2.7 (0.6)	2.9 (0.8)	0.264
Mean recession (mm)	0.5 (0.6)	0.5 (0.6)	0.6 (0.8)	0.515
IOTN/AC >5 (%)	23.5	26.9	16.2	0.245
Mean contact point deviation (mm)	8.2 (3.1)	7.8 (2.9)	8.9 (3.4)	0.075
Mean Little's index	7.5 (3.4)	7.7 (3.3)	7.0 (3.5)	0.268
Self-perceived impaired dental esthetics (%)	40	27	68	<0.001

\* comparing those interested in orthodontics VS not. Values are mean and (SD), unless otherwise stated. Abbreviations: IOTN = Index of Orthodontic Treatment Need, IOTN/AC esthetic component of the IOTN, IOT = interest in orthodontic treatment.






Figure 3

Journal of Periodontology Page 28 of 28

