

Age should not be a barrier for pulmonary endarterectomy in carefully selected patients

Newnham, Michael; Hernandez-Sanchez, Jules; Dunning, John; Ng, Choo; Tsui, Steven; Bunclark, Katherine; Sheares, Karen; Taboada, Dolores; Toshner, Mark; Pepke-Zaba, Joanna; Jenkins, David; Cannon, John

DOI:

[10.1183/13993003.01804-2017](https://doi.org/10.1183/13993003.01804-2017)

License:

None: All rights reserved

Document Version

Peer reviewed version

Citation for published version (Harvard):

Newnham, M, Hernandez-Sanchez, J, Dunning, J, Ng, C, Tsui, S, Bunclark, K, Sheares, K, Taboada, D, Toshner, M, Pepke-Zaba, J, Jenkins, D & Cannon, J 2017, 'Age should not be a barrier for pulmonary endarterectomy in carefully selected patients', *European Respiratory Journal*, vol. 50, no. 6, 1701804. <https://doi.org/10.1183/13993003.01804-2017>

[Link to publication on Research at Birmingham portal](#)

Publisher Rights Statement:

Checked for eligibility: 21/03/2019

This is an author-submitted, peer-reviewed version of a manuscript that has been accepted for publication in the *European Respiratory Journal*, prior to copy-editing, formatting and typesetting. This version of the manuscript may not be duplicated or reproduced without prior permission from the copyright owner, the European Respiratory Society. The publisher is not responsible or liable for any errors or omissions in this version of the manuscript or in any version derived from it by any other parties. The final, copy-edited, published article, which is the version of record, is available without a subscription 18 months after the date of issue publication.

The final version of record can be found at: <https://doi.org/10.1183/13993003.01804-2017>

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

Title

Age should not be a barrier to pulmonary endarterectomy in carefully selected patients

Authors and affiliations

Michael Newnham^{1,2}, Jules Hernández-Sánchez¹, John Dunning¹, Choo Ng¹, Steven Tsui¹, Katherine Bunclark¹, Karen Sheares¹, Dolores Taboada¹, Mark Toshner^{1,2}, Joanna Pepke-Zaba¹, David Jenkins¹, John Cannon¹

¹ Papworth Hospital NHS Foundation Trust, Cambridge, UK

² University of Cambridge, Department of Medicine, Cambridge, UK

Corresponding author

Dr John Cannon
Pulmonary Vascular Disease Unit
Papworth Hospital NHS Foundation Trust
Papworth Everard
Cambridge
UK
CB23 3RE

Email: john.cannon1@nhs.net

“Take home” message / Abstract

CTEPH patients over 80 years old undergoing pulmonary endarterectomy have similar outcomes to those under 80.

To the Editor:

Pulmonary endarterectomy (PEA) is the treatment of choice in operable chronic thromboembolic pulmonary hypertension (CTEPH) with excellent long term outcomes [1]. It is a complex surgical procedure requiring cardiopulmonary bypass and removal of obstructive thromboembolic material during periods of deep hypothermic circulatory arrest [1]. We have observed an increase in the number of older CTEPH patients referred for consideration of PEA, which is consistent with other cardiothoracic surgeries. The UK population is aging with a projected 3% increase in those over 85 years in the next 20 years [2]. This may be mirrored by patients with CTEPH getting older, as the incidence of pulmonary embolism, which frequently precedes CTEPH, increases markedly with age [3,4]. Furthermore, an epidemiological analysis by Gall *et al*, has projected that the annual incidence of CTEPH will increase over the next 10 years [5]. Therefore, the management of CTEPH in older patients is a pertinent topic for investigation.

We previously reported that hospital survival in patients over 70 years undergoing PEA was comparable to those under 70, albeit with longer hospital and intensive care unit (ICU) stays [6]. For selected cardiac surgeries (coronary artery bypass and aortic valve surgery) octogenarians have equivalent or improved long-term mortality compared to an age- and sex-matched population [7,8]. Furthermore, patient reported outcomes (PROs) including quality of life improve post-operatively, and are equivalent or better than a matched general population of octogenarians [7,9]. However, increased in-hospital mortality and prolonged hospital / ICU length of stay have also been reported, which may translate to increased health utilisation costs [10,11]. Therefore, in the current study, we aimed to assess the outcomes of CTEPH patients over 80 years old who underwent PEA.

Consecutive CTEPH patients undergoing PEA from June 2006 to August 2016 at the UK National PEA centre (Papworth, UK) were included in a retrospective analysis. The diagnosis of CTEPH was based on international criteria [12]. Suitability for PEA was discussed by a multidisciplinary team comprising pulmonary hypertension physicians, specialist cardiothoracic radiologists and pulmonary endarterectomy surgeons. The cohort was dichotomized into over and under 80 years according to the age at time of surgery. Pre-operative baseline and post-operative 3-6-month follow-up data were recorded, with follow-up data included until 3 months after the end of the census period. PROs were assessed using the Cambridge Pulmonary Hypertension Outcome Review (CAMPHOR) score, a PH specific quality of life measure [13]. Groups were compared using a chi-squared test for categorical data, Wilcoxon rank-sum test for continuous data and log-rank test for survival data. A false discovery rate adjusted p-value was used to account for multiple testing.

1152 individuals underwent PEA (under 80 years: 1115 (97%); over 80: 37 (3%)) during the study period. Baseline and 3-6 month follow-up variables and outcomes are summarised in table 1. Overall survival at 1, 3 and 5 years was 91.8%, 88.2% and 84.4% in the under 80 group, and 83.5%, 76.4% and 69.4% in the over 80 group. Whilst survival was lower in the over 80 group (Log-rank test, $p=0.020$), it was no different from an age- and sex-matched UK reference population ($p=0.500$) [15].

There were significant improvements in World Health Organisation (WHO) functional class, 6-minute walk distance (6mwd) and haemodynamics for both under and over 80s post-PEA. Whilst the 6mwd was lower in the over 80s post-PEA (286 ± 164 vs. 360 ± 92 ; $p=0.020$) there was an equivalent change from baseline ($p=0.676$) reflecting the same magnitude of improvement. Furthermore, there was no difference in the median change from baseline for WHO functional class, haemodynamics or PROs between under and over 80s, indicating an equivalent improvement in outcomes.

There were more concomitant cardiac surgical procedures in the over 80 group (11% vs. 27% $p=0.042$), predominantly due to more coronary artery bypass grafts (7% vs. 19%). There was a shorter cardiopulmonary bypass time (323 ± 305 minutes vs. 305 ± 65 ; $p=0.020$) in the over 80s but no difference in the total deep hypothermic circulatory arrest time or type of surgical disease (Jamieson classification based on location and morphology) ($p=0.230$ and $p=1.000$). The hospital length of stay was longer in those over 80 (19 ± 7 vs. 14 ± 10 days; $p=0.020$) but there was no difference in time spent on the ICU (4 ± 3 vs. 5 ± 5 days; $p=0.310$). There was also no difference in post-PEA complications ($p=1.000$) or in hospital mortality (4% vs. 8%; $p=0.510$).

Despite the study limitations (small group size of over 80s, retrospective single-centre analysis), we found similar outcomes in patients under and over 80 years old undergoing PEA, except for a prolonged hospital length of stay in octogenarians. Whilst survival is reduced in the over 80 group compared to the under 80s, it is no different to a reference age- and sex-matched UK population. The greater number of concomitant cardiac surgeries in octogenarians could indicate that their improvement is multi-factorial. Future research should consider the health utilization and cost implications of older patients undergoing PEA, given they have a prolonged hospital length of stay.

We acknowledge that CTEPH patients over 80 were highly selected to undergo PEA and therefore our results may not apply to “all comers”. However, it reinforces the effectiveness of the PEA selection process at expert centres. Age alone should not be a contraindication for PEA and individuals with suspected CTEPH should be referred for specialist evaluation.

Acknowledgements: The National Pulmonary Hypertension Centres UK and Ireland for referring patients that were considered for PEA.

References:

1. Jenkins D, Madani M, Fadel E, et al. Pulmonary endarterectomy in the management of chronic thromboembolic pulmonary hypertension. *Eur Respir Rev* 2017; **26**: 160111.
2. Office for National Statistics. Overview of the UK population: July 2017. <https://www.ons.gov.uk/peoplepopulationandcommunity/populationandmigration/populationestimates/articles/overviewoftheukpopulation/july2017>. Date last updated: July 21 2017. Date last accessed: Aug 1 2017.

3. Silverstein MD, Heit JA, Mohr DN, et al. Trends in the incidence of deep vein thrombosis and pulmonary embolism: a 25-year population-based study. *Arch Intern Med* 1998; **158**: 585-593.
4. Pepke-Zaba J, Delcroix M, Lang I, et al. Chronic thromboembolic pulmonary hypertension (CTEPH): results from an international prospective registry. *Circulation* 2011; **124**: 1973-1981.
5. Gall H, Hoeper MM, Richter MJ, et al. An epidemiological analysis of the burden of chronic thromboembolic pulmonary hypertension in the USA, Europe and Japan. *Eur Respir Rev* 2017; **26**: 160121.
6. Berman M, Hardman G, Sharples L, et al. Pulmonary endarterectomy: outcomes in patients aged >70. *Eur J Cardiothorac Surg* 2012; **41**: 154-160
7. Krane M, Voss B, Hiebinger A, et al. Twenty years of cardiac surgery in patients aged 80 years and older: risks and benefits. *Ann Thorac Surg* 2011; **91**: 506-513.
8. Likosky DS, Sorensen MJ, Dacey LJ, et al. Long-term survival of the very elderly undergoing aortic valve surgery. *Circulation* 2009; **120**: S127-S133.
9. Ghanta RK, Shekar PS, McGurk S, et al. Long-term survival and quality of life justify cardiac surgery in the very elderly patient. *Ann Thorac Surg* 2011; **92**: 851-857.
10. Kuo K, Shah P, Hiebert B, et al. Predictors of survival, functional survival, and hospital readmission in octogenarians after surgical aortic valve replacement. *J Thorac Cardiovasc Surg* 2017; in press.
11. Gelsomino S, Lorusso R, Livi U, et al. Cost and cost-effectiveness of cardiac surgery in elderly patients. *J Thorac Cardiovasc Surg* 2011; **142**: 1062-1073.
12. Galiè N, Hoeper MM, Humbert M, et al. Guidelines for the diagnosis and treatment of pulmonary hypertension. *Eur Respir J* 2009; **34**: 1219-1263.
13. McKenna SP, Doughty N, Meads DM, et al. The Cambridge Pulmonary Hypertension Outcome Review (CAMPHOR): a measure of health-related quality of life and quality of life for patients with pulmonary hypertension. *Qual Life Res* 2006; **15**: 103-115.
14. Cannon JE, Su L, Kiely DG, et al. Dynamic Risk Stratification of Patient Long-Term Outcome After Pulmonary Endarterectomy: Results from the United Kingdom National Cohort. *Circulation* 2016; **133**: 1761-1771.
15. Office for National Statistics. National Life Tables, 2011-2013. <http://webarchive.nationalarchives.gov.uk/20160106164026/http://www.ons.gov.uk/ons/publications/re-reference-tables.html?edition=tcm%3A77-365199>. Date last updated: Archived on Jan 6 2016. Date last accessed: Feb 1 2004.

TABLE 1

Baseline variables and post-PEA outcomes subdivided by age cohorts.

	Baseline			Follow-up		
	Under 80	Over 80	p-value	Under 80	Over 80	p-value
Subjects (n)	1115	37				
Age at PEA [years]	61 (22)	81 (2)				
Gender [female]	517 (46)	17 (46)	1.000			
Comorbidities						
COPD	78 (7)	2 (5)	1.000			
Diabetes	94 (9)	2 (6)	0.950			
Atrial fibrillation/flutter	91 (9)	5 (14)	0.810			
Ischaemic heart disease	124 (12)	7 (19)	0.560			
Malignancy	85 (8)	4 (11)	1.000			
Systemic hypertension	307 (30)	10 (28)	1.000			
CAMPBOR score^a:						
Symptoms	13 ± 10	12 ± 9	1.000	4 ± 8	4 ± 7	0.860
Activity	11 ± 9	14 ± 8	0.260	6 ± 9	9 ± 6	0.300
Quality of life	11 ± 11	11 ± 11	0.870	4 ± 10	6 ± 8	1.000
WHO functional class [1/2/3/4 (%)]^b	0 / 14/ 73/ 13	0/ 3/ 70/ 26	0.046	28/ 47/ 23/ 1	22/ 40/ 33/ 4	0.450
6mwd [metres]	290 ± 216	227 ± 210	0.051	360 ± 164	286 ± 92	0.020
mPAP [mmHg]	45 ± 15	42 ± 12	0.560	25 ± 13	28 ± 12	0.560
PVR [dynes.s.cm⁻⁵]	666 ± 479	743 ± 462	0.310	235 ± 212	297 ± 294	0.081
CI [L/min/m²]	2.2 ± 0.8	1.9 ± 0.8	0.020	2.3 ± 0.7	2.3 ± 0.8	0.810
Type of surgical disease^c [1/2/3/4 (%)]	14/ 58/ 26/ 3	14/ 59/ 27/ 0	1.000			
Bypass time [mins]	323 ± 65	305 ± 51	0.020			
Arrest time [mins]	37 ± 15	33 ± 13	0.230			
Concomitant surgery: Total	123 (11)	10 (27)	0.042			
CABG	76 (7)	7 (19)				
PFO/ASD closure	34 (3)	1 (3)				
MVR	12 (2)	0 (0)				
AVR	8 (1)	2 (5)				
Complications: Total^d	277 (32)	12 (35)	1.000			
ECMO	60 (6)	2 (6)				
Pneumonia	110 (12)	6 (18)				
Re-intubation	91 (10)	3 (9)				
Renal replacement therapy	52 (6)	0 (0)				
Return to theatre	80 (9)	2 (6)				
Tracheotomy	61 (6)	1 (3)				
Length of stay [days]						
Intensive care unit	4 ± 3	5 ± 5	0.310			
Total hospital	14 ± 10	19 ± 7	0.020			
In-hospital mortality [%]	4	8	0.510			
Residual PH^e						
mPAP≥25mmHg				463 (51)	18 (62)	1.000
mPAP≥30mmHg				309 (34)	13 (44)	0.560
mPAP≥38mmHg and PVR≥425dynes.s.cm ⁻⁵				101 (11)	2 (7)	0.930
Survival [%]						
1 year				91.8	83.5	0.020
3 years				88.2	76.4	
5 years				84.4	69.4	

Data presented as median ± Interquartile range or number of patients (%) unless otherwise stated. 6mwt: 6-minute walk distance; AVR: aortic valve replacement; CABG: coronary artery bypass graft; CAMPBOR: Cambridge Pulmonary Hypertension Outcome Review score; CI: cardiac index; ECMO: Extracorporeal membrane oxygenation; mPAP: mean pulmonary arterial pressure; MVR: mitral valve replacement; PFO/ASD: patent foramen ovale / atrial septal defect; PH: pulmonary hypertension; PVR: pulmonary vascular resistance; WHO: World Health Organisation.

^a CAMPBOR score: a higher score denotes a worse patient reported outcome

^b Total percentage may not add up to 100 due to rounding

^c Jamieson classification

^d Total: any of the listed complications

^e The higher residual PH thresholds have been associated with post-PEA outcomes [14]