

## The local structure theorem

Parker, Chris; Stroth, Gernot

DOI:

[10.1016/j.jalgebra.2019.08.013](https://doi.org/10.1016/j.jalgebra.2019.08.013)

License:

Creative Commons: Attribution-NonCommercial-NoDerivs (CC BY-NC-ND)

*Document Version*

Peer reviewed version

*Citation for published version (Harvard):*

Parker, C & Stroth, G 2019, 'The local structure theorem: the wreath product case', *Journal of Algebra*.  
<https://doi.org/10.1016/j.jalgebra.2019.08.013>

[Link to publication on Research at Birmingham portal](#)

### General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

### Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact [UBIRA@lists.bham.ac.uk](mailto:UBIRA@lists.bham.ac.uk) providing details and we will remove access to the work immediately and investigate.

# THE LOCAL STRUCTURE THEOREM: THE WREATH PRODUCT CASE

CHRIS PARKER AND GERNOT STROTH

Dedicated to the memory of Kay Magaard

ABSTRACT. Groups with a large  $p$ -subgroup,  $p$  a prime, include almost all of the groups of Lie type in characteristic  $p$  and so the study of such groups adds to our understanding of the finite simple groups. In this article we study a special class of such groups which appear as wreath product cases of the Local Structure Theorem [MSS2].

## 1. INTRODUCTION

Throughout this article  $p$  is a prime and  $G$  is a finite group. We say that  $L \leq G$  has *characteristic  $p$*  if

$$C_G(O_p(L)) \leq O_p(L).$$

For  $T$  a non-trivial  $p$ -subgroup of  $G$ , the subgroup  $N_G(T)$  is called a  *$p$ -local* subgroup of  $G$ . By definition  $G$  has *local characteristic  $p$*  if all  $p$ -local subgroups of  $G$  have characteristic  $p$  and  $G$  has *parabolic characteristic  $p$*  if all  $p$ -local subgroups containing a Sylow  $p$ -subgroup of  $G$  have characteristic  $p$ .

A group  $K$  is called a  *$\mathcal{K}$ -group* if all its composition factors are from the known finite simple groups. So, if  $K$  is a simple  $\mathcal{K}$ -group, then  $K$  is a cyclic group of prime order, an alternating group, a simple group of Lie type or one of the 26 sporadic simple groups. A group  $G$  is a  *$\mathcal{K}_p$ -group*, provided all subgroups of all  $p$ -local subgroups of  $G$  are  $\mathcal{K}$ -groups. This paper is part of a programme to investigate the structure of certain  $\mathcal{K}_p$ -groups. See [MSS1, MSS2] for an overview of the project.

Of fundamental importance to the development of the programme are large subgroups of  $G$ : a  $p$ -subgroup  $Q$  of  $G$  is *large* if

- (i)  $C_G(Q) \leq Q$ ; and
- (ii)  $N_G(U) \leq N_G(Q)$  for all  $1 \neq U \leq C_G(Q)$ .

For example, if  $G$  is a simple group of Lie type defined in characteristic  $p$ ,  $S \in \text{Syl}_p(G)$  and  $Q = O_p(C_G(Z(S)))$ , then  $Q$  is a large subgroup of  $G$  unless there is some degeneracy in the Chevalley commutator relations which define  $G$ . This means that  $Q$  is a large subgroup of  $G$  unless  $G$  is one of  $\text{Sp}_{2n}(2^k)$ ,  $n \geq 2$ ,  $F_4(2^k)$  or  $G_2(3^k)$ .

If  $Q$  is a large subgroup of  $G$ , then it is easy to see that  $O_p(N_G(Q))$  is also a large  $p$ -subgroup of  $G$ . Thus we also assume that

- (iii)  $Q = O_p(N_G(Q))$ .

One of the consequences of  $G$  having a large  $p$ -subgroup is that  $G$  has parabolic characteristic  $p$ . In fact any  $p$ -local subgroup of  $G$  containing  $Q$  is

of characteristic  $p$  [MSS2, Lemma 1.5.5 (e)]. Further, if  $Q \leq S \in \text{Syl}_p(G)$ , then  $Q$  is weakly closed in  $S$  with respect to  $G$  ( $Q$  is the unique  $G$ -conjugate of  $Q$  in  $S$ ) [MSS2, Lemma 1.5.2 (e)]. A significant part of the programme described in [MSS1] aims to determine the groups which possess a large  $p$ -subgroup. This endeavour extends and generalizes earlier work of Timmesfeld and others in the original proof of the classification theorem where groups with a so-called large extraspecial 2-subgroup were investigated. The state of play at the moment is that the Local Structure Theorem has been completed and published [MSS2]. To describe this result we need some further notation.

For a finite group  $L$ ,  $Y_L$  denotes the unique maximal elementary abelian normal  $p$ -subgroup of  $L$  with  $O_p(L/C_L(Y_L)) = 1$ . Such a subgroup exists [MSS1, Lemma 2.0.1(a)]. From now on assume that  $G$  is a finite  $\mathcal{K}_p$ -group,  $S$  a Sylow  $p$ -subgroup of  $G$  and  $Q$  a large  $p$ -subgroup of  $G$  with  $Q \leq S$  and  $Q = O_p(N_G(Q))$ . We define

$$\mathcal{L}_G(S) = \{L \leq G \mid S \leq L, O_p(L) \neq 1, C_G(O_p(L)) \leq O_p(L)\}.$$

Under the assumption that  $S$  is contained in at least two maximal  $p$ -local subgroups, for  $L \in \mathcal{L}_G(S)$  with  $L \not\leq N_G(Q)$ , the Local Structure Theorem provides information about  $L/C_L(Y_L)$  and its action on  $Y_L$ . Given the Local Structure Theorem there are two cases to treat in order to fully understand groups with a large  $p$ -subgroup. Either there exists  $L \in \mathcal{L}_G(S)$  with  $Y_L \not\leq Q$  or, for all  $L \in \mathcal{L}_G(S)$ ,  $Y_L \leq Q$ . Research in the first case has just started and, for this situation, this paper addresses the wreath product scenario in the Local Structure Theorem [MSS2, Theorem A (3)]. This case is separated from the rest because of the special structure of  $L$  and  $Y_L$ . This structure allows us to use arguments measuring the size of certain subgroups to reduce to three exceptional configurations and has a distinct flavour from the remaining cases. For instance, the groups which are examples in the wreath product case typically have  $Q$  of class 3 whereas in the more typical cases it has class at most 2. The configurations in the Local Structure Theorem which are not in the wreath product case and have  $Y_L \not\leq Q$  will be examined in a separate publication as there are methods which apply uniformly to cover many possibilities at once. Contributions to the  $Y_L \leq Q$  for all  $L \in \mathcal{L}_G(S)$  are the subject of [PPS].

For  $L \in \mathcal{L}_G(S)$  with  $Q$  not normal in  $L$  we set

$$L^\circ = \langle Q^L \rangle, \bar{L} = L/C_L(Y_L) \text{ and } V_L = [Y_L, L^\circ]$$

and use this notation throughout the paper. Set  $q = p^a$ . We recall from [MSS2, Remark A.25] the definition of a *natural wreath  $\text{SL}_2(q)$ -module* for the group  $X$  with respect to  $\mathcal{K}$ : suppose that  $X$  is a group,  $V$  is a faithful  $X$ -module and  $\mathcal{K}$  is a non-empty  $X$ -invariant set of subgroups of  $X$ . Then  $V$  is a *natural  $\text{SL}_2(q)$ -wreath product module* for  $X$  with respect to  $\mathcal{K}$  if and only if

$$V = \bigoplus_{K \in \mathcal{K}} [V, K] \text{ and } \langle \mathcal{K} \rangle = \times_{K \in \mathcal{K}} K,$$

and, for each  $K \in \mathcal{K}$ ,  $K \cong \text{SL}_2(q)$  and  $[V, K]$  is the natural  $\text{SL}_2(q)$ -module for  $K$ .

We now describe the wreath product case in [MSS2, Theorem A (3)]. For  $L \in \mathcal{L}_G(S)$  with  $L \not\leq N_G(Q)$ ,  $L$  is in the *wreath product case* provided

- there exists a unique  $\bar{L}$ -invariant set  $\mathcal{K}$  of subgroups of  $\bar{L}$  such that  $V_L$  is a natural  $\mathrm{SL}_2(q)$ -wreath product module for  $\bar{L}$  with respect to  $\mathcal{K}$ .
- $\bar{L}^\circ = O^p(\langle \mathcal{K} \rangle \bar{Q})$  and  $Q$  acts transitively on  $\mathcal{K}$  by conjugation.
- $Y_L = V_L$  or  $p = 2$ ,  $|Y_L : V_L| = 2$ ,  $\bar{L}^\circ \cong \mathrm{SL}_2(4)$  or  $\Gamma\mathrm{SL}_2(4)$  and  $V_L \not\leq Q$ .

We say that  $\bar{L}$  is *properly wreathed* if  $|\mathcal{K}| > 1$ .

There are overlaps between the wreath product case and some other divisions in the Local Structure Theorem.

If  $\bar{L}^\circ \cong \mathrm{SL}_2(q)$  with  $V_L = Y_L$ , then this situation can be inserted in the linear case of [MSS2, Theorem A (1)] by including  $n = 2$  is that case. Suppose that  $|\mathcal{K}| = 2$  and  $K \cong \mathrm{SL}_2(2)$ . If  $\bar{Q}$  is a fours group, then, as  $\bar{Q}$  conjugates  $\bar{K}_1$  to  $\bar{K}_2$ ,

$$\bar{L}^\circ \cong \Omega_4^+(2) \cong \mathrm{SL}_2(2) \times \mathrm{SL}_2(2)$$

and  $Y_L$  is the tensor product module. This is an example in the tensor product case of [MSS2, Theorem A (6)]. We declare  $L$  to be in the *unambiguous wreath product case* if these two *ambiguous* configurations do not occur. The ambiguous cases will be handled in a more general setting in a forthcoming paper mentioned earlier.

**Main Theorem.** *Suppose that  $p$  is a prime,  $G$  is a finite group,  $S$  a Sylow  $p$ -subgroup of  $G$  and  $Q \leq S$  is a large  $p$ -subgroup of  $G$  with  $Q = O_p(N_G(Q))$ . If there exists  $L \in \mathcal{L}_G(S)$  with  $L$  in the unambiguous wreath product case and  $V_L \not\leq Q$ , then  $G \cong \mathrm{Mat}(22)$ ,  $\mathrm{Aut}(\mathrm{Mat}(22))$ ,  $\mathrm{Sym}(8)$ ,  $\mathrm{Sym}(9)$  or  $\mathrm{Alt}(10)$ .*

The proof of this theorem splits into four parts. First, in Section 3, we show that in the properly wreathed case we must have  $q = |\mathcal{K}| = 2$  and, as  $L$  is unambiguous,  $\bar{S} = \bar{Q} \cong \mathrm{Dih}(8)$  and  $\bar{L}^\circ \cong \mathrm{O}_4^+(2)$ . If  $|\mathcal{K}| = 1$ , we show that  $\bar{L}^\circ \cong \Gamma\mathrm{SL}_2(4)$  or  $\mathrm{SL}_2(4)$  and  $V_L$  is the natural module with  $|Y_L : V_L| \leq 2$ , where, if  $\bar{L}^\circ \cong \mathrm{SL}_2(4)$ ,  $|Y_L : V_L| = 2$  holds. In the following three sections, we determine the groups corresponding to these three cases. Finally the Main Theorem follows by combining Propositions 3.5, 4.1, 5.1 and 6.2.

In [PPS] the authors proved that the unambiguous wreath product case does not lead to examples if for all  $L \in \mathcal{L}_G(S)$  we have  $Y_L \leq Q$ , with the additional assumption that  $G$  is of local characteristic  $p$ . In this paper we do not make the assumption that  $G$  is of local characteristic  $p$ .

In the Local Structure Theorem there is also a possibility that  $L \in \mathcal{L}_G(S)$  is of weak wreath type. Any such group is contained in one, which is of unambiguous wreath type. A corollary of our theorem is

**Corollary.** *Suppose that  $p$  is a prime,  $G$  is a finite group,  $S$  a Sylow  $p$ -subgroup of  $G$  and  $Q \leq S$  is a large  $p$ -subgroup of  $G$  with  $Q = O_p(N_G(Q))$ . If  $L \in \mathcal{L}_G(S)$  is of weak wreath product type, then either  $G$  is as in the Main Theorem or  $V_L \leq Q$ .*

In addition to the notation already introduced, we will use the following

**Notation.** For  $p$  a prime,  $G$  a group with a large  $p$ -subgroup  $Q = O_p(N_G(Q))$  and  $L \in \mathcal{L}_G(S)$ , we set  $Q_L = O_p(L)$  and assume that  $V_L \not\leq Q$ . Define  $D = \langle V_L^{N_G(Q)} \rangle (L \cap N_G(Q)) \in \mathcal{L}_G(S)$ . Furthermore, set

$$W = \langle (V_L \cap Q)^D \rangle,$$

$$U_L = \langle (W \cap Q_L)^L \rangle$$

and

$$Z = C_{V_L}(Q).$$

Notice that for  $L_0 = N_L(S \cap C_L(Y_L))$ , we have  $L = C_L(Y_L)L_0$  and  $C_L(Y_L) \leq D$ . Further

$$Y_{L_0} = Y_L = \Omega_1(Z(O_p(L_0)))$$

by [MSS2, Lemma 1.2.4 (i)]. Since  $C_L(Y_L)$  normalizes  $Q$ ,

$$L^\circ = \langle Q^L \rangle = \langle Q^{C_L(Y_L)L_0} \rangle = \langle Q^{L_0} \rangle = L_0^\circ.$$

Therefore, if  $L$  is in the unambiguous wreath product case, then so is  $L_0$ . Hence we also assume that  $L = L_0$  and so

$$Y_L = \Omega_1(Z(Q_L)).$$

## 2. PRELIMINARIES

In this section we present some lemmas which will be used in the forthcoming sections.

**Lemma 2.1.** *Suppose that  $X$  is a group,  $E = O_2(X)$  is elementary abelian of order 16 and  $X/E \cong \text{Alt}(6)$  induces the non-trivial irreducible part of the 6-point permutation module on  $E$ . Then  $X$  splits over  $E$ .*

*Proof.* Choose  $R \leq X$  such that  $R/E \cong \text{Sym}(4)$  and  $Z(R) = 1$ . Let  $T \in \text{Syl}_3(R)$ . As  $T$  acts fixed-point freely on  $O_2(R)$ ,  $N_R(T) \cong \text{Sym}(3)$  and so there are involutions in  $X/E$ . Hence, as  $X/E$  has one conjugacy class of involutions, there are involutions in  $O_2(R) \setminus E$ . Therefore  $O_2(R)/Z(O_2(R))$  is elementary abelian of order 16. Now we consider  $O_2(R)$ . The fixed-point free action of  $T$  on  $O_2(R)/Z(O_2(R))$  implies there is partition of this group into five  $T$ -invariant subgroups of order 4. As  $T$  acts fixed-point freely on  $O_2(R)$  the preimages of all these four groups are abelian. As there are involutions in  $O_2(R) \setminus E$ , there is a  $T$ -invariant fours group  $F^* \leq O_2(R)/Z(O_2(R))$  with  $F^* \neq E/Z(O_2(R))$  and such that the preimage  $F$  of  $F^*$  is elementary abelian of order 16. Now the action of  $X$  on  $E$  shows that for any involution  $i \in R \setminus E$  all involutions in the coset  $Ei$  are conjugate to  $i$  by an element of  $E$ . Hence all involutions in  $O_2(R) \setminus E$  are in  $F$ . This shows that  $F$  is invariant under  $N_R(T)$ .

Again there is a partition of  $F$  into five groups of order four invariant under  $T$ . Let  $t$  be an involution in  $N_R(T)$ . Then  $|C_F(t)| = 4$ , where  $|C_{E \cap F}(t)| = 2$ . Hence there is some fours group  $F_1 \leq F$ ,  $F_1 \neq E \cap F$  and  $C_{F_1}(t) \neq 1$ . This shows that  $F_1$  is normalized by  $t$ . Then  $F_1 \langle t \rangle \cong \text{Dih}(8)$  is a complement to  $E$ . Using a result of Gaschütz [GLS2, Theorem 9.26],  $X$  splits over  $E$ .  $\square$

The next lemma is well-known.

**Lemma 2.2.** *Suppose that  $X \cong \text{Sym}(5)$ ,  $F_1$  and  $F_2$  are four groups of  $X$  with  $F_1 \leq \text{Alt}(5)$  and  $V$  is a non-trivial irreducible  $\text{GF}(2)X$ -module. Then*

- (i)  *$V$  is either the non-trivial irreducible part of the permutation module, which is the same as the natural  $O_4^-(2)$ -module, or  $V$  is the natural  $\Gamma\text{L}_2(4)$ -module.*
- (ii)  *$F_1$  acts quadratically on  $V$  if and only if  $V$  is the natural  $\Gamma\text{L}_2(4)$ -module.*
- (iii)  *$F_2$  acts quadratically on  $V$  if and only if  $V$  is the natural  $O_4^-(2)$ -module.*

**Lemma 2.3.** *Suppose that  $p$  is a prime,  $X$  is a group of characteristic  $p$  and  $U$  is a normal  $p$ -subgroup of  $X$ . Let  $R$  be a normal subgroup of  $X$  with  $R \leq C_X(U/[U, O_p(X)])$ . If  $[O_p(X), O^p(R)] \leq U$ , then  $R \leq O_p(X)$ .*

*Proof.* It suffices to prove that  $O^p(R) = 1$ . Suppose that  $n \geq 1$  is such that  $[U, O^p(R)] \leq [U, O_p(X); n]$ . Then

$$[O_p(X), O^p(R)] = [O_p(X), O^p(R), O^p(R)] \leq [U, O^p(R)] \leq [U, O_p(X); n]$$

and so

$$[O_p(X), O^p(R), U] \leq [[U, O_p(X); n], O_p(X)] = [U, O_p(X); n+1].$$

We also have

$$[U, O^p(R), O_p(X)] \leq [[U, O_p(X); n], O_p(X)] = [U, O_p(X); n+1]$$

and thus the Three Subgroups Lemma implies

$$[U, O_p(X), O^p(R)] \leq [U, O_p(X); n+1].$$

This yields

$$[U, O^p(R)] = [U, O^p(R), O^p(R)] \leq [U, O_p(X), O^p(R)] \leq [U, O_p(X); n+1].$$

Since  $O_p(X)$  is nilpotent, we deduce  $[U, O^p(R)] = 1$ . Hence

$$[O_p(X), O^p(R)] = [O_p(X), O^p(R), O^p(R)] \leq [U, O^p(R)] = 1.$$

As  $X$  has characteristic  $p$ ,  $O^p(R) = 1$  and so  $R \leq O_p(X)$  as claimed.  $\square$

**Lemma 2.4.** *Assume that  $X$  is a group,  $Y$  is a normal subgroup of  $X$  and  $xC_X(Y) \in Z(X/C_X(Y))$ . If  $[Y, x] \leq Z(Y)$ , then  $Y/C_Y(x) \cong [Y, x]$  as  $X$ -groups.*

*Proof.* Define

$$\begin{aligned} \theta : Y &\rightarrow [Y, x] \\ y &\mapsto [y, x]. \end{aligned}$$

Then  $\theta$  is independent of the choice of the coset representative in  $xC_X(Y)$ .

For  $y, z \in Y$ ,

$$(yz)\theta = [yz, x] = [y, x]^z[z, x] = [y, x][z, x] = (y)\theta(z)\theta,$$

and, for  $y \in Y$  and  $\ell \in X$ , as  $[x, \ell] \in C_R(Y)$ ,  $x^\ell = xc$  for some  $c \in C_X(Y)$ , and so

$$(y\theta)^\ell = [y, x]^\ell = [y^\ell, x^\ell] = [y^\ell, xc] = [y^\ell, c][y^\ell, x]^c = [y^\ell, x] = (y^\ell)\theta.$$

Thus  $\theta$  is an  $X$ -invariant homomorphism from  $Y$  to  $[Y, x]$ . As  $\ker \theta = C_Y(x)$ , we have  $Y/C_Y(x) \cong [Y, x]$  as  $X$ -groups.  $\square$

**Lemma 2.5.** *Assume that  $p$  is a prime,  $X$  is a group,  $Y$  is an abelian normal  $p$ -subgroup of  $X$  and  $R$  is a normal  $p$ -subgroup of  $X$  which contains  $Y$ . Suppose that  $Y = [Y, O^p(X)]$ ,  $[R, O^p(X)] \leq C_R(Y)$  and  $R$  acts quadratically or trivially on  $Y$ . Suppose that no non-central  $X$ -chief factor of  $Y/C_Y(R)$  is isomorphic to an  $X$ -chief factor of  $[Y, R]$ . Then  $Y \leq Z(R)$ .*

*Proof.* Assume that  $R > C_R(Y)$ . Using  $[R, O^p(X)] \leq C_R(Y)$ , we may select  $x \in R \setminus C_R(Y)$  such that  $xC_X(Y) \in Z(X/C_X(Y))^\#$ . As  $Y$  is abelian,  $[Y, x] \leq Z(Y)$  and so Lemma 2.4 applies to give  $Y/C_Y(x) \cong [Y, x]$  as  $X$ -groups. As  $R$  acts quadratically on  $Y$ ,

$$C_Y(x) \geq C_Y(R) \geq [Y, R] \geq [Y, x]$$

and so the hypothesis on non-central  $X$ -chief factors now gives  $Y/C_Y(x)$  and  $[Y, x]$  only have central  $X$ -chief factors. In particular,  $Y = [Y, O^p(X)] \leq C_Y(x)$  and this contradicts the initial choice of  $x \in R \setminus C_R(Y)$ . Hence  $Y \leq Z(R)$ .  $\square$

**Lemma 2.6.** *Suppose that  $p$  is a prime,  $X$  is a group,  $V \leq U$  are normal  $p$ -subgroups of  $X$ , and  $Q$  is a large  $p$ -subgroup of  $X$  which is not normal in  $X$ . Assume that  $V$  is a non-trivial irreducible  $\text{GF}(p)X$ -module and  $U/V$  is centralized by  $O^p(X)$ . Then*

- (i)  $U$  is elementary abelian; and
- (ii) if  $U \not\leq \Omega_1(Z(O_p(X)))$ , then  $O_p(X)/C_{O_p(X)}(U)$  contains a non-central chief factor isomorphic to  $V$  as a  $\text{GF}(p)X$ -module.

*Proof.* Set  $Z_X = \Omega_1(Z(O_p(X)))$ . We have  $[U, O^p(X)] \leq V \leq Z_X$  as  $V$  is irreducible. As  $O^p(X)$  does not centralize  $U/\Phi(U)$  by Burnside's Lemma [GLS2, Proposition 11.1] and  $V$  is a non-trivial irreducible  $X$ -module,  $V \not\leq \Phi(U)$  and  $\Phi(U)$  is centralized by  $O^p(X)$ . Therefore  $\Phi(U) \cap Z_X$  is centralized by  $O^p(X)$  and is normalized by  $Q$ . Since  $Q$  is large and  $O^p(X) \not\leq N_X(Q)$ , we deduce  $\Phi(U) \cap Z_X = 1$ . Thus  $\Phi(U) = 1$  and so  $U$  is elementary abelian. Hence (i) holds.

Set  $Y = O_p(X)$  and assume that  $U \not\leq Z_X$ . Select  $x \in U \setminus Z_X$  such that  $[X, x] \leq U \cap Z_X \leq Z(Y)$ . Then  $xC_X(Y) \in Z(X/C_X(Y))$ . Thus Lemma 2.4 implies  $Y/C_Y(x) \cong [Y, x] \leq U \cap Z_X$  and this isomorphism is as  $X$ -groups. Since  $[Y, x]$  is normalized by  $Q$ ,  $[Y, x] \neq 1$  and  $Q$  is large,  $O^p(X)$  does not centralize  $[Y, x]$ . Thus  $[Y, x] \geq V$  as  $[U, O^p(X)] \leq V$ . This proves (ii).  $\square$

**Lemma 2.7.** *Assume that  $p$  is a prime,  $X$  is a group,  $U$  is an elementary abelian normal subgroup of  $X$ ,  $U = [U, O^p(X)]$  and  $O_p(X)$  acts quadratically and non-trivially on  $U$ . Set  $R = O_p(X)$ ,  $W = R/C_R(U)$ , and  $Z = [U, R]$ . Then  $W$ ,  $U/Z$  and  $Z$  are  $X/R$ -modules and  $W$  is isomorphic to an  $X/R$ -submodule of  $\text{Hom}(U/Z, Z)$ . In particular, if  $Z$  is centralized by  $X$ , then the set of  $X$ -chief factors of  $W$  can be identified with a subset of the  $\text{GF}(p)$ -duals of the  $X$ -chief factors of  $U/Z$ .*

*Proof.* Since  $R$  acts quadratically on  $U$ ,  $W$  is elementary abelian. Furthermore,  $R$  centralizes  $W$ ,  $U/Z$  and  $Z$ . Hence all of these groups can be regarded

as  $\text{GF}(p)X/R$ -modules. For  $w \in R$ , define

$$\begin{aligned} \theta : R &\rightarrow \text{Hom}(U/Z, Z) \\ w &\mapsto \begin{array}{ccc} \theta_w : U/Z &\rightarrow & Z \\ uZ &\mapsto & [u, w] \end{array} \end{aligned}$$

The calculation in the proof of Lemma 2.4 shows that the commutator  $[u, w]$  defines a homomorphism from  $U$  to  $Z$  and, as  $w$  centralizes  $Z$ ,  $\theta_w$  is a well-defined homomorphism from  $U/Z$  to  $Z$ . Thus  $\theta$  is a well-defined map. Consider  $w_1, w_2 \in R$ ,  $uZ \in U/Z$  and  $\ell \in X$ . Then

$(uZ)\theta_{w_1w_2} = [u, w_1w_2] = [u, w_2]^{w_1}[u, w_1] = [u, w_1][u, w_2] = (uZ)\theta_{w_1}(uZ)\theta_{w_2}$  which means  $\theta_{w_1w_2} = \theta_{w_1}\theta_{w_2}$  and so  $\theta$  is a group homomorphism. We show that  $\theta$  is an  $X$ -module homomorphism. So let  $\ell \in X$ ,  $uZ \in U/Z$  and  $w \in R$ . Then  $(w^\ell)\theta = \theta_{w^\ell}$  and

$$(uZ)\theta_{w^\ell} = [u, w^\ell] = [u^{\ell^{-1}}, w]^\ell = (u)(\theta_w \cdot \ell).$$

Since  $\ker \theta = C_R(U)$ , this completes the proof of the main claim.

If  $Z$  is centralized by  $X$ , then

$$\text{Hom}(U/Z, Z) \cong (U/Z)^* \otimes Z = \bigoplus_{i=1}^n (U/Z)^*$$

where  $n$  is such that  $|Z| = p^n$ . This completes the proof of the lemma.  $\square$

**Lemma 2.8.** *Suppose that  $V$  is a  $p$ -group and  $X$  is a group which acts faithfully on  $V$  with  $O_p(X) = 1$ . Assume  $A \leq X$  is an elementary abelian  $p$ -subgroup of order at least  $p^2$  which has the property  $C_V(A) = C_V(a)$  for all  $a \in A^\#$ . If  $L$  is a non-trivial subgroup of  $X$  and  $L = [L, A]$ , then  $A$  acts faithfully on  $L$ .*

*In particular,  $A$  centralizes every  $p'$ -subgroup which it normalizes,  $[A, F(X)] = 1$ ,  $E(X) \neq 1$  and, if  $L$  is a component of  $X$  which is normalized but not centralized by  $A$ , then  $A$  acts faithfully on  $L$ .*

*Proof.* Suppose that  $L = [L, A]$  is a non-trivial subgroup of  $X$ . Assume that there is  $b \in A^\#$  with  $[L, b] = 1$ . Then  $L$  normalizes  $C_V(b)$  and so, as  $C_V(b) = C_V(A)$ ,  $L = [L, A]$  centralizes  $C_V(b)$ . Since  $L = [L, A]$ ,  $L = O^p(L)$  and the Thompson  $A \times B$ -Lemma implies  $[L, V] = 1$ , a contradiction. Hence  $A$  acts faithfully on  $L$ .

Let  $F$  be a  $p'$ -subgroup of  $X$  which is normalized by  $A$ . Then  $F = \langle C_F(a) \mid a \in A^\# \rangle$ . If  $A$  does not centralizes  $F$ , then there exists  $a \in A^\#$  such that  $1 \neq [C_F(a), A] = [C_F(a), A, A]$ . Hence, taking  $L = [C_F(a), A]$ , we have  $L = [L, A]$  and  $a \in C_A(L)$ , a contradiction. Hence  $[F, A] = 1$ . Now  $A$  centralizes  $F(X)$  and therefore  $E(X) \neq 1$ .

If  $L$  is a component of  $X$  which is normalized by  $A$ , then either  $[L, A] = L$  or  $[L, A] = 1$ . If  $[L, A] \neq 1$ , then we have  $A$  acts faithfully on  $L$ .  $\square$

**Lemma 2.9.** *Let  $X$  be a group,  $N$  a normal subgroup of  $G$  and  $T \in \text{Syl}_p(X)$ . Assume that  $X = NT$ ,  $C_T(N) = 1$ ,  $q = p^a$  and*

$$N = N_1 \times N_2 \cdots \times N_s,$$

*where  $N_i \cong \text{SL}_2(q)$  for  $1 \leq i \leq s$ . Then the  $p$ -rank of  $G$  is  $sa$ .*



*Proof.* Assume first that  $q = 2$ . Then  $T$  acts faithfully on  $O_3(N)$ . As the 2-rank of  $\mathrm{GL}_s(3)$  is  $s$ , we are done. Similarly, if  $q = 3$ , then  $T$  acts faithfully on  $O_2(N)/Z(N)$ , which is elementary abelian of order  $2^{2s}$  we are done as  $\mathrm{GL}_{2s}(2)$  has 3-rank  $s$ .

Thus we may assume that  $q > 3$ . In particular, the subgroups  $N_i$  are quasisimple and  $T$  permutes the set  $\{N_i \mid 1 \leq i \leq s\}$ .

Assume that  $p$  is odd. Let  $A$  be an elementary abelian subgroup in  $T$  of maximal rank and assume that  $A \not\leq N$ . Then by Thompson replacement [GLS2, Theorem 25.2] we may assume that  $A$  acts quadratically on  $T \cap N$ . This shows that  $A$  has to normalize each  $N_i$ . As non-trivial field automorphisms are not quadratic on  $T \cap N_i$ , we get that  $A$  centralizes  $T \cap N$  and so  $A \leq T \cap N$ , the assertion.

Assume that  $q = 2^a$  with  $a \geq 2$ . Let  $B = N_N(T \cap N)$ . We have that  $T$  normalizes  $B$  and  $T/(T \cap N)$  acts faithfully on  $B/(T \cap N)$ . Now the Thompson dihedral Lemma [GLS2, Lemma 24.1] says that for any elementary abelian subgroup  $A$  of  $T$  we have a  $B$ -conjugate  $A^g$  such that  $U = \langle A, A^g \rangle (T \cap N) / (T \cap N)$  is a direct product of  $r$  dihedral groups where  $2^r = |A/(A \cap N)| \leq 2^s$  and  $A(T \cap N)/(T \cap N)$  is a Sylow 2-subgroup of  $U$ . Set  $T_1 = [O_{2'}(U), T \cap N]$ . As  $U$  is generated by two conjugates of  $A$  we see that  $|T_1| = |C_{T_1}(A/A \cap N)|^2$ . This now shows that  $|A| \leq |T \cap N|$ , the assertion again. This proves the lemma.  $\square$

In the next two lemmas we use the notation presented in the introduction though we do not assume that  $L$  is unambiguous.

**Lemma 2.10.** *Suppose that  $L \in \mathcal{L}_G(S)$ ,  $L \not\leq N_G(Q)$  and  $V_L = [Y_L, L^\circ]$ . Then*

- (i)  $C_{Y_L}(L^\circ) = 1$ .
- (ii)  $\Omega_1(Z(S)) \leq V_L$ .
- (iii) *If  $V_L$  is an irreducible  $L$ -module,  $V_L \not\leq Q$  and  $\Omega_1(Z(Q_L)) < Q_L$ , then  $V_L \leq Q'_L \leq \Phi(Q_L)$ .*

*Proof.* As  $C_{Y_L}(L^\circ) \leq C_G(Q)$  is normalized by  $L$ , (i) is a consequence of  $Q$  being large.

By [MSS2, Lemma 1.24 (g)],  $\Omega_1(Z(S)) \leq Y_L$  now Gaschütz Theorem [GLS2, Theorem 9.26] and (i) give (ii).

Assume that  $N$  is a non-trivial normal  $p$ -subgroup of  $L$ . Then  $\Omega_1(Z(S)) \cap N \neq 1$ . Since  $V_L$  is irreducible as a  $L$ -module, (ii) gives  $V_L \leq N$ . In particular, as  $V_L \not\leq Q$ ,  $N \not\leq Q$ .

Suppose that  $Q_L$  is abelian. Then, as  $Q = O_p(N_G(Q))$  and  $[Q, Q_L, Q_L] \leq Q'_L = 1$ ,  $Q_L$  is quadratic on  $Q$ , and hence  $Q_L Q/Q$  is elementary abelian and so  $\Phi(Q_L) \leq Q$ . By the remark earlier taking  $N = \Phi(Q_L)$  we obtain  $\Phi(Q_L) = 1$ , contrary to  $\Omega_1(Z(Q_L)) < Q_L$ . Hence  $Q_L$  is non-abelian. Thus  $Q'_L \neq 1$  and so, as  $V_L$  is irreducible,  $V_L \leq Q'_L \leq \Phi(Q_L)$ . This proves (iii).  $\square$

**Lemma 2.11.** *Suppose that  $L \in \mathcal{L}_G(S)$ ,  $L \not\leq N_G(Q)$  and  $V_L = [Y_L, L^\circ]$ . Assume that  $Y_L = \Omega_1(Z(Q_L))$ ,  $m \in L$  and  $O^p(L)Q_L \leq KQ_L$ , where  $K = \langle W, W^m \rangle$ . Then  $O^p(L) \leq K$  and the following hold*

- (i)  $[O^p(L), Q_L] \leq [W, Q_L][W^m, Q_L] \leq (W \cap Q_L)(W^m \cap Q_L) = U_L$ .
- (ii) *If  $[W, W] \leq V_L$ , then  $W$  acts quadratically on the non-central chief factors of  $Q_L/V_L$ .*

Assume, in addition, that  $V_L$  is irreducible as a  $K$ -module,  $[V_L, W, W] \neq 1$ , and  $[W, W] \leq V_L$ . Then

- (iii)  $W \cap W^m \cap Q_L \leq Y_L$ ;
- (iv)  $U_L/Y_L$  is elementary abelian or trivial; and
- (v) either  $Q_L = Y_L$  or  $U'_L \geq V_L$ .

*Proof.* Since  $W$  and  $W^m$  are normalized by  $Q_L$ ,  $K = \langle W, W^m \rangle$  is normalized by  $Q_L K$  and so  $O^p(L) \leq K$ . Since  $W$ ,  $W^m$ ,  $[Q_L, W]$  and  $[Q_L, W^m]$  are normalized by  $Q_L$ , we have

$$[Q_L, O^p(L)] \leq [Q_L, \langle W, W^m \rangle] = [Q_L, W][Q_L, W^m] \leq (W \cap Q_L)(W^m \cap Q_L).$$

In particular,  $A = (W \cap Q_L)(W^m \cap Q_L)$  is normalized by  $O^p(L)$ . Since  $(W \cap Q_L)^L = (W \cap Q_L)^{SO^p(L)} = (W \cap Q_L)^{O^p(L)}$ , we have  $A = U_L$ . Thus (i) holds.

By the additional hypothesis,

$$[Q_L, W, W] \leq [W, W] \leq V_L$$

and so  $W$  acts quadratically on all the non-central  $L$ -chief factors in  $Q_L/V_L$ , which is (ii).

Notice that part (ii),  $V_L$  irreducible as a  $K$ -module and  $[V_L, W, W] \neq 1$  together imply that the non-central  $K$ -chief factors in  $Q_L/V_L$  are not isomorphic to  $V_L$ .

Set  $I = W \cap W^m \cap Q_L$ . Then  $I \leq W \cap W^m$  and so

$$[I, W] \leq [W, W] \leq V_L$$

and

$$[I, W^m] \leq [W^m, W^m] \leq V_L^m = V_L.$$

Hence  $IV_L/V_L$  is centralized by  $\langle W, W^m \rangle = K$ . As  $W$  acts quadratically on all the non-central chief factors of  $K$  in  $Q_L/V_L$  by (ii) and by assumption,  $W$  does not act quadratically on  $V_L$ , Lemma 2.6 implies that  $I \leq \Omega_1(Z(Q_L)) = Y_L$ . This proves (iii).

Since  $W$  is generated by elements of order  $p$ ,  $W/[W, W]$  is elementary abelian and therefore, as  $[W, W] \leq V_L$ ,  $WV_L/V_L$  is also elementary abelian. Since  $W \cap Q_L$  and  $Q_L \cap W^m$  normalize each other parts (i) and (iii) give (iv).

If  $V_L \not\leq U'_L$  and  $Q_L \neq Y_L$ , then, as  $U_L/Y_L$  is elementary abelian by (iv), Lemma 2.10 (ii) implies  $U_L$  is elementary abelian. Select  $E$  with  $Q_L \geq E > V_L$  of minimal order such that  $E = [E, O^p(L)]$  and  $E/V_L$  has a non-central  $K$ -chief factor. Then

$$E \leq [Q_L, O^p(L)] \leq [Q_L, W][Q_L, W^m] \leq U_L \leq C_L(E).$$

Furthermore,  $V_L[E, Q_L] < E$  and so  $[[E, Q_L], O^p(L)] \leq V_L$ . Therefore Lemma 2.6 implies that  $[E, Q_L] \leq Y_L$  and so  $Q_L$  acts quadratically on  $E$ . Hence Lemma 2.5 implies that  $E \leq Y_L$ , a contradiction. Hence  $U'_L$  is non-trivial and it follows that  $V_L \leq U'_L$ .  $\square$

### 3. THE REDUCTION

We use the notation presented in the introduction. For the rest of this article we have  $L \in \mathcal{L}_G(S)$  with  $Q$  not normal in  $L$  and  $L$  is in the unambiguous wreath product case. This means that  $Y_L = V_L$  unless we are in the special case that  $\overline{L}^\circ \cong \mathrm{SL}_2(4)$  or  $\Gamma\mathrm{SL}_2(4)$ ,  $|Y_L : V_L| = 2$  and

$$V_L \not\leq Q.$$

We start with a general result which just requires  $V_L \not\leq Q$ .

**Lemma 3.1.** *The following hold.*

- (i)  $\langle V_L^D \rangle$  is not a  $p$ -group;
- (ii)  $[Q, \langle V_L^D \rangle] \leq W$ ; and
- (iii)  $W \not\leq C_G(V_L)$ .

*Proof.* Let  $\tilde{C} = N_G(Q)$  and  $K = \langle V_L^{\tilde{C}} \rangle$ . As  $D = KN_L(Q)$  and  $N_L(Q)$  acts on  $V_L$  we have  $\langle V_L^D \rangle = \langle V_L^K \rangle$  is subnormal in  $H$ . If  $\langle V_L^D \rangle$  is a  $p$ -group, we obtain  $V_L \leq O_p(N_G(Q)) = Q$  which is a contradiction. This proves (i).

We have  $[Q, V_L] \leq Q \cap V_L \leq W$ . As  $W$  and  $Q$  are normalized by  $D$ , (ii) holds.

Assume  $W \leq C_G(V_L)$ . Then  $[W, V_L] = 1$  and so  $[W, \langle V_L^D \rangle] = 1$ . Hence  $X = O^p(\langle V_L^D \rangle)$  centralizes  $Q$  by (ii). Since  $C_G(Q) \leq Q$ , we have  $X \leq Q$ . Thus  $X = 1$  and  $\langle V_L^D \rangle$  is a  $p$ -group, which contradicts (i). Hence  $W \not\leq C_G(V_L)$ .  $\square$

We adopt the following notation. Let  $B \geq C_L(V_L)$  be such that  $\overline{B} = \langle \mathcal{K} \rangle$  and let  $S_0 = S \cap B$ . We write  $B = K_1 \dots K_s$  where  $K_i \geq C_L(V_L)$ ,  $\overline{K}_i \in \mathcal{K}$ ,  $\overline{K}_i \cong \mathrm{SL}_2(q)$  and, for  $1 \leq i \leq s$ , put

$$\begin{aligned} S_i &= S \cap K_i \\ V_L^i &= [V_L, K_i], \\ Z_i &= C_{V_L^i}(S_i) = C_{V_L^i}(S_0) \end{aligned}$$

and

$$Z_0 = Z_1 \dots Z_s = C_{V_L}(S_0).$$

We begin by showing that  $\overline{W}$  is not contained in the base group  $\overline{B}$ .

**Lemma 3.2.** *Suppose that  $\overline{L}$  is either properly wreathed, or  $q = p^a$  (where  $p$  divides  $a$ ) and some element of  $\overline{L}^\circ$  induces a non-trivial field automorphism on  $O^p(\overline{L}^\circ) \cong \mathrm{SL}_2(q)$ . Then  $W$  is not contained in  $S_0$ . In particular, if  $\overline{L}$  is properly wreathed with  $q = s = 2$ , then  $\overline{Q}$  is not cyclic of order 4.*

*Proof.* Set  $F = \bigcap_{g \in D} C_Q(V_L)^g$ .

Suppose that  $W$  is contained in  $S_0$ . As  $\overline{Q}$  normalizes  $\overline{W}$  and acts transitively on  $\mathcal{K}$  when  $\overline{L}$  is properly wreathed and, as  $V_L$  is the natural  $\mathrm{SL}_2(q)$ -module when  $s = 1$ , and field automorphisms are present, the structure of  $V_L$  yields

$$[V_L, S_0] = [V_L, W] = C_{V_L}(W) = Z_0.$$

Suppose that  $g \in D$ . Then using Lemma 3.1(ii) and  $(V_L)^g = V_{L^g}$  yields

$$(3.2.1) \quad [Z_0, [V_{L^g}, Q]] \leq [Z_0, W] = 1.$$

We also remark that as  $W \leq Q$ ,  $Z_0 \leq [V_L, Q] \leq W = W^g \leq S_0^g$  and  $Z_0 \leq Z(W)$ . In particular, as  $S_0^g$  normalizes every element of  $\mathcal{K}^g$ , so does  $Z_0$ . Therefore, for  $1 \leq i \leq s$ ,  $Z_0$  also normalizes each  $K_i^g$  and so also  $[Y_L^g, K_i^g] = (V_L^i)^g$ .

If  $s = 1$  and we have field automorphisms in  $\overline{L}^\circ$ , then  $[V_L, Q] > Z_0$  and so (3.2.1) provides  $Z_0 \leq C_Q([V_L, Q]) = C_Q(V_L)$ . Thus

$$[V_L, W] = Z_0 \leq F$$

in this case.

We will show that the same holds in the properly wreathed case. Because  $Q$  acts transitively on  $\mathcal{K}^g$ ,

$$V_{L^g} = V_{L^g}^1[V_{L^g}, Q] = V_{L^g}^2[V_{L^g}, Q].$$

As  $[Z_0, [V_{L^g}, Q]] = 1$  by (3.2.1),

$$\begin{aligned} [V_{L^g}, Z_0] &= [V_{L^g}^1[V_{L^g}, Q], Z_0] \cap [V_{L^g}^2[V_{L^g}, Q], Z_0] \\ &= [V_{L^g}^1, Z_0] \cap [V_{L^g}^2, Z_0] \leq V_{L^g}^1 \cap V_{L^g}^2 = 1. \end{aligned}$$

Hence  $Z_0 \leq C_Q(V_{L^g})$  and this implies that

$$[V_L, W] = Z_0 \leq F$$

in the properly wreathed case too. Therefore,

$$\begin{aligned} [Q, V_L] &\leq W \\ [W, V_L] &= Z_0 \leq F \cap W \\ [F \cap W, V_L] &= 1. \end{aligned}$$

Hence  $V_L$  stabilizes the normal series  $Q \geq W \geq W \cap F \geq 1$  in  $D$  and so  $V_L \leq O_p(D)$ . But then  $\langle V_L^D \rangle$  is a  $p$ -group contrary to Lemma 3.1 (i). We conclude that  $W \not\leq S_0$  as claimed.

If  $q = s = 2$  and  $\overline{Q}$  is cyclic of order four, then, as  $\overline{W}$  is generated by involutions,  $\overline{W} = \overline{Q} \cap \overline{S}_0$ , a contradiction. Thus  $\overline{Q}$  is not cyclic of order 4 in this case.  $\square$

We now reduce the properly wreathed case to one specific configuration which will be handled in detail in Section 4.

**Proposition 3.3.** *Assume that  $\overline{L}$  is properly wreathed and unambiguous. Then  $|\mathcal{K}| = 2$ ,  $q = 2$ , and  $\overline{W}$  permutes  $\mathcal{K}$  transitively by conjugation. Furthermore,  $\overline{Q} = \overline{S} \cong \text{Dih}(8)$ ,  $\overline{L}^\circ \cong \text{O}_4^+(2)$  and  $Y_L = V_L$  is the natural  $\text{O}_4^+(2)$ -module.*

*Proof.* Since  $Q$  permutes  $\mathcal{K}$  transitively by conjugation and  $S_0$  normalizes  $Q$ , we have

**(3.3.1)**

- (i)  $\overline{Q \cap S_0}$  contains  $[\overline{Q}, \overline{S_0}]$ ;
- (ii)  $|\overline{S_0} : \overline{Q \cap S_0}| \leq |\overline{S_0} : [\overline{Q}, \overline{S_0}]| \leq q$ ; and
- (iii)  $[\overline{Q}, \overline{S_0}]C_{\overline{L}}(\overline{K_i})/C_{\overline{L}}(\overline{K_i}) \in \text{Syl}_p(\overline{K_i}C_{\overline{L}}(\overline{K_i})/C_{\overline{L}}(\overline{K_i}))$ .  $\blacksquare$

As  $W = \langle V_{L^g} \cap Q \mid g \in D \rangle$ , Lemma 3.2 implies there exists  $g \in D$  such that  $V_{L^g} \cap Q \not\leq S_0$ . We fix this  $g$ .

**(3.3.2)** We have  $\overline{V_{L^g} \cap Q} \cap \overline{S_0} \neq 1$ .

Suppose that  $\overline{V_{L^g} \cap Q} \cap \overline{S_0} = 1$ . Then, as  $\overline{Q \cap S_0}$  and  $\overline{V_{L^g} \cap Q}$  normalize each other,  $\overline{V_{L^g} \cap Q}$  centralizes  $\overline{Q \cap S_0}$ . If  $\overline{V_{L^g} \cap Q}$  normalizes some  $\overline{K_i} \in \mathcal{K}$ , then, as  $\overline{Q}$  acts transitively on  $\mathcal{K}$  and normalizes  $\overline{V_{L^g} \cap Q}$ ,  $\overline{V_{L^g} \cap Q}$  normalizes every member of  $\mathcal{K}$ . As  $\overline{V_{L^g} \cap Q}$  centralizes  $\overline{[Q, S_0]}$ , (3.3.1) (iii) implies that

$$\overline{V_{L^g} \cap Q} \leq \overline{[Q, S_0]} C_L(\overline{K_i}).$$

Since  $Q$  acts transitively on  $\mathcal{K}$ , this is true for each  $\overline{K_i} \in \mathcal{K}$ . Thus

$$\overline{V_{L^g} \cap Q} \leq \bigcap_{i=1}^s \overline{[Q, S_0]} C_L(\overline{K_i}) = \bigcap_{i=1}^s \overline{S_i} C_L(\overline{K_i}) = \overline{S_0},$$

which contradicts the choice of  $g \in D$ .

Hence  $\overline{V_{L^g} \cap Q}$  does not normalize any member of  $\mathcal{K}$ . As  $\overline{B}$  is a direct product we calculate that  $C_{\overline{S_0}}(\overline{V_{L^g} \cap Q})$  has index at least  $q^{p-1}$  in  $\overline{S_0}$ . However (3.3.1) (ii) states that  $\overline{Q \cap S_0}$  has index at most  $q$  in  $\overline{S_0}$  and, as this subgroup is centralized by  $\overline{V_{L^g} \cap Q}$ , we deduce that

$$p = 2.$$

Furthermore, as  $\overline{V_{L^g} \cap Q}$  does not normalize any member of  $\mathcal{K}$ , if  $s > 2$ , we have  $C_{\overline{S_0}}(\overline{V_{L^g} \cap Q})$  has index at least  $q^2$  in  $\overline{S_0}$ , and so we must have

$$s = 2.$$

Since  $\overline{V_{L^g} \cap Q}$  centralizes  $\overline{[S_0, Q]}$  by (3.3.1)(iii), no element in  $\overline{V_{L^g} \cap Q}$  can act as a non-trivial field automorphism on  $\overline{K_1}$  and so we infer from  $\overline{V_{L^g} \cap Q} \cap \overline{S_0} = 1$ , that  $|\overline{V_{L^g} \cap Q}| = 2$ . In particular,  $|C_{V_L}(V_{L^g} \cap Q)| = q^2$  as  $V_{L^g} \cap Q$  exchanges  $V_L^1$  and  $V_L^2$ .

We know that  $|V_{L^g}| = q^4$ . As  $|[V_{L^g}, Q]| \geq q^3$ , we have

$$|V_{L^g} : V_{L^g} \cap Q| \leq q,$$

and we have just determined that

$$|V_{L^g} \cap Q : V_{L^g} \cap Q \cap C_G(V_L)| = |\overline{V_{L^g} \cap Q}| = 2.$$

Hence  $V_{L^g} \cap Q \cap C_G(V_L)$  has order at least  $2^{3a-1}$ , where  $q = 2^a$ .

Assume that  $a \neq 1$ . Then, as  $V_{L^g}^1$  has order  $q^2$ ,

$$V_{L^g} \cap Q \cap C_G(V_L) \cap V_{L^g}^1 \neq 1.$$

It follows that  $V_L \cap Q$  normalizes both  $K_1^g$  and  $K_2^g$ . As  $(V_L \cap Q)C_{L^g}(V_{L^g})/C_{L^g}(V_{L^g})$  is normalized by  $Q$  and  $Q$  permutes  $\{K_1^g, K_2^g\}$  transitively,  $(V_L \cap Q)C_{L^g}(V_{L^g})/C_{L^g}(V_{L^g})$  does not centralize  $K_i^g/C_{L^g}(V_{L^g})$  for  $i = 1, 2$ . Thus  $|C_{V_{L^g}^i}(V_L \cap Q)| \leq q$  for  $i = 1, 2$ . But then

$$2^{3a-1} \leq |V_{L^g} \cap Q \cap C_G(V_L)| \leq |C_{V_{L^g}^i}(V_L \cap Q)| \leq 2^{2a},$$

which contradicts  $a \neq 1$ . We conclude that  $q = s = 2$  and  $|\overline{V_{L^g} \cap Q}| = 2$ . Furthermore,  $\overline{V_{L^g} \cap Q}$  is centralized by  $\overline{Q}$  and so  $\overline{Q}$  is elementary abelian of order 4. It follows that  $\overline{L^\circ} \cong \Omega_4^+(2)$  and  $V_L$  is the natural module. Hence  $L$

is ambiguous and we conclude that  $\overline{V_{L^g} \cap Q} \cap \overline{S_0} \neq 1$ . ■

**(3.3.3)** We have  $|C_{V_L}(V_{L^g} \cap Q)| \leq q^{s/p}$ .

We know  $\overline{V_{L^g} \cap Q} \not\leq \overline{S_0}$  and  $\overline{V_{L^g} \cap Q} \cap \overline{S_0} \neq 1$  by (3.3.2). As  $\overline{V_{L^g} \cap Q}$  is normalized by  $\overline{Q}$ ,  $\overline{V_{L^g} \cap Q} \cap \overline{S_0} \neq 1$  implies that

$$C_{V_L}(\overline{V_{L^g} \cap Q}) = C_{Z_0}(\overline{V_{L^g} \cap Q}).$$

If some element  $d \in V_{L^g} \cap Q$  induces a non-trivial field automorphism on  $\overline{K_i}$  for some  $\overline{K_i} \in \mathcal{K}$ , then  $C_{V_L^i}(V_{L^g} \cap Q) \leq C_{Z_i}(d)$  has order  $q^{1/p}$  and the result follows by transitivity of  $\overline{Q}$  on  $\mathcal{K}$ . On the other hand, if  $d \in V_{L^g} \cap Q$  has an orbit of length  $p$  on  $\mathcal{K}$ , then  $C_{\langle (V_L^1)^{(d)} \rangle}(V_{L^g} \cap Q) \leq C_{\langle Z_1^{(d)} \rangle}(d)$  which has order  $q$ . Using the transitivity of  $Q$  on  $\mathcal{K}$ , we deduce  $|C_{V_L}(V_{L^g} \cap Q)| \leq q^{s/p}$ . This proves the result. ■

As  $Q$  acts transitively on the  $\{V_i \mid 1 \leq i \leq s\}$ , we have  $V_L = [V_L, Q]V_1$ . By (3.3.2)  $\overline{Q} \cap \overline{S_0} \neq 1$  and so  $|[V_1, Q]| \geq q$ . In particular

$$|V_L : [V_L, Q]| \leq q.$$

Since  $V_L \cap Q \cap C_{L^g}(V_{L^g}) \leq C_{V_L}(V_{L^g} \cap Q)$ , (3.3.3) and  $|V_L| = q^{2s}$  together give

$$|(V_L \cap Q)C_{L^g}(V_{L^g})/C_{L^g}(V_{L^g})| \geq q^{2s-1-s/p}.$$

On the other hand, by Lemma 2.9 the  $p$ -rank of  $\overline{L}$  is  $as$  where  $q = p^a$ . Hence

$$s \geq 2s - 1 - s/p$$

and so

$$s = p = 2.$$

In particular, Lemma 2.9 implies

**(3.3.4)**  $|(V_L \cap Q)C_{L^g}(V_{L^g})/C_{L^g}(V_{L^g})| = q^2 = 2^{2a}$ .

Assume that  $q > 2$ . Since  $S^g/S_0^g$  has 2-rank 2 and  $V_L \cap Q$  is elementary abelian,  $(V_L \cap Q \cap S_0^g)C_{L^g}(V_{L^g})/C_{L^g}(V_{L^g})$  has rank at least  $2a - 2 \neq 1$ . Since  $V_L \cap Q \cap S_0^g$  is normalized by  $Q$  and  $Q$  permutes  $\{K_1^g, K_2^g\}$  transitively,  $V_L \cap Q \cap S_0^g$  contains an element which projects non-trivially on to both  $S_1^g C_{L^g}(V_{L^g})/C_{L^g}(V_{L^g})$  and  $S_2^g C_{L^g}(V_{L^g})/C_{L^g}(V_{L^g})$ . Thus  $V_L \geq [V_L \cap Q, [V_{L^g}, Q]] \geq Z_0^g$ . But then, using (3.3.3) yields the contradiction

$$q^2 = |Z_0^g| \leq |C_{V_L}(V_{L^g} \cap Q)| \leq q.$$

Thus  $q = s = 2$ . It follows from Lemma 3.2 that  $W$  is transitive on  $\mathcal{K}$  and  $\overline{Q} \cong \text{Dih}(8)$  or  $\overline{Q}$  is elementary abelian of order 4. The second possibility gives  $\overline{L}^\circ \cong \Omega_4^+(2)$ , which is ambiguous. This proves Proposition 3.3. □

Next we deal with the case  $s = 1$ .

**Proposition 3.4.** *Suppose that  $OP(\overline{L}^\circ) \cong \text{SL}_2(q)$  where  $q = p^a = r^p$ ,  $V_L = Y_L$  is the natural  $OP(\overline{L}^\circ)$ -module and that some element of  $\overline{L}^\circ$  induces a non-trivial field automorphism on  $OP(\overline{L}^\circ)$ . Then  $p = 2 = r$ .*

*Proof.* We may assume that  $r^p > 4$ . By Lemma 3.2 we have that  $W \not\leq S_0$  and, as  $W$  is generated by elements of order  $p$ , we have that  $|S_0W : S_0| = p$ . As  $Q$  is normal in  $S$ ,  $1 \neq \overline{Q} \cap \overline{S_0}$ , so  $Z_0 \leq Q \cap Y_L$ . Furthermore, as  $\overline{Q}$  contains elements which act as field automorphisms on  $O^p(\overline{L}^\circ)$ ,

$$|V_L \cap Q : Z_0| \geq |[V_L, Q] : Z_0| \geq r^{p-1} > p,$$

by assumption. Thus no element in  $S \setminus Q_L$  centralizes a subgroup of index  $p$  in  $V_L \cap Q$ .

Set  $W_1 = \langle Z_0^D \rangle$ . As  $Z_0$  centralizes  $W \cap S_0$ , every element of  $Z_0$  centralizes a subgroup of index at most  $p$  in  $W$ . As  $W_1$  is generated by conjugates of  $Z_0$ , and these conjugates all contain elements which centralize a subgroup of index at most  $p$  in  $W$ ,  $W_1$  is generated by elements which centralize a subgroup of index at most  $p$  in  $V_L \cap Q$ . As no element in  $S \setminus Q_L$  has this property, we conclude that  $W_1 \leq Q_L$ . Hence  $[V_L, W_1] = 1$ . In particular  $[V_L \cap Q, W_1] = 1$  and so also  $[W, Z_0] = [W, W_1] = 1$ . This shows  $W \leq S_0$  and contradicts Lemma 3.2.  $\square$

We collect the results of this section in the following proposition:

**Proposition 3.5.** *Suppose that  $L \in \mathcal{L}_G(S)$ ,  $L \not\leq N_G(Q)$ ,  $V_L \not\leq Q$  and  $L$  is in the unambiguous wreath product case. Then one of the following holds:*

- (i)  $\overline{L}^\circ \cong O_4^+(2)$ ,  $\overline{Q} = \overline{S} \cong \text{Dih}(8)$  and  $Y_L = V_L$  is the natural module.
- (ii)  $\overline{L}^\circ \cong \Gamma\text{SL}_2(4)$ ,  $V_L$  is the natural  $\text{SL}_2(4)$ -module and  $|Y_L : V_L| \leq 2$ .
- (iii)  $\overline{L}^\circ \cong \text{SL}_2(4)$ ,  $V_L$  is the natural module and  $|Y_L : V_L| = 2$ .

*Proof.* If  $|\mathcal{K}| > 1$ , then (i) holds by Proposition 3.3, so we may assume that  $|\mathcal{K}| = 1$ . As  $L$  is unambiguous, either  $Y_L \neq V_L$  or  $\overline{L}^\circ \not\cong \text{SL}_2(q)$ . If  $Y_L \neq V_L$ , then by definition of the wreath product case, (ii) or (iii) holds. So we may assume  $Y_L = V_L$  and  $\overline{L}^\circ \not\cong \text{SL}_2(q)$ . Now (ii) holds by Proposition 3.4.  $\square$

#### 4. $\overline{L}^\circ \cong O_4^+(2)$

In this section we analyse the configuration from Proposition 3.5(i). We prove

**Proposition 4.1.** *Suppose that  $L \in \mathcal{L}_G(S)$ ,  $L \not\leq N_G(Q)$  and  $L$  is in the unambiguous wreath product case. If  $Y_L \not\leq Q$  and  $\overline{L}^\circ \cong O_4^+(2)$ , then  $G \cong \text{Sym}(8)$ ,  $\text{Sym}(9)$  or  $\text{Alt}(10)$ .*

*Proof.* By Proposition 3.5 we have  $\overline{Q} \cong \text{Dih}(8)$ . Since  $Y_L$  is the natural  $O_4^+(2)$ -module for  $L/C_L(Y_L)$  and  $V_L$  is also the wreath product module for  $L/C_L(Y_L)$  with respect to  $\{\overline{K}_1, \overline{K}_2\}$ , we have the following well known facts.

##### (4.1.1)

- (i)  $|[Y_L, Q]| = 2^3$ ,  $|[Y_L, Q, Q]| = 2^2$  and  $C_{Y_L}(Q) = [Y_L, Q, Q, Q]$  has order 2.
- (ii)  $[Y_L, S_0] = C_{Y_L}(S_0)$  has order  $2^2$ ;
- (iii)  $|[Y_L, Q']| = 2^2$ ;
- (iv)  $C_L([Y_L, Q]) \leq C_L(Y_L)$ .

Our first aim is to prove

**(4.1.2)**  $\overline{W}$  is elementary abelian of order  $2^2$ ,  $[Y_L, W] = [Y_L, Q] = Y_L \cap Q$  and  $[Y_L, W, W] = C_{Y_L}(W) = C_{Y_L}(Q) = Z$ .

Applying Lemma 3.1, we consider  $x \in D$  such that  $Y_{L^x} \cap Q \not\leq C_L(Y_L)$ . Then  $Y_{L^x} \cap Q$  is normalized by  $Q$  and so

$$\overline{Y_{L^x} \cap Q} \text{ contains a 2-central involution in } \overline{Q}.$$

In particular, (4.1.1)(iii) gives

$$|[Y_L, Y_{L^x} \cap Q]| \geq 2^2.$$

As  $Y_L$  is elementary abelian,  $\overline{Y_{L^x} \cap Q}$  is elementary abelian.

Suppose that  $[Y_L, Y_{L^x} \cap Q, Y_{L^x} \cap Q] = 1$ . Then

$$[Y_L, Y_{L^x} \cap Q] \leq C_{S^x}([Y_{L^x}, Q]) = Q_{L^x}$$

by (4.1.1) (iv). Hence  $[Y_L, Y_{L^x} \cap Q, Y_{L^x}] = 1$ . Then as  $|[Y_L, Y_{L^x} \cap Q]| = 2^2$  and  $|Y_L \cap Q| = 2^3$ , we conclude that  $(Y_L \cap Q)C_{L^x}(Y_{L^x})/C_{L^x}(Y_{L^x})$  has order 2. Thus  $[Y_{L^x}, Y_L \cap Q, Y_L \cap Q] = 1$ . Now the argument just presented implies that  $|\overline{Y_{L^x} \cap Q}| = 2$  and so, as  $Q$  normalizes  $Y_{L^x} \cap Q$ ,  $\overline{Y_{L^x} \cap Q} = Z(\overline{Q})$ . In particular, as  $[Y_L, S_0, S_0] = 1$ , we have proved that

$$\text{if } \overline{Y_{L^x} \cap Q} \leq \overline{S_0}, \text{ then } \overline{Y_{L^x} \cap Q} = Z(\overline{Q}).$$

For a moment let  $\overline{Q_1}$  be the four subgroups of  $\overline{Q}$  not equal to  $\overline{S_0}$ . Then as  $\Phi(Y_{L^x} \cap Q) = 1$  the displayed line implies that  $\overline{W} \leq \overline{Q_1}$  and Lemma 3.2 and  $\overline{Q}' \leq \overline{Y_{L^x} \cap Q}$  imply  $\overline{W} = \overline{Q_1}$ . The remaining statements in (4.1.2) now follow from the action of  $L$  on  $Y_L$ .  $\blacksquare$

We have that  $Z(Q)$  centralizes  $[Y_L, Q]$  and so  $Z(Q) \leq S \cap C_L(Y_L) = Q_L$ . Hence using (4.1.2) we obtain

$$\begin{aligned} [W, W] &= [\langle [Y_L, Q]^D \rangle, W] = \langle [[Y_L, Q], W]^D \rangle \\ &= \langle Z^D \rangle = Z[Z, \langle V_L^{N_G(Q)} \rangle] \leq Z[Z(Q), \langle V_L^{N_G(Q)} \rangle] \\ &= Z[\langle [Z(Q), V_L]^{N_G(Q)} \rangle] = Z. \end{aligned}$$

**(4.1.3)** We have  $Q_L = Y_L$ .

Suppose that  $Q_L > Y_L$ . Let  $m \in L$  be such that  $\overline{K} \cong \text{SL}_2(2) \times \text{SL}_2(2)$ , where  $K = \langle W, W^m \rangle$ . Recall that by the choice of  $L$  in the Notation at the end of the introduction, we have  $Y_L = \Omega_1(Z(Q_L))$  and by Proposition 3.5 and (4.1.2),  $K$  acts irreducibly on  $Y_L = V_L$ . Hence we may apply Lemma 2.11 (iii), (iv) and (v) which combined yield  $U_L/Y_L$  is elementary abelian and

$$U'_L = Y_L.$$

Since  $[Q_L, W, W] \leq [W, W] = Z \leq Y_L$ , we have  $W$  acts quadratically on every chief factor of  $L$  in  $Q_L/Y_L$ . In particular, no non-central  $L$ -chief factor of  $Q_L/Y_L$  is isomorphic to  $Y_L$ .

Let  $E$  be the preimage of  $C_{U_L/Y_L}(K)$ . Then  $E$  is normal in  $L$  and application of Lemma 2.6 implies that  $E = Y_L$ . Let  $X \in \text{Syl}_3(K)$ . By Lemma 2.11(i),  $[K, C_L(Y_L)] \leq U_L$ , so  $XU_L$  is normal in  $L$ . As  $L$  is solvable,  $C_L(Y_L) = C_X(Y_L)Q_L$  and either  $C_X(Y_L) = 1$  or  $X \cong 3_+^{1+2}$ . The latter case is impossible as  $W$  is quadratic on  $U_L/Y_L$ . Hence  $U_L = [U_L, O^2(L)]$  and  $U_L/Y_L$  contains no central  $L$ -chief factors. We know that every  $L$ -chief factor in



$U_L/Y_L$  is a wreath product module for  $\mathrm{SL}_2(2) \wr 2$  with  $\overline{W}$  acting quadratically. In particular, for every non-central chief factor  $F$  of  $L$  in  $U_L/Y_L$  we have  $[F, \overline{W}] = [F, Z(\overline{Q})]$ . Set  $W_1 = [W, D]$ . Then

$$\overline{W}_1 \geq [\overline{W}, \overline{Q}] = Z(\overline{Q}).$$

Hence  $[F, W] = [F, W_1]$  for every non-central chief factor  $F$  of  $L$  in  $U_L/Y_L$ . Set  $\tilde{L} = L/Y_L$  and let  $z \in Q$  with  $Z(\overline{Q}) = \langle \bar{z} \rangle$ . As  $C_F(Z(\overline{Q})) = [F, Z(\overline{Q})]$  for each  $F$ , we have  $C_{\tilde{U}_L}(z) = [\tilde{U}_L, z]$ ; then as  $W$  acts quadratically on  $\tilde{U}_L$ , we have  $[W, \tilde{U}_L] = C_{\tilde{U}_L}(W)$ . Thus  $[U_L, W]Y_L = [U_L, W_1]Y_L$ . In particular,

$[W/W_1, U_L] = [U_L, W]W_1/W_1 = (Y_L \cap Q)[U_L, W_1]W_1/W_1 = (Y_L \cap Q)W_1/W_1$  and so  $U_L$  acts quadratically on  $W/W_1$ . Therefore  $U_L C_D(W/W_1)/C_D(W/W_1)$  is elementary abelian. Hence

$$Y_L = U'_L \leq C_D(W/W_1).$$

Set  $R = \langle Y_L^D \rangle$ . Then, as  $Y_L \not\leq O_2(D)$  by Lemma 3.1 (i),  $Y_L \cap O_2(D) = Y_L \cap Q \leq W$  and so  $R$  centralizes  $O_2(D)/W$  and  $W/W_1$ . Lemma 2.3 yields  $Y_L \leq O_2(D)$  and this contradicts Lemma 3.1 (i). We have shown  $Q_L = Y_L$ . ■

**(4.1.4)**  $|S| = 2^7$  and  $N_G(Q)/Q \cong \mathrm{Sym}(3)$ .

Since  $Q_L = Y_L = V_L$  and  $\overline{Q} \cong \mathrm{Dih}(8)$ ,  $|S| = 2^7$  and  $|Q| = 2^6$ . Then  $N_G(Q) = SX$ , where  $X$  is a Hall  $2'$ -subgroup of  $N_G(Q)$  and  $QX$  is normal in  $N_G(Q)$ . Furthermore  $W$  is extraspecial of order  $2^5$ . As  $W/Z = J(Q/Z)$ , we have  $W$  is normal in  $N_G(Q)$ . Hence  $X$  acts faithfully on  $W$  and embeds in  $O_4^+(2)$ . As  $[\overline{W}, \overline{Q}] = Z(\overline{Q})$ ,  $S/W$  is faithful on  $W/Z$ , so  $N_G(Q)/W$  embeds into  $O_4^+(2)$ . Because  $O_4^+(2) \cong \mathrm{Sym}(3) \wr 2$ , and  $O_2(N_G(Q)/W) \neq 1$ , we get the claim. ■

Taking  $T \in \mathrm{Syl}_3(L)$ , we have  $N_L(T)$  is a complement to  $Q_L$  and so  $L = Q_L N_L(T)$  is a split extension of  $Q_L$  by  $O_4^+(2)$ . In particular, the isomorphism type of  $S$  is uniquely determined. As  $\mathrm{Sym}(8)$  has a subgroup isomorphic to  $L$  and  $\mathrm{Sym}(8)$  has odd index in  $\mathrm{Alt}(10)$ , we have  $S$  is isomorphic to a Sylow 2-subgroup of  $\mathrm{Alt}(10)$ .

Let  $z \in C_{Y_L}(Q)^\#$ , then as  $Y_L$  is a  $+$ -type space for  $L$ , there is a fours group  $A$  of  $Y_L$  which has all non-trivial elements  $L$ -conjugate to  $z$ . Since  $C_G(z)$  has characteristic 2,  $C_{O(G)}(z) = 1$  and so by coprime action

$$O(G) = \langle C_{O(G)}(a) \mid a \in A^\# \rangle = 1.$$

Assume that  $G$  has no subgroup of index two. Then  $S$  is isomorphic to a Sylow 2-subgroup of  $\mathrm{Alt}(10)$ . Therefore [Mas, Theorem 3.15] implies that  $F^*(G) \cong \mathrm{Alt}(10)$ ,  $\mathrm{Alt}(11)$ ,  $\mathrm{PSL}_4(r)$ ,  $r \equiv 3 \pmod{4}$ , or  $\mathrm{PSU}_4(r)$ ,  $r \equiv 1 \pmod{4}$ . Notice that  $Z(Q) = C_{Y_L}(Q) = \langle z \rangle$  and so  $C_G(z) = N_G(Q)$  has characteristic 2. In  $\mathrm{Alt}(11)$ ,  $z$  corresponds to  $(12)(34)(56)(78)$  and so  $C_G(z) \leq (\mathrm{Alt}(8) \times Z_3) : 2$ , which implies that  $C_G(z)$  is not of characteristic 2. In the linear and unitary groups  $C_G(z)$  has a normal subgroup isomorphic to  $\mathrm{SL}_2(r) \circ \mathrm{SL}_2(r)$ , and this contradicts (4.1.4). Hence  $G \cong \mathrm{Alt}(10)$ .

Assume now that  $G$  has a subgroup of index two. As  $V_L \leq G'$  we also have  $W \leq G'$ . Therefore  $(G' \cap L)/Y_L \cong \Omega_4^+(2)$  and so  $G'$  has Sylow 2-subgroups isomorphic to those of  $\text{Alt}(8)$ . Applying [GH, Corollary A\*] we have  $F^*(G) \cong \text{Alt}(8)$ ,  $\text{Alt}(9)$  or  $\text{PSP}_4(3)$ . Again in  $G' \cong \text{PSP}_4(3)$ , we have that  $G'$  contains a subgroup of shape  $\text{SL}_2(3) \circ \text{SL}_2(3)$ . This contradicts (4.1.4) and proves the proposition.  $\square$

### 5. $\overline{L}^\circ \cong \Gamma\text{SL}_2(4)$

In this section we attend to the case from Proposition 3.5(ii). Hence we have  $p = 2$ ,  $\overline{L}^\circ \cong \Gamma\text{SL}_2(4)$ ,  $V_L$  is the natural  $\text{SL}_2(4)$ -module and either  $Y_L = V_L$  or  $|Y_L/V_L| = 2$ . Notice that as  $L \not\leq N_G(Q)$  and  $L$  centralizes  $Y_L/V_L$ , if  $Y_L > V_L$ ,  $Y_L$  does not split over  $V_L$  and  $C_{Y_L}(Q) = C_{V_L}(Q)$  has order 2. Furthermore,  $C_S([Y_L, Q]) = Q_L$ .

Our aim is to prove

**Proposition 5.1.** *Suppose  $L \in \mathcal{L}_G(S)$  and  $L \not\leq N_G(Q)$  with  $\overline{L}$  in the unambiguous wreath product case. If  $Y_L \not\leq Q$  and  $\overline{L}^\circ \cong \Gamma\text{SL}_2(4)$ , then  $G \cong \text{Mat}(22)$  or  $\text{Aut}(\text{Mat}(22))$ .*

Notice that as  $Q_L \in \text{Syl}_2(C_L(Y_L))$ ,  $C_L(Y_L)/Q_L$  is centralized by  $L^\circ$ , and so  $C_{L^\circ}(Y_L) = Q_L \cap L^\circ$  as the Schur multiplier of  $\text{SL}_2(4)$  has order 2. We also have  $|\overline{Q}| \geq 4$  and  $|Z(Q) \cap V_L| = 2$ .

**Lemma 5.2.** *For  $N = N_G(Q_L)$  we have  $(Z(Q) \cap V_L)^N \cap Y_L \subseteq V_L$ . In particular,  $N$  normalizes  $V_L$ .*

*Proof.* If  $V_L = Y_L$ , there is nothing to prove. Assume that  $|Y_L : V_L| = 2$ . Choose  $g \in N$ , put  $U = (Z(Q) \cap V_L)^g$  and assume that  $U \not\leq V_L$ . Recall that  $Y_L = \Omega_1(Z(Q_L))$  and so  $U \leq Y_L$  and  $Y_L$  is normalized by  $N$ . Then  $C_L(U)C_N(Y_L)/C_N(Y_L) \cong 5 : 4$  or  $2 \times \text{Sym}(3)$ . As  $C_N(U^{g^{-1}})$  normalizes  $Q \cap Y_L$ ,  $C_N(U^{g^{-1}})$  is not irreducible on  $Y_L/U^{g^{-1}}$ . This excludes the possibility  $C_L(U)C_N(Y_L)/C_N(Y_L) \cong 5 : 4$  which is irreducible on  $Y_L/U$ . Hence we see that  $Z(Q) \cap V_L$  has exactly  $15 + 10 = 25$  conjugates under  $N$ , but 25 does not divide the order of  $\text{SL}_5(2) = \text{Aut}(Y_L)$ . This contradiction proves the lemma.  $\square$

**Lemma 5.3.** *We have  $Q_L = Y_L$  and either*

- (i)  $|S| = 2^7$ ,  $L/Q_L \cong \Gamma\text{SL}_2(4)$ ,  $N_G(Q)/Q \cong \text{SL}_2(2)$ , there exists a subgroup  $E \leq S$  of order  $2^4$  which is normalized by  $N_G(Q)$  such that  $N_G(E)/E \cong \text{Alt}(6)$  and  $N_L(E)$  has index 5 in  $L$ . Furthermore all the involutions in  $\langle N_G(E), L \rangle$  are conjugate.
- (ii)  $G$  has a subgroup of index 2 which satisfies the conditions in (i).

*Proof.* We have  $\overline{S} \cong \text{Dih}(8)$  and  $\overline{Q} \not\leq \overline{S}_0$  as  $\overline{L}^\circ \cong \Gamma\text{SL}_2(4)$ . Lemma 3.2 implies that  $\overline{W} \not\leq \overline{S}_0$ . By assumption, we either have  $Y_L = V_L$  or  $|Y_L : V_L| = 2$ . In particular,  $2^4 \leq |Y_L| \leq 2^5$ . Since  $\overline{Q}$  is normal in  $\overline{S}$  and contains  $\overline{W}$  we know

**(5.3.1)** Either  $\overline{Q}$  is elementary abelian of order 4 or  $\overline{Q} = \overline{S}$

As  $V_L$  is a natural  $\text{SL}_2(4)$ -module and  $L \not\leq N_G(Q)$ , we have  $C_{Y_L}(Q) = C_{Y_L}(S)$  has order 2 and  $[Y_L, Q] = [V_L, Q]$  has order 8. Furthermore, as  $W$

is normal in  $S$  and is not contained in  $S_0$ , we have  $[Y_L, Q, W] = Z$  where  $Z = C_{V_L}(S)$  has order 2. Thus, arguing exactly as before (4.1.3) and in the proof of (4.1.2) we obtain

**(5.3.2)**  $|\overline{W}| = 4$ ,  $[W, W] = Z$  and  $[Q_L, W, W] \leq Y_L$ .

**(5.3.3)** Assume that  $Q_L > Y_L$ . Then  $[Q_L, O^2(L)] \not\leq Y_L$ .

Suppose that  $[Q_L, O^2(L)] \leq Y_L$ . Then  $V_L \not\leq \Phi(Q_L)$  by Burnside's Lemma [GLS2, Proposition 11.1], which contradicts Lemma 2.10(iii). This proves the claim  $\blacksquare$

**(5.3.4)** If  $V_L < Y_L$ , then  $\overline{Q} = \overline{S}$ .

If  $\overline{Q}$  has order 4, then  $\overline{Q} = \overline{W}$  by (5.3.2), so  $\overline{Q}$  normalizes a Sylow 3-subgroup  $\overline{T}$  of  $\overline{L}$  and so  $Q$  normalizes  $C_{Y_L}(T)$  which has order 2 and complements  $V_L$ . Hence  $C_{Y_L}(T) \leq Z(Q)$ , so  $T \leq N_G(Q)$  and therefore  $L = \langle T, S \rangle \leq N_G(Q)$ , a contradiction. Thus  $\overline{Q} = \overline{S}$  has order 8.  $\blacksquare$

**(5.3.5)** We have  $Q_L = Y_L$ .

Suppose false. By (5.3.2)  $W$  acts quadratically on  $Q_L/Y_L$  and  $|\overline{W}| = 4$ . Also  $\overline{W} \not\leq \overline{S}_0$ , so Lemma 2.2 implies that the non-central  $L$ -chief factors in  $Q_L/Y_L$  are orthogonal modules for  $\overline{L} \cong O_4^-(2)$ . In particular, as  $L$ -modules, the non-central  $L$ -chief factors of  $Q_L/Y_L$  are not isomorphic to  $V_L$ .

Choose  $E \leq Q_L$  normal in  $L$  and minimal so that  $E/Y_L$  contains a non-central  $L$ -chief factor and let  $F$  be the preimage of  $C_{E/Y_L}(O^2(L))$ . Then  $[F, O^2(L)] \leq Y_L$  and Lemma 2.6 applies to yield  $F \leq Y_L$ . In particular,  $[E, E] \leq Y_L$ .

We claim  $E' \leq V_L$ . This is obviously the case if  $V_L = Y_L$ . So suppose that  $|Y_L : V_L| = 2$ . If  $E' \not\leq V_L$ . Then the minimal choice of  $E$  and  $E'V_L = Y_L$  implies that  $E/V_L$  is extraspecial of order  $2^5$ . Notice that  $[E, W] \leq W$  and  $W/Z$  is elementary abelian as  $[W, W] = Z$  by (5.3.2). Hence, as  $[E, W]Y_L/V_L$  has order  $2^3$ , we infer that  $E/V_L$  has  $+$ -type contrary to  $\overline{L} \cong \Gamma\text{SL}_2(4)$ . Hence  $E/V_L$  is elementary abelian. If  $[Q_L, E] = V_L$ , then  $E/V_L$  has order  $2^4$  by Lemma 2.2 and so  $Q_L/C_{Q_L}(E)$  embeds into

$$\text{Hom}_L(E/V_L, V_L) \cong (E/V_L)^* \otimes V_L \cong (E/V_L) \otimes V_L$$

by Lemma 2.7. Since  $Q_L/C_{Q_L}(E)$  involves only trivial and orthogonal modules this contradicts [Pr, Lemma 2.2].

Thus  $[E, Q_L] = Y_L > V_L$ .

By (5.3.4)

$$\overline{Q} = \overline{S} \text{ has order 8.}$$

In summary we now know  $|\overline{W}| = 4$  and  $[\overline{W}, \overline{Q}] = [\overline{W}, \overline{S}] = Z(\overline{S})$ .

We calculate using  $Z$  is normal in  $D$  by (5.3.2) that

$$[W, Q, Q] = \langle [V_L, Q, Q, Q]^D \rangle = \langle Z^D \rangle = Z.$$

Therefore

$$[E, [W, Q], Q] \leq E \cap [[W, Q], Q] \leq Z \leq Y_L.$$

As  $||Z(\bar{S}), E/Y_L|| = 4$  and  $\bar{Q} = \bar{S}$ , this implies that  $|C_{E/Y_L}(\bar{S})| = 4$ . As  $E/Y_L$  is the orthogonal  $O_4^-(2)$ -module for  $L$ , this is impossible. We have proved the claim.  $\blacksquare$

**(5.3.6)** Suppose that  $Y_L = V_L$ . Then  $L$  is a maximal 2-local subgroup of  $G$ ,  $N_G(Q)/Q \cong \text{SL}_2(2)$ , there exists a subgroup  $E \leq S$  of order  $2^4$  which is normalized by  $N_G(Q)$  such that  $N_G(E)/E \cong \text{Alt}(6)$  and  $N_L(E)$  has index 5 in  $L$ .

By (5.3.5) we have  $|S| = 2^7$ , and  $|\bar{W}| = 2^2$ . Also  $||W, Y_L|| = 8$  and  $Y_L \not\leq Q$ , so  $Q \cap Y_L = [W, Y_L] = W \cap Y_L$ , Therefore  $|W| = 2^5$ . Set  $C = C_Q(W)$ . Then  $C$  centralizes  $[Y_L, Q]$  which has order  $2^3$  and so  $C \leq C_L([Y_L, Q]) = Y_L$ . Thus  $C \leq C_{Y_L}(W)$  which has order 2. Then, by (5.3.2),  $W' = Z = C$  and, as  $W$  is generated by involutions, we have  $W$  is extraspecial. Since  $[Y_L, Q] \leq W$ ,  $W$  has  $+$ -type.

Observe  $W/Z = J(Q/Z)$ , so  $W$  is normal in  $N_G(Q)$  and  $N_G(Q)/Z$  embeds into  $\text{Aut}(W) \cong 2^4:\text{O}_4^+(2)$ .

Assume that  $Y_L Q/Q$  normalizes a subgroup  $T$  of  $O_3(N_G(Q))/Q$  which has fixed points on  $W/Z$ . Then  $W = [W, T]C_W(T)$  and  $[W, T] \cong C_W(T) \cong \text{Q}_8$  and these subgroups are normalized by  $Y_L$ . But then

$$[W, Y_L] = [C_W(T), Y_L][W, T, Y_L].$$

Since  $[W, Y_L]$  is elementary abelian and  $\Omega_1(P) = Z(P)$  for  $P \cong \text{Q}_8$ , we conclude that

$$[C_W(T), Y_L] = [W, T, Y_L] = Z$$

and then  $[W, Y_L]$  has order 2 which is nonsense as  $Y_L$  is the natural module. Therefore  $Y_L$  normalizes no such subgroup.

Let  $F = O_{2,3}(N_G(Q))$ . Assume that  $|F/Q| = 9$ . Then the previous argument implies that  $C_{F/Q}(Y_L) \neq 1$ . Let  $T_1$  be the preimage of this subgroup. Then  $[Y_L, Q]$  is normalized by  $T_1$ . Hence  $Y_L = C_{Y_L Q}([Y_L, Q])$  is normalized by  $T_1$ . Using the fact that  $Q$  is weakly closed in any 2-group which contains it, for  $w \in Y_L^\#$ , we let  $Q_w$  be the unique conjugate of  $Q$  in  $O_2(C_G(w))$ . Then  $T_1$  permutes the elements of  $Y_L$  and so  $T_1$  normalizes  $L^\circ = \langle Q_w \mid w \in Y_L^\# \rangle$ . Since  $L = L^\circ Y_L$ , we have that  $T_1$  normalizes  $L$ . On the other hand,  $WY_L$  is normalized by  $T_1$  and, as  $T_1$  acts fixed-point freely on  $W/Z$ ,  $T_1$  acts transitively on  $WY_L/Y_L \cong W/[Y_L, Q] \cong 2^2$  and this is impossible as  $W \cap O^2(L)$  is a maximal subgroup of  $W$  and is normalized by  $T_1$ .

Hence  $|F/Q| = 3$ ,  $N_G(Q) = FS$  and  $N_G(Q)/Q \cong \text{SL}_2(2)$ . In particular,  $|Q| = 2^6$ ,  $S = Y_L Q$ , and  $FY_L/W \cong 2 \times \text{SL}_2(2)$ . It follows that

$$[W, Q] \text{ is elementary abelian of order } 8.$$

et  $E = C_S([W, Q])$ . As  $W$  is normal in  $N_G(Q)$ , so is  $E$ . As  $|S| = 2^7$  and  $|\text{GL}_3(2)_2| = 2^3$ , we have  $|E| \geq 2^4$ . Since  $F$  acts fixed-point freely on  $W/Z$  (being normalized by  $Y_L$ ), we have  $E \leq Q$  and then  $E$  is normal in  $N_G(Q)$ . Since  $E \cap W = [W, Q]$ , we find  $|E| = 2^4$ . Let  $S \leq L_1 \leq L$  be such that  $L_1/Q_L \cong \text{Sym}(4)$  has index 5 in  $L$ . Notice that  $O_2(L_1) = S_0$ . Then  $E \leq C_L([Y_L, Q, Q]) = Y_L S_0$ . Also  $Y_L \leq S_0$ , so  $S_0 = Y_L S_0$ . Therefore  $E \leq S_0$ . Now  $EY_L/Y_L$  acts as a Sylow 2-subgroup of  $\text{SL}_2(4)$  on the natural module. In particular for any involution  $e \in E \setminus Y_L$  we have that  $C_{Y_L}(e) =$

$E \cap Y_L$ . This implies that all involutions in  $EY_L$  are contained in  $Y_L \cup E$  and therefore  $E$  and  $Y_L$  are the only elementary abelian subgroups of  $S_0$  of order  $2^4$ . In particular,  $L_1$  normalizes  $E$ . Now  $N_G(E) \geq \langle L_1, N_G(Q) \rangle \in \mathcal{L}_G(S)$ . Notice that  $L_1$  has orbits of lengths 3, and 12 on  $E$  and that  $N_G(Q)$  does not preserve these orbits. Hence  $N_G(E)$  acts transitively on  $E^\#$ . As  $N_G(Q) = C_G(Z)$ , we now have that  $|N_G(E)| = 15|N_G(Q)| = 2^7 \cdot 3^2 \cdot 5$ . We have that  $X = N_G(E)/E$  is isomorphic to a subgroup of  $\text{GL}_4(2) \cong \text{Alt}(8)$  of order  $2^3 \cdot 3^2 \cdot 5$ . We consider the action of  $X$  on a set of size 8. As  $\text{Alt}(8)$  has no subgroups of order 45,  $X$  is not transitive. Hence  $X$  is isomorphic to a subgroup of  $\text{Alt}(7)$ ,  $\text{Sym}(6)$  or  $X \cong (\text{Alt}(5) \times 3):2$ . Suppose that  $X \cong (\text{Alt}(5) \times 3):2$ . As  $N_G(Q)/Q \cong \text{Sym}(4)$ , we see that  $EQ/E \leq \text{Alt}(5)$ . Since  $E$  is the natural  $\text{SL}_2(4)$ -module, we get that  $|Z(Q)| = 4$ . But, by (5.3.2),  $|Z(Q)| = 2$ . Hence we have one of the first two possibilities and then obviously  $X = N_G(E)/E \cong \text{Alt}(6)$ .

We just have to show that  $L$  is a maximal 2-local subgroup. Let  $M$  be a 2-local subgroup with  $L \leq M$ . As  $Q \leq M$ , we have that  $M$  is of characteristic 2. Then  $Y_L = Y_M$  and  $C_G(Y_L) = Y_L$ . As  $|N_G(Q) : S| = 3$  and  $Y_L$  is not normal in  $N_G(Q)$ , we have  $N_M(Q) = S = N_L(Q)$ . As  $L$  acts transitively on  $Y_L^\#$ , we conclude  $M = N_M(Q)L = N_L(Q)L = L$ . ■

**(5.3.7)** If  $Y_L = V_L$ , then  $G$  has just one conjugacy class of involutions.

By (5.3.6)  $N_G(E)/E \cong \text{Alt}(6)$ . As  $Y_L \not\leq E$ , there is an involution  $y \in Y_L \setminus E$ . Now  $y$  inverts an element of order 5 in  $N_G(E)$  and so  $|[E, y]| = |C_E(y)| = 4$ . This shows that all involutions in  $Ey$  are conjugate. As all involutions in  $S/E$  are conjugate in  $\text{Alt}(6)$  and all the involutions in  $Y_L$  are  $L$ -conjugate, this proves the claim. ■

We have now proved that (i) holds when  $Y_L = V_L$ .

**(5.3.8)** Suppose that  $Y_L > V_L$ . Then  $G$  has a subgroup of index 2.

We have that  $|S| = 2^8$ . By (5.3.4) and (5.3.5),  $S = QY_L$ . We are going to show that  $J(S) = Y_L$ . For this let  $A \leq S$  be elementary abelian of maximal order and assume that  $A \neq Y_L$ . Then  $|AY_L/Y_L| \leq 4$ . As there are no transvections on  $V_L$ , we get  $|AY_L/Y_L| = 4$  and we may assume that  $A$  acts quadratically on  $Y_L$  by [GLS2, Theorem 25.2]. As  $W \not\leq S_0$  by Lemma 3.2 and  $|\bar{W}| = 4$  by (5.3.2),  $W$  does not act quadratically on  $Y_L$ ,  $AY_L/Y_L \leq S_0/Y_L$  and  $S_0 = AY_L$ . Now  $A \cap Y_L$  has order 8 and so  $|C_{Y_L}(S_0)| = 8$ . But  $(L^\circ)'$  is generated by two conjugates of  $S_0$ , which gives  $C_{Y_L}(L^\circ) \neq 1$  a contradiction to Lemma 2.10(i). Thus  $Y_L = J(S)$  is the Thompson subgroup of  $S$ . In particular,  $N_G(Y_L)$  controls  $G$ -fusion of elements in  $Y_L$ . As  $S \in \text{Syl}_2(G)$  and  $C_S(Y_L) = Q_L$ ,  $Q_L \in \text{Syl}_2(C_G(Y_L))$  and we have  $N_G(Y_L) = C_G(Y_L)N_{N_G(Y_L)}(Q_L)$ . By Lemma 5.2

$V_L$  is normal in  $N_G(Y_L)$ .

Suppose that  $O^2(L) \geq Y_L$ . Then  $O^2(L)/V_L \cong \text{SL}_2(5)$  has quaternion Sylow 2-subgroups and  $|L : O^2(L)| = 2$ . On the other hand, there exists  $g \in N_G(Q) \setminus N_G(Y_L)$  with  $WY_L \geq (Y_L^g \cap Q)Y_L \neq Y_L$  and  $(Y_L^g \cap Q)V_L/V_L$  is elementary abelian, which is a contradiction. Therefore  $O^2(L)/V_L \cong \text{SL}_2(4)$

and, as  $W$  does not act quadratically on  $Y_L$ , we see that  $|W : W \cap O^2(L)| = 2$  and thus  $O^2(L)W/V_L \cong \Gamma\text{SL}_2(4)$ . Hence  $L$  has a subgroup  $L_0 = O^2(L)W$  of index 2 with  $Y_L \cap L_0 = V_L$ .

Let  $T \in \text{Syl}_2(L_0)$  and  $w \in Y_L \setminus T$ . Suppose that for some  $x \in G$ ,  $w^x \in T$  and  $|C_S(w^x)| \geq |C_S(w)|$ . As  $L^\circ$  has orbits of length 6 and 10 on  $Y_L \setminus V_L$ , we may assume  $|C_S(w^x)| \geq |S|/2$ . But then as  $V_L$  is the natural module, it does not admit transvections and so  $w^x \in V_L$ . As  $N_G(Y_L) = N_G(V_L)$  and  $N_G(Y_L)$  controls fusion in  $Y_L$ , this is not possible. Hence the supposed condition cannot hold. Application of [GLS2, Proposition 15.15], shows that  $G$  has a subgroup of index 2. This proves (5.3.8).  $\blacksquare$

Let  $G_0$  be a subgroup of  $G$  of index 2, and set  $Q_0 = Q \cap G_0$ . We have  $V_L \leq L^\circ \leq G_0$ . Hence  $W = \langle [V_L, Q]^D \rangle \leq G_0$ . In particular,  $W \leq Q_0$  and so  $Z(Q_0) = Z$  and  $Q_0$  is large in  $G_0$ . Set  $L_0 = O^2(L)Q_0 = O^2(L)W$ . Then  $L_0^\circ/V_L \cong \Gamma\text{SL}_2(4)$  and  $Y_{L_0} = V_{L_0} = V_L \not\leq Q_0$ . Thus  $(G_0, L_0)$  satisfies the hypotheses of (i). This proves (ii) holds if  $V_L \neq Y_L$ .  $\square$

*Proof of Proposition 5.1:* By Lemma 5.3 we just have to examine the structure in Lemma 5.3(i), so we may assume that Lemma 5.3(i) holds.

By Lemma 2.1

$N_G(E)$  splits over  $E$ .

As  $N_G(Q) \leq N_G(E)$ , for a 2-central involution  $z$  we have that  $C_G(z)$  is a split extension of  $E$  by  $\text{Sym}(4)$ . As  $O(C_G(z)) = 1$  coprime action yields  $O(G) = \langle C_{O(G)}(e) \mid e \in E^\# \rangle = 1$ . In particular  $F(G) = 1$  and  $E(G) \neq 1$ . Suppose that  $J^*$  is a non-trivial subnormal subgroup of  $G$  normalized by  $\langle L, N_G(E) \rangle$ . Then  $S \cap J^* \neq 1$ . Since  $1 \neq J^* \cap N_G(E)$  is normal in  $N_G(E)$  and  $1 \neq J^* \cap L$  is normal in  $L$ , it follows that  $J^* \cap N_G(E) \geq J^* \cap S \geq EY_L$ . Hence  $J^* \geq \langle Y_L^{N_G(E)} \rangle = N_G(E) \geq S$  and  $J^* \geq \langle S^L \rangle = L$ . Therefore there is a unique non-trivial subnormal subgroup of  $G$  of minimal order normalized by  $\langle L, N_G(E) \rangle$ . It follows that  $\langle L, N_G(E) \rangle$  is contained in a component  $J$  of  $G$ . Since  $O(G) = 1$  and  $S \leq J$ ,  $J = E(G)$ . As  $J$  has just one conjugacy class of involutions by Lemma 5.3(i) and, for  $z \in E^\#$ ,  $C_G(z) \leq N_G(E)$ , it follows that  $G = J$  is simple. Using  $G$  has just one conjugacy class of involutions and applying [J, Theorem] yields  $G \cong \text{Mat}(22)$ . This proves the proposition when Lemma 5.3(i) holds. If Lemma 5.3(ii) holds, then  $G \cong \text{Aut}(\text{Mat}(22))$ .  $\square$

## 6. $\overline{L}^\circ \cong \text{SL}_2(4)$

In this section we investigate the configuration in Proposition 3.5(iii). Thus  $\overline{L}^\circ \cong \text{SL}_2(4)$ ,  $|Y_L : V_L| = 2$  and  $V_L$  is the natural  $\text{SL}_2(4)$ -module.

As  $Q \leq L^\circ$ ,  $C_{V_L}(S_0) = C_{V_L}(Q) \leq Z(Q)$ , so  $Q$  is normal in  $N_{L^\circ}(C_{V_L}(S_0))$  and hence  $\overline{Q} = \overline{S_0}$  is a Sylow 2-subgroup of  $\overline{L}^\circ$ . In particular  $Z(Q) \cap Y_L = Z(Q) \cap V_L$  is of order 4.

**Lemma 6.1.** *The subgroup  $Q$  is elementary abelian. In particular,  $Q \cap Y_L = Q \cap V_L = C_{Y_L}(Q) = Z$ ,  $|Y_L Q/Q| = 2^3$  and  $|V_L Q/Q| = 2^2$ .*

*Proof.* We know that  $[Q, V_L] = C_{V_L}(Q) = Q \cap V_L$  and, as  $\overline{Q}$  is elementary abelian,  $\Phi(Q) \leq Q_L$ . If  $\Phi(Q) \neq 1$ , then, since  $Z(S) \cap \Phi(Q) \neq 1$ , we deduce

$\Phi(Q) \cap V_L \neq 1$ . As  $N_L(QQ_L)$  normalizes  $Q$  and is irreducible on  $[V_L, Q]$ ,  $[V_L, Q] \leq \Phi(Q)$ . But then  $V_L$  centralizes  $Q/\Phi(Q)$ , so  $V_L \leq O_p(N_G(Q)) = Q$ , a contradiction. This shows  $Q$  is elementary abelian and then also  $Y_L \cap Q = V_L \cap Q = C_{Y_L}(Q)$ .  $\square$

**Proposition 6.2.** *Suppose  $L \in \mathcal{L}_G(S)$  and  $L \not\leq N_G(Q)$  with  $\bar{L}$  in the unambiguous wreath product case. If  $Y_L \not\leq Q$ ,  $\bar{L}^\circ \cong \text{SL}_2(4)$  and  $|Y_L : V_L| = 2$ , then  $G$  is  $\text{Aut}(\text{Mat}(22))$ .*

*Proof.* We start by observing that the action of  $L$  on  $Y_L$  gives

(6.2.1)

- (i)  $|V_L Q/Q| = |Q : C_Q(V_L)| = 2^2$ ;
- (ii) for all  $v \in V_L \setminus Q$ ,  $C_Q(v) = C_Q(V_L)$ ; and
- (iii) for all  $w \in Q \setminus Q_L$ ,  $[w, V_L] = [Q, V_L]$ .

Let  $B = N_L(QQ_L)$ . Then  $B$  contains an element  $\beta$  of order 3 which acts fixed-point freely on  $V_L$  and irreducibly on  $[V_L, Q] = C_{Y_L}(Q)$ .

Using (6.2.1) (ii) and Lemma 2.8 yields  $[V_L, F(N_G(Q)/Q)] = 1$ . Let  $K \geq Q$  be the preimage of

$$[E(N_G(Q)/Q), V_L Q/Q].$$

Then  $K$  is non-trivial, normalized by  $B$  and Lemma 2.8 implies  $V_L Q/Q$  acts faithfully on  $K/Q$ .

The three involutions of  $QQ_L/Q_L$  each centralize a subgroup of  $Y_L$  of order  $2^3$  and by Lemma 2.10(i), there are three elements of  $Y_L Q/Q$  which act on  $Q$  as  $\text{GF}(2)$ -transvections, they generate  $Y_L Q/Q$  and are permuted transitively by  $B/Q$ . As  $B$  normalizes  $K$  and as  $V_L Q/Q$  acts faithfully on  $K/Q$ , at least one and hence all of the transvections in  $Y_L Q/Q$  act faithfully on  $K/Q$ .

If  $C_Q(K) \neq 1$ , then  $C_{C_Q(K)}(S) \neq 1$ . As  $\Omega_1(Z(S)) = C_{V_L}(S)$  by Lemma 2.10 (ii), and  $C_Q(K)$  is normalized by  $B$ , we have  $[Q, V_L] \leq C_Q(K)$ . But then  $K = \langle V_L^K \rangle Q$  centralizes  $Q/C_Q(K)$  contrary to  $C_K(Q) = Q$ . Hence  $C_Q(K) = 1$ .

Let  $V$  be a non-trivial minimal  $KY_L$ -invariant subgroup of  $Q$ . Then  $KY_L$  acts irreducibly on  $V$ . Moreover, as  $Y_L$  does not centralize  $V$ ,  $V \not\leq Q_L$  and, as  $V_L$  is the natural  $\bar{L}^\circ$ -module we have  $[Y_L, V] = [Y_L, Q] = Y_L \cap Q \leq V$ . It follows that  $K$  centralizes  $Q/V$  and so  $K/Q$  acts faithfully on  $V = [Q, K]$  which is normalized by  $B$ . Hence  $C_{Y_L}(V) = Y_L \cap V = Y_L \cap Q$  and  $Y_L Q/Q$  acts faithfully on  $V$ . Recall that  $Y_L Q/Q$  is generated by elements which operate as transvections on  $Q$  and hence on  $V$ . Therefore [McL, Theorem] applies to give  $KY_L/Q \cong \text{SL}_m(2)$  with  $m \geq 3$ ,  $\text{Sp}_{2m}(2)$  with  $m \geq 2$ ,  $\text{O}_{2m}^\pm(2)$  with  $m \geq 2$ , or  $\text{Sym}(m)$  with  $m \geq 7$ . Furthermore,  $V = [Q, K]$  is the natural module in each case.

Since  $C_{Y_L Q/Q}(S/Q)$  contains a transvection and has order  $2^2$ ,  $KY_L/Q \not\cong \text{SL}_m(2)$  with  $m \geq 3$  or  $\text{O}_{2m}^\pm(2)$  with  $m \geq 2$ . Suppose that  $KY_L/Q \cong \text{Sym}(m)$  with  $m \geq 7$ . Then, as  $Y_L Q/Q$  is generated by three transvections, we see that  $Y_L Q/Q$  is generated by three commuting transpositions in  $KY_L/Q$ . Let  $t$  be the product of these transpositions. Then, as  $m \geq 7$ ,  $|[V, t]| = 2^3$ . However,  $|[V, Y_L]| = 2^2$ , and so we have a contradiction. We have demonstrated

**(6.2.2)**  $KY_L/Q \cong \text{Sp}_{2m}(2)$ ,  $m \geq 2$  and  $[Q, K] = [Q, KY_L]$  is the natural module.

Since  $[Q, K]$  is the natural  $KY_L/Q$ -module and  $[V_L, Q] \leq [Q, K]$  has order  $2^2$ , we have  $[[V_L, Q], S] \neq 1$ . In particular,  $QQ_L/Q_L < S/Q_L \cong \text{Dih}(8)$  and  $SQ/Q \cap K/Q$  acts non-trivially on  $[Q, V_L]$ .

Consider  $Q^* = O_2(KS)$ . Since  $Q^*$  centralizes  $[Q, K]$ ,  $Q^*$  centralizes  $[V_L, Q]$  and so  $Q^*Q_L = QQ_L$ . Hence  $\Phi(Q^*) \leq Q_L$ . If  $\Phi(Q^*) \neq 1$ , then

$$[Q, K] = \langle \Omega_1(Z(S))^K \rangle \leq \Phi(Q^*)$$

and so also  $[Q^*, K] = [Q^*, K, K] \leq [Q, K] \leq \Phi(Q^*)$  which is impossible. Hence  $Q^*$  is elementary abelian and it follows that  $Q \leq Q^* = C_{Q^*}(Q) \leq Q$ . Since  $KS$  acts on  $[Q, K]$  and  $KY_L/Q \cong \text{Sp}_{2m}(2)$ , we now deduce  $S \leq KY_L$  from the structure of  $\text{Out}(K/Q)$ . Hence  $B = \langle S^B \rangle \leq KY_L$  as  $B$  normalizes  $KV_L$ . It follows that  $B/Q$  is the minimal parabolic subgroup  $P$  of  $K/Q$  irreducible on  $[Y_L, V]$  and with  $O^2(P)$  centralizing  $[Y_L, V]^\perp/[Y_L, V] = C_{Y_L}(V)/[Y_L, V]$ . Therefore there is  $\beta \in K$  of order three such that  $\langle \beta \rangle$  is transitive on the transvections in  $Y_LQ/Q$  and normalizes  $Q_LQ/Q$  which has index 2 in  $S/Q$ . In particular, from the structure of the natural  $\text{Sp}_{2m}(2)$ -module  $\beta$  centralizes

$$C_V(Y_L)/[V, Y_L] = (V \cap Q_L)/(V \cap Y_L) = (V \cap Q_L)Y_L/Y_L \leq [Q_L, V]Y_L/Y_L.$$

As  $V$  is abelian,  $V$  acts quadratically on  $Q_L/V_L$ . By Lemma 2.2,  $Q_L/V_L$  involves only natural  $\text{SL}_2(4)$ -modules and trivial modules as  $L$ -chief factors. We know  $\beta$  acts fixed-point freely on the natural module and so, as  $\beta$  centralizes  $[Q_L, V]Y_L/Y_L$ , all the  $L$ -chief factors of  $Q_L/V_L$  are centralized by  $L$ . In particular,  $V_L$  is the unique non-central  $L$ -chief factor in  $Q$  and so  $Y_L \cap \Phi(Q_L) = 1$ . As  $\Omega_1(Z(S)) \leq V_L$  by Lemma 2.10 (ii),  $\Phi(Q_L) = 1$ , so  $Q_L = \Omega_1(Z(Q_L)) = Y_L$ , which together with  $S/Q_L \cong \text{Dih}(8)$  implies

**(6.2.3)**  $Y_L = Q_L$  has order  $2^5$  and  $|S| = 2^8$ .

Together (6.2.2) and (6.2.3) give

**(6.2.4)**  $|Q| = 2^4$  and  $N_G(Q)/Q \cong \text{Sym}(6)$ .

We next show that  $G$  has a subgroup of index two. In  $N_G(Q)$  we have a subgroup  $U$  of index 2 of shape  $2^4.\text{Alt}(6)$ . Furthermore  $Y_L \not\leq U$  and  $V_L \leq U$ . Since  $[v, Q] = C_Q(v)$  for  $v \in V_L \setminus Q$  and  $U/Q$  has one conjugacy class of involutions, all the involutions in  $U \setminus Q$  are  $U$ -conjugate. Since  $L$  acts transitively on  $V_L$  and  $U$  is transitive on  $Q^\#$ , we have that all the involutions in  $U$  are  $G$ -conjugate. As  $Q$  is large, we have  $C_G(z) \leq N_G(Q)$  for  $z \in Q^\#$ . Hence all the involutions in  $U$  have centralizer which is a  $\{2, 3\}$ -group. There is an involution  $t$  in  $Y_L \setminus V_L$ , which is not in  $U$  and centralized by an element of order 5 in  $L$ . Hence  $t$  is not conjugate to any involution of  $U$ . Application of [GLS2, Proposition 15.15] gives a subgroup  $G_1$  of index two in  $G$ . We have  $N_{G_1}(Q)/Q \cong \text{Alt}(6)$ . By Lemma 2.1 this extension splits and we have that the centralizer of a 2-central involution  $z \in G_1$  is a split extension of an elementary abelian group of order 16 by  $\text{Sym}(4)$ . In particular  $O(C_G(z)) = 1$  and so by coprime action  $O(G) = \langle C_{O(G)}(e) \mid e \in Q^\# \rangle = 1$ . As  $Y_L \not\leq Q$ , there is an involution  $y \in N_{G_1}(Q) \setminus Q$ . Since all involutions in  $Qy$  and in  $N_{G_1}(Q)/Q$  are conjugate,  $G_1$  has just one conjugacy class of involutions. In particular



$F^*(G_1)$  is simple. Application of [J, Theorem] gives that  $F^*(G_1) \cong \text{Mat}(22)$  and so  $G \cong \text{Aut}(\text{Mat}(22))$ .  $\square$

#### ACKNOWLEDGMENT

We thank the referee for numerous comments which have improved the readability and clarity of our work. The second author was partially supported by the DFG.

#### REFERENCES

- [GH] D. Gorenstein, K. Harada, On finite groups with Sylow 2-subgroups of type  $A_n$ ,  $n = 8, 9, 10, 11$ , Math. Z., 117 (1970), 207–238.
- [GLS2] D. Gorenstein, R. Lyons, R. Solomon, The classification of the finite simple groups, Amer. Math. Soc. Surveys and Monographs 40(2), (1996).
- [J] Z. Janko, A characterization of the Mathieu simple groups, I, J. Algebra 9, 1968, 1–19.
- [Mas] D. Mason, Finite simple groups with Sylow 2-subgroup dihedral wreath  $Z_2$ , J. Algebra 26, (1973), 10–68.
- [MSS1] U. Meierfrankenfeld, B. Stellmacher, G. Stroth, Finite groups of local characteristic  $p$ : an overview in *Groups, combinatorics and geometry*, Durham 2001 (eds. A. Ivanov, M. Liebeck, J. Saxl), Cambridge Univ. Press, 155–191.
- [MSS2] U. Meierfrankenfeld, B. Stellmacher and G. Stroth, *The local structure theorem for finite groups with a large  $p$ -subgroup*, Mem. Amer. Math. Soc. 242, Nr. 1147 (2016).
- [McL] J. McLaughlin, Some subgroups of  $\text{SL}_n(F_2)$ , Illinois J. Math 13, 1969, 105–115.
- [PPS] Chr. Parker, G. Parmeggiani, B. Stellmacher, The  $P!$ -Theorem, Journal of Algebra 263 (2003), 17–58.
- [Pr] A. R. Prince, On 2-groups admitting  $A_5$  or  $A_6$  with an element of order 5 acting fixed point freely. J. Algebra 49 (1977), no. 2, 374–386.

CHRIS PARKER, SCHOOL OF MATHEMATICS, UNIVERSITY OF BIRMINGHAM, EDGBASTON, BIRMINGHAM B15 2TT, UNITED KINGDOM  
*Email address:* `c.w.parker@bham.ac.uk`

GERNOT STROTH, INSTITUT FÜR MATHEMATIK, UNIVERSITÄT HALLE - WITTENBERG, THEODOR LIESER STR. 5, 06099 HALLE, GERMANY  
*Email address:* `gernot.stroth@mathematik.uni-halle.de`